

## The ISOPHOT - MAMBO survey of 3CR radio sources: Further evidence for the unified schemes<sup>★,★★</sup>

M. Haas<sup>1,2</sup>, S. A. H. Müller<sup>2</sup>, F. Bertoldi<sup>3</sup>, R. Chini<sup>2</sup>, S. Egner<sup>1</sup>, W. Freudling<sup>4</sup>, U. Klaas<sup>1</sup>,  
O. Krause<sup>1</sup>, D. Lemke<sup>1</sup>, K. Meisenheimer<sup>1</sup>, R. Siebenmorgen<sup>4</sup>, and I. van Bemmel<sup>5</sup>

<sup>1</sup> Max-Planck-Institut für Astronomie (MPIA), Königstuhl 17, 69117 Heidelberg, Germany

<sup>2</sup> Astronomisches Institut, Ruhr-Universität Bochum (AIRUB), Universitätsstr. 150/NA7, 44780 Bochum, Germany  
e-mail: haas@astro.rub.de

<sup>3</sup> Max-Planck-Institut für Radioastronomie (MPIfR), Auf dem Hügel 69, 53121 Bonn, Germany

<sup>4</sup> European Southern Observatory (ESO), Karl-Schwarzschild-Str. 2, 85748 Garching, Germany

<sup>5</sup> Space Telescope Science Institute (STScI), 3700 San Martin Drive, Baltimore, MD 21218, USA

Received 1 December 2003 / Accepted 1 June 2004

**Abstract.** We present the complete set of ISOPHOT observations of 3CR radio galaxies and quasars, which are contained in the ISO Data Archive, providing 75 mid- and far-infrared spectral energy distributions (SEDs) between 5 and 200  $\mu\text{m}$ . For 28 sources they are supplemented with MAMBO 1.2 mm observations and for 15 sources with new submillimetre data from the SCUBA archive. The sample includes flat and steep spectrum quasars, broad and narrow line radio galaxies, as well as Fanaroff-Riley FR 1 and FR 2 types. The SED shapes exhibit a diversity in the infrared (IR), ranging from a smooth dominating synchrotron component in flat spectrum sources to a thermal dust bump around 60–100  $\mu\text{m}$  in steep spectrum sources. The detection rate of a thermal bump in more than 50% of the cases suggests that dust emission may be a general phenomenon in these sources. We check the orientation-dependent unified scheme, in which the powerful FR 2 narrow line galaxies are quasars viewed at high inclination, so that their nuclei are hidden behind a dust torus intercepting the optical-ultraviolet AGN radiation and reemitting it in the infrared. We find that (1) both the quasars and the galaxies show a high mid- to far-infrared luminosity ratio typical for powerful AGNs and (2) – when matched in 178 MHz luminosity – both show the same ratio of isotropic far-infrared to isotropic 178 MHz lobe power. Therefore, from our large sample investigated here we find strong evidence for the orientation-dependent unification of the powerful FR 2 galaxies with the quasars. The distribution of the dust-to-lobe luminosity ratio shows a dispersion which we suggest to be most likely due to the additional influence of evolution and environment superposed on the orientation-dependent unification. We discuss our data also in the frame of the receding torus model. At the high 178 MHz luminosities of our sources above  $10^{26.5}$  W/Hz we do not find any support for this model in its original formulation and therefore we propose a refinement: The scale height of the torus might not be independent of luminosity, rather it may increase at high luminosities due to the impact of supernovae from starbursts accompanying the AGN phenomena.

**Key words.** Galaxy: fundamental parameters – galaxies: photometry – galaxies: quasars: general – infrared: galaxies

### 1. Introduction

#### 1.1. Unified schemes

While observationally at least a dozen classes of active galactic nuclei (AGN) are discernible, on the conceptional side consensus is growing that in most, if not all, of these different objects the accretion onto a massive black hole ( $M_{\text{bh}} \geq 10^6 M_{\odot}$ ) provides the main source of energy. It seems natural to explain their apparent differences by observational effects such

as (1) the *circumgalactic environment*, (2) the *evolutionary phase*, or (3) the *aspect angle*. To disentangle these effects is one of the major challenges of the current AGN research, as reviewed by e.g. Urry & Padovani (1995).

The Revised Third Cambridge Catalogue of radio galaxies and quasars at 178 MHz by Laing et al. (1983, abbreviated 3CRR) and – applying softer criteria – by Spinrad et al. (1985, for short 3CR) contains 178 and 298 sources, respectively, which fall into several classes of radio-loud AGN:

- 1) flat spectrum quasars, which are almost core dominated;
- 2) steep spectrum broad line radio galaxies (BLRGs) and quasars (here collectively called *quasars*), with either compact or lobe dominated morphology;
- 3) powerful narrow line radio galaxies (NLRGs) of Fanaroff-Riley type FR 2 with – often asymmetric – jets

\* Based on observations with the Infrared Space Observatory ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, The Netherlands and the UK) and with the participation of ISAS and NASA.

\*\* Table 2 and Fig. 3 are only available in electronic form at <http://www.edpsciences.org>

ending in edge-brightened double lobes extending far beyond the host galaxy;

- 4) low power narrow line radio galaxies of Fanaroff-Riley type FR 1, with double jets feeding edge-darkened radio lobes often contained within the host galaxy.

The diversity and the brightness of the sources makes the 3CR catalogue a well suited database for studying the relationship between these AGN classes. The aspect angle relation between flat and steep spectrum quasars, with the flat spectrum sources being steep spectrum sources seen pole-on (as proposed by Orr & Browne 1982), is now widely accepted. The relation between steep spectrum quasars and FR 2 radio galaxies and that between FR 1 and FR 2 sources is still a matter of debate.

While the quasars show the optical signatures of a powerful AGN (like broad emission lines and high excitation and optical luminosity), the galaxies do not. In an early attempt, physically different central engines were proposed as reviewed by Begelman et al. (1984): quasars have a high mass accretion rate but a low black hole spin, and galaxies have a low mass accretion rate but a high black hole spin. Both engines should be able to create the radio jets. Although a viable explanation, from the conceptional view point it appears somewhat astonishing that two intrinsically different AGN types should lead to the *same* radio jet and lobe phenomena. Alternatively, in the picture of orientation-dependent unification, galaxies could be quasars in which the AGN is hidden behind a dust torus seen “edge-on”, as proposed by Barthel (1989). Further debate arises from the fact that galaxies are distributed over the entire redshift range of the 3CR catalogue from the nearby local universe at  $z \approx 0.01$  up to  $z \approx 2$ , but the quasars are preferentially found at larger distances at  $z \gtrsim 0.3$  and 178 MHz luminosities above  $10^{26.5}$  W/Hz (e.g. Singal 1996; Willott et al. 2000). The increase of the quasar fraction with distance and luminosity may find an explanation by the *receding torus model* (Lawrence 1991; Hill et al. 1996): due to dust sublimation the inner wall of the torus recedes with increasing luminosity of the central engine and – under the assumption that the torus scale height is independent of luminosity – the cone opening angle increases, leading to a higher chance to see the broad line region. However, it is still under debate how far this elegant model is valid for the powerful radio galaxies with 178 MHz luminosities above  $10^{26.5}$  W/Hz. Alternatively, also physical or cosmic evolution could play a role: in this picture galaxies or a subset of them could be old quasars with a starving black hole, but still with luminous radio phenomena, possibly due to a favourable circumgalactic environment. Support for an environmental or evolutionary relationship comes from the fact that many FR 2s and most FR 1s fall into the low ionisation emission line category (Hine & Longair 1979), with little evidence for thick obscuring dust tori on HST images (Chiaberge et al. 1999, 2000).

In order to test the orientation-dependent unification between FR 2 galaxies and quasars, henceforth simply called *unification*, the task is to reveal a hidden powerful AGN in a (powerful) galaxy. Clear evidence for an AGN hidden behind a dust torus can be obtained by spectropolarimetry, when broad lines

show up as light scattered in the bipolar cones with the polarisation angle being perpendicular to the polar axis (Antonucci & Miller 1985). Despite a number of cases with broad lines revealed by spectropolarimetry, in numerous cases huge extinction or dust lanes extended over kpc could have prevented the view into the central region via a “suitable mirror” (Antonucci & Barvainis 1990; Hill et al. 1996; Cohen et al. 1999). Note that the optical signatures of dust extinction only indicate that something *could* be hidden, but it remains to be shown that there is actually a powerful energy source behind the dust lanes. A robust check of the *unification* is to look for the mid- and far-infrared reemission of the absorbed light from the AGN. A great advantage is that at wavelengths  $\lambda \gtrsim 25 \mu\text{m}$  the IR emission is largely optically thin, hence isotropic and independent of the aspect angle.

### 1.2. Infrared observations

Observations of the 3CR sample with IRAS did not allow a substantial conclusion about the *unification* (Heckman et al. 1992; Heckman et al. 1994; Hes et al. 1995; Hoekstra et al. 1997). The wavelength coverage and sensitivity of the IRAS observations were not sufficient to discriminate between dust and synchrotron emission in the individual sources, hence to correct for the contribution by a beamed synchrotron component to the mid- and far-infrared spectra of the sources.

With the Infrared Space Observatory ISO (Kessler et al. 1996) several small 3CR sub-samples were successfully observed: on the basis of a few ISOPHOT-MAMBO detections Haas et al. (1998) demonstrated that the IR emission of quasars and galaxies is a mixture of thermal and synchrotron radiation, dominated by thermal emission in radio-quiet quasars and synchrotron emission in flat spectrum radio-loud ones. Remarkably, steep spectrum quasars as well as galaxies showed powerful infrared dust emission.

In order to test the *unification*, a dedicated strategy compared galaxy – quasar pairs, which match in isotropic 178 MHz radio luminosity as well as redshift (in order to minimize possible cosmological effects). However, the four FR 2 galaxy – quasar pairs studied by van Bemmelen et al. (2000) yielded a lower detection rate of thermal dust emission and a lower dust luminosity for the galaxies than for the quasars, contrary to the expectations from the *unification*. On the other hand, ISOPHOT-MAMBO observations of ten FR 2 galaxy – quasar pairs with high radio power by Meisenheimer et al. (2001) yielded a balanced detection statistics and a similar dust luminosity for the quasars and galaxies, as predicted by the *unification*. Furthermore, the extrapolation of the synchrotron spectra of the quasars from the cm and mm regime into the far- and mid-IR has on average a more than ten times higher level than that for the radio galaxies. This is also consistent with the unified schemes, which predict that the jet axis of quasars is oriented more toward the line-of-sight, resulting in a stronger beamed component. Further ISOPHOT-MAMBO studies of small 3CR samples confirmed that many galaxies as well as quasars produce luminous dust emission (Andreani et al. 2002), and that the synchrotron component in quasars is higher than in

galaxies (van Bemmel & Bertoldi 2001). A closer quantitative look at the dust-to-radio luminosity ratio, however, reveals also differences between galaxies and quasars which suggest the additional role of evolution and environment superposed on the *unification* (Meisenheimer et al. 2001; Haas 2001).

Due to the limited detection of individual 3CR sources by IRAS, no definite conclusions about the *FR 1 – FR 2 relation* could be drawn; to our knowledge neither ISO results on that topic have been reported so far.

### 1.3. This paper

In order to overcome the small number statistics and derive results from a broader statistical basis, we present here the remaining unpublished ISOPHOT SEDs of the 3CR sources in the ISO archive. We discuss the statistical properties of the complete mid- to far-infrared data base of 75 galaxies and quasars from the 3CR catalogue observed by ISOPHOT with regard to unification, evolution and environment as well as the applicability of the receding torus model. We focus here on the *unification*; the discussion of the *FR 1 – FR 2 relation* will follow in a subsequent paper. Since common selection criteria for the sources were low cirrus foreground and good visibility to the satellite, this sample provides a fairly representative subsample of the 3CR catalogue, with a redshift range from nearby to distant ( $z \lesssim 2$ ) objects. Throughout this paper we use blackbodies modified with a dust emissivity index of  $\beta = 2$ . We adopt a  $\Lambda$  cosmology with  $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_{\text{matter}} = 0.27$  and  $\Omega_{\Lambda} = 0.73$  (Bennett et al. 2003).

## 2. The data

The ISO Data Archive (Kessler et al. 2000) contains 5 to 200  $\mu\text{m}$  photometry for 75 3CR sources obtained with ISOPHOT (Lemke et al. 1996), the photometer on board ISO. About half of them have also been observed between 5 and 15  $\mu\text{m}$  with ISOCAM. Our sample does not contain those 3CR sources, for which only ISOCAM mid-infrared data are available, without any far-infrared complement by ISOPHOT data or IRAS detections. The full set of 85 ISOCAM observations is published by Siebenmorgen et al. (2004). Our sample does not include the two FR 1 sources 3C 71 (NGC 1068) and 3C 231 (M 82), which do not belong to the classical FR 1s with elliptical hosts, and which are nearby and therefore too extended to be observed in ISOPHOT photometry modes. In order to characterize the (sub)-mm part of the SEDs for the full sample, we complemented the available literature data: for 15 sources we found submm photometry in the JCMT-SCUBA archive, and performed additional mm observations for 28 sources with the Max-Planck Millimeter bolometer array MAMBO (Kreysa et al. 1998) at the IRAM 30 m telescope.

### 2.1. ISOPHOT

The observing modes (Laureijs et al. 2002) comprise chopped measurements, including those with the spectrometer

ISOPHOT-S, from which we determined broad band fluxes, and small maps.

We have reduced the data using the ISOPHOT Interactive Analysis tool (PIA<sup>1</sup> V10), together with the calibration data set V7.0. Using the latest versions of the data reduction tools, there are no systematic photometric offsets with regard to PIA V7 and V8 used for most of the results published earlier by various authors. The accuracy of the absolute photometric calibration depends mainly on systematic errors as described in Laureijs & Klaas (1999), and it is currently known to be better than 30% for faint sources. The relative filter-to-filter calibration agrees within 15% (Klaas et al. 2002).

### 2.2. MAMBO

With the MAMBO array at the IRAM 30-m telescope on Pico Veleta, Spain, during the pooled observation campaigns between January and March 2003, we obtained 1.2 mm (250 GHz) continuum flux densities for 28 of our sources. We used the standard on-off photometry observing mode, chopping between the target and sky at 2 Hz, and nodding the telescope every 10 s. The total on sky integration time varied between 2.5 and 23 min per source, and each object was observed at least twice, on different days. The absolute flux calibration was established by observations of Mars and Uranus, resulting in a flux calibration uncertainty of approx. 20%. The data were reduced using the MOPSI software package.

### 2.3. SCUBA

Yet unpublished submm continuum data obtained with SCUBA were retrieved from the JCMT archive for 15 sources. The observations were performed in photometry mode, and for one source (3C 274 = M 87) as jiggle maps. We have reduced them with the SCUBA User Reduction Facility (SURF), including identification of noisy bolometer pixels and removal of sky noise. The atmospheric transmission was determined from skydips and water radiometer data at the CSO. IRC+10206 and HL-Tau served as standard calibrators. The integration time per source was 10–20 min. In case of very good atmospheric transmission ( $\tau_{\text{CSO}} \approx 0.05$ ) we also included the 450  $\mu\text{m}$  data. The photometric accuracy at 450  $\mu\text{m}$  and 850  $\mu\text{m}$  is about 30% and 25%, respectively.

### 2.4. Literature data

ISO photometry of 49 3CR sources has so far been published by various authors, as labelled in col. “publ.” of Table 1. The SED plots shown here include 4.5 to 14.3  $\mu\text{m}$  data with ISOCAM (Cesarsky et al. 1996) for 33 sources, which are published by Siebenmorgen et al. (2004) and for a subset by Freudling et al. (2003).

Additional literature data were retrieved from the NED and SIMBAD, the 2MASS archive and numerous

<sup>1</sup> PIA is a joint development by the ESA Astrophysics Division and the ISOPHOT consortium led by the MPI für Astronomie, Heidelberg. Contributing institutes are DIAS, RAL, AIP, MPIK, and MPIA.



Table 1. continued.

| 3C                 | Publ.    | Map/<br>Chop | 4.8<br>23"     | 7.3<br>23"      | 10<br>23" | 11.5<br>23"     | 12.8<br>23" | 16<br>23" | 20<br>52" | 25<br>52" | Map/<br>Chop | 60<br>45"         | 80<br>45" | 90<br>45" | 100<br>45" | Map/<br>Chop | 120  | 150  | 170 | 180  | 200  | 450<br>8"         | 850<br>15"        | 1200<br>11"       |
|--------------------|----------|--------------|----------------|-----------------|-----------|-----------------|-------------|-----------|-----------|-----------|--------------|-------------------|-----------|-----------|------------|--------------|------|------|-----|------|------|-------------------|-------------------|-------------------|
| 336.0              |          | c            | <15            | <6              | <12       |                 | <36         | <75       | <42       |           | c            | <96               |           | <168      |            | c            |      |      |     |      |      | <47               | <5.5              |                   |
| 337.0              | A, F     |              |                |                 |           |                 |             |           |           |           | m            | <63               |           | <78       |            | m            |      |      |     |      |      | <102              |                   |                   |
| 340.0              |          |              |                |                 |           |                 |             |           |           |           | m            |                   |           | <99       |            | m            |      |      |     |      |      | <168              |                   |                   |
| 343.1              | A, F     |              |                |                 |           |                 |             |           |           |           | c            | <108              |           | <462      |            | c            |      |      |     |      |      | <411              |                   | 7.6               |
| 351.0              | vB, H2   | c            | 18             | 31              |           | 50              |             |           |           | 112       | m            | 201               |           | 168       | 151        | m            |      | 94   |     | 66   |      |                   | 48                | <3.3              |
| 352.0              | A        |              |                |                 |           |                 |             |           |           |           | m            |                   |           | <75       |            | m            |      |      |     |      |      | <156              |                   |                   |
| 356.0              | A        |              |                |                 |           |                 |             |           |           |           | m            |                   |           | 27        |            | m            |      |      |     |      |      | <84               |                   |                   |
| 368.0              | M        | c            | 4 <sup>x</sup> | 16 <sup>x</sup> |           |                 | <120        |           | <162      |           | c            | 94                |           | <144      |            | c            |      |      |     |      |      | <168              | <60 <sup>A</sup>  | <3.1 <sup>A</sup> |
| 371.0 <sup>*</sup> |          |              |                |                 |           |                 |             |           |           |           | c            |                   |           | 397       |            | c            |      |      |     |      |      | 275               | <45 <sup>A</sup>  | 4.1 <sup>A</sup>  |
| var.               |          |              |                |                 |           |                 |             |           |           |           | c            |                   |           | 226       |            | c            |      |      |     |      |      | 215               |                   |                   |
| 380.0              | M        | c            | <9             | 7               |           | <18             |             |           |           | <66       | m            | 54                |           | 56        |            | m            |      |      |     |      |      | 83                |                   | 659               |
| 381.0              |          | m            |                |                 |           | 19              |             |           |           | 52        | m            | 74                |           | <78       |            | m            |      |      |     |      |      | 42                |                   | <2.1              |
| 390.3              |          | c            |                |                 |           | 55 <sup>s</sup> |             |           |           | 139       | c            | 191               |           | <1233     |            | c            | <963 |      |     | <984 |      | <385              | <14               |                   |
| 401.0              | F        |              |                |                 |           |                 |             |           |           |           | c            | <78               |           | <99       |            | c            |      |      |     |      | <351 | <270              |                   | 12.2              |
| 405.0 <sup>*</sup> | H1, M, P | c            | <12            | 40              |           |                 | 250         |           | 590       |           | c            | 3034              |           | 2155      |            | m            | 994  |      |     | 419  |      | ~230 <sup>R</sup> | ~490 <sup>R</sup> | 628               |
| 427.1              | M        | c            | <48            | <15             |           |                 | <114        |           | <75       |           | c            | <135              |           | <204      |            | c            |      |      |     |      |      | <180              |                   |                   |
| 437.0              | M        | c            | <21            | <12             |           |                 | <102        |           | <138      |           | c            | <126              |           | <246      |            | c            |      |      |     |      |      | <189              |                   | <52 <sup>W</sup>  |
| 441.0              | F        |              |                |                 |           |                 |             |           |           |           | c            | <174              |           | <153      |            | c            |      |      |     |      |      | <234              | <324              | <3.0              |
| 445.0 <sup>*</sup> |          | c            | 46             | 73              |           |                 |             |           |           | 360       | c            | 435               |           | <357      |            | c            |      | <849 |     |      |      | <1647             |                   | <4.8              |
| 454.3              |          | c            | <12            | 16              |           |                 | 34          |           | <291      |           | c            | 706               |           | 324       |            | m            | 187  |      |     | 248  |      | 1370              | 2780              | 3716.0            |
| 459.0 <sup>*</sup> | F        |              |                |                 |           |                 |             |           |           | <450      | c            | 1035 <sup>+</sup> |           | 1375      |            | c            |      | 1110 | 815 |      |      | <1413             |                   |                   |
| 460.0              | vB       |              |                |                 |           |                 |             |           |           |           | m            | <57               |           | <54       |            | m            |      |      |     |      |      | <351              |                   |                   |

\* Contained in Spinrad et al. (1985), but not in Laing et al. (1983); <sup>c</sup> contaminated by NGC 4138; <sup>x</sup> possibly contaminated by an asteroid; <sup>s</sup> integrated over spectral continuum obtained with ISOPHOT-S; <sup>+</sup> also: 1060 mJy at 65  $\mu$ m.

papers: Lilly & Longair (1984), Spinrad et al. (1985), Laing et al. (1983), Lilly et al. (1985), Neugebauer et al. (1987), Simpson et al. (2000) and de Vaucouleurs et al. (1991) at optical and NIR wavelengths, Chini et al. (1989a,b), Steppe et al. (1988, 1992, 1993), Robson et al. (1998), Hughes et al. (1993), Best et al. (1998), van Bemmell et al. (2000), Polletta et al. (2000), Archibald et al. (2001), Willott et al. (2002), Andreani et al. (2002) and Stevens et al. (1998) at mm wavelengths, and at radio wavelengths from Jodrell-Bank Web pages (at <http://www.jb.man.ac.uk/atlas/>), Kühr et al. (1981), Kellermann et al. (1989) and Akujor et al. (1994).

### 3. Results

#### 3.1. General properties

The IR, submm and mm fluxes of the 3CR sources are listed in Table 1. ISOPHOT achieved a detection rate of about 70% in the MIR ( $\leq 25 \mu$ m) and 60% in the FIR ( $\geq 60 \mu$ m). Even at the long wavelengths between 120 and 200  $\mu$ m the detection rate is about 20%. The fluxes of the brightest non-variable sources, also detected by IRAS (e.g. 3C 272.1, 3C 405, 3C 459), agree within 15%.


14 out of 28 sources observed with MAMBO at 1.2 mm were detected, and 3- $\sigma$  upper flux limits are provided for the others. From the 15 sources in the SCUBA archive eight were detected at 850  $\mu$ m and three at 450  $\mu$ m.

Supplementing our data with further photometry from the literature as listed in Sect. 2.4, we display the observed spectral energy distributions (SEDs) in Figs. 1–3, whereby each figure contains a subset of sources suitably chosen with regard to the illustration of the discussion items. Figure 1 shows only

those quasars, BLRGs and FR 2 NLRGs which exhibit a clear thermal bump and are used for investigating the *unification*. Figure 2 shows the FR 1 SEDs. Finally, Fig. 3 shows the remaining SEDs which are either those of flat spectrum sources or which exhibit poorly constrained thermal bumps due to upper limits only – nevertheless illustrating the high IR luminosity limits. Note that the SEDs in Figs. 1 and 2 run smoothly and the photometric uncertainties represented by the size of the symbols are small.

The most remarkable feature of the SEDs in Fig. 1 is the steady increase in flux density from near- to far-IR wavelengths with a peak typically at 60–100  $\mu$ m, followed by a steep drop at (sub)-mm wavelengths in many cases. Where it could be measured, in particular for 3C 48, 3C 298, 3C 318, 3C 405, this drop follows exactly the Rayleigh-Jeans tail of the modified blackbody (with dust emissivity index  $\beta = 2$ ). This strongly suggests a thermal nature of the FIR emission.


In order to discuss the SEDs in the framework of thermal emission, we have eyeball fitted a modified blackbody to the *peak* of the flux distribution, as shown in Figs. 1 and 2. The peak blackbody is mainly determined by the maximum in the FIR and, if available, also by the sub-millimetre points. However, this greybody fit does not always match the (sub)-millimetre measurements, since they may often be dominated by the synchrotron emission (for example in 3C 109 or 3C 234 in Fig. 1). Note that our “peak blackbody” corresponds in a  $\nu F_\nu$  versus  $\nu$  plot to the “break” of the thermal spectrum, where the slope equals unity. Its temperature  $T_p$  (corrected for redshift) lies between  $\sim 30$  and 100 K, with higher  $T_p$  values preferentially found in the most luminous sources at high redshift – probably a natural consequence for a flux limited sample like



**Fig. 1.** Spectral energy distributions of those quasars, BLRGs and FR 2 NLRGs which show a clear thermal bump and are used in the discussion of the unification in Sect. 4.2. The measurement errors are of the size of the symbols. The wavelength and frequency ranges are as observed and not corrected to the rest frame of the objects. The MIR (10–40  $\mu\text{m}$ ) and FIR (40–1000  $\mu\text{m}$ ) wavelength ranges, as well as the temperature of the modified blackbody (solid line curve) fitted to the thermal FIR bump are shown in the object’s restframe. Where the data coverage is sparse, the straight lines indicate the “envelopes” used for computing the restframe luminosities.

ours. It is clear from Fig. 1 that also dust which is colder than the peak blackbody may be present, but if so, then at lower flux levels. Also, as being evident from the MIR data points above

the peak blackbody curve, warmer dust components up to the dust sublimation temperature of about 1500 K are present, but for lucidity of the SED plots they are not shown explicitly here.



**Fig. 2.** Spectral energy distributions of the FR 1 sources. The meaning of the lines and labels is as in Fig. 1. Where the ISOPHOT data coverage is sparse, IRAS flux values from the NED are shown, if available (e.g. for 3C 293).

Note that for the cases of upper limits and poorly determined dust bumps shown in Fig. 3 we tentatively adopt an average  $T_p \sim 40$  K determined from the good cases shown in Fig. 1, if not constrained sufficiently.

The restframe thermal mid- (10–40  $\mu\text{m}$ ) and far-IR (40–1000  $\mu\text{m}$ ) luminosities are computed by integrating the SED “envelopes”. These curves are either sufficiently sampled by observations, as e.g. in the case of 3C 48, or they are partly sampled between 10 and 100  $\mu\text{m}$  and follow the Rayleigh-Jeans tail of the peak blackbody between 100 and 1000  $\mu\text{m}$ , as e.g. in the case of 3C 47. For a few sources in Fig. 1 (3C 33, 3C 49, 3C 245 and 3C 284) the SEDs are sparsely sampled in the MIR-FIR regime; in these cases they are interpolated linearly in the log-log SED plots between NIR and FIR flux points as indicated by the solid lines. Since the short wavelength part of the integration interval dominates the luminosity calculation, for poorly sampled SEDs the MIR luminosity may be overestimated by up to  $\sim 50\%$ , but the FIR luminosity is not significantly ( $< 15\%$ ) affected by the choice of  $T_p$ .

For all sources Table 2 lists the derived parameters like luminosities, dust temperatures and masses, and other values used in the discussion.

### 3.2. Notes on some individual sources

**3C 274 (M 87):** The SED of this FR 1 source clearly exhibits as three basic features (1) the optical-NIR bump from the host galaxy, (2) the thermal dust bump in the mid- and far-IR, and (3) the radio synchrotron component. Table 1 lists the total fluxes at 450 and 850  $\mu\text{m}$ , nevertheless we were able to resolve the one-sided jet and the core in the SCUBA jiggle maps. The core fluxes assuming a point source with  $FWHM$  8'' and 15'' are  $0.391 \pm 0.10$  Jy and  $1.085 \pm 0.10$  Jy at 450 and 850  $\mu\text{m}$ , respectively.

**3C 268.4:** Our FIR fluxes at 90 and 170  $\mu\text{m}$  are similar to those reported by Andreani et al. (2002), but they are surprisingly high for a quasar at  $z \approx 1.4$ . Therefore we checked for the presence of other nearby galaxies. In fact, the galaxy NGC 4138 lies less than 200'' apart and has an extent of more than 100''. In the FIR it is also bright with  $F(170 \mu\text{m}) > 6$  Jy, as we found with the help of the ISOPHOT Serendipity Survey (Stickel et al. 2000). Hence the large ISOPHOT beams of 45''

and 180'', respectively, have seen mainly the contamination by NGC 4138. Therefore, we do not consider 3C 268.4 any further.

**3C 313:** The FIR flux reported by Andreani et al. (2002) is due to a misidentification, actually the ISO observations refer to IRAS F15086+0801 ( $\neq$  3C 313).

**3C 368:** The source was observed three times in the MIR 5–15  $\mu\text{m}$  range, once with ISOPHOT on Oct. 18, 1996 (at 4.8 and 7.3  $\mu\text{m}$ ) and twice at 12  $\mu\text{m}$  with ISOCAM on Feb. 27, 1996, and on Oct. 27, 1997. The detections on Oct. 18, 1996 exceed by far the  $3\sigma$  upper limits from Feb. 27, 1996 and Oct. 27, 1997. The ISOPHOT chopped series show a perfectly oscillating on-off pattern leaving no doubt about the reality of the source detection. Since the explanation of strong MIR variability of 3C 368 seems unlikely to us, we suggest that the “MIR flux excess” during the observations on Oct. 18, 1996 is due to the passage of an asteroid (or comet), seen in the ISOPHOT aperture. However, no so far known asteroid has crossed the ISOPHOT observations (private communication by Thomas Müller). Nevertheless, an asteroid (or comet) seems to be the best explanation for the spurious 3C 368 detection and non-detections. Furthermore, at a temperature of 230 K as indicated by the MIR fluxes, an asteroid superposed on the 3C 368 field of view does not affect the FIR fluxes of 3C 368 at  $\lambda \geq 60 \mu\text{m}$ . Since 3C 368 was also detected with SCUBA at 450  $\mu\text{m}$  (Archibald et al. 2001), we ascribe the FIR-submm dust bump to 3C 368 and use the ISOCAM upper limit for MIR luminosity estimates. Nevertheless, we do not include 3C 368 in the discussion here.

**3C 390.3:** This broad line radio galaxy was also detected by IRAS, but both the ISOPHOT and the ISOCAM MIR data do not confirm the exceptionally high 12 and 25  $\mu\text{m}$  IRAS fluxes (Miley et al. 1984), thus making the mid- to far-IR colours of 3C 390.3 more similar to those of other BLRGs. The high IRAS fluxes could be contaminated by an asteroid or comet.

**3C 454.3:** This is one of the few examples of a flat spectrum quasar with a strong FIR dust bump sticking out of the bright synchrotron continuum. Other such examples are 3C 273 (Fig. 3), 3C 279 (Haas et al. 1998) and PG 1302-102 (Haas et al. 2000). We checked, whether the bump in 3C 454.3 could be due to another galaxy, but the only candidate on DSS and 2MASS images is an unresolved object about 15'' NNE with a K-star like optical-NIR SED. Therefore we conclude that the bright FIR dust bump is in fact caused by 3C 454.3.


## 4. Discussion

Figures 1 and 2 demonstrate that at least half ( $>40/75$ ) of the 3CR SEDs display a mid- to far-infrared bump due to thermal emission by dust. Even in the remaining cases the upper limits allow for considerable dust emission (Fig. 3, Table 2). In the following we consider the different classes (e.g. flat and steep spectrum quasars, BLRGs, FR 2 NLRGs and FR 1s), and investigate how the dust emission compares to the properties at other wavelengths. First, we discuss the SED shapes for representative class members. Then we study the statistical properties of the FR 2 NLRGs, quasars and BLRGs with respect to the *unification*, whereby we also consider other effects like those predicted by the receding torus model as well as evolution and environment.

### 4.1. SED shapes

The SEDs shown in Figs. 1 and 2 exhibit a striking diversity of shapes. Figure 4 shows five representative examples among the different AGN classes, i.e. radio and optical types:

- 1) Flat spectrum core dominated quasars (“FSQs”) like 3C 454.3: the synchrotron spectrum extends at a nearly constant flux level from the cm to the mm range, and declines at shorter wavelengths. In the FIR, the SEDs may show a thermal dust bump, as in the powerful FSQ 3C 454.3, or not, as in the less luminous BL Lac object 3C 371 (Fig. 3). In the context of unified schemes these flat spectrum sources are viewed nearly pole-on as proposed by Orr & Browne (1982).
- 2) Steep spectrum quasars like 3C 351: the synchrotron spectrum drops from the cm to the mm wavelengths, so that the thermal FIR dust emission bump becomes clearly visible. The optical spectra are quite blue, typical for an AGN outshining the host galaxy. Also, the SEDs show a power-law like flux increase, from the NIR over the MIR and peaking in the FIR, as seen also in optically identified PG quasars (Haas et al. 2003). With respect to the optical flux, the FIR dust bump may be quite pronounced with a steep slope from the NIR to the FIR as in 3C 48 at  $z \sim 0.3$  and 3C 318 at  $z \sim 1.5$  or more shallow as in 3C 249.1 at  $z \sim 0.3$  and 3C 298 at  $z \sim 1.5$  (Fig. 1). We do not see any difference between the SEDs of FR 2, lobe dominated and CSS types.
- 3) Broad line radio galaxies (BLRG) like 3C 33.1: the *infrared* and *radio* spectra are similar to those of the quasars, showing in the IR also the powerlaw rise between 5 and  $60 \mu\text{m}$ . In contrast, the *optical* spectrum is redder than that of quasars, probably due to a stronger contribution of the host galaxy relative to that of the AGN, and possible obscuration of the intrinsically blue central spectrum.
- 4) Narrow line radio galaxies (NLRG) of class FR 2 like 3C 405: the far-infrared and radio spectra are similar to those of quasars and BLRGs. Relative to the FIR, however, the emission in the optical and also in the NIR is weaker. This suppression at short wavelengths shows some range: while for 3C 405 the optical-NIR SEDs are only “moderately” reddened, the optical to IR slope is very steep for both 3C 65 and 3C 234 (Fig. 1). In the unified scheme the



**Fig. 4.** SED shapes of 3CR sources as representative examples of the different optical and radio classes, having good spectral data coverage. The symbols are as in Fig. 1.

NLRGs of class FR 2 represent edge-on quasars. 3C 234 shows polarised broad  $H\alpha$  but narrow  $H\beta$  lines (Antonucci 1984); hence it might be seen at the grazing angle to the edge of the dust torus. Some NLRGs of class FR 2 like 3C 321 and 3C 459 (Fig. 1) show a strong FIR bump and only a moderate MIR contribution, more reminiscent of the SEDs of starburst galaxies.

- 5) Narrow line and low-excitation radio galaxies (LERG) of class FR 1 like 3C 272.1, showing three basic SED features:
- 1) in the optical-NIR a bright bump due to the elliptical hosts, represented by a blackbody of  $T \approx 3500$  K peaking at about  $1 \mu\text{m}$ ;
  - 2) in the cm-mm wavelength range a steep synchrotron spectrum; and
  - 3) in the MIR-FIR a well discernible dust emission bump.


The examples shown in Fig. 4 illustrate the basic SED features. In the optical they arise from the host galaxy and the blue AGN continuum, in some cases reddened by extinction toward the central region. In the IR and radio ranges the emission from dust and from the jet and lobes can well be discerned. Remarkably, the SEDs exhibit also a diversity of shapes even within each AGN class, for example in the strength of the FIR dust bump or the slope between the NIR and FIR fluxes. For the optically selected PG quasars, which are thought to be seen nearly pole-on, such differences in the IR emission could be interpreted as evolution of the dust distribution around the heating sources (Haas et al. 2003). Although the diversity of SED shapes within each 3CR AGN class may also indicate such intrinsic differences, the orientation with respect to the line of sight is not yet known for our 3CR sample and it is still under debate, how far the observed near- and mid-IR emission is influenced by different viewing angles. Therefore, we will restrict our investigation to the orientation problem. Furthermore, average SEDs for each AGN class could be computed, but they will smear out the details recognisable in the individual SEDs. Therefore, instead of using averages, in the next subsection we prefer to analyse the properties from the individual SEDs.

#### 4.2. On the FR 2 unification

In this section we consider the following two basic classes: i) the steep spectrum quasars and the BLRGs, henceforth for short denoted as *quasars*, and ii) the FR 2 NLRGs, henceforth denoted as *galaxies*. The strategy to check the *unification* includes two steps: (1) to show that both the *quasars* and the *galaxies* exhibit a high mid- to far-IR luminosity ratio typical for AGNs, and (2) that the isotropic FIR-to-radio luminosity ratio is the same for *quasars* and *galaxies* at matched isotropic 178 MHz radio power. We consider only those sources shown in Fig. 1, which have sufficiently well sampled SEDs. The sample investigated here consists of 17 *galaxies* and 18 *quasars*, hence it can be considered as balanced. The set of sources shown in Fig. 3 and not included here comprises 20 *galaxies* and 15 – partly flat spectrum – *quasars*, and we will see below that their exclusion does not introduce a strong bias.

##### 4.2.1. AGN typical dust emission

The thermal 10–1000  $\mu\text{m}$  infrared luminosities of our sample span four orders of magnitude, from moderate  $10^{10} L_{\odot}$  over luminous  $10^{11} L_{\odot}$  and ultraluminous  $10^{12} L_{\odot}$  to hyperluminous  $10^{13}$ – $10^{14} L_{\odot}$  objects (Table 2 and abscissa of Fig. 5). The most luminous objects lie at the highest redshift, as is typical for flux limited samples like the 3CR catalogue.




**Fig. 5.** Mid- to far-infrared dust luminosity ratio versus 10–1000  $\mu\text{m}$  luminosity. The rectangular box shows the range of local starburst ULIRGs (from Klaas et al. 2001), each plotted with a “+” sign, and the dotted line indicates the upper end of their  $L_{\text{MIR}} / L_{\text{FIR}}$  distribution. The only 3CR source falling below this line is 3C 459.

The dust masses derived from the peak blackbodies shown in Fig. 1 range between  $10^6$  and  $10^9 M_{\odot}$  (Table 2), which is comparable to those found in PG quasars (Haas et al. 2003) and in ULIRGs (Klaas et al. 2001). Although the derived dust masses are possibly uncertain by a factor of ten, they do still provide a rough estimate of interstellar material associated with these objects.

In order to compare the 3CR sources with other samples of known properties, we have decomposed the IR emission into the MIR (10–40  $\mu\text{m}$ ) and FIR (40–1000  $\mu\text{m}$ ) luminosities (Table 2). For the *quasars* as well as for the *galaxies* the mid- to far-infrared luminosity ratio ranges between 0.4 and 5 as shown in Fig. 5. This is comparable to what is found for optically selected PG quasars (Haas et al. 2003). In contrast, those ultraluminous infrared galaxies (ULIRGs), in which the dust emission is powered mainly by starbursts and *not* by an AGN, populate the  $L_{\text{MIR}}/L_{\text{FIR}}$  range from 0.15 to 0.5 (Klaas et al. 2001). This is significantly lower than that of both the *quasars* and the *galaxies*. Therefore, the high mid- to far-infrared luminosity ratio above 0.5 together with the high dust luminosity ( $L_{\text{IR}} \geq 10^{11} L_{\odot}$ ) provides evidence for the presence of a powerful AGN in the *quasars* and the *galaxies* as well. Their AGN may be accompanied by starbursts, but for our purpose here the presence of a powerful buried AGN in the *galaxies* is relevant. It argues in favour of the *unification*. It should be noted here, that the orientation-dependent unification picture may not apply for the FR 1 and the low-power FR 2 sources, which have relatively low dust luminosities and masses.

A closer look on the mid-to-far-infrared luminosity ratio shows that the *quasars* lie above unity, while many of the *galaxies* lie below unity (Fig. 5). Such a difference between *quasars* and *galaxies* is consistent with unified models, if the proposed dust torus becomes optically thick at MIR wavelengths (Pier & Krolik 1992, 1993; van Bemmell & Dullemond 2003). In this case, if seen edge-on, our view towards the hot MIR emitting dust at the inner walls of the torus will suffer from extinction even at MIR wavelengths by the outer parts of the torus or by more extended dust. The actual degree of MIR opacity is not yet known. A value of  $A_{\text{MIR}} \sim 1$ –2 is




**Fig. 6.** *Top:* ratio of far-infrared dust power to radio lobe power  $R_{\text{dr}}$  versus radio lobe power, at restframes  $70 \mu\text{m}$  and 178 MHz. *Bottom:*  $R_{\text{dr}}$  versus  $L_{\text{IR}}$ . The dotted lines indicate the range of the dispersion discussed in the text, and the extreme sources 3C 321, 3C 405 and 3C 459 are labelled explicitly.

needed to explain the lower  $L_{\text{MIR}}/L_{\text{FIR}}$  ratios for the *galaxies*. Using standard extinction curves this value corresponds to  $A_V \sim 100\text{--}200$ , hence not exceptional with regard to that found in ULIRGs (e.g. Haas et al. 2001). Therefore we do not yet see any conflict with the *unification* and further investigations with more detailed observations as well as models may clarify that issue.

#### 4.2.2. Radio-infrared-comparison

So far we have found evidence for a powerful AGN in both the *quasars* and the *galaxies*. In a strict sense, however, the concept of *unification* requires that for an object drawn from the parent population any isotropic emission remains the same while rotating the viewing angle to its axis. Thus, for an ideal sample of parent objects the emitted isotropic FIR dust power should be the same for objects of identical isotropic lobe power. Therefore, following the ideas by Meisenheimer et al. (2001), we consider  $R_{\text{dr}}$ , the ratio of  $P_v$  at FIR wavelength  $70 \mu\text{m}$  ( $=4.3 \text{ THz}$ ) and  $P_v$  at radio frequency 178 MHz. Figure 6 shows  $R_{\text{dr}}$  versus the 178 MHz radio lobe power. All along the range of the 178 MHz radio lobe power, the distribution of  $R_{\text{dr}}$  for the *quasars* is strikingly similar to that of the *galaxies*. There is a marginal impression of a decline of  $R_{\text{dr}}$  with 178 MHz radio lobe power. This trend, however, is most likely caused by the



**Fig. 7.** Same as Fig. 6, but for the sources from Fig. 3 with IR upper limits only.

mathematical dependency “ $a/x$  versus  $x$ ”. It disappears when plotting  $R_{\text{dr}}$  versus  $L_{\text{IR}}$  (Fig. 6 bottom). As a quantitative check, we divide the luminosity range into two bins, one for luminosity below  $2 \times 10^{12} L_{\odot}$  and one above, and calculate for each bin the mean  $R_{\text{dr}}$  values. They lie in the range of  $10^{2.0 \pm 0.3}$  (Table 3). Again, a possible marginal decline of the mean  $R_{\text{dr}}$  with increasing luminosity is not yet statistically significant and we discuss possible reasons further below (Sect. 4.2.3).

With regard to the “matching strategy” applied by van Bemmell et al. (2000) and Meisenheimer et al. (2001) this means that, if we choose a *galaxy – quasar* pair matching in 178 MHz radio lobe power, i.e. lying in a vertical bin of Fig. 6, they exhibit a similar  $R_{\text{dr}}$ , exactly as predicted by the unified schemes. Some exceptional cases are discussed below. Also, if we plot  $R_{\text{dr}}$  versus redshift, or if we use the MIR power at  $15 \mu\text{m}$  (with values given in Table 2) instead of the FIR power at  $70 \mu\text{m}$ , all along the abscissa the  $R_{\text{dr}}$  distributions look quite similar for *galaxies* and *quasars*.

As illustrated in Fig. 7 with values given in Table 3, the FR 2 NLRGs and quasars from Fig. 3, for which we obtained only upper limits for the dust luminosity, populate the same range of  $R_{\text{dr}}$  as the detected sources from Fig. 1 do. Obviously the upper limits are so high that their constraints are consistent with our conclusions on the *unification*. Future data with the Spitzer and HERSCHEL satellites may consolidate this.

The range of  $R_{\text{dr}}$  values spans a factor of about ten around the average  $R_{\text{dr}} \sim 100$  as indicated by the dotted lines in Fig. 6. This appears large at first glance, but it becomes actually small with regard to the range of more than four decades of power. In view of the complexity of the objects, with a diversity of the strength of the thermal IR dust bump, evolving giant radio lobes (Kaiser et al. 1997; Blundell & Rawlings 1999) and various circumgalactic environments (Barthel & Miley 1988), such a dispersion is certainly not surprising. Therefore, the striking similarity of the  $R_{\text{dr}}$  distributions for *quasars* and *galaxies* provides strong evidence in favour of the basic role of the orientation-dependent *unification*.

#### 4.2.3. Other influences

When looking at the radio maps, it becomes clear that the jet and lobe phenomena are of course also influenced by the

**Table 3.** Average values of  $\log(R_{\text{dr}})$  for galaxies and quasars shown in Figs. 6 and 7. The means  $\pm$  standard deviations are given for two bins in luminosity of  $L_{\text{IR}}$  and  $P_{178 \text{ MHz}}$ .  $N$  denotes the number of sources in the bins. The averages are also calculated excluding extreme outliers.

| Bins                             | Quasars           | $N$            | Galaxies          | $N$            |
|----------------------------------|-------------------|----------------|-------------------|----------------|
| Detections:                      |                   |                |                   |                |
| $L_{\text{IR}}$                  |                   |                |                   |                |
| $<2 \times 10^{12} L_{\odot}$    | $1.946 \pm 0.290$ | 6              | $1.907 \pm 0.620$ | 11             |
|                                  |                   |                | $1.900 \pm 0.375$ | 9 <sup>a</sup> |
| $>2 \times 10^{12} L_{\odot}$    | $1.818 \pm 0.327$ | 12             | $1.859 \pm 0.650$ | 6              |
|                                  |                   |                | $1.640 \pm 0.408$ | 5 <sup>b</sup> |
| Detections:                      |                   |                |                   |                |
| $P_{178 \text{ MHz}}$            |                   |                |                   |                |
| $<2 \times 10^{28} \text{ W/Hz}$ | $1.970 \pm 0.301$ | 8              | $2.174 \pm 0.548$ | 10             |
|                                  |                   |                | $1.960 \pm 0.351$ | 8 <sup>c</sup> |
| $>2 \times 10^{28} \text{ W/Hz}$ | $1.774 \pm 0.308$ | 10             | $1.484 \pm 0.466$ | 7              |
|                                  |                   |                | $1.603 \pm 0.376$ | 6 <sup>d</sup> |
| Upper limits:                    |                   |                |                   |                |
| $P_{178 \text{ MHz}}$            |                   |                |                   |                |
| $<2 \times 10^{28} \text{ W/Hz}$ | $2.197 \pm 0.533$ | 7              | $2.117 \pm 0.382$ | 8              |
|                                  | $2.040 \pm 0.367$ | 6 <sup>e</sup> |                   |                |
| $>2 \times 10^{28} \text{ W/Hz}$ | $1.819 \pm 0.434$ | 8              | $1.935 \pm 0.337$ | 12             |
|                                  | $1.957 \pm 0.203$ | 7 <sup>f</sup> |                   |                |

<sup>a</sup> Excluding the extremes 3C 321 and 3C405.

<sup>b</sup> Excluding the extreme 3C 459.

<sup>c</sup> Excluding the extremes 3C 321 and 3C459.

<sup>d</sup> Excluding the extreme 3C 405.

<sup>e</sup> Excluding the extreme 3C 371 (BL Lac type, outside figure ranges).

<sup>f</sup> Excluding the extreme 3C 380.

circumgalactic environment and by evolution. Therefore it appears not appropriate, to consider and apply the three “principles” *unification*, *environment* and *evolution* on an exclusive either-or basis, rather the interplay of all three principles will help to understand the commonalities and differences of *galaxies* and *quasars*. The task now is to find strategies for disentangling the effects of the three principles. Firstly, we discuss our data in view of the receding torus model – this topic may be considered as *evolution* of the torus with luminosity.

#### 4.2.3.1 Receding torus model

In this modification of the general orientation-unifying model (Lawrence 1991; Hill et al. 1996; review by Simpson 2003, and references therein) both the *galaxies* and the *quasars* are basically surrounded by a dusty torus, but the inner wall of the torus recedes with increasing luminosity of the central engine due to dust sublimation ( $r_{\text{sub}} \propto L^{0.5}$ , e.g. Netzer & Laor 1993). Under the assumption, that the scale height of the dust torus is independent of luminosity, the opening angle of the bi-cone increases with luminosity.

When defining those objects with direct view to the broad emission line region simply as *quasars* and all others as *galaxies*, then the receding torus model predicts that the *quasar* fraction increases with luminosity. In fact, such a transition of

the *quasar* fraction from about 20% to higher values is observed at a 151 MHz luminosity threshold of about  $10^{26.5} \text{ W/Hz}$  (Willott et al. 2000). However, in all the luminosity bins above  $10^{26.5} \text{ W/Hz}$  the *quasar* fraction is 40% at the  $1-\sigma$  level (Willott et al. 2000). This constancy of the *quasar* fraction at high luminosities is surprising in the frame of a pure receding torus model. It suggests that above a luminosity threshold – possibly around  $10^{26.5} \text{ W/Hz}$  – the geometry of the receding torus does not change significantly and that the bi-cone opening angle has reached a maximum value compatible with other constraints. All of our FR 2 sources have 178 MHz luminosities higher than  $10^{26.5} \text{ W/Hz}$  (Fig. 6), hence lie above this threshold. With regard to our 3CR data we emphasize two points:

- 1) If *galaxies* and *quasars* belong to the same parent population, then – irrespective of whether a change of the source geometry with luminosity occurs or not – any orientation-dependent *unification* predicts that for a given luminosity bin the dust-to-lobe luminosity ratio  $R_{\text{dr}}$  should be the same for the *galaxy* and the *quasar*. If the *galaxy* had a lower  $R_{\text{dr}}$  than the *quasar*, this would point to a weaker central engine in the *galaxy*. In such a case, however, we cannot determine from the current data, whether this difference between *galaxy* and *quasar* is accompanied by different torus properties – and even if we could, this information would be only of moderate relevance with regard to other effects like lobe evolution and circumgalactic environment discussed in the next subsection. Nevertheless, our 3CR data show a striking similarity of the  $R_{\text{dr}}$  distributions for the *quasars* and the *galaxies* (Fig. 6) and any difference or decline with luminosity can be established at most at the  $1-\sigma$  level or less (Table 3).
- 2) If the bi-cone opening angle increases significantly with luminosity, then the covering angle of the dust, which absorbs the radiation from the central engine, decreases. Then one would expect that – relative to another measure of the AGN strength – the IR luminosity declines. For example an increase of the half-opening angle of the bi-cone from  $30^\circ$  to  $60-65^\circ$  results in a decrease of the dusty solid angle seen from the central source by a factor of about two ( $\cos 30^\circ / \cos 65^\circ = 2.05$ ). We searched for such a relative decline of the IR luminosity in two data samples:
  - a) if in *optically selected quasars* the blue continuum from the central engine is seen largely free of extinction, then the ratio of blue to IR luminosity should decline. We examined the sample of 64 Palomar Green quasars with good near- and mid-IR data and covering four decades in luminosity (Fig. 4 and Table 2 in Haas et al. 2003). We find that for all bins above  $L_B = 10^{11} L_{\odot}$  (corresponding to  $M_B < -24 \text{ mag}$ ) the mean ratios of  $L_{\text{NIR}}/L_B$  and  $L_{\text{MIR}}/L_B$  are clearly constant at the  $1-\sigma$  level with values about  $1.1 \pm 0.5$ . But for the low luminosity bins with  $L_B < 10^{11} L_{\odot}$  the ratios are higher by a factor of about 1.5 and 1.7, respectively, although the differences are only marginally significant;
  - b) for the *radio-loud 3CR objects* one may expect a decline of  $R_{\text{dr}}$  with luminosity, but our data do not show

a clear trend at a statistically significant level (Fig. 6, Table 3).

Thus we conclude that the effects of the receding torus with increasing bi-cone opening angle may be significant at lower luminosity levels for Seyfert galaxies and radio galaxies having 178 MHz luminosities below  $10^{26.5}$  W/Hz, but that at the high luminosity level of quasars the receding torus model in its originally proposed rigid form plays only a minor role and needs some further refinement.

One refinement should focus on the torus scale height: for the receding torus model the assumption of a constant torus scale height is most critical. It may not be justified if the AGN phenomenon is accompanied by nuclear starbursts and if the star formation rate (SFR) grows with AGN luminosity. Then supernovae will blow-up the scale height  $h$  of the torus (and the disk) by injection of kinetic energy. For example, three-dimensional numerical simulations of the interstellar matter located in the combined gravitational potential of both a central black hole and a stellar system yield  $h \propto \text{SFR}^{0.5} \propto L^{0.5}$  (Wada & Norman 2002). If this is the case, then both  $h$  and  $r_{\text{sub}}$  scale with  $L^{0.5}$  and the opening angle of the bi-cone does not change with luminosity. This refinement would explain (1) that the quasar fraction remains constant at 151 MHz luminosities above  $10^{26.5}$  W/Hz, and (2) that the ratios  $L_{\text{NIR}}/L_B$  and  $L_{\text{MIR}}/L_B$  do not decline at luminosities above  $L_B = 10^{11} L_{\odot}$  ( $M_B < -24$  mag).

#### 4.2.3.2 Environment and lobe evolution

The ISO data suggest, that the FIR and MIR luminosities actually yield good estimates for the strength of the AGN. Now, under the premise that the *unification* is basically valid, sensitive infrared measurements offer to pursue the following strategy proposed by Barthel & Arnaud (1996): we adopt the IR luminosity as primary measure for the AGN power and examine the strength of the radio lobes. Then deviations in the dust-to-lobe luminosity ratio allow, free of orientation bias, to explore the additional influence of environment and evolution onto the extended radio lobes. In the following, we consider the extreme cases of high and low  $R_{\text{dr}}$  values in Fig. 6.

High values of  $R_{\text{dr}}$  are expected for sources, which are either rather dust-rich or do not show strong radio lobe activity. Examples are the “cool” FR 2 galaxies 3C 321 and 3C 459 both with a high FIR contribution possibly enhanced by starbursts. Very extreme examples for high  $R_{\text{dr}}$  sources are radio-quiet quasars lying even above the upper plot range of Fig. 6.

Low values of  $R_{\text{dr}}$  are expected for sources, which are either rather dust-poor or do show exceptional radio lobe activity – possibly caused by a special circumgalactic environment or by a high evolutionary state of the lobes. For example, 3C 405 (Cygnus A) has exceptionally bright radio lobes (Barthel & Arnaud 1996) and shows a low  $R_{\text{dr}} < 10$ , confirming the suggestion by Barthel & Arnaud, that radio luminosity is not generally a good measure of AGN power. The ISO data show that the dust emission may serve as a better isotropic indicator for the AGN strength. Also the galaxy 3C 295 lies at the low end of the FIR-to-radio distribution, indicating high efficiency in

creating the radio lobe luminosity. Note that 3C 295 is located in a dense X-ray bright cluster, suggesting the influence of the environment in enhancing the lobe power. Also 3C 380 is peculiar in that it shows both a relatively flat radio spectrum probably seen pole-on and bright FR 2 lobes. A possible way out of this dilemma is that there is an intrinsic bend in the jet (Wilkinson et al. 1991). In this case a strong interaction with the circumgalactic medium may provide an extraordinarily high radio luminosity, too.

The extreme cases with high and low  $R_{\text{dr}}$  show that the dispersion of the  $R_{\text{dr}}$  distribution in general may be well understood in terms of evolution of the radio lobes and by effects of the circumgalactic environment.

If evolutionary and environmental effects are superposed on the pure orientation-dependent unification, then it is a priori not clear whether the increased dispersion affects *galaxies* and *quasars* in the same manner, or whether a bias is introduced. A detailed investigation of the possible selection effects (Chapter 3.5.3 in Haas 2001) shows that there is a bias, which limits the strategies to check the *unification*: a radio source, which has already been recognised as a *quasar* due to some criteria, will on average always exhibit a higher IR luminosity than a source with matching radio luminosity, which has not yet been identified as a *quasar*. Larger data samples are required in order to overcome this “strategical bias”.

## 5. Conclusions

We have obtained sensitive Infrared SEDs for a large sample of 3CR radio galaxies and quasars. In order to check the orientation-dependent *unification* of powerful FR 2 sources we considered 17 *galaxies* and 18 *quasars* with good IR detections. The main conclusions for this sample are:

- 1) Both the *quasars* and the *galaxies* exhibit a high mid- to far-IR luminosity ratio typical for AGNs.
- 2) The ratio of IR dust to radio lobe luminosity  $R_{\text{dr}}$  is similar for the *quasars* and the *galaxies*. Therefore, this result provides strong evidence in favour of the orientation-dependent *unification* of luminous ( $L_{\text{IR}} \gtrsim 10^{11} L_{\odot}$ ) narrow line FR 2 radio *galaxies* with *quasars*.
- 3) The dispersion of the  $R_{\text{dr}}$  distribution might be caused by evolution by the radio lobes as well as effects of the circumgalactic environment.
- 4) The receding torus model in its original formulation is not valid at the high luminosity of our sources ( $L_{\text{IR}} \gtrsim 10^{11} L_{\odot}$ ). Therefore we propose as a refinement that the scale height of the torus increases also with luminosity due to supernovae from starbursts accompanying the AGN.

Compared with the detections, the upper limits of the remaining 20 *galaxies* and 15 *quasars* are so high that they do not provide any deep constraints or conflicts with the conclusions above.

*Acknowledgements.* We thank the anonymous referee for constructive suggestions and for encouraging us to discuss the receding torus model. The ISOPHOT Data Centre at MPIA is supported by Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR) with funds of

Bundesministerium für Bildung und Forschung, grant No. 50 QI 0201. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). Martin Haas thanks for grants from the Nordrhein-Westfälische Akademie der Wissenschaften, funded by the Federal State Nordrhein-Westfalen and the Federal Republic of Germany. This research is essentially based on data from the ISO and JCMT/SCUBA Archives. For literature and photometry search NED and SIMBAD were used. It is a pleasure to thank IRAM for discretionary observing time with the 30 m telescope at Pico Veleta.

## References

- Akujor, C. E., Lüdke, E., Browne, I. W. A., et al. 1994, *A&AS*, 105, 247
- Andreani, P., Fosbury, R. A. E., van Bemmell, I., & Freudling, W. 2002, *A&A*, 381, 389
- Antonucci, R. R. J. 1984, *ApJ*, 278, 499
- Antonucci, R. R. J., & Barvainis, R. 1990, *ApJ*, 363, L17
- Antonucci, R. R. J., & Miller, J. S. 1985, *ApJ*, 297, 621
- Archibald, E. N., Dunlop, J. S., Hughes, D. H., et al. 2001, *MNRAS*, 323, 417
- Barthel, P. D. 1989, *ApJ*, 336, 606
- Barthel, P. D., & Arnaud, K. A. 1996, *MNRAS*, 283, L45
- Barthel, P. D., & Miley, G. K. 1988, *Nature*, 333, 319
- Begelman, M. C., Blandford, R. D., & Rees, M. J. 1984, *RvMP*, 56, 255
- Bennett, C. L., Halpern, M., Hinshaw, G., et al. 2003, *ApJS*, 148, 1
- Best, P. N., Röttgering, H. J. A., Bremer, M. N., et al. 1998, *MNRAS*, 301, L15
- Blundell, K., & Rawlings, S. 1999, *Nature*, 399, 330
- Cesarsky, C., Abergel, A., Agnese, P., et al. 1996, *A&A*, 315, L32
- Chiaberge, M., Capetti, A., & Celotti, A. 1999, *A&A*, 349, 77
- Chiaberge, M., Capetti, A., & Celotti, A. 2000, *A&A*, 355, 873
- Chini, R., Kreysa, E., & Biermann, P. L. 1989a, *A&A*, 219, 87
- Chini, R., Biermann, P. L., Kreysa, E., & Gmünd H.-P. 1989b, *A&A*, 221, L3
- Cohen, M. H., Ogle, P. M., Tran, H. D., Goodrich, R. W., & Miller, J. S. 1999, *AJ*, 118, 1963
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, Jr. H. G., et al. 1991, *Third reference catalog of bright galaxies, version 3.9* (Berlin: Springer-Verlag)
- Fanti, C., Pozzi, F., Fanti, R., et al. 2000, *A&A*, 358, 499
- Freudling, W., Siebenmorgen, R., & Haas, M. 2003, *ApJ*, 599, L13
- Haas, M. 2001, *Habilitationschrift Universität Heidelberg*, available from <http://www.astro.ruhr-uni-bochum.de/haas/>
- Haas, M., Chini, R., Meisenheimer, K., et al. 1998, *ApJ*, 503, L109
- Haas, M., Müller, S. A. H., Chini, R., et al. 2000, *A&A*, 354, 453
- Haas, M., Klaas, U., Müller, S. A. H., Chini, R., & Coulson, I. 2001, *A&A*, 367, L9
- Haas, M., Klaas, U., Müller, S. A. H., et al. 2003, *A&A*, 402, 87
- Heckman, T. M., Chambers, K. C., & Postman, M. 1992, *ApJ*, 391, 39
- Heckman, T. M., O’Dea, Ch. P., Baum, S. A., & Laurikainen, E. 1994, *ApJ*, 428, 65
- Hes, R., Barthel, P. D., & Hoekstra, H. 1995, *A&A*, 303, 8
- Hildebrand, R. H. 1983, *QJRAS*, 24, 267
- Hill, G. J., Goodrich, R. W., & Depoy, D. L. 1996, *ApJ*, 462, 163
- Hine, R. G., & Longair, M. S. 1979, *MNRAS*, 188, 111
- Hoekstra, H., Barthel, P. D., & Hes, R. 1997, *A&A*, 319, 757
- Hughes, D. H., Robson, E. I., Dunlop, J. S., & Gear, W. K. 1993, *MNRAS*, 263, 607
- Kaiser, C. R., Dennett-Thorpe, J., & Alexander, P. 1997, *MNRAS*, 292, 723
- Kellermann, K. I., Sramek, R., Schmidt, M., et al. 1989, *AJ*, 98, 1195
- Kessler, M. F., Steinz, J. A., Anderegg, M. E., et al. 1996, *A&A*, 315, L27
- Kessler, M. F., Müller, T. G., Arviset, C., García-Lario, P., & Prusti, T. 2000, *The ISO Handbook, SAI/2000-035/Dc*, ESA publ.
- Klaas, U., Haas, M., Müller, S. A. H., et al. 2001, *A&A*, 379, 823
- Klaas, U., Laureijs, R., Radovich, M., Schulz, B., & Wilke, K. 2002, *ISOPHOT calibration accuracies, V5.0* (adapted to OLP10), *ISO Explanatory Library Doc. SAI/02-xxx/rp*, ESA publ.
- Kreysa, E., Gemünd, H.-P., Gromke, J., et al. 1998, *SPIE*, 3357, 319
- Kühr, H., Witzel, A., Pauliny-Toth, I. I. K., & Nauber, U. 1981, *A&AS*, 45, 367
- Laing, R. A., Riley, J. M., & Longair, M. S. 1983, *MNRAS*, 204, 151
- Laureijs, R., & Klaas, U. 1999, *ISOPHOT Error Budgets V1.0*, *ISO Explanatory Library Doc. SAI/98-091/dc*, ESA publ.
- Laureijs, R., Klaas, U., Richards, P. J., Schulz, B., & Abraham, P. 2002, *The ISO Handbook, Vol. IV: PHT – the Imaging Photopolarimeter, version 2.0*, *SAI/99-069/dc*, ESA publications
- Lawrence, A. 1991, *MNRAS*, 252, 586
- Lemke, D., Klaas, U., Abolins, J., et al. 1996, *A&A*, 315, L64
- Lilly, S. J., & Longair, M. S. 1984, *MNRAS*, 211, 833
- Lilly, S. J., Longair, M. S., & Miller, L. 1985, *MNRAS*, 214, 109
- Meisenheimer, K., Haas, M., Müller, S. A. H., et al. 2001, *A&A*, 372, 719
- Miley, G., Neugebauer, G., Soifer, B. T., et al. 1984, *ApJ*, 278, L79
- Neugebauer, G., Green, R. F., Matthews, K., et al. 1987, *ApJS*, 63, 615
- Netzer, H., & Laor, A. 1993, *ApJ*, 404, L51
- Orr, M. J. L., & Browne, I. W. A. 1982, *MNRAS*, 200, 1067
- Pier, E. A., & Krolik, J. H. 1992, *ApJ*, 401, 99
- Pier, E. A., & Krolik, J. H. 1993, *ApJ*, 418, 673
- Polletta, M., Courvoisier, Th., Hooper, E. J., & Wilkes, B. J. 2000, *A&A*, 362, 75
- Robson, E. I., Leeuw, L. L. Stevens, J. A., & Holland, W. S. 1998, *MNRAS*, 301, 935
- Saunders, R., Baldwin, J. E., Rawlings, S., Warner, P. J., & Miller, L. 1989, *MNRAS*, 238, 777
- Siebenmorgen, R., Freudling, W., Krügel, E., & Haas, M. 2004, *A&A*, 421, 129
- Simpson, C. 2003, *New Astron. Rev.*, 47, 211
- Simpson, C., Ward, M., & Wall, J. V. 2000, *MNRAS*, 319, 963
- Singal, A. K. 1996, *MNRAS*, 278, 1069
- Spinrad, H., Djorgowski, S., Marr, J., & Aguilar, L. 1985, *PASP*, 97, 932
- Steppe, H., Salter, C. J., Chini, R., et al. 1988, *A&AS*, 75, 317
- Steppe, H., Liechti, S., Mauersberger, R., et al. 1992, *A&AS*, 96, 441
- Steppe, H., Paubert, G., Sievers, A., et al. 1993, *A&AS*, 102, 611
- Stevens, J. A., Robson, E. I., Gear, W. K., et al. 1998, *ApJ*, 502
- Stickel, M., Lemke, D., Klaas, U., et al. 2000, *A&A*, 359, 865
- Urry, C. M., & Padovani, P. 1995, *PASP*, 107, 803
- van Bemmell, I. M., Barthel, P., & de Graauw, Th. 2000, *A&A*, 359, 523
- van Bemmell, I. M., & Bertoldi, F. 2001, *A&A*, 368, 414
- van Bemmell, I. M., & Dullemond, C. P. 2003, *A&A*, 404, 1
- Wada, K., & Norman, C. 2002, *ApJ*, 566, L21
- Wilkinson, P. N., Akujor, C. E., Cornwell, T. J., & Saikia, D. J. 1991, *MNRAS*, 248, 86
- Willott, C. J., Rawlings, S., Blundell, K. M., & Lacy, M. 2000, *MNRAS*, 316, 449
- Willott, C. J., Rawlings, S., Archibald, E. N., & Dunlop, J. S. 2002, *MNRAS*, 331, 435

# Online Material

**Table 2.** Types, redshifts and luminosity distances, restframe luminosities, dust temperatures and masses. The luminosity distance  $D_L$  is calculated adopting a  $\Lambda$  cosmology with  $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$ ,  $\Omega_{\text{matter}} = 0.27$  and  $\Omega_{\Lambda} = 0.73$ . The luminosities  $L_{\text{opt}}$  ( $0.5\text{--}1 \mu\text{m}$ ),  $L_{\text{NIR}}$  ( $3\text{--}10 \mu\text{m}$ ),  $L_{\text{MIR}}$  ( $10\text{--}40 \mu\text{m}$ ) and  $L_{\text{FIR}}$  ( $40\text{--}1000 \mu\text{m}$ ) are computed in the restframes of the objects from the “envelopes” of the SEDs. The luminosities give the total emission from host galaxy and the core. The IR luminosities largely represent the thermal dust luminosities, except for some sources (like 3C 371 or 3C 380), which do not exhibit any prominent thermal bump and are excluded from the discussion. The thermal dust emission power  $\nu P_{\nu}$  has been computed at restframe wavelengths 15 and  $70 \mu\text{m}$ , respectively. The 178 MHz radio lobe power is computed from the observed 178 MHz fluxes, applying a K-correction with a spectral index  $\alpha = 0.5$ . The choice of  $\alpha$  has only little impact on the resulting lobe power and the relations discussed.  $R_{\text{dr}}$  is the ratio of dust power to radio lobe power at 15 and  $70 \mu\text{m}$ , respectively. The values of  $T_p$  refer to the peak blackbody shown in the SED plots, the uncertainty is 5–10 K and can raise to 50% in cases of poor FIR upper limits. The dust mass  $M_{\text{dust}}$  is estimated according to Formula 6 in Stickett et al. (2000) following Hildebrand (1983), with uncertainties up to an order of magnitude. A “~” marks highly uncertain  $L_{\text{MIR}}/L_{\text{FIR}}$ ,  $T_{\text{dust}}$  and  $M_{\text{dust}}$  values due to poor dust bump determinations or upper limit constraints, and these sources are not used in the discussion.

| 3CR   | Type* |          | Redshift | $D_L$<br>Mpc | $L_{\text{opt}}$   | $L_{\text{NIR}}$   | $L_{\text{MIR}}$   | $L_{\text{FIR}}$   | $L_{\text{MIR}}/$ | $P(15 \mu\text{m})$ | $P(70 \mu\text{m})$ | $P(178 \text{ MHz})$ | $R_{\text{dr}}$ | $R_{\text{dr}}$ | $T_p$ | $M_d$ |
|-------|-------|----------|----------|--------------|--------------------|--------------------|--------------------|--------------------|-------------------|---------------------|---------------------|----------------------|-----------------|-----------------|-------|-------|
|       |       |          |          |              | $\log [L_{\odot}]$ | $\log [L_{\odot}]$ | $\log [L_{\odot}]$ | $\log [L_{\odot}]$ | $L_{\text{FIR}}$  |                     |                     |                      |                 |                 |       |       |
| 2.0   | QSR   | C CSS    | 1.037    | 6943         | 11.39              | <12.00             | 12.80              | 12.61              | 1.54              | 39.11               | 39.09               | 37.34                | 59              | 57              | 55    | 7.92  |
| 9.0   | QSR   | L        | 2.012    | 15852        | 12.76              | 13.85              | <13.87             | <13.62             | ~1.79             | <40.28              | <40.16              | 38.22                | <115            | <87             | ~50   | ~8.99 |
| 16.0  | LERG  | FR 2     | 0.405    | 2193         | 10.38              | <11.31             | <11.85             | <11.90             | ~0.89             | <38.11              | <38.45              | 36.13                | <95             | <206            | ~40   | ~7.81 |
| 17.0  | BLRG  | ?        | 0.220    | 1081         | 10.37              | 10.67              | <10.65             | <10.96             | ~0.49             | <37.06              | <37.54              | 35.74                | <21             | <63             | ~40   | ~6.91 |
| 19.0  | NLRG  | FR 2     | 0.482    | 2701         | 10.80              | <11.29             | <11.50             | <11.47             | ~1.06             | <37.87              | <38.05              | 36.36                | <32             | <49             | ~40   | ~7.41 |
| 20.0  | NLRG  | FR 2     | 0.174    | 830          | 9.93               | 10.52              | 10.95              | 11.18              | 0.58              | 37.21               | 37.68               | 35.83                | 24              | 70              | 35    | 7.45  |
| 33.0  | NLRG  | FR 2     | 0.059    | 261          | 10.31              | 9.93               | <10.06             | 10.14              | 0.83              | <36.45              | 36.72               | 34.92                | <34             | 63              | 35    | 6.40  |
| 33.1  | BLRG  | FR 2     | 0.181    | 869          | 10.15              | 10.96              | 11.00              | 10.61              | 2.47              | 37.47               | 37.14               | 35.36                | 129             | 60              | 50    | 6.08  |
| 34.0  | NLRG  | FR 2     | 0.689    | 4177         | 11.01              | <11.70             | <12.09             | <12.03             | ~1.16             | <38.41              | <38.58              | 36.76                | <45             | <67             | ~40   | ~7.95 |
| 42.0  | NLRG  | FR 2     | 0.395    | 2128         | 10.60              | <10.69             | <11.12             | <11.28             | ~0.70             | <37.42              | <37.86              | 36.14                | <19             | <53             | ~40   | ~7.23 |
| 46.0  | NLRG  | FR 2     | 0.437    | 2400         | 10.91              | <11.37             | <11.63             | <11.61             | ~1.06             | <37.98              | <38.17              | 36.18                | <64             | <99             | ~40   | ~7.54 |
| 47.0  | QSR   | L        | 0.425    | 2322         | 11.09              | 11.62              | 12.01              | 11.91              | 1.24              | 38.38               | 38.48               | 36.56                | 65              | 84              | 35    | 8.18  |
| 48.0  | QSR   | C CSS    | 0.367    | 1950         | 11.64              | 12.21              | 12.44              | 12.47              | 0.93              | 38.79               | 39.02               | 36.72                | 117             | 203             | 35    | 8.70  |
| 49.0  | NLRG  | FR 2 CSS | 0.621    | 3675         | 10.48              | <11.65             | 12.28              | 12.34              | 0.88              | 38.52               | 38.90               | 36.58                | 87              | 209             | 40    | 8.26  |
| 61.1  | NLRG  | FR 2     | 0.186    | 895          | 9.89               | 10.02              | 10.28              | 10.59              | 0.49              | 36.47               | 37.16               | 35.76                | 5               | 25              | 40    | 6.54  |
| 65.0  | NLRG  | FR 2     | 1.176    | 8125         | 11.11              | 12.71              | 13.10              | 12.51              | 3.94              | 39.42               | 38.95               | 37.50                | 83              | 29              | 80    | 7.17  |
| 67.0  | BLRG  | FR 2 CSS | 0.310    | 1600         | 10.50              | <10.96             | <11.17             | <11.17             | ~1.01             | <37.54              | <37.75              | 35.80                | <55             | <90             | ~40   | ~7.11 |
| 68.1  | QSR   | L        | 1.238    | 8664         | 12.13              | <12.31             | <12.65             | <13.06             | ~0.39             | <38.96              | <39.63              | 37.49                | <30             | <139            | ~40   | ~8.01 |
| 79.0  | NLRG  | FR 2     | 0.256    | 1283         | 10.66              | 11.08              | 11.58              | 11.38              | 1.60              | 37.94               | 37.94               | 36.08                | 73              | 73              | 55    | 6.63  |
| 84.0  | NLRG  | FR 1     | 0.018    | 74           | 10.61              | 10.46              | 10.87              | 10.80              | 1.18              | 37.11               | 37.30               | 33.85                | 1831            | 2779            | 35    | 7.01  |
| 109.0 | BLRG  | FR 2     | 0.306    | 1577         | 11.28              | 12.06              | 12.22              | 11.69              | 3.38              | 38.67               | 38.21               | 36.12                | 362             | 126             | 55    | 6.95  |
| 138.0 | QSR   | C CSS    | 0.759    | 4707         | 11.45              | 12.24              | 12.38              | <12.29             | ~1.23             | 38.81               | <38.86              | 37.14                | 47              | <52             | ~40   | ~8.20 |
| 234.0 | NLRG  | FR 2     | 0.185    | 890          | 10.56              | 11.64              | 11.95              | 11.50              | 2.81              | 38.39               | 38.04               | 35.76                | 426             | 189             | 45    | 7.17  |
| 245.0 | QSR   | L        | 1.029    | 6876         | 11.83              | 12.49              | 13.09              | 12.60              | 3.03              | 39.38               | 38.94               | 37.32                | 115             | 42              | 85    | 7.08  |
| 249.1 | QSR   | L        | 0.311    | 1607         | 11.60              | 11.70              | 11.66              | 11.16              | 3.14              | 38.11               | 37.57               | 35.83                | 191             | 55              | 60    | 6.26  |
| 268.3 | BLRG  | FR 2 CSS | 0.371    | 1975         | 10.27              | <11.22             | <11.57             | <11.54             | ~1.08             | <37.90              | <38.09              | 36.02                | <77             | <119            | ~40   | ~7.46 |
| 270.0 | NLRG  | FR 1     | 0.007    | 31           | 10.28              | 9.50               | 8.84               | 9.05               | 0.61              | 35.10               | 35.51               | 33.03                | <120            | 305             | 28    | 5.81  |
| 272.1 | LERG  | FR 1     | 0.003    | 12           | 10.25              | 9.43               | 8.13               | 8.33               | 0.63              | 34.46               | 34.77               | 31.90                | 361             | 736             | 27    | 5.13  |
| 273.0 | QSR   | FR 2     | 0.158    | 747          | 12.13              | 12.28              | 12.13              | 12.00              | 1.35              | 38.56               | 38.47               | 35.91                | 453             | 368             | 30    | 8.53  |
| 274.0 | NLRG  | FR 1     | 0.004    | 16           | 10.29              | 9.56               | 8.43               | 8.50               | 0.86              | 34.72               | 35.01               | 32.56                | 144             | 286             | 35    | 4.75  |
| 274.1 | NELR  | FR 2     | 0.422    | 2303         | 10.67              | <11.56             | <12.12             | <12.34             | ~0.60             | <38.38              | <38.93              | 36.35                | <108            | <382            | ~40   | ~8.30 |
| 277.1 | QSR   | FR 2     | 0.321    | 1667         | 10.90              | <11.23             | <11.60             | <11.16             | ~2.78             | <37.93              | <37.59              | 35.76                | <147            | <67             | ~40   | ~6.95 |
| 277.2 | NLRG  | FR 2     | 0.766    | 4761         | 10.89              | <11.86             | 12.71              | 12.13              | 3.82              | 38.89               | 38.51               | 36.89                | 101             | 42              | 90    | 6.58  |
| 280.0 | NLRG  | FR 2     | 0.996    | 6602         | 10.98              | <12.23             | 12.97              | 12.36              | 4.02              | 39.19               | 38.73               | 37.49                | 49              | 17              | 90    | 6.81  |
| 280.1 | QSR   | FR 2     | 1.659    | 12481        | 11.95              | <12.67             | <13.03             | <13.12             | ~0.82             | <39.38              | <39.61              | 37.70                | <48             | <82             | ~40   | ~8.28 |
| 284.0 | NLRG  | FR 2     | 0.239    | 1187         | 10.82              | <11.19             | <11.49             | 11.55              | 0.89              | <37.84              | 38.11               | 35.58                | <182            | 344             | 40    | 7.49  |
| 286.0 | QSR   | C CSS    | 0.849    | 5410         | 11.94              | 12.01              | 12.66              | 12.28              | 2.42              | 38.87               | 38.71               | 37.33                | 35              | 24              | 75    | 7.00  |
| 287.0 | QSR   | C CSS    | 1.055    | 7094         | 11.92              | <12.18             | <12.51             | <12.54             | ~0.93             | <38.84              | <39.10              | 37.40                | <28             | <50             | ~40   | ~8.47 |
| 288.1 | QSR   | L        | 0.961    | 6313         | 11.75              | 12.01              | 12.66              | 12.51              | 1.43              | 38.97               | 39.02               | 37.03                | 87              | 98              | 55    | 7.78  |
| 289.0 | NLRG  | FR 2     | 0.967    | 6362         | 11.21              | <11.82             | <12.16             | <12.15             | ~1.02             | <38.49              | <38.71              | 37.16                | <21             | <36             | ~40   | ~8.08 |
| 293.0 | LERG  | FR 1     | 0.045    | 197          | 10.52              | 10.07              | 9.98               | 10.43              | 0.36              | 36.32               | 36.85               | 34.04                | 191             | 655             | 26    | 7.41  |
| 295.0 | NLRG  | FR 2     | 0.461    | 2559         | 11.09              | 11.29              | 11.81              | 11.99              | 0.67              | 38.07               | 38.57               | 37.15                | 8               | <26             | 40    | 7.94  |
| 298.0 | QSR   | FR 2     | 1.436    | 10427        | 12.77              | 13.32              | 13.52              | 13.19              | 2.17              | 39.88               | 39.70               | 38.24                | 44              | 29              | 40    | 9.06  |
| 299.0 | NLRG  | FR 2 CSS | 0.367    | 1950         | 10.48              | <11.05             | <11.68             | <12.35             | ~0.21             | <37.78              | <38.94              | 36.05                | <53             | <780            | ~40   | ~8.30 |
| 303.1 | NLRG  | FR 2 CSS | 0.267    | 1347         | 10.61              | 10.56              | 10.83              | 11.11              | 0.52              | 37.10               | 37.71               | 35.55                | 36              | 145             | 40    | 7.07  |
| 309.1 | QSR   | C CSS    | 0.905    | 5858         | 12.05              | 12.40              | 12.79              | 12.39              | 2.51              | 39.10               | 38.82               | 37.36                | 55              | 29              | 70    | 7.22  |
| 318.0 | QSR   | FR 2 CSS | 1.574    | 11690        | 11.62              | <12.86             | 13.56              | 13.38              | 1.51              | 39.80               | 39.89               | 37.76                | 111             | 134             | 55    | 8.64  |
| 321.0 | NLRG  | FR 2     | 0.096    | 435          | 10.60              | 10.60              | 11.33              | 11.36              | 0.92              | 37.45               | 37.86               | 34.76                | 497             | 1258            | 35    | 7.49  |
| 323.1 | QSR   | FR 2     | 0.264    | 1329         | 11.42              | 11.62              | 11.53              | 10.88              | 4.49              | 38.03               | 37.38               | 35.61                | 261             | 59              | 60    | 5.98  |
| 324.0 | NLRG  | FR 2     | 1.206    | 8385         | 11.25              | 12.12              | <13.00             | <13.01             | ~0.98             | <39.16              | <39.56              | 37.55                | <41             | <103            | ~50   | ~8.42 |

**Table 2.** continued

| 3CR                | Type <sup>a</sup> |          | Redshift | $D_L$<br>Mpc | $L_{\text{opt}}$    | $L_{\text{NIR}}$    | $L_{\text{MIR}}$    | $L_{\text{FIR}}$    | $L_{\text{MIR}}/$ | $P(15 \mu\text{m})$ | $P(70 \mu\text{m})$ | $P(178 \text{ MHz})$ | $R_{\text{dr}}$  | $R_{\text{dr}}$  | $T_{\text{d}}$ | $M_{\text{d}}$  |
|--------------------|-------------------|----------|----------|--------------|---------------------|---------------------|---------------------|---------------------|-------------------|---------------------|---------------------|----------------------|------------------|------------------|----------------|-----------------|
|                    |                   |          |          |              | log [ $L_{\odot}$ ] | $L_{\text{FIR}}$  | log [W]             | log [W]             | log [W]              | 15 $\mu\text{m}$ | 70 $\mu\text{m}$ | [K]            | log $M_{\odot}$ |
| 325.0              | QSR               | L        | 0.860    | 5498         | 11.43               | <13.10              | 13.07               | 12.42               | 4.54              | 39.54               | 38.84               | 37.14                | 252              | 50               | 75             | 7.14            |
| 330.0              | NLRG              | FR 2     | 0.550    | 3168         | 11.08               | 11.29               | <11.54              | <11.62              | -0.83             | <37.86              | <38.20              | 36.87                | <10              | <21              | ~40            | ~7.57           |
| 332.0              | BLRG              | FR 2     | 0.152    | 716          | 10.87               | 10.70               | 10.78               | <10.80              | -0.93             | 37.14               | <37.38              | 35.05                | 122              | <211             | ~40            | ~6.75           |
| 334.0              | QSR               | L        | 0.555    | 3203         | 11.54               | <12.03              | 12.20               | 11.68               | 3.30              | 38.58               | 38.11               | 36.47                | 127              | 44               | 75             | 6.39            |
| 336.0              | QSR               | L        | 0.927    | 6036         | 11.79               | 11.93               | <12.64              | <12.75              | -0.77             | <38.83              | <39.32              | 37.09                | <55              | <168             | ~40            | ~8.69           |
| 337.0              | NLRG              | FR 2     | 0.635    | 3777         | 10.66               | <11.60              | <12.08              | <12.04              | -1.09             | <38.36              | <38.60              | 36.66                | <50              | <86              | ~40            | ~7.96           |
| 340.0              | NLRG              | FR 2     | 0.775    | 4830         | 10.77               | <11.75              | <12.22              | <12.37              | -0.70             | <38.51              | <38.94              | 36.83                | <48              | <131             | ~40            | ~8.32           |
| 343.1              | NLRG              | FR 2 CSS | 0.750    | 4638         | 11.13               | <11.83              | <12.46              | <12.89              | -0.37             | <38.65              | <39.48              | 36.84                | <64              | <431             | ~40            | ~8.85           |
| 351.0              | QSR               | L        | 0.371    | 1975         | 11.80               | 12.22               | 12.27               | 11.87               | 2.50              | 38.68               | 38.39               | 36.13                | 360              | 185              | 50             | 7.30            |
| 352.0              | NLRG              | FR 2     | 0.806    | 5071         | 10.57               | <11.56              | <12.10              | <12.27              | -0.68             | <38.37              | <38.84              | 36.92                | <28              | <84              | ~40            | ~8.22           |
| 356.0              | NLRG              | FR 2     | 1.079    | 7296         | 11.23               | 11.52               | 12.02               | 12.25               | 0.59              | 38.28               | 38.83               | 37.27                | 10               | 37               | 40             | 8.20            |
| 368.0              | NLRG              | FR 2     | 1.131    | 7738         | 11.35               | <12.39              | 12.92               | <12.82              | -1.26             | 39.21               | <39.36              | 37.41                | 63               | <89              | ~40            | ~8.73           |
| 371.0              | BL Lac            |          | 0.051    | 224          | 10.77               | <10.70              | <10.63              | <10.23              | -2.49             | <37.10              | <36.73              | 33.59                | <3227            | <1373            | ~40            | ~6.11           |
| 380.0              | QSR               | C CSS    | 0.692    | 4199         | 11.88               | <12.28              | <12.33              | <11.85              | -3.01             | <38.76              | <38.31              | 37.46                | <20              | <7               | ~65            | ~6.81           |
| 381.0 <sup>x</sup> | NLRG              | FR 2     | 0.160    | 758          | 10.43               | 10.91               | 11.10               | 10.65               | 2.82              | 37.51               | 37.15               | 35.34                | 147              | 64               | 40             | 6.70            |
| 390.3              | BLRG              | FR 2     | 0.056    | 246          | 10.30               | 10.64               | 10.65               | 10.23               | 2.61              | 37.13               | 36.76               | 34.80                | 211              | 90               | 40             | 6.13            |
| 401.0              | LERG              | FR 2     | 0.201    | 976          | 10.72               | <10.15              | <10.51              | <11.00              | -0.33             | <36.52              | <37.56              | 35.67                | <7               | <78              | ~40            | ~6.94           |
| 405.0              | NLRG              | FR 2     | 0.056    | 246          | 9.95                | 10.55               | 11.38               | 11.30               | 1.19              | 37.64               | 37.84               | 37.07                | 4                | 6                | 50             | 6.78            |
| 427.1              | LERG              | FR 2     | 0.572    | 3324         | 10.59               | <10.90              | <11.99              | <12.29              | -0.50             | <37.92              | <38.85              | 36.89                | <11              | <89              | ~40            | ~8.21           |
| 437.0              | NLRG              | FR 2     | 1.480    | 10827        | 11.15               | <12.35              | <13.33              | <13.12              | -1.63             | <39.50              | <39.60              | 37.76                | <55              | <70              | ~60            | ~8.21           |
| 441.0              | NLRG              | FR 2     | 0.707    | 4311         | 10.72               | <11.72              | <12.26              | <12.42              | -0.68             | <38.52              | <38.99              | 36.81                | <51              | <150             | ~40            | ~8.36           |
| 445.0              | BLRG              | FR 2     | 0.056    | 247          | 10.26               | 10.87               | 10.98               | 10.50               | 3.02              | 37.46               | 37.03               | 34.52                | 855              | 323              | 50             | 5.96            |
| 454.3              | FSQ               | C OVV    | 0.860    | 5498         | 12.50               | <12.92              | 13.53               | 12.79               | 5.59              | 39.81               | 39.13               | 37.06                | 566              | 118              | 100            | 7.10            |
| 459.0              | NLRG              | FR 2     | 0.220    | 1081         | 10.89               | 11.07               | 11.83               | 12.24               | 0.39              | 37.93               | 38.81               | 35.85                | 120              | 908              | 40             | 8.18            |
| 460.0              | NLRG              | FR 2     | 0.268    | 1353         | 10.39               | <10.15              | <10.43              | <10.98              | -0.28             | <36.55              | <37.58              | 35.56                | <10              | <104             | ~40            | ~6.94           |


\* The types are abbreviated as:

QSR = quasar, BLRG = broad line radio galaxy, NLRG = narrow line radio galaxy, LERG = low-excitation radio galaxy;

C = core dominated, L = lobe dominated, FR 1, FR 2 = Fanaroff-Riley classes 1 and 2;

CSS = compact steep spectrum source, OVV = optically violent variable.

<sup>x</sup> 3C 381 could be classified as BLRG based on a noisy  $H_{\beta}$  line, which appears to be broader than the [OIII] $\lambda$ 5007 line, but with considerable uncertainty (Saunders et al. 1989). A new spectrum obtained at the KECK telescope shows that 3C 381 should be classified as NLRG (van Bemmel, private communication).



**Fig. 3.** Spectral energy distributions of the remaining sources not used in the discussion, which are either flat spectrum sources or have only poorly constrained thermal bumps. The meaning of the lines and labels is as in Fig. 1. For the poorly constrained cases, however, the blackbody curves and the envelopes are shown here as *dotted* lines in order to emphasize that they represent only a maximum possible dust contribution. For the blackbody, if not sufficiently constrained by the data, we adopted the average of  $T_p \approx 40$  K derived from the good cases shown in Fig. 1. The long dashed lines for 3C 138, 3C 273 and 3C 380 indicate the extrapolation of the synchrotron emission towards shorter wavelengths.