

LETTER TO THE EDITOR

A remarkable recurrent nova in M 31: The X-ray observations[★]

M. Henze¹, J.-U. Ness¹, M. J. Darnley², M. F. Bode², S. C. Williams², A. W. Shafter³, M. Kato⁴, and I. Hachisu⁵

¹ European Space Astronomy Centre, PO Box 78, 28692 Villanueva de la Cañada, Madrid, Spain
e-mail: mhenze@sciops.esa.int

² Astrophysics Research Institute, Liverpool John Moores University, IC2 Liverpool Science Park, Liverpool, L3 5RF, UK

³ Department of Astronomy, San Diego State University, San Diego, CA 92182, USA

⁴ Department of Astronomy, Keio University, Hiyoshi, Yokohama 223-8521, Japan

⁵ Department of Earth Science and Astronomy, College of Arts and Sciences, The University of Tokyo, Komaba, Meguro-ku, 153-8902 Tokyo, Japan

Received 13 January 2014 / Accepted 7 February 2014

ABSTRACT

Context. Another outburst of the recurrent M 31 nova M31N 2008-12a was announced in late November 2013. Optical data suggest an unprecedentedly short recurrence time of approximately one year.

Aims. In this Letter we address the X-ray properties of M31N 2008-12a.

Methods. We requested *Swift* monitoring observations shortly after the optical discovery. We estimated source count rates and extracted X-ray spectra from the resulting data. The corresponding ultraviolet (UV) data were also analysed.

Results. The nova M31N 2008-12a was clearly detected as a bright supersoft X-ray source (SSS) only six days after the well-constrained optical discovery. It displayed a short SSS phase of two weeks' duration and an exceptionally hot X-ray spectrum with an effective black-body temperature of ~ 97 eV. During the SSS phase the X-ray light curve displayed significant variability that might have been accompanied by spectral variations. The very early X-ray variability was found to be anti-correlated with simultaneous variations in the UV flux.

Conclusions. The X-ray properties of M31N 2008-12a coherently point towards a high-mass white dwarf in the nova system. This object might be a promising Type Ia supernova progenitor. We rediscovered additional X-ray detections of M31N 2008-12a that are consistent with our data and increase the number of known nova outbursts to seven. This nova is an exceptional object that merits further attention in the future.

Key words. galaxies: individual: M 31 – novae, cataclysmic variables – X-rays: binaries – stars: individual: M31N 2008-12a

1. Introduction

Recurrent novae (RNe) are those for which more than one nova outburst has been observed. As in the case for classical novae, the outburst is caused by a thermonuclear runaway in an accreted envelope of matter on top of a white dwarf (WD) in a close binary system (for recent reviews see Bode & Evans 2008). A proportion of the accreted matter is ejected during the outburst and forms a drastically enlarged pseudo-photosphere, thereby leading to the optical nova. The matter remaining on the WD hosts stable hydrogen burning soon after the outburst. This process powers a supersoft X-ray source (SSS; photon energies below 1 keV, see Parmar et al. 1998) that can be observed once the ejected matter becomes optically thin to soft X-rays.

While the optical characteristics of a nova depend on the properties of the ejected envelope, only X-ray measurements allow us to observe the WD directly and study its physics. Novae with a short SSS phase are expected to host a massive WD (e.g. Hachisu & Kato 2006, 2010). In addition, a high effective temperature of the SSS indicates a high-mass WD (Sala & Hernanz 2005). A massive WD is required to produce recurrence times that are short enough to be observed (Truran & Livio 1986; Yaron et al. 2005). The shortest time between two confirmed nova outbursts was eight years for the Galactic RN U Sco

(Schaefer 2010). In the single-degenerate scenario of type Ia supernova (SN Ia) progenitors, RNe correspond to the final stage of the binary evolution (see Hachisu et al. 1999b,a).

Novae in our neighbour galaxy M 31 (distance 780 kpc; Holland 1998; Stanek & Garnavich 1998) have been observed for almost a century. Currently, almost a thousand nova outbursts are known in M 31 (see the online catalogue¹ of Pietsch et al. 2007). Dedicated X-ray monitoring observations of M 31 novae and archival studies produced a sample of 79 novae with SSS counterparts (see Henze et al. 2014, hereafter HPH2014, and references therein). A statistical analysis of this sample revealed strong correlations between optical and X-ray parameters, showing that novae that decline fast in the optical are hot in X-rays with short SSS durations (HPH2014).

Nova M31N 2008-12a was first discovered in December 2008 by Kabashima & Nishiyama². Two additional eruptions were found in October 2011 and October 2012; the second was classified as the outburst of a He/N nova by Shafter et al. (2012). For a comprehensive description of the optical properties and the identification of the 2011 and 2012 outbursts with M31N 2008-12a see the accompanying Letter by Darnley et al. (2014, hereafter DWB2014). In November 2013, another

¹ <http://www.mpe.mpg.de/~m31novae/opt/m31/index.php>

² http://www.cbat.eps.harvard.edu/CBAT_M31.html#2008-12a

[★] Tables 1 and 2 are available in electronic form at <http://www.aanda.org>

outburst was detected by [Tang et al. \(2013\)](#) and this Letter focusses on the results from the subsequent X-ray monitoring.

2. Observations and data analysis

Soon after the announcement of the November 2013 outburst of M31N 2008-12a we requested a target of opportunity (ToO) X-ray monitoring with *Swift* ([Gehrels et al. 2004](#)). The extremely short recurrence time suggested a high-mass WD, the SSS phase of which we detected and characterised successfully. All observations are summarised in Table 1. Additionally, in Table 2 we list all observations of a *Swift* monitoring programme after the 2012 outburst that did not detect an X-ray source.

The *Swift* X-ray telescope (XRT; [Burrows et al. 2005](#)) data analysis was based on the cleaned event files. We derived the count rates given in Table 1 using the source statistics (*sosta*) tool within the HEASoft XIMAGE package (version 4.5.1.). This method takes into account the background, estimated in a source-free region around the object, as well as corrections for sampling dead time, vignetting and point spread function (PSF). The upper limits in Tables 1 and 2 were computed assuming a PSF constructed from the merged data of all detections.

The source and background photons were extracted from the event files within the HEASoft Xselect environment (version v2.4c). The spectral analysis used XSPEC ([Arnaud 1996](#), version 12.8.1g) with the Tübingen-Boulder ISM absorption model (TBabs in XSPEC), the photoelectric absorption cross-sections from [Balucinska-Church & McCammon \(1992\)](#), and the ISM abundances from [Wilms et al. \(2000\)](#). All spectra were binned to include at least one photon per bin and fitted in XSPEC assuming Poisson statistics according to [Cash \(1979\)](#).

The ultraviolet (UV) magnitudes are given in the Vega system and were determined using the *uvot* source tool. All magnitudes assume the *Swift* UV/optical telescope (UVOT, [Roming et al. 2005](#)) photometric system ([Poole et al. 2008](#)) and have not been corrected for extinction.

All upper limits correspond to 3σ confidence and all error ranges to 1σ confidence unless otherwise noted.

3. Results

Nova M31N 2008-12a was clearly detected by *Swift* as a bright X-ray source following its 2013 outburst (first announced by [Henze et al. 2013a](#)). Based on the unprecedentedly short recurrence time, we had expected a massive WD that should display a fast SSS appearance (according to [Hachisu & Kato 2010](#), HPH2014). Consequently, the *Swift* monitoring was designed to start as soon as possible to constrain the beginning of the SSS phase. Surprisingly, the SSS turned on even faster than expected and the first observation, approximately six days after discovery, detected a bright SSS. The source was monitored with *Swift* until the SSS turned off two weeks later (see Table 1).

The source position, as determined from the merged XRT event files of all detections, is RA = 00h45m29.29s, Dec = +41°54′08″.5 (J2000; 90% confidence error 3″.6). Although this is 4″.8 away from the optical coordinates reported by [Tang et al. \(2013\)](#) (RA = 00h45m28.89s, Dec = +41°54′10″.2) both positions are consistent ($\sim 2\sigma$ level) because of the large uncertainties. Moreover, the appearance of the X-ray source soon after the optical nova outburst as well as its characteristics (see below) leave little doubt that both events are related. Therefore, we identify the transient X-ray source as the counterpart of M31N 2008-12a.

The best-fit parameters for a simultaneous fit of all X-ray spectra with a black-body model are $kT = (97_{-4}^{+5})$ eV and




Fig. 1. *Swift* XRT spectrum extracted from merged event files. This is shown only for visualising the high effective temperature. Model fitting was performed on the individual spectra. In grey we show a scaled spectrum of the bright and hot nova M31N 2007-12b ([Pietsch et al. 2011](#)), which is a suspected RN ([Bode et al. 2009](#)), for comparison.




Fig. 2. Evolution of X-ray count rate **a)** and effective black-body temperature **b)** of M31N 2008-12a during the 2013 *Swift* monitoring. This assumes an outburst on November 26.6, 2013 UT (see also Table 1). The error bars in time represent the duration of the individual observation **a)** or the time between the grouped observations **b)**. Panel **a)** Upper limits are indicated by black open triangles. A blue asterisk marks the first upper limit estimated from the *Swift* monitoring after the 2012 outburst (see Table 2). Panel **b)** Groups of observations have been fitted simultaneously with the N_{H} fixed to the global best fit. The red dashed line shows the overall best-fit temperature derived in Sect. 3.

$N_{\text{H}} = (1.4_{-0.3}^{+0.2}) \times 10^{21} \text{ cm}^{-2}$. A χ^2 test statistic for this fit gave a reduced χ^2 of 1.19 for 574 degrees of freedom. The estimated kT is exceptionally high for an M 31 nova (cf. HPH2014). From the M 31 reddening maps of [Montalto et al. \(2009\)](#) we derived an approximate $E(B - V) \sim 0.26$ for the position of M31N 2008-12a. This corresponds to $N_{\text{H}} \sim 1.8 \times 10^{21} \text{ cm}^{-2}$ (using the relation of [Güver & Özel 2009](#)), which is consistent with the best-fit value.

A spectrum extracted from the combined event files of all detections is shown in Fig. 1. This spectrum was grouped to include at least 20 counts per bin and is only used for visualisation. Practically all source photons have energies below 1 keV. This source is clearly a SSS. Because the Wien tail of the spectrum extends to relatively high energies we can infer that the SSS is unusually hot even without the use of specific spectral models (which are controversial, see e.g. [Ness et al. 2013](#)).

In Fig. 2a we show the X-ray light curve of the nova during the 2013 outburst. This plot and Table 1 assume that the optical outburst occurred between the last upper limit and the first detection reported by [Tang et al. \(2013\)](#), i.e. on November 26.6,




Fig. 3. Short-term X-ray and UV light curves for the five snapshots of *Swift* observation 00032613005 on day six after outburst (see Table 1).

2013 UT (with an uncertainty of ± 0.5 d). Following the first detection of the SSS on day six after the outburst its count rate increased and subsequently showed significant variability (see Fig. 2a). This variation was not due to instrumental vignetting or bad columns. The count rate started dropping on day 16 and the source disappeared a few days later. The last detection in an individual observation on day 18 showed a count rate that was approximately an order of magnitude below the peak level (see also Table 1).

The SSS might still have been detected at the 2σ level in the merged 14.3 ks of the last five upper limit observations. However, with $(7.7 \pm 3.4) \times 10^{-4}$ ct s $^{-1}$ its count rate would have been more than an order of magnitude below the bright phase. Therefore, this bright SSS phase which indicates stable hydrogen burning (see e.g. Sala & Hernanz 2005), was definitely over at this point. Consequently, we assume a SSS turn-off time (t_{off}) of (19 ± 1) d. This is consistent with the results of the low-cadence *Swift* monitoring following the 2012 eruption that failed to detect a SSS ~ 20 d after outburst (see Table 2 and Fig. 2a).

The X-ray variability might have been accompanied by spectral variation. In Fig. 2b we show the results of a simultaneous fitting (with N_{H} fixed to the global best fit) of separate groups of consecutive observations with similar count rate. Figure 2b shows indications for lower effective temperatures at the very beginning and end of the SSS phase. We also fitted the second and sixth temperature bin (low count rates) and the third and fifth bin (high count rates) in Fig. 2b simultaneously with fixed absorption. These groups of observations represent the high- and low-luminosity states of M31N 2008-12a, with the fourth temperature bin possibly belonging to a transitory stage. The resulting best-fit temperatures were found to be different, albeit with weak significance: $kT = (105 \pm 3)$ eV (high) vs. (91 ± 4) eV (low).

Furthermore, we found significant X-ray variability within the first (6 ks) *Swift* observation of the 2013 outburst (see also Henze et al. 2013b). This count rate variation was accompanied by variability in the UV. Subsequently, no significant UV variations were found against a gradual decline (see Table 1). In Fig. 3 we show both short-term light curves based on the individual *Swift* snapshots during the first detection. They appear to be anti-correlated. This process could indicate the gradual emergence of the SSS. Therefore, we estimate a SSS turn-on time (t_{on}) of (6 ± 1) d.

4. Discussion

4.1. Variability

Significant X-ray variability during the early stages of the SSS phase has been observed in several Galactic novae (e.g. KT Eri,

RS Oph; Bode et al. 2010; Osborne et al. 2011). One interpretation of this effect is a variable absorption column during the emergence of the SSS from the expanding ejecta and/or optically thick wind (see e.g. Hachisu & Kato 2006). Short-term X-ray variability of various kinds has been observed in the M 31 nova sample (see HPH2014).

The anti-correlation between X-ray and UV flux during the early SSS phase (Fig. 3) and the later possible temperature variations (Fig. 2b) might point towards intrinsic variations in the WD temperature. Schwarz et al. (2011) suggested a similar scenario to explain the anti-correlated X-ray and UV variability during the SSS phase of the Galactic nova V458 Vul (see also Ness et al. 2009). They drew an interesting parallel to the variability in persistent SSSs which might originate in accretion-rate variations that cause the WD photosphere to expand or shrink (see Reinsch et al. 2000; Greiner & Di Stefano 2002). Based on its frequent outbursts and evidence from the optical data (DWB2014), M31N 2008-12a is likely to have a high accretion rate. The observed variability might indicate that accretion had been re-established as early as six days after outburst.

4.2. Theoretical implications of the short recurrence time

Theoretically, a recurrence period (τ_r) of the order of one year is only expected for extremely massive WDs ($\geq 1.35 M_{\odot}$) with a high mass accretion rate exceeding $10^{-7} M_{\odot} \text{ yr}^{-1}$ (Wolf et al. 2013; Saio et al., in prep.). A massive WD is consistent with the high temperature suggested in our X-ray spectra (Sect. 3, Fig. 1). For such a short τ_r , however, the accreted mass is too small (several $10^{-7} M_{\odot}$) for the pseudo-photosphere to expand to a red-giant size. Therefore, we would expect a relatively high effective surface temperature even at the optical maximum that results in a faint peak magnitude. The shorter the τ_r , the fainter the peak magnitude.

The low peak brightness of M31N 2008-12a (never above 18th mag, see DWB2014) seems consistent with this scenario. While high accretion rates can also trigger hydrogen shell flashes on less massive WDs without causing a nova (see Jose et al. 1993), the observed evidence suggests that M31N 2008-12a was experiencing actual nova outbursts. Optically faint novae with a fast and hot SSS phase might generally be good RN candidates.

An extremely massive WD can be born as an ONe WD in the final stage of stellar evolution for a narrow range of zero-age main-sequence mass ($\sim 8\text{--}10 M_{\odot}$). Another possibility is mass-accretion onto an initially less massive WD in a binary system. If a (CO) WD was born with $M_{\text{WD}} < 1.07 M_{\odot}$ and accumulates mass until reaching the Chandrasekhar mass it will explode as a SN Ia; RNe with short τ_r correspond to immediate progenitors of SNe Ia in the single degenerate scenario (Hachisu et al. 1999b,a).

4.3. Population picture

In Fig. 4 we show the position of M31N 2008-12a in the five-parameter space of the correlations presented by HPH2014 for the M 31 nova sample. The SSS time scales were unprecedentedly short, also compared to Galactic novae where, for example, the fastest t_{on} was displayed by the RN U Sco (≥ 8 d; Schlegel et al. 2010). The average SSS temperature seems considerably higher than measured for any other M 31 nova and might have been even hotter during the high count rate observations (see Sect. 3 and Fig. 2b). Overall, the X-ray parameter relations (Figs. 4a and b) are broadly consistent with the population trends. Nova M31N 2008-12a therefore provides an important extension of these relations into previously unpopulated regions of the parameter space.




Fig. 4. Double-logarithmic plots of the nova parameter correlations from HPH2014. The solid grey lines indicate a robust power-law fit with corresponding 95% confidence regions in light grey. The correlations displayed are **a)** t_{on} vs. t_{off} , **b)** black-body kT vs. t_{off} , **c)** R -band optical decay time $t_{2,R}$ vs. t_{on} , and **d)** expansion velocity vs. t_{on} . All timescales are in units of days after outburst. Overplotted in red are the parameters of M31N 2008-12a as derived in Sect. 3 or by DWB2014.

The two optical parameters, $t_{2,R}$ and v_{exp} (see DWB2014), show stronger deviations from the rest of the nova sample. This behaviour might be consistent with a fainter nova outburst, as was suggested by theoretical models (Sect. 4.2). However, for the two correlations in Figs. 4c and d the scatter is generally larger. At this stage, it is difficult to say whether the connection between the optical and X-ray parameters of M31N 2008-12a is significantly different from the overall population trend.

4.4. Previous X-ray outbursts

During a search of the literature we found that M31N 2008-12a had actually been detected in X-rays before the first optical outburst was reported. It was initially discovered as the “recurrent supersoft X-ray transient” RX J0045.4+4154 by White et al. (1995) based on archival ROSAT data. Because no optical nova was known before 2008, this source was not recognised as a nova counterpart by Pietsch et al. (2005) in the course of their archival analysis. White et al. (1995) reported two X-ray outbursts in February 1992 and January 1993. The recurrence time of approximately one year as well as the short duration of the outbursts and the high black-body temperature (~ 90 eV) estimated by White et al. (1995) agree well with our results.

Furthermore, another X-ray outburst was reported by Williams et al. (2004) in *Chandra* data taken in September 2001. Because this observation used the HRC detector no X-ray spectrum exists. The outburst was shorter than five months. Overall, there is strong evidence that M31N 2008-12a had at least three outbursts before 2008. A follow-up paper will study the optical and X-ray archives in more detail.

5. Conclusions

The X-ray properties of M31N 2008-12a clearly suggest that the binary system hosts a high-mass WD. The fast recurrence

times and optical evidence indicate a high accretion rate, the re-establishing of which could explain the observed early X-ray and UV variability. The M 31 galaxy remains a treasure island for nova science and might hold the key to the understanding of the (potentially numerous) optically faint RNe and their impact on SN Ia progenitor theories. The position of M31N 2008-12a will be watched closely in particular in late 2014.

Acknowledgements. We thank the referee, Eric Schlegel, for a careful reading of the manuscript. We wish to thank Margarita Hernanz for constructive comments on an early version of this Letter. We are grateful to the *Swift* Team for making the ToO observations possible, in particular N. Gehrels, the duty scientists, as well as the science planners. M.H. acknowledges support from an ESA fellowship. S.C.W. acknowledges PhD funding from STFC. A.W.S. acknowledges support from NSF grant AST1009566.

References

- Arnaud, K. A. 1996, in *Astronomical Data Analysis Software and Systems V*, eds. G. H. Jacoby, & J. Barnes, ASP Conf. Ser., 101, 17
- Balucinska-Church, M., & McCammon, D. 1992, *ApJ*, 400, 699
- Bode, M. F., & Evans, A. 2008, *Classical Novae*, eds. M. F. Bode, & A. Evans (Cambridge University Press)
- Bode, M. F., Darnley, M. J., Shafter, A. W., et al. 2009, *ApJ*, 705, 1056
- Bode, M. F., Osborne, J. P., Page, K. L., et al. 2010, *ATel*, 2392, 1
- Burrows, D. N., Hill, J. E., Nousek, J. A., et al. 2005, *Space Sci. Rev.*, 120, 165
- Cash, W. 1979, *ApJ*, 228, 939
- Darnley, M. J., Williams, S. C., Bode, M. F., et al. 2014, *A&A*, 563, L9 [DWB2014]
- Gehrels, N., Chincarini, G., Giommi, P., et al. 2004, *ApJ*, 611, 1005
- Greiner, J., & Di Stefano, R. 2002, *A&A*, 387, 944
- Güver, T., & Özel, F. 2009, *MNRAS*, 400, 2050
- Hachisu, I., & Kato, M. 2006, *ApJS*, 167, 59
- Hachisu, I., & Kato, M. 2010, *ApJ*, 709, 680
- Hachisu, I., Kato, M., & Nomoto, K. 1999a, *ApJ*, 522, 487
- Hachisu, I., Kato, M., Nomoto, K., & Umeda, H. 1999b, *ApJ*, 519, 314
- Henze, M., Ness, J., Bode, M. F., Darnley, M. J., & Williams, S. C. 2013a, *ATel*, 5627, 1
- Henze, M., Ness, J., Bode, M. F., et al. 2013b, *ATel*, 5633, 1
- Henze, M., Pietsch, W., Haberl, F., et al. 2014, *A&A*, 563, A2 [HPH2014]
- Holland, S. 1998, *AJ*, 115, 1916
- Jose, J., Hernanz, M., & Isern, J. 1993, *A&A*, 269, 291
- Montalto, M., Seitz, S., Riffeser, A., et al. 2009, *A&A*, 507, 283
- Ness, J.-U., Drake, J. J., Beardmore, A. P., et al. 2009, *AJ*, 137, 4160
- Ness, J.-U., Osborne, J. P., Henze, M., et al. 2013, *A&A*, 559, A50
- Osborne, J. P., Page, K. L., Beardmore, A. P., et al. 2011, *ApJ*, 727, 124
- Parmar, A. N., Kahabka, P., Hartmann, H. W., Heise, J., & Taylor, B. G. 1998, *A&A*, 332, 199
- Pietsch, W., Fliri, J., Freyberg, M. J., et al. 2005, *A&A*, 442, 879
- Pietsch, W., Haberl, F., Sala, G., et al. 2007, *A&A*, 465, 375
- Pietsch, W., Henze, M., Haberl, F., et al. 2011, *A&A*, 531, A22
- Poole, T. S., Breeveld, A. A., Page, M. J., et al. 2008, *MNRAS*, 383, 627
- Reinsch, K., van Teeseling, A., King, A. R., & Beuermann, K. 2000, *A&A*, 354, L37
- Roming, P. W. A., Kennedy, T. E., Mason, K. O., et al. 2005, *Space Sci. Rev.*, 120, 95
- Sala, G., & Hernanz, M. 2005, *A&A*, 439, 1061
- Schaefer, B. E. 2010, *ApJS*, 187, 275
- Schlegel, E. M., Schaefer, B., Pagnotta, A., et al. 2010, *ATel*, 2430, 1
- Schwarz, G. J., Ness, J.-U., Osborne, J. P., et al. 2011, *ApJS*, 197, 31
- Shafter, A. W., Hornoch, K., Ciardullo, J. V. R., Darnley, M. J., & Bode, M. F. 2012, *ATel*, 4503, 1
- Stanek, K. Z., & Garnavich, P. M. 1998, *ApJ*, 503, L131
- Tang, S., Cao, Y., & Kasliwal, M. M. 2013, *ATel*, 5607
- Truran, J. W., & Livio, M. 1986, *ApJ*, 308, 721
- White, N. E., Giommi, P., Heise, J., Angelini, L., & Fantasia, S. 1995, *ApJ*, 445, L125
- Williams, B. F., Garcia, M. R., Kong, A. K. H., et al. 2004, *ApJ*, 609, 735
- Wilms, J., Allen, A., & McCray, R. 2000, *ApJ*, 542, 914
- Wolf, W. M., Bildsten, L., Brooks, J., & Paxton, B. 2013, *ApJ*, 777, 136
- Yaron, O., Prialnik, D., Shara, M. M., & Kovetz, A. 2005, *ApJ*, 623, 398

Table 1. *Swift* observations of nova M31N 2008-12a.

ObsID	Exp ^a [ks]	Date ^b [UT]	MJD ^b [d]	Δt^c [d]	UV ^d [mag]			Rate [10 ⁻² ct s ⁻¹]	$L_{0.2-1.0}^e$ [10 ³⁸ erg s ⁻¹]
					<i>uvm2</i>	<i>uvw1</i>	<i>uvw2</i>		
00032613005	6.0	2013-12-03.03	56 629.03	6.43	–	–	19.4 ± 0.1	1.3 ± 0.2	1.0 ± 0.1
00032613006	2.0	2013-12-04.09	56 630.09	7.49	19.8 ± 0.3	>19.5	19.7 ± 0.2	1.8 ± 0.4	1.3 ± 0.3
00032613007	2.0	2013-12-04.70	56 630.70	8.10	19.4 ± 0.2	19.4 ± 0.2	19.3 ± 0.2	1.7 ± 0.3	1.2 ± 0.3
00032613008	1.7	2013-12-05.23	56 631.23	8.63	20.0 ± 0.4	19.6 ± 0.2	19.8 ± 0.3	2.2 ± 0.4	1.7 ± 0.3
00032613009	1.9	2013-12-05.70	56 631.70	9.10	19.8 ± 0.3	>19.9	19.9 ± 0.3	4.0 ± 0.5	3.0 ± 0.4
00032613010	2.0	2013-12-06.10	56 632.10	9.50	19.9 ± 0.3	19.8 ± 0.3	19.5 ± 0.2	4.0 ± 0.5	3.0 ± 0.4
00032613011	2.0	2013-12-06.63	56 632.63	10.03	>20.2	19.7 ± 0.3	20.0 ± 0.3	3.8 ± 0.5	2.8 ± 0.4
00032613012	3.3	2013-12-07.10	56 633.10	10.50	20.1 ± 0.2	>19.8	20.2 ± 0.3	1.3 ± 0.2	1.0 ± 0.2
00032613013	1.8	2013-12-07.57	56 633.57	10.97	20.1 ± 0.4	19.4 ± 0.2	19.6 ± 0.2	1.7 ± 0.4	1.3 ± 0.3
00032613014	1.5	2013-12-08.03	56 634.03	11.43	19.6 ± 0.3	>20.0	19.9 ± 0.3	3.0 ± 0.5	2.2 ± 0.4
00032613015	1.8	2013-12-08.57	56 634.57	11.97	19.9 ± 0.4	20.0 ± 0.3	>20.3	3.2 ± 0.5	2.4 ± 0.4
00032613016	4.4	2013-12-09.03	56 635.04	12.44	19.8 ± 0.2	19.7 ± 0.3	20.0 ± 0.2	2.5 ± 0.3	1.9 ± 0.2
00032613017	3.8	2013-12-10.44	56 636.44	13.84	>20.6	19.9 ± 0.2	20.0 ± 0.2	3.2 ± 0.3	2.4 ± 0.2
00032613018	2.1	2013-12-11.04	56 637.04	14.44	20.0 ± 0.4	>20.2	>20.4	1.6 ± 0.3	1.2 ± 0.2
00032613019	3.0	2013-12-11.31	56 637.30	14.70	20.1 ± 0.3	19.6 ± 0.3	20.2 ± 0.3	1.7 ± 0.3	1.3 ± 0.2
00032613020	1.9	2013-12-11.57	56 637.57	14.97	>20.0	19.6 ± 0.3	19.9 ± 0.3	1.9 ± 0.4	1.4 ± 0.3
00032613021	1.9	2013-12-12.04	56 638.04	15.44	19.9 ± 0.2	–	–	1.3 ± 0.3	1.0 ± 0.2
00032613022	1.9	2013-12-12.32	56 638.32	15.72	20.4 ± 0.3	–	–	1.3 ± 0.3	1.0 ± 0.2
00032613023	1.9	2013-12-12.51	56 638.52	15.92	20.2 ± 0.4	19.8 ± 0.3	20.0 ± 0.3	1.8 ± 0.4	1.4 ± 0.3
00032613024	2.1	2013-12-12.78	56 638.78	16.18	>20.9	–	–	1.7 ± 0.3	1.3 ± 0.2
00032613025	2.0	2013-12-13.10	56 639.11	16.51	–	19.8 ± 0.2	–	0.9 ± 0.2	0.7 ± 0.2
00032613026	1.9	2013-12-13.31	56 639.31	16.71	–	20.4 ± 0.3	–	0.6 ± 0.2	0.5 ± 0.2
00032613027	1.9	2013-12-13.58	56 639.58	16.98	–	20.2 ± 0.2	–	0.5 ± 0.2	0.4 ± 0.1
00032613028	2.0	2013-12-13.78	56 639.78	17.18	>20.2	>20.2	>20.5	0.3 ± 0.1	0.2 ± 0.1
00032613029	0.3	2013-12-14.24	56 640.24	17.64	–	–	–	<3.4	<2.5
00032613030	0.9	2013-12-14.38	56 640.38	17.78	–	–	–	0.9 ± 0.4	0.7 ± 0.3
00032613031	1.9	2013-12-14.52	56 640.52	17.92	–	–	–	<3.4	<2.5
00032613032	1.9	2013-12-14.78	56 640.78	18.18	>20.2	20.1 ± 0.4	20.3 ± 0.3	0.5 ± 0.2	0.4 ± 0.1
00032613036	1.6	2013-12-15.06	56 641.06	18.46	>20.1	>20.0	>20.3	<1.2	<0.9
00032613033	1.4	2013-12-15.64	56 641.64	19.04	–	–	20.8 ± 0.3	<0.9	<0.7
00032613034	1.9	2013-12-16.05	56 642.05	19.45	>20.2	>20.1	>20.4	<0.7	<0.5
00032613035	1.7	2013-12-16.52	56 642.52	19.92	20.7 ± 0.3	–	–	<0.5	<0.3
00032613037	7.6	2013-12-17.06	56 643.06	20.46	>19.9	20.4 ± 0.2	>20.3	<0.2	<0.2
00032613038	7.8	2013-12-20.66	56 646.66	24.06	20.4 ± 0.2	>20.6	20.2 ± 0.3	<0.1	<0.1
00032613039	8.0	2013-12-23.26	56 649.26	26.66	20.0 ± 0.4	>19.6	20.5 ± 0.2	<0.2	<0.1
00032613040	3.8	2013-12-23.59	56 649.59	26.99	–	–	–	<0.2	<0.2

Notes. ^(a) Dead-time corrected exposure time; ^(b) start date of the observation; ^(c) time in days after the outburst of nova M31N 2008-12a in the optical on 2013-11-26.60 UT (MJD = 56 622.60; see [Tang et al. 2013](#)); ^(d) *Swift* UVOT filters were *UVM2* (166–268 nm), *UVW1* (181–321 nm), and *UVW2* (112–264 nm); ^(e) X-ray luminosities (unabsorbed, black-body fit, 0.2–1.0 keV) and upper limits were estimated according to Sect. 3.

Table 2. *Swift* observations of the 2012 outburst of nova M31N 2008-12a.

ObsID	Exp ^a [ks]	Date ^b [UT]	MJD ^b [d]	Δt^c [d]	UV ^d [mag]			Rate [10 ⁻² ct s ⁻¹]	$L_{0.2-1.0}^e$ [10 ³⁸ erg s ⁻¹]
					U	<i>uvw1</i>	<i>uvw2</i>		
00032613001	4.0	2012-11-06.45	56 237.45	20.35	–	–	20.2 ± 0.2	<0.3	<0.2
00032613002	4.2	2012-11-16.34	56 247.34	30.24	–	19.8 ± 0.1	–	<0.3	<0.2
00032613003	3.5	2012-12-16.13	56 277.13	60.03	–	–	20.4 ± 0.2	<0.3	<0.2
00032613004	3.7	2013-02-05.36	56 328.36	111.26	19.8 ± 0.2	–	–	<0.4	<0.3

Notes. As for Table 1. The time ^(c) is in days after the outburst of nova M31N 2008-12a in the optical on 2012-10-17.10 UT (MJD = 56 217.10; see Sect. 3).