

Bands of solid CO₂ in the 2–3 μm spectrum of S 140:IRS1[★]

J. V. Keane¹, A. C. A. Boogert², A. G. G. M. Tielens^{1,3}, P. Ehrenfreund⁴, and W. A. Schutte⁴

¹ Kapteyn Astronomical Institute, PO Box 800, 9700 AV Groningen, The Netherlands

² California Institute of Technology, Downs Laboratory of Physics 320-47, Pasadena CA 91125, USA

³ SRON, PO Box 800, 9700 AV Groningen, The Netherlands

⁴ Leiden Observatory, PO Box 9513, 2300 RA Leiden, The Netherlands

Received 14 November 2000 / Accepted 6 July 2001

Abstract. We investigate the 2–3 μm ISO–SWS spectrum of the luminous protostellar object S 140:IRS1. Two narrow absorption features are detected at 2.70 and 2.77 μm, which are well fitted with laboratory spectra of the $\nu_1 + \nu_3$ and the $2\nu_2 + \nu_3$ combination modes of solid CO₂. The ice in this line of sight must have been subjected to significant heating, in agreement with previously studied CO₂ bands. A combined laboratory fit to all CO₂ bands detected toward S 140:IRS1 shows, among others, the need for particle shape calculations for the CO₂ stretch mode. Finally, we discuss the absence of features of isolated H₂O and dangling OH groups in the spectrum of S 140:IRS1.

Key words. ISM: molecules – ISM: abundances – stars: S 140:IRS1 – infrared: stars

1. Introduction

Ice species toward massive protostars were discovered by their characteristic stretching and bending modes at infrared wavelengths (e.g. Willner et al. 1982; Ehrenfreund & Schutte 2000). From band profile analyses, it is now well established that in specific interstellar environments the most abundant species (H₂O, CH₃OH and CO₂) reside on the grain mantle mixed with highly diluted volatiles (CO, CH₄, OCS) (e.g. Tielens et al. 1991; Gibb et al. 2000). A spectral region that received little attention so far is the 2–3 μm region, as wavelengths longer than 2.6 μm are not observable from the ground, a problem remedied by the launch of the Infrared Space Observatory (ISO). Laboratory studies show that this region is awash with a wide variety of combination modes and isolated bands of ices including those of CO₂ and H₂O (Hagen & Tielens 1981; Sandford & Allamandola 1990, 1993; Ehrenfreund et al. 1996).

The abundant presence of CO₂ ice in molecular clouds and star forming regions has been well established (d’Hendecourt & de Muizon 1989; de Graauw et al. 1996; d’Hendecourt et al. 1996; Gürtler et al. 1996; Whittet et al.

1998). Recently, detailed studies of the ¹²CO₂ (Gerakines et al. 1999; Boogert 1999) interstellar absorption bands have revealed that variations in the profile of these bands reflect different degrees of thermal processing of the ice mantles and variation in their chemical composition. The ¹²CO₂ bands are due to strong transitions and hence the detailed profiles and peak positions will depend on the grain shape and shape (Ehrenfreund et al. 1997; Boogert et al. 1999; Baratta et al. 2000). The conclusions based on the ¹²CO₂ bands are confirmed by an analysis of the much weaker ¹³CO₂ band (Boogert et al. 2000), for which grain shape effects are unimportant. The CO₂ combination modes, which absorb weakly in the 2–3 μm spectral region, are also not affected by particle shape effects, and provide a new and independent test of the physical and chemical properties of interstellar CO₂-ices.

Isolated molecules have been studied extensively in the laboratory in the 2–3 μm spectral (Sandford et al. 1990, 1993; Ehrenfreund et al. 1996). Thermal and energetic processes in the interstellar medium destroy sites in the ice matrix where such pockets could reside and lead to conversion of isolated molecules over relatively short time-scales into polymers. Therefore the detection of features due to isolated molecules requires cold and UV shielded astronomical environments.

In this letter, we identify for the first time the CO₂ combination modes along the line of sight toward an embedded object and perform a search for bands of isolated molecules.

Send offprint requests to: J. V. Keane,
e-mail: jacquie@astro.rug.nl

[★] Based on observations with ISO, an ESA project with instruments funded by ESA Member States (especially the PI countries: France, Germany, The Netherlands and the UK) and with the participation of ISAS and NASA.




Fig. 1. ISO-SWS AOT 6 spectrum of S 140:IRS1 **a)**. For comparison the detector response, scaled to the brightness of this source, is shown **b)**.

2. Observations

The protostellar object S 140:IRS1 was observed with ISO-SWS in the high resolution “SWS06” mode (de Graauw et al. 1996). The spectrum was reduced with version 9.5 of the SWS pipeline using the latest calibration files. All detector scans were checked for excessive noise levels, deviating continuum slopes, flux levels and dark current jumps. Clipping all points that deviate by 3σ or more ensured that hits due to cosmic rays were also removed. Bad scans were cut out of the data and the remaining scans were averaged and rebinned to two points per resolution element at the maximum resolving power of band 1b ($R \sim 2500$). Figure 1 shows the final reduced spectrum. Also plotted is the detector response curve to check the reliability of the observed spectrum. No obvious residuals of the detector responsivity are apparent in the interstellar spectrum, that might affect our study of narrow absorption features.

3. Results

The spectrum of S 140:IRS1 shows two narrow absorption features near $2.70 \mu\text{m}$ and $2.78 \mu\text{m}$ and perhaps a broader absorption feature around $2.73 \mu\text{m}$. The independent up and down scans agree well for the $2.70 \mu\text{m}$ band and this feature is detected at the $>6\sigma$ level. The $2.78 \mu\text{m}$ band differs in the individual scans and this feature is only detected at the 3σ level. The $2.73 \mu\text{m}$ feature occurs only at the 2σ level and will require further confirmation. The spectrum was converted to optical depth scale by adopting the polynomial continuum shown in Fig. 1. The peak positions, widths ($FWHM$) and central optical depths are summarized in Table 1.

Table 1. Spectroscopic properties of the absorption features. Errors are shown in the parenthesis.

λ (μm)	Feature	ν cm^{-1}	$FWHM$ cm^{-1}	τ^a
2.70	$\nu_1 + \nu_3$	3707.8 (0.5)	4 (1)	0.076
2.78	$2\nu_2 + \nu_3$	3600.3 (1.15)	5 (1.5)	0.026
2.73		3660.5 (3)	10 (1.5)	0.014

^a Error on τ is 0.007.

4. Identification of CO₂ combination modes

The 2.70 and $2.78 \mu\text{m}$ bands seen in the astronomical observations match the $\nu_1 + \nu_3$ and $2\nu_2 + \nu_3$ combination bands of CO₂, measured in the laboratory. These bands have been studied in detail by Sandford & Allamandola (1990) and Ehrenfreund et al. (1997) in various ice mixtures at different temperatures. The $2\nu_2 + \nu_3$ combination mode has a similar behaviour as the $\nu_1 + \nu_3$ mode at 10 K. Astronomical observations indicate the existence of different types of ices, such as hydrogen-rich (H₂O dominated, polar ices) or hydrogen-poor (apolar ices, as well as thermally processed ices). In pure 10 K CO₂ ice or CO dominant ice mixtures, the peak position of the $\nu_1 + \nu_3$ band is centred at higher frequencies and has a slightly narrower width than the observed interstellar feature. When H₂O is added a large broadening and substantial shift to lower frequencies is observed. A slight narrowing, but an even greater shift to lower frequencies, occurs when CH₃OH is added (Boogert et al. 2000). This difference between polar mixtures (H₂O or CH₃OH dominated) and apolar mixtures (e.g. CO dominated) is similar to that observed by Ehrenfreund et al. (1998) and Boogert et al. (2000) for the fundamental ¹²CO₂ and ¹³CO₂ bands.

After an initial broadening, the widths of both combination bands narrow dramatically with increasing temperature. The CO₂ is then in a segregated, pure CO₂ state (Ehrenfreund et al. 1999; Boogert et al. 2000). A comparison of the interstellar CO₂ combination bands with laboratory CO₂ profiles in different mixtures reveals that both the positions and the widths of the interstellar features can well matched by various mixtures provided that the laboratory ices are heated (Figs. 2a,b,c,d) in order that “segregated boundary phases of pure CO₂” are formed. Thus, the observed interstellar CO₂ combination bands in S 140:IRS1 reflect a pure segregated CO₂ phase. The broader 3660 cm^{-1} ($2.73 \mu\text{m}$) interstellar feature is not matched by any of the CO₂ laboratory mixtures.

5. Alternative candidates

5.1. Isolated water

Isolated H₂O, in low temperature matrices, has been extensively studied in the laboratory and displays a set of bands between 2.5 and $2.9 \mu\text{m}$, with both position and width sensitive to matrix composition and concentration




Fig. 2. A comparison of the S 140:IRS1 CO₂ combination modes with various heated laboratory ice spectra **a–d**). Additionally, isolated H₂O in a CO matrix is compared to the ISO-SWS spectrum **e**).

(Hagen & Tielens 1981; Ehrenfreund et al. 1996). Monomers and dimers show sharp bands (ν_3 and ν_1) between 3800 cm⁻¹ and 3400 cm⁻¹. The position and the width of the interstellar 3700 cm⁻¹ feature can be well matched by isolated monomeric H₂O in a CO dominated matrix (Fig. 2e). The ν_1 mode of monomeric H₂O, on the other hand, is at a higher frequency than the interstellar 3600 cm⁻¹ feature. An assignment of the 2.70 μm band with monomeric H₂O requires a significant amount of CO (CO/H₂O > 100) but the amount of CO detected along the line sight toward S 140:IRS1 is very small (<1%). Hence, we conclude that the observed interstellar 2.70 μm band is not due to isolated H₂O.

5.2. Dangling OH-groups

Water molecules link to each other in clusters through hydrogen bonds and the last OH group is referred to as a dangling bond. The relative intensity and position of the band depends strongly on the matrix composition (Hagen & Tielens 1981). CO₂ will act as a weak base resulting in a broad, weak OH-stretching band lying between 3700 cm⁻¹ and the main OH-stretching band of completely bonded H₂O molecules (Hagen et al. 1983). The terminal OH-band may account for the interstellar “3660 cm⁻¹” feature (if it is real).




Fig. 3. Simultaneous fits to all the detected solid CO₂ bands toward S 140:IRS1 by different laboratory mixtures. The dot-dashed line denotes a fit based on the observed abundances, the dashed line represents the polar + annealed fit of Gerakines et al. (1999), the solid line denotes the annealed mixture of Boogert et al. (2000), and the dotted line represents the particle shape corrected spectrum of the annealed mixture (see Sect. 6).

6. Discussion

Extensive analyses of the CO₂ bands toward different lines of sight (Gerakines et al. 1999) yield inconclusive results about the exact composition of ice mantles. In the case of S 140:IRS1, the line of sight is dominated by H₂O and the only other major species present is CO₂ which is 20% relative to H₂O. CH₃OH is a very minor constituent of the ice mantle toward S 140:IRS1 (<3% of H₂O; Allamandola et al. 1992). Furthermore, the lack of volatile species, such as CO, indicate that this line of sight is warm. Fitting the interstellar CO₂ bands by a heated laboratory mixture constrained by the observed H₂O, CO₂, and CH₃OH abundances (Fig. 3; dot-dashed line) reveals that the combination modes are the only bands that are reasonably matched. The spectral signatures of the other CO₂ bands are poorly matched by this mixture and in particular the stretching mode is significantly narrower than the interstellar band. Gerakines et al. (1999) have shown that the CO₂ stretching and bending modes can be well matched by a combination of the heated mixture of H₂O:CO₂:CH₃OH = 1:1:1 and an H₂O:CO₂ = 100:14 polar mixture at 10 K (Fig. 3; dashed line). The CO₂ combination bands are well fitted by this model since the heated component produces a segregated boundary pure CO₂ phase required to match these bands (Sect. 4). However, due to the large contribution by the apolar mixture (50%), a broad feature appears around 3650 cm⁻¹ that is not apparent in the interstellar spectrum. Alternatively, Boogert et al. (2000) have shown that the heated H₂O:CO₂:CH₃OH = 1:1:1 mixture alone provides an excellent fit to the CO₂ bending mode (Fig. 3;

solid line). The combination modes and the ¹³CO₂ band are also well matched. However, the profile of the CO₂ stretching mode is not as successfully reproduced. One possible reason may be that the absorption profile is very sensitive to the shape and size of the grains. Ehrenfreund et al. (1997) showed that particle shape effects are important for strong transitions and that substantial profile changes occur. Assuming a distribution of ellipsoidally shaped particles and adopting the method of grain shape corrections applied by Ehrenfreund et al. (1997), we find that the blue side of the CO₂ stretching mode is now better fitted (Fig. 3; dotted line). However, there is still considerable uncertainty regarding the optical constants (Barrata et al. 2000) and which type of particles should be used. Due to the inherent weakness of the combination bands and the ¹³CO₂ band, they are insensitive to grain shape and hence there are no variations in their spectral profiles.

Two crucial results emerge from fitting the various mixtures to all the solid CO₂ bands observed toward S 140:IRS1. Firstly, the combination modes are only fitted if the various mixtures are heated so that a segregated boundary phase of pure CO₂ is formed. Hence, the CO₂ combination bands are not useful in constraining the ice mantle composition. Secondly, a mixture based on the observed interstellar abundances fails to match the observed solid CO₂ bands. The best fitting mixtures of Gerakines et al. (1999) and Boogert et al. (2000) reveal that the CO₂ bands are only successfully matched by laboratory mixtures that contain significant amounts of CH₃OH which contradict the observations that show CH₃OH is a minor constituent of the ice toward S 140:IRS1. The current laboratory database does not contain a sufficiently extensive range of CH₃OH mixtures and therefore the effects of varying CH₃OH intensities in H₂O:CO₂ mixtures are not readily assessable. Thus, further laboratory studies of CH₃OH in H₂O:CO₂ matrices are required.

The CO₂ column density can be determined from the central optical depth and *FWHM*, using the band strength of pure solid CO₂ determined by Gerakines et al. (1995). We find that the column densities derived, from both combination bands, agree with the CO₂ stretching and bending mode column densities ($4.2 \times 10^{17} \text{ cm}^{-2}$; Gerakines et al. 1999). This agreement also shows that over this wide wavelength range (2.7–16 μm) the same ice column is probed. Hence, in contrast to previous discussions – which centred on the methanol column density derived from the 3.5 μm and 6.8 μm absorption features (Allamandola et al. 1992) –, radiation transfer effects do not mar the column density derivations, at least for S 140:IRS1.

7. Conclusions

We have presented ISO-SWS observations of the 2–3 μm region toward the embedded object S 140:IRS1. The spectrum shows for the first time two narrow absorption features near 2.7 and 2.77 μm attributed to the combination modes ($\nu_1 + \nu_3$ and $2\nu_2 + \nu_3$) of solid CO₂ ice.

The derived column densities agree with previous estimates derived from the CO₂ stretching and bending modes. The main results presented here show that simultaneous fits to the solid CO₂ stretching, bending, combination and ¹³CO₂ bands are achieved using an H₂O:CO₂:CH₃OH = 1:1:1 laboratory mixture. This illustrates that the relative strengths and profiles of the combination bands are in good agreement with the other CO₂ bands. Applying particle shape corrections significantly improved the fit to the peak of the fundamental CO₂ stretching mode. Furthermore, isolated water can be excluded as a cause of these bands, however multimers near 2.7 μm may be present.

Acknowledgements. This research has made use of the Leiden Observatory Laboratory database of spectra (www.strw.LeidenUniv.nl/~lab/ISO-WWW3).

References

- Allamandola, L. J., Sandford, S. A., Tielens, A. G. G. M., & Herbst, T. M. 1992, *ApJ*, 399, 134
- Baratta, G. A., Palumbo, M. E., & Strazzulla, G. 2000, *A&A*, in press
- Boogert, A. C. A. 1999, Ph.D. Thesis, University of Groningen
- Boogert, A. C. A., Ehrenfreund, P., Gerakines, P. A., et al. 2000, *A&A*, 353, 349
- Chiar, J. E., Gerakines, P. A., Whittet, D. C. B., et al. 1998, *ApJ*, 498, 716
- de Graauw, Th., Haser, L. N., Beintema, D. A., et al. 1996, *A&A*, 315, L49
- d'Hendecourt, L. B., & Jourdain de Muizon, M. 1989, *A&A*, 223, L5
- d'Hendecourt, L. B., Jourdain de Muizon, M., Dartois, E., et al. 1996, *A&A*, 315, L341
- Ehrenfreund, P., Gerakines, P. A., Schutte, W. A., van Hemert, M. C., & van Dishoeck, E. F. 1996, *A&A*, 312, 263
- Ehrenfreund, P., Boogert, A. C. A., Gerakines, P. A., Tielens, A. G. G. M., & van Dishoeck, E. F. 1997, *A&A*, 328, 649
- Ehrenfreund, P., Dartois, E., Demyk, K., & d'Hendecourt, L. 1998, *A&A*, 339, L17
- Ehrenfreund, P., Kerkhof, O., Schutte, W. A., et al. 1999, *A&A*, 350, 240
- Ehrenfreund, P., & Schutte, W. A. 2000, in *Astrochemistry: From molecular clouds to planetary systems*, IAU Symp., 197, 135
- Gerakines, P. A., Schutte, W. A., Greenberg, J. M., & van Dishoeck, E. F. 1995, *A&A*, 296, 810
- Gerakines, P. A., Whittet, D. C. B., Ehrenfreund, P., et al. 1999, *ApJ*, 522, 357
- Gibb, E. L., Whittet, D. C. B., Schutte, W. A., et al. 2000, *ApJ*, 536, 347
- Gürtler, J., Henning, Th., Kömpe, C., et al. 1996, *A&A*, 315, L189
- Hagen, W., & Tielens, A. G. G. M. 1981, *J. Chem. Phys.*, 75, 4198
- Hagen, W., Tielens, A. G. G. M., & Greenberg, J. M. 1983, *A&AS*, 51, 389
- Sandford, S. A., & Allamandola, L. J. 1990, *ApJ*, 355, 357
- Sandford, S. A., & Allamandola, L. J. 1993, *ApJ*, 417, 815
- Tielens, A. G. G. M., Tokunaga, A. T., Geballe, T. R., & Bass, F. 1991, *ApJ*, 381, 181
- Willner, S. P., Gilletet, F. C., Herter, T. L., et al. 1982, *ApJ*, 253, 174