

## The radio counter-jet of the QSO 3C 48

W. X. Feng<sup>1,2</sup>, T. An<sup>1,3</sup>, X. Y. Hong<sup>1,3</sup>, Jun-Hui Zhao<sup>4</sup>, T. Venturi<sup>5</sup>,  
Z.-Q. Shen<sup>1</sup>, and W. H. Wang<sup>1,3</sup>

<sup>1</sup> Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai 200030, PR China  
e-mail: wfeng@shao.ac.cn

<sup>2</sup> Graduate School of Chinese Academy of Sciences, Beijing 100039, PR China

<sup>3</sup> National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, PR China

<sup>4</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, MS 78, Cambridge, MA 02138, USA

<sup>5</sup> Istituto di Radioastronomia del CNR, via Gobetti 101, 4129 Bologna, Italy

Received 5 November 2004 / Accepted 9 December 2004

**Abstract.** We present multi-frequency radio observational results of the quasar 3C 48. The observations were carried out with the Very Large Array (VLA) at five frequencies, 0.33, 1.5, 4.8, 8.4, and 22.5 GHz, and with the Multi-Element Radio Linked Interferometer Network (MERLIN) at the two frequencies of 1.6 and 5 GHz. The source shows a one-sided jet to the north within 1'', which then extends to the northeast and becomes diffuse. Two bright components (N2 and N3), containing most of the flux density, are present in the northern jet. The spectral index of the two components is  $\alpha_{N2} \sim -0.99 \pm 0.12$  and  $\alpha_{N3} \sim -0.84 \pm 0.23$  ( $S \propto \nu^\alpha$ ). Our images show the presence of an extended structure surrounding component N2, suggestive of strong interaction between the jet and the interstellar medium (ISM) of the host galaxy. A steep-spectrum component, labelled S, located 0.25'' southwest to the flat-spectrum component which could be the core of 3C 48, is detected at a significance of  $>15\sigma$ . Both the location and the steepness of the spectrum of component S suggest the presence of a counter-jet in 3C 48.

**Key words.** galaxies: quasars: individual: 3C 48 – galaxies: jets – galaxies: interactions

### 1. Introduction

The radio source 3C 48 (B0134+329) is a 16th mag quasar, with redshift  $z = 0.368$  (Spinrad et al. 1985). It was the first QSO identified at optical wavelengths (Smith et al. 1961), and the second identified one (after 3C 273) at radio wavelengths (Schmidt 1963). The source shows unexpectedly strong emission in the far IR bands, suggesting that it is located in a gas-rich environment (Chatzichristou et al. 1999). The variability measured with ROSAT suggests that at least part of the X-ray emission (principally the softest) arises from the region of the accretion disc or from a beamed blazar component (Worrall et al. 2004). The image obtained with the University of Hawaii's 2.2 m optical telescope at Mauna Kea Observatory is dominated by a previously unsuspected high surface brightness region, referred to as "3C 48A", centered  $\sim 1''$  ( $1'' \sim 4.1$  kpc projected linear size<sup>1</sup>) to the north of the QSO nucleus (Stockton & Ridgway 1991; Hook et al. 1994). This region was identified as the potential second nucleus of a galaxy in the process of merging with the host galaxy of 3C 48 (Zuther et al. 2003). There is evidence that the nuclear activity in 3C 48 is triggered by the merger process.

3C 48 is classified as a Compact Steep Spectrum (CSS) source based on its steep radio spectrum and its small angular size (Fanti et al. 1985). The spectrum is reasonably straight from  $\sim 178$  MHz up to 2.7 GHz, with an integrated radio index  $\alpha \sim -0.83$  (Peacock et al. 1982). It shows a low-frequency peak at about 100 MHz (<http://nedwww.ipac.caltech.edu>). The quasar 3C 48 does not show significant variations with time, and has a simple spectrum at MHz/GHz wavelengths (Baars et al. 1977), and it is usually used as the primary flux density calibrator in interferometer observations owing to its compactness and brightness. While Ott et al. (1994) found that the source could have become brighter from the 1960s to the 1990s, the increase in flux density is enhanced at shorter wavelengths, as is consistent with the activity in the core (Wilkinson et al. 1990).

Most of the radio emission occurs on a scale smaller than about  $0.6''$  and additional emission extends to  $\sim 6''$  northwest to the nucleus, with a clumpy morphology (Hartas et al. 1983; Woan 1992). At sub-arcsecond resolution, the source exhibits an elongated structure in the north–south direction. VLBI observations showed a disrupted radio morphology (Wilkinson et al. 1991). The active nucleus, which is identified from its inverted radio spectrum (Simon et al. 1990), is weak at radio frequencies and is buried deeply within the host galaxy. A one-sided jet stretches out of the core towards the north  $\sim 1''$  within

<sup>1</sup> Hubble constant  $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ , deceleration parameter  $q_0 = 0.5$  and zero cosmological constant.

**Table 1.** Observational information and image parameters.

No.	Epoch	Array	Freq. (GHz)	$BW^a$ (MHz)	$\tau^b$ (min)	$S_{\text{peak}}$ (Jy/b)	rms (mJy/b)	Maj×Min, PA (arcsec, °)	Figure
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)
1	1992.45	MERLIN	5.0	13	850	2.61	0.3	$0.04 \times 0.04$ , 34.4	Fig. 1d
2	1993.64	MERLIN	1.6	15	900	4.65	1.3	$0.17 \times 0.14$ , 33.2	Fig. 1b
3	1999.08	VLA-C	8.4	50	2	3.17	0.6	$2.37 \times 2.26$ , 7.9	
4			22.5	50	7	1.02	0.5	$1.07 \times 1.03$ , 10.6	
5	2000.92	VLA-A	8.4	50	2	1.66	0.3	$0.32 \times 0.20$ , -73	
6			22.5	50	5.5	0.34	0.4	$0.12 \times 0.07$ , -72	Fig. 1c
7	2003.46	VLA-A	0.33	0.003	4.5	43.0	15	$11.3 \times 6.2$ , -77.0	
8			1.5	0.012	4	15.3	1.8	$2.1 \times 1.3$ , -81.8	
9			4.8	50	3	3.69	1.0	$0.46 \times 0.33$ , -63.3	
10			8.4	50	4	1.60	0.3	$0.27 \times 0.21$ , -82.5	Fig. 1a

<sup>a</sup> The VLA observations at 0.33 and 1.5 GHz were carried out in spectral line mode, and recorded with narrow bandwidth.

<sup>b</sup> On-source integration time.

the nucleus of the host galaxy. The northern jet is highly dispersed at  $0.05''$ – $0.5''$ , and its diffused morphology has been interpreted as the result of the jet outflow interaction with the dense ISM in the host galaxy (Wilkinson et al. 1991). There is also morphological evidence for a recent merger event, such as a possible double nucleus and a tidal tail extending several arcseconds to the northwest (Chatzichristou et al. 1999; Canalizo et al. 2000; Zuther et al. 2003).

In this paper, we present results of the multi-frequency VLA and MERLIN observations of 3C 48. The observations and data reduction are described in the next section. The analysis of the images is given in Sect. 3. Section 4 is a discussion. A summary is given in the last section.

## 2. Observation and data reduction

Quasar 3C 48 was observed as a flux density calibrator with the VLA on Jan. 29, 1999 (observational code AH635) at 8.4 and 22.5 GHz, on Dec. 1, 2000 (AH721) at 8.4 and 22.5 GHz, as well as on June 17, 2003 (AZ143) at 0.33, 1.5, 4.8 and 8.4 GHz. The observations were carried out in snapshot mode. 3C 48 was observed at 4.8, 8.4 and 22.5 GHz with the VLA in the C configuration with a bandwidth of 50 MHz in two IFs, while the observations at 0.33 and 1.5 GHz were done in the spectral line mode in order to reject radio frequency interference (RFI). Two epochs of archival MERLIN data at 1.6 and 5.0 GHz were also used in this paper. The observational information of the VLA and the MERLIN is listed in Table 1 in a time series. Column 1 is the serial number. Columns 2 and 3 are the epoch and the array configuration. The observing frequency, bandwidth, and on-source time are listed in Cols. 4–6, respectively.


The raw VLA data were calibrated and corrected for the phase and amplitude errors using the Astronomical Image Processing System (AIPS) of the National Radio Astronomy Observatory (NRAO). The MERLIN data were calibrated with the suite of D-programs (Thomasson 1986).

The further process in the reduction of the VLA and MERLIN data, including editing, self-calibration and image analysis, was performed using both the AIPS and the DIFMAP packages (Shepherd et al. 1994). In order to do analysis of the jet structure, we fitted Gaussian models to the self-calibrated  $uv$  data sets. The task MODELFIT in the DIFMAP package was used for all data sets except for the 5 GHz MERLIN, where we used the task JMFIT in AIPS to take out the extended emission prior to the model fitting. We found that 80–90% of the total flux density of the source is dominated by a few compact components. Elliptical Gaussian models were preferentially used. If elliptical Gaussian models failed in the least-squares fitting, then a circular Gaussian was used. The errors in the model fitting parameters are estimated based on the method described by Fomalont (1999). The uncertainties of the flux density, originating from the calibration process, are estimated within a range between 5% (for VLA) and 10% (for MERLIN). The uncertainty of the component position is proportional to the FWHM of the synthesized beam and is inversely proportional to the ratio of peak signal to noise (Fomalont 1999).

## 3. Analysis

### 3.1. Images

We imaged the source using the data sets listed in Table 1. The results show that the source is resolved when the resolution is higher than  $0.5''$ . Figure 1 shows four images at 1.6, 5, 8.4 and 22.5 GHz with sub-arcsec resolutions. The detailed structure of the source is shown in the images from Figs. 1a to 1d in order of increasing resolution. These images have been made using uniform weighting. Since most of the observations lack nearby phase calibrators, the bright, compact component N2 was used as the phase reference in the self-calibration and imaging process. The parameters of each image are summarized in Table 1. Columns 7 and 8 give the peak flux density and the off-source rms noise in each of the images; Col. 9 gives



**Fig. 1.** The multi-frequency total intensity images of 3C 48. The peak flux density, beam size, off source rms level are referred to in Cols. 7–9 in Table 1. Contour levels for a), b), and c) are  $1.1, 5.0,$  and  $1.2 \text{ mJy}/b \times (-1, 1, 2, \dots, 1024)$ , respectively, and those for d) are  $1.1 \text{ mJy}/b \times (-1, 1, 4, 16, 32, 64, 128, 256, 512)$ . The lowest contour levels are generally 3 times the rms levels.

the size of the synthesized beam; Col. 10 indicates the label of the images that are present in Fig. 1. Each of the major components in the images is fitted in least squares to a Gaussian model. The results of the model-fit to the calibrated data are listed in Table 2. Column 1 is the serial number of the observations, which is the same as that in Table 1; Col. 2 is the observing epoch; Col. 3 is the observing frequency; Col. 4 is the component label; Col. 5 is the integral flux density of the Gaussian component; Cols. 6 and 7 are the peak position of the Gaussian component in  $RA_{B1950}$  and  $Dec_{B1950}$  coordinates; Cols. 8–10 are the sizes of major and minor axes and the position angle. Previous observations show that the source is dominated by steep spectrum components and that the radiation from the core is rather weak and difficult to identify at radio frequencies. We note that the phase center in each image is closer to the brightest jet component N2 ( $RA_{B1950}: 01^{\text{h}}34^{\text{m}}49.826^{\text{s}}$ ,  $Dec_{B1950}: +32^{\circ}54'20.260''$ ) than to the radio core. The core component and its location will be discussed in Sect. 4.1.

Figure 1a is the VLA image at 8.4 GHz observed at the epoch of 2003. The image shows an elongated structure in

the NE–SW direction, which includes 4 components (N1, N2, N3, and S). The main jet is dominated by the components N1, N2, and N3. Component S has been detected for the first time at a significance level above  $15\sigma$ , at  $RA \sim 01^{\text{h}}34^{\text{m}}49.810^{\text{s}}$  and  $Dec \sim 32^{\circ}54'19.626''$ . An independent VLA image at 8.4 GHz, made from the observation at epoch 2000, confirms the detection. The epoch 2000 image at 8.4 GHz is not shown, while the parameters derived from model fitting are listed in Table 2.

The two major components (N2 and N3), the northern extended component (N1), and the weak southern component (S) have also been detected in the 1.6 GHz MERLIN image (Fig. 1b). A Gaussian fit failed to fit the diffuse component N1, visible in the MERLIN 1.6 GHz image. Component S is confirmed at  $12\sigma$ , and can be fitted with a Gaussian of  $0.28'' \times 0.23''$  ( $PA = 37.6^{\circ}$ ) with peak brightness of 26 mJy per beam.

Both components N1 and S appear to be completely resolved in the high resolution images at 22.5 GHz (Fig. 1c) observed with the VLA and at 5 GHz observed with MERLIN (Fig. 1d). At the high resolutions, the components N2

**Table 2.** Parameters of Gaussian components.

No.	Freq. (GHz)	Comp.	$S_{\text{int}}$ (Jy)	RA ( <sup>h</sup> , <sup>m</sup> , <sup>s</sup> )	Dec ( <sup>°</sup> , <sup>'</sup> , <sup>''</sup> )	$a$ (mas)	$b$ (mas)	$\Phi$ ( <sup>°</sup> )	Figure
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	
1	4.8	N2a	1.549	01 3449.826 ± 0.005	32 5420.264 ± 0.004	86.0	53.6	168.1	Fig. 1d
		N2b	0.799	49.826 ± 0.004	20.175 ± 0.003	65.0	57.3	71.8	
		N3a	1.423	49.821 ± 0.006	20.025 ± 0.003	99.7	47.4	11.0	
		N3b	0.164	49.821 ± 0.003	19.936 ± 0.002	57.6	43.5	1.1	
2	1.6	N1	3.29	49840 ± 0.034	20.296 ± 0.036	512.0	342.5	4.9	Fig. 1b
		N2	10.33	49.827 ± 0.030	20.264 ± 0.017	255.5	224.71	179.8	
		N3	3.13	49.821 ± 0.006	20.075 ± 0.006	264.0	205.3		
		S	0.039	49.812 ± 0.026	19.574 ± 0.014	187.2	99.4	172.0	
3	8.4	N	3.20	49.826 ± 0.036	20.264 ± 0.010	317.2	105.4	25.4	
4	22.5	N	1.06	49.827 ± 0.055	20.260 ± 0.021	262.0	84.2	18.9	
5	8.4	N1	0.211	49.843 ± 0.030	20.512 ± 0.030	330.0	330.0		
		N2	1.997	49.827 ± 0.012	20.281 ± 0.012	131.3	131.3		
		N3	1.025	49.821 ± 0.011	20.075 ± 0.011	118.3	118.3		
		S	0.013	49.807 ± 0.033	19.619 ± 0.026	356.0	280.0	112.9	
6	22.5	N2a	0.55	49.827 ± 0.038	20.263 ± 0.038	64.1	64.1		Fig. 1c
		N2b	0.35	49.825 ± 0.045	20.172 ± 0.045	75.7	75.7		
		N3a	0.30	49.821 ± 0.033	20.029 ± 0.033	56.5	56.5		
		N3b	0.12	49.821 ± 0.028	19.940 ± 0.028	48.4	48.4		
7	0.33	N	43.14	49.827 ± 0.081	20.257 ± 0.058	466.4	332.7	-40.9	
8	1.5	N	15.88	49.826 ± 0.022	20.260 ± 0.006	373.5	108.0	28.4	
9	4.8	N	5.35	49.826 ± 0.014	20.254 ± 0.020	322.7	146.1	24.6	
		S	0.015	49.814 ± 0.018	19420 ± 0.016	705.5	341.1	80.6	
10	8.4	N1	0.250	49.839 ± 0.031	20.453 ± 0.031	330.0	330.0		Fig. 1a
		N2	1.848	49.827 ± 0.011	20.280 ± 0.011	118.3	118.3		
		N3	1.011	49.821 ± 0.010	20.079 ± 0.010	107.5	107.5		
		S	0.016	49.810 ± 0.027	19.633 ± 0.027	292.0	282.0	72.7	

and N3 are further resolved into subcomponents, labelled N2a and N2b, N3a and N3b, respectively.


### 3.2. Spectral index

The multi-frequency observations allow us to study the spectral index distribution of 3C 48 at sub-arcsecond resolutions. Figure 2a shows the image of the spectral index distribution (grey scale) derived from the VLA images at 8.4 and 22.5 GHz at epoch 2000.92, with the contours of the radiation intensity at 8.4 GHz superimposed. The uv-data at 22.5 GHz have been tapered to match the synthesized beam of the data at 8.4 GHz. Both images at 8.4 and 22.5 GHz are convolved to the same resolution and are aligned with respect to the common phase center. With the same method, we obtained the spectral index image (Fig. 2b) with the MERLIN data sets. Figure 2b shows the spectral index image (grey scale) between 1.6 and 5 GHz, with the MERLIN 5 GHz contours overplotted. The dark end

of the grey scale corresponds to a flat spectrum. Most of the source shows a steep spectrum. The spectrum is steepest in the northern jet (component N2), while it becomes flatter to the south.

In order to image the distribution of the spectral index across the source, we plotted the spectral index along the jet axis at position angle 8.5°. The slice spectral indices are shown in Figs. 3a and 3b, corresponding to Figs. 2a and 2b, respectively. There is a component centered at RA  $\sim 01^{\text{h}}34^{\text{m}}49.820^{\text{s}} \pm 0.020^{\text{s}}$ , Dec  $\sim 32^{\circ}54'19.740'' \pm 0.013''$ , marked with a cross in Figs. 2a and b, showing a rather flat spectrum.

The spectral indices of components N2, N3 and S are estimated based on the model fitting results. Figure 4 shows the power-law fitting in logarithmic coordinates. The spectrum of component N2 ( $\alpha_{\text{N2}} \sim -0.99 \pm 0.12$ ) is slightly steeper than that of component N3 ( $\alpha_{\text{N3}} \sim -0.84 \pm 0.23$ ); and component S also shows a steep spectrum with  $\alpha_{\text{S}} \sim -0.70 \pm 0.14$  between 1.6 and 8.4 GHz. The results are consistent with those shown in



**Fig. 2.** **a)** Simultaneous spectral index distribution (grey) between 8.4 and 22.5 GHz at epoch 2000.92, superimposed on the 22.5 GHz VLA image. **b)** Spectral index distribution (grey) between 1.6 and 5 GHz, superimposed on the 5 GHz MERLIN image at the epoch 1992.45; Restored beams of the contour images are  $0.20'' \times 0.20''$  and  $0.23'' \times 0.19''$ , PA  $45^\circ$ ; The contour levels are  $1.3 \text{ mJy}/b \times (-1, 1, 4, 64, 256)$  and  $3.5 \text{ mJy}/b \times (-1, 1, 4, 64, 256)$ , respectively. The scale of the spectral index is shown in each panel. The darkest region corresponds to the flattest spectrum.

Figs. 2 and 3. A mean spectral index  $\alpha \sim -0.87 \pm 0.10$  for the overall source (3C 48) is estimated from the total flux densities at the frequencies between 0.33 and 22.5 GHz (Fig. 4).

## 4. Discussion

### 4.1. The position of the nucleus

In Sect. 3.2 we showed that the flat-spectrum component centered at RA  $\sim 01^{\text{h}}34^{\text{m}}49.820^{\text{s}} \pm 0.020^{\text{s}}$ , Dec  $\sim 32^\circ 54' 19.740'' \pm 0.013''$ , is most likely the nuclear core in 3C 48. The flat-spectrum core is likely to be buried in the strong emission at the centimeter and longer wavelengths. A detection of the active core was also claimed by Simon et al. (1990). The core was identified at the southern end of the source in their 22.5 GHz image. Wilkinson et al. (1991) suggested that the radio core detected by Simon et al. (1990) is associated with the southernmost component in their high resolution image. Based on the analysis of our data, we find that the position of the flat-spectrum core in our image is  $0.15''$  south of that identified by Simon et al. (1990) and Wilkinson et al. (1991). Further higher resolution and higher dynamics range images at millimeter wavelengths are needed to unambiguously identify the nucleus in 3C 48.


### 4.2. A counter-jet in 3C 48

We have distinctly detected an extended component  $S \sim 0.25''$  southwest of the flat-spectrum core, which is the putative nucleus of 3C 48. Component  $S$  shows a steep spectrum, i.e.  $\alpha_s \sim -0.70 \pm 0.14$ . Because of its location and the steep spectrum nature, we suggest that  $S$  is a counter-jet.

Alternatively, a multi-particle model has been successfully developed and has been applied to the host galactic nucleus of 3C 48 (Scharwächter et al. 2004), suggesting a counter tidal tail extending to the southwest in front of the main body of the 3C 48 host. The detection of component  $S$  could be evidence supporting the counter tidal tail in this quasar.

### 4.3. Interaction with ISM

CSS sources usually have sub-kpc sizes with complex radio morphologies. The nature of CSS sources is not completely understood yet and two scenarios have been suggested. In the *youth scenario*, CSS sources are small and could reflect an early stage in the evolution of radio sources (Phillips & Mutel 1982; Fanti et al. 1995). According to the *frustration scenario*, the unusual conditions in the interstellar medium of their host galaxies, such as a higher density and/or the presence of turbulence, inhibit the radio source from growing to larger sizes (van Breugel et al. 1984). The youth scenario is very much preferred in statistical research (Saikia et al. 1995, 2001; Murgia et al. 1999, 2001). Despite this statistical interpretation, the frustration scenario may still apply to individual sources which can be interpreted in terms of confinement and interaction between the jets and ISM in the host galaxies. Numerical simulations have also shown that a dense ISM may substantially confine CSS sources (De Young 1993). The radio structure of 3C 48 shows evidence for strong interaction between the radio jet and the ISM in the host galaxy. Our scenario is as follows: a supersonic collimated jet stretches out from the core to the north where a shock is produced in the interaction between jet flow and the dense medium. The interaction converts the




**Fig. 3.** Slice plot of the spectral index at PA  $8.5^\circ$  corresponding to Figs. 2a and 2b, respectively. The slice center for **a**) is RA  $01^{\text{h}}34^{\text{m}}49.827^{\text{s}}$ , Dec  $+32^\circ54'20.250''$  and for **b**) RA  $01^{\text{h}}34^{\text{m}}49.827^{\text{s}}$ , Dec  $+32^\circ54'20.268''$ , respectively.

kinetic energy carried by the jet to internal energy or turbulence which could enhance the acceleration of particles and then the radiation field, causing the jet to brighten. The observed brightening of the northern jet (hot spot N2 in Figs. 1c and 1d) after losing collimation appears to be consistent with this picture. Based on the study of the kinematics and physical conditions of the extended emission line gas and their relations to the nuclear star formation, Chatzichristou (2001) found that the coupling between radio jet and gas is the kinematic signature of a recent interaction. Both Canalizo & Stockton (2000) and Chatzichristou et al. (1999) suggested that the unusually high stellar velocity dispersion as well as very young stellar population are possibly related to the interaction of the nuclear radio jets with the ambient media. The NIR (near infrared) images and spectra also suggest that reddening by several magnitudes in the nuclear emission can be explained by the interactions of the 3C 48 radio jet with the circum-nuclear medium (Zuther et al. 2003). The ratio of the recombination lines  $P_{\alpha\beta}$  and  $P_{\alpha\gamma}$  determined from 3C 48 (Zuther et al. 2003) is about  $1.6 (\pm 0.5)$ , which is consistent with the value typical for the ISM of active galactic nuclei being excited by the central-engine continuum emission.

## 5. Summary

In this paper, we carried out radio multi-frequency observations of 3C 48. The high dynamic range images exhibit the detailed structure of the source on the sub-arcsec scale. The position of the nucleus with a flat spectrum was determined based on the spectral index distribution images. The jet components show steep spectra. We confirm the existence of a counter-jet in 3C 48, which shows evidence for the first time that 3C 48 actually has a double-sided jet. The northern jet appears to be disrupted at  $0.5''$  northeast of the core. Our analysis suggests



**Fig. 4.** Total spectrum of 3C 48 (solid line) and of the various components in the source.

that the initially collimated jet can be disrupted by the interaction between jet flow and the circum-nuclear medium.

*Acknowledgements.* This research is supported by the National Science Foundation of PR China (10328306, 10333020, and 10473018). The authors are grateful for the technique support by the staff of the VLA and the MERLIN. The VLA is a facility of the National Radio Astronomy Observatory, which is operated by Associated Universities Inc. under cooperative agreement with the National Science Foundation. MERLIN is a National Facility operated

by the University of Manchester at Jodrell Bank Observatory on behalf of PPARC. W. X. Feng thanks the MERLIN staff (Peter Thomasson) for providing the archive data. It is a great pleasure to thank the referee Paul Wiita who suggested some improvements, and also thanks Prof. D. R. Jiang for his valuable advice in this paper. This research has made use of the NASA/IPAC Extragalactic Database (NED).

## References

- Baars, J. W. M., Genzel, R., Pauliny-Toth, I. I. K., & Witzel, A. 1977, *A&A*, 61, 99
- van Breugel, W. J. M., Miley, G. K., & Heckman, T. A. 1984, *AJ*, 89, 5
- Canalizo, G., & Stockton, A. 2000, *ApJ*, 528, 201
- Chatzichristou, E. T., Vanderriest, C., & Jaffe, W. 1999, *A&A*, 343, 407
- Chatzichristou, E. T. 2001, *IAUS*, 205, 162
- De Young, D. 1993, *ApJ*, 402, 95
- Fanti, C., Fanti, R., Parma, P., et al. 1985, *A&A*, 143, 295
- Fanti, C., Fanti, R., Dallacasa, D., et al. 1995, *A&A*, 302, 317
- Fomalont, E. B. 1999, in *Synthesis Imaging in Radio Astronomy II*, ed. G. B. Taylor, C. L. Carilli, & R. A. Perley, 301
- Hartas, J. S., Ree, W. G., Scott, P. F., et al. 1983, *MNRAS*, 205, 625
- Hook, R., Lucy, L., Stockton, A., & Ridgway, S. 1994, *ST-ECF Newsletter*, 21, 16
- Murgia, M., Fanti, C., Fanti, R., et al. 1999, *A&A*, 345, 769
- Murgia, M., Fanti, C., Fanti, R., et al. 2001, *NewAR.*, 46, 307
- Ott, M., Witzel, A., Quirrenbach, A., et al. 1994, *A&A*, 284, 331
- Peacock, J. A., & Wall, J. V. 1982, *MNRAS*, 198, 843
- Phillips, T. J., & Mutel, R. L. 1982, *A&A*, 106, 21
- Saikia, D. J., Jeyakumar, S., Wiita, P. J., et al. 1995, *MNRAS*, 276, 1215
- Saikia, D. J., Jeyakumar, S., Salter, C. J., et al. 2001, *MNRAS*, 321, 37
- Scharwächter, J., Eckart, A., Pfalzner, S., et al. 2004, *A&A*, 414, 497
- Schmidt, M. 1963, *Nature*, 197, 1040
- Shepherd, M. C., Pearson, T. J., & Taylor, G. B. 1994, *BAAS*, 26, 987
- Smith, H. J., & Hoffleit, D. 1961, *PASP*, 73, 292S
- Simon, R. S., Readhead, A. C. S., Moffet, A. T., et al. 1990, *ApJ*, 354, 140
- Spinrad, H., Djorgovski, S., Marr, J., et al. 1985, *PASP*, 97, 932
- Stockton, A., & Ridgway, S. E. 1991, *AJ*, 102, 2
- Thomasson, P. 1986, *QJRAS*, 27, 413
- Wilkinson, P. N., Tzioumis, A. K., Akujor, C. E., et al. 1990, in *Parsec-Scale Radio Jets*, ed. J. A. Zensus, & T. J. Pearson (Cambridge University Press), 152
- Wilkinson, P. N., Tzioumis, A. K., Benson, J. M., et al. 1991, *Nature*, 352, 313
- Woan, G. 1992, *A&A*, 352, 313
- Worrall, D. M., Hardcastle, M. J., Pearson, T. J., et al. 2004, *MNRAS*, 347, 632
- Zuther, J., Eckart, A., Scharwächter, J., et al. 2003, *A&A*, 414, 919