

Time lag between prompt optical emission and γ -rays in GRBs (Research Note)

S. M. Tang¹ and S. N. Zhang^{1,2,3,4}

¹ Department of Physics and Center for Astrophysics, Tsinghua University, Beijing 100084, PR China
e-mail: tangsm99@mails.tsinghua.edu.cn

² Department of Physics, University of Alabama in Huntsville, Optics Building 201C, Huntsville, AL 35899, USA

³ Space Science Laboratory, NASA Marshall Space Flight Center, SD50, Huntsville, AL 35812, USA

⁴ Institute of High Energy Physics, Chinese Academy of Sciences, PO Box 918-3, Beijing 100039, PR China

Received 5 May 2006 / Accepted 30 June 2006

ABSTRACT

Aims. The prompt optical emission contemporaneous with the γ -rays from γ -ray bursts (GRBs) carries important information on the central engine and explosion mechanism. We study the time lag between prompt optical emission and γ -rays in GRB 990123 and GRB 041219a, which are the only two GRBs detected at optical wavelengths during the ascending burst phase.

Methods. Assuming profiles of prompt optical light curves are the same as the prompt γ -rays, we simulate optical light curves with different time lags and compare them with the observed optical flux. Then the best fit time lag and its error are determined by chi-squared values.

Results. We find that time lags between prompt optical emission and γ -rays in GRB host galaxy rest-frames are consistent in the two GRBs, which is 5 ~ 7 s for GRB 990123 and 1 ~ 5 s for GRB 041219a. This result is consistent with a common origin of prompt optical and γ -ray emissions in the two GRBs. Based on synchrotron cooling model, we also derive the parameters for the two GRBs.

Key words. gamma rays: bursts – methods: data analysis

1. Introduction

The prompt optical emission contemporaneous with the γ -rays from γ -ray bursts (GRBs) carries crucial information on the central engine and explosion mechanism. Internal shock models and external shock models have been proposed to explain the prompt optical emission (see e.g. Meszaros & Rees 1999; Liang et al. 1999). However, the origin of the prompt optical emission remains an open issue.

Prompt optical emission has been found for several GRBs during the brief durations of bursts (e.g. Akerlof et al. 1999; Blake et al. 2005; Vestrand et al. 2005; Rykoff et al. 2005; Klotz et al. 2006). For two of them, i.e. GRB 990123 (Akerlof et al. 1999; Kippen et al. 1999) and GRB 041219a (Vestrand et al. 2005; Barthelmy et al. 2004), early-time optical observations were carried out during their bursting phase, including both ascending and descending parts. Hence, it is possible to study the time lag between prompt optical emission and γ -rays in these two GRBs, which could provide important clues to the origin of prompt optical emission.

2. Data analysis

As shown in Fig. 1, in each GRB, there are only three points with a positive optical detection (crosses) during the burst phase and the profile of optical light curve remains unknown. To avoid redundant parameters, we simply assume that the prompt optical light curve profile is exactly the same as the γ -rays, but with a time lag to be determined. Then we simulate optical light curves with different time lags and compare them with the observed optical flux. The redshift of GRB 990123 is 1.6 (Andersen et al. 1999; Kulkarni et al. 1999). The redshift of GRB 041219a is




Fig. 1. Observed and simulated optical and γ -ray light curves. The upper panel is for GRB 990123 with time indicated in the top axis, and the lower panel is for GRB 041219a with time indicated in the bottom axis. The optical flux measured by Akerlof et al. (GRB 990123) or Vestrand et al. (GRB 041219a) are indicated by crosses, where error bars for detections are given as 1σ values and non-detections are plotted as 2σ upper limits. The corresponding best-fitting simulated values for model 3 described in the paper are plotted as circles. The γ -ray light curves measured by the BASTE (GRB 990123) or Swift BAT (GRB 041219a) are plotted as solid lines. The best-fitting simulated optical light curves for model 3 are plotted as dashed lines.

unknown. Barkov & Bisnovatyi-Kogan (2005) gave a redshift upper limit as $z \leq 0.12$ based on a model of a dust reradiation

Table 1. Results of χ^2 tests for different optical lag models.

model ^a	χ^2	ν	$P(>\chi^2)^b$	Lag ₉₉ ^c (s)	Lag ₀₄ ^d (s)
1	7.22	4	0.12	–	0
2	5150	6	$<10^{-20}$	0	0
3	6.36	4	0.17	12.2	3.0
4	7.17	5	0.21	12.2	5.2

^a Refer to text for details.

^b The probability that a single observation from a χ^2 distribution with degrees of freedom ν will be larger than the χ^2 values in Col. 2.

^c Time lag for GRB 990123.

^d Time lag for GRB 041219a.




Fig. 2. χ^2 value of simulated optical flux as a function time lag for model 3 described in the paper. The upper panel is for GRB 990123 and the lower panel is for GRB 041219a. The solid, dashed and dash-dotted horizontal lines indicate 1σ , 2σ and 3σ errors, respectively.

of IR afterglow in the envelope surrounding the GRB source. In this note, we assume $z = 0.1$ for GRB 041219a.

We consider the following four models for time lag between optical and gamma-ray emissions:

1. No time lag between optical and gamma-ray emissions in GRB 041219a.
2. No time lag between optical and gamma-ray emissions in both GRB 990123 and GRB 041219a.
3. Time lags between optical and gamma-ray emissions in GRB 990123 and GRB 041219a are independent.
4. Time lags between optical and gamma-ray emissions in GRB 990123 and GRB 041219a are the same in their host galaxy rest-frames.

Results of χ^2 tests for the four models are given in Table 1. For model 3, χ^2 values of different time lags are shown in Fig. 2, and simulated light curves with corresponding lags of minimum χ^2 values are shown in Fig. 1. χ^2 values of different time lags for model 4 are shown in Fig. 3.

As shown in Table 1, model 2 is rejected. Models 4 and 3 are better than model 1, indicating that a common time lag between prompt optical emission and γ -rays in both GRBs is quite possible. In model 3 (Fig. 2), for GRB 990123, the best-fitting time lag is 12.2 s, and there is another local minimum of χ^2 values around 18.2 s. The 12 s lag corresponds to moving the second pulse to the second optical point, while 18 s corresponds to moving the first pulse to the second optical point. Therefore the host galaxy rest-frame time lag should be 4.6 ~ 7.2 s. For




Fig. 3. χ^2 value of simulated optical flux as a function time lag for both GRB 990123 and GRB 041219a in their host galaxy rest-frames, for model 4 described in the paper. The solid, dashed and dash-dotted horizontal lines indicate 1σ , 2σ and 3σ errors, respectively.

GRB 041219a, the best-fitting time lag is $3.0^{+2.5}_{-2.3}$ s with 1σ error, and the host galaxy rest-frame time lag will be 0.6 ~ 5.0 s. The χ^2 value is very sensitive to the time lag in GRB 990123, thus in model 4 (Fig. 3), the best-fitting lag value is dominated by the contribution of GRB 990123.

3. Discussion and conclusions

It was reported in some previous studies that the behaviors and origins of the prompt optical emission in GRB 990123 and GRB 041219a are totally different: In GRB 990123, the optical emission was uncorrelated with the prompt gamma-rays, suggesting that the optical emission was generated by a reverse external shock arising from the ejecta's collision with surrounding material; while in GRB 041219a, the optical emission was correlated with the prompt gamma-rays, indicating optical emission was generated by internal shocks driven into the burst ejecta by variations of the inner engine (Vestrand et al. 2005). However, our results show that time correlation between prompt optical and gamma-ray emission are quite consistent in both GRBs. If GRB 041219a is a low-redshift object, the rest-frame time lag of prompt optical emission behind γ -rays in the two GRBs is similar, indicating a common mechanism to produce the prompt optical emission in the two GRBs.

If both prompt optical and γ -ray emissions come from cooling e^\pm pairs in the moving ejecta with a Lorentz factor Γ through synchrotron radiation, assuming velocities of e^\pm are isotropic, the frequency of emitted photon is around the peak frequency of synchrotron photons, which in the observer's frame is

$$\nu \approx \nu_m \approx 2 \times 10^6 \gamma^2 B \Gamma / (1+z) \text{ Hz}, \quad (1)$$

where γ is the Lorentz factor of e^\pm and B is the magnetic field strength in Gauss.

The power radiated per one electron or positron in the observation frame is

$$P = 1.1 \times 10^{-15} \gamma^2 B^2 \Gamma^2 / (1+z)^2 \text{ erg/s}. \quad (2)$$

The time lag between prompt optical emission and γ -rays should be the typical lifetime of e^\pm which emit optical photons

$$t_{\text{lag}} \approx \frac{E}{P} \approx \frac{3.3 \times 10^8 (1+z)}{\gamma B^2 \Gamma} \text{ s}, \quad (3)$$

Table 2. Parameters of GRB 990123 and GRB 041219a based on synchrotron cooling model.

Parameter	GRB 990123	GRB 041219a	Ref. ^a
Observation			
Redshift	1.6	0.1	1, 2, 3
Time lag (s)	12.2	3.0	
$L_{\text{peak},\gamma}$ (erg/s)	1.5×10^{53}	1.2×10^{52}	2, 4
$L_{\text{peak,opt}}$ (erg/s)	5×10^{49}	1.5×10^{47}	5, 6
$\nu_{100 \text{ keV}}$ (Hz)	2.4×10^{19}	2.4×10^{19}	
$\nu_{5000 \text{ \AA}}$ (Hz)	6×10^{14}	6×10^{14}	
Assumption			
Bulk Lorentz factor	300	300	7
Radius (cm)	2×10^{16}	2×10^{16}	7, 8
Derived parameters			
B (Gs)	28	53	
γ_γ	6.1×10^4	2.9×10^4	
γ_{opt}	310	140	
N_e^b	3.6×10^{51}	6.3×10^{49}	
$\nu_{a,\gamma}$ (Hz)	6.5×10^{14}	8.2×10^{14}	9
$\nu_{a,\text{opt}}$ (Hz)	6.6×10^{13}	3.2×10^{13}	9
Σ (cm ⁻²)	2.6×10^{23}	4.5×10^{21}	

^a References: 1. Andersen et al. (1999); 2. Kulkarni et al. (1999); 3. Barkov & Bisnovatyi-Kogan (2005); 4. Barthelmy et al. (2004); 5. Akerlof et al. (1999); 6. Vestrand et al. (2004); 7. Liang et al. (1999); 8. Li et al. (2003); 9. Li et al. (2004).

^b Calculated from $L_{\text{peak},\gamma}$.

Assuming $\Gamma = 300$, which is a typical value in GRBs (see e.g. Liang et al. 1999), we can derive B , γ and the total e^\pm number N_e based on synchrotron cooling, as listed in Table 2. Further assuming an emission radius $R = 2 \times 10^{16}$ cm (see e.g. Liang et al. 1999; Li et al. 2003), we can derive the synchrotron self-absorption frequency ν_a (Li et al. 2004) and the e^\pm column density Σ in the observer's frame

$$\nu_a \approx 1 \times 10^{16} L_{50}^{2/7} \Gamma^{3/7} R_{14}^{-4/7} B_4^{1/7} / (1+z) \text{ Hz}, \quad (4)$$

and

$$\Sigma = \frac{N_e}{\pi(R/\Gamma)^2} \text{ cm}^{-2}. \quad (5)$$

As shown in Table 2, the self-absorption frequencies are less than the observed frequencies, thus prompt emissions can be observed. When absorption is negligible, the optical to γ -ray flux ratio from synchrotron cooling with constant Γ and B should

be $F_{5000 \text{ \AA}}/F_{100 \text{ keV}} = \nu_{5000 \text{ \AA}}/\nu_{100 \text{ keV}} = 2.5 \times 10^{-5}$. This is consistent with the observed value of GRB 041219a, which is 1.2×10^{-5} when there is zero lag between prompt optical emission and γ -rays (Vestrand et al. 2005) and about 1.4×10^{-5} for our best-fitting model. However, in GRB 990123, the peak optical/peak γ -ray flux ratio is 3.3×10^{-4} (Kulkarni et al. 1999; Akerlof et al. 1999) and the flux ratio changes to be 7.1×10^{-4} for our best-fitting model when the time lag is 12.2 s, which is one magnitude larger than the synchrotron-cooling model predicted value, indicating that such a simplified non-absorption synchrotron cooling model alone could not account for all the observed properties of GRB 990123. It is possible that another mechanism may be operating simultaneously which can modify the flux ratio. For example, down-Comptonization of gamma-ray photons in intervening electron clouds may be able to reduce the flux ratio substantially if some gamma-ray photons are converted into optical photons which are also delayed by several seconds with respect to the gamma-ray emission, as proposed independently by Zheng et al. (2006). In this model, different covering factors of high density region around the central engine could lead to different optical to gamma-ray flux ratios.

Acknowledgements. We thank the anonymous referee for helpful comments that improved the manuscript. SNZ acknowledges partial funding support by the Ministry of Education of China, Directional Research Project of the Chinese Academy of Sciences and by the National Natural Science Foundation of China under project no. 10233010 and 10521001.

References

- Akerlof, C., Balsano, R., Barthelemy, S., et al. 1999, *Nature*, 398, 400
Andersen, M. I., Castro-Tirado, A. J., Hjorth, J., et al. 1999, *Science*, 283, 2075
Barthelmy, S., et al. 2004, *GCN Circ.* 2874
<http://gcn.gsfc.nasa.gov/gcn/gcn3/2874.gcn3>
Barkov, M. V., & Bisnovatyi-Kogan, G. S. 2005, *Astrophysics*, 48, 369
Blake, C. H., Bloom, J. S., Starr, D. L., et al. 2005, *Nature*, 435, 181
Kippen, R. M., et al. 1999, *GCN Circ.* 224
<http://gcn.gsfc.nasa.gov/gcn/gcn3/224.gcn3>
Klotz, A., Gendre, B., Stratta, G., et al. 2006, preprint
[arXiv:astro-ph/0604061]
Kulkarni, S. R., Djorgovski, S. G., Odewahn, S. C., et al. 1999, *Nature*, 398, 389
Li, Z., & Song, L. M. 2004, *ApJ*, 608, L17
Li, Z., Dai, Z. G., Lu, T., & Song, L. M. 2003, *ApJ*, 599, 380
Liang, E. P., Crider, A., Böttcher, M., & Smith, I. A. 1999, *ApJ* 519, L21
Meszaros, P., & Rees, M. 1999, *MNRAS*, 306, L39
Rykoff, E. S., Yost, S. A., Krimm, H. A., et al. 2005, *ApJ*, 631, L121
Vestrand, W. T., Wozniak, P. R., Wren, J. A., et al. 2005, *Nature*, 435, 178
Zheng, Z., Lu, Y., & Zhao, Y. H. 2006, *ApJ*, accepted
[arXiv:astro-ph/0606312]