

The bright spiky emission of pulsar B0656+14

P. Weltevrede¹, G. A. E. Wright^{2,1}, B. W. Stappers^{3,1}, and J. M. Rankin^{4,1}

¹ Astronomical Institute “Anton Pannekoek”, University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands
e-mail: wltvrede@science.uva.nl

² Astronomy Centre, University of Sussex, Falmer, BN1 9QJ, UK
e-mail: G.Wright@sussex.ac.uk

³ Stichting ASTRON, Postbus 2, 7990 AA Dwingeloo, The Netherlands
e-mail: stappers@astron.nl

⁴ Physics Department, 405 Cook Physical Science building, University of Vermont, Burlington 05405, USA
e-mail: Joanna.Rankin@uvm.edu

A&A 458, 269–283 (2006), DOI: 10.1051/0004-6361:20065572

Key words. stars: pulsars: individual: PSR B0656+14 – stars: pulsars: general – radiation mechanisms: non-thermal – errata, addenda

A misprint was inserted in Table 2 during the printing process ([2mm] was wrongly placed before AO-P1). The correct table is as follows:

Table 2. The parameters of the best fits to the pulse-energy distributions of the four observations. Besides the parameters of the fitted distribution (where $f_{\text{weak}} = N_{\text{weak}}/N_{\text{tot}}$ is the fraction of weak pulses that are added to the distribution), also the total χ^2 , the number of degrees of freedom and the significance probability are tabulated.

REF	σ	μ	f_{weak}	χ^2	$N_{\text{d.o.f}}$	$P(\chi^2)$
AO-P1	0.99	-0.34	13%	105	32	1×10^{-9}
AO-P2	0.79	-0.34	0%	146	24	2×10^{-19}
AO-L	0.56	-0.13	0%	240	44	8×10^{-29}
WSRT-L	0.48	-0.03	8%	148	44	4×10^{-13}