

Erratum

Thermal and non-thermal components of the interstellar medium at sub-kiloparsec scales in galaxies

R. Paladino^{1,2}, M. Murgia^{1,3}, T. T. Helfer⁴, T. Wong^{5,6}, R. Ekers⁵, L. Blitz⁴, L. Gregorini^{3,7}, and L. Moscadelli¹

¹ INAF – Osservatorio Astronomico di Cagliari, Loc. Poggio dei Pini, Strada 54, 09012 Capoterra (CA), Italy
e-mail: rpaladin@ca.astro.it

² Dipartimento di Fisica, Università di Cagliari, Cittadella Universitaria, 09042 Monserrato (CA), Italy

³ Istituto di Radioastronomia – INAF, Via Gobetti 101, 40129 Bologna, Italy

⁴ Radio Astronomy Laboratory, University of California, Berkeley, CA 94720, USA

⁵ Australia Telescope National Facility, CSIRO, PO Box 76, Epping, N.S.W. 1710, Australia

⁶ School of Physics, University of New South Wales, Sydney, NSW 2052, Australia

⁷ Dipartimento di Fisica, Università di Bologna, Via Irnerio 46, 40126 Bologna, Italy

A&A 456, 847–859 (2006), DOI: 10.1051/0004-6361:20065002

Key words. radio continuum: galaxies – galaxies: spiral – ISM: molecules – stars: formation – errata, addenda

Figure 7 published in A&A, 456 (2006) p. 856 is a duplicate of Fig. 8. Here, we provide the correct version of the figure. The change has no consequences for the discussion in the paper.




Fig. 1. RC-CO, RC-24 μm and 24 μm -CO correlations of NGC 7331, NGC 3627 and NGC 3521 at 6 arcsec resolution. The linear resolution achieved for each galaxy is indicated in the lower-right of the right-most panels.