

LETTER TO THE EDITOR

## ***INTEGRAL* observation of hard X-ray variability of the TeV binary LS 5039/RX J1826.2-1450<sup>★</sup>**

A. D. Hoffmann<sup>1</sup>, D. Klochkov<sup>1</sup>, A. Santangelo<sup>1</sup>, D. Horns<sup>2</sup>, A. Segreto<sup>3</sup>, R. Staubert<sup>1</sup>, and G. Pühlhofer<sup>1</sup>

<sup>1</sup> Kepler Zentrum für Astro- und Teilchenphysik, Institut für Astronomie und Astrophysik, University of Tübingen, Sand 1, 72076 Tübingen, Germany

e-mail: hoffmann@astro.uni-tuebingen.de

<sup>2</sup> Universität Hamburg, Institut für Experimentalphysik, Hamburg, Germany

<sup>3</sup> INAF IFC-Pa, via U. La Malfa 153, 90146 Palermo, Italy

Received 21 November 2008 / Accepted 26 December 2008

### **ABSTRACT**

**Context.** LS 5039/RX J1826.2-1450 is one of the few High Mass X-ray binary systems from which radio and high energy TeV emission has been observed. Moreover, variability of the TeV emission with orbital period was detected.

**Aims.** We investigate the hard X-ray (25–200 keV) spectral and timing properties of the source with the monitoring IBIS/ISGRI instrument on-board the *INTEGRAL* satellite.

**Methods.** We present the analysis of *INTEGRAL* observations for a total of about 3 Msec exposure time, including both public data and data from the Key Programme. We search for flux and spectral variability related to the orbital phase.

**Results.** The source is observed to emit from 25 up to 200 keV and the emission is concentrated around inferior conjunction. Orbital variability in the hard X-ray band is detected and established to be in phase with the orbitally modulated TeV emission observed with HESS. For this energy range we determine an average flux for the inferior conjunction phase interval of  $(3.54 \pm 2.30) \times 10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup>, and a flux upper limit for the superior conjunction phase interval of  $1.45 \times 10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup> (90% conf. level respectively). The spectrum for the inferior conjunction phase interval follows a power law with an index  $\Gamma = 2.0_{-0.2}^{+0.2}$  (90% conf. level).

**Key words.** X-ray: binaries – stars: individual: LS 5039/RX J1826.2-1450

### **1. Introduction**

LS 5039/RX J1826.2-1450 is a High Mass X-ray Binary (HMXRB) system, which was discovered in the *ROSAT* all-sky survey and identified with optical follow-up observations by Motch et al. (1997). The compact companion is estimated to have a mass of  $M \sim 3.7 M_{\odot}$ , orbiting a bright,  $V = 11.2$ , O6.5V(f) star with an orbital period  $P_{\text{orb}} \sim 3.9060 \pm 0.00017$  d, an eccentricity  $e = 0.35 \pm 0.04$  and a distance of  $d = 2.5 \pm 0.1$  kpc (Casares et al. 2005). Based on the observation of persistent asymmetric milliarsecond radio outflows (extending up to  $\sim 1000$  AU), Martí et al. (1998) and Paredes et al. (2000) suggested the presence of a mildly relativistic ( $v \sim 0.2 c$ ) jet. Paredes et al. (2002) and Ribó et al. (2008) have therefore proposed the source to be a microquasar.

LS 5039 has been observed by several X-ray satellites. A summary of all observations can be found in Bosch-Ramon et al. (2005). Flux variations on timescales of days as well as mini-flares on shorter timescales have been observed. RXTE observations, covering the whole orbital period, show variability with a maximum of emission at orbital phase  $\phi = 0.8$ , interpreted due to periastron passage. Moreover, the power-law shaped X-ray


spectrum appears harder in the high flux state (Bosch-Ramon et al. 2005).

Because inhomogeneous jets could emit  $\gamma$ -rays via inverse Compton scattering, Paredes et al. (2000, 2002) have proposed the source to be the counterpart of the unidentified high energy EGRET source 3EG J1824-1514 (Hartman et al. 1999) and also a component of the MeV emitter GRO J1823-12, which is a superposition of three EGRET sources (Collmar 2003).

The source has likewise been discovered to emit at TeV energies with the High Energy Stereoscopic System (HESS) (Aharonian et al. 2005). This demonstrated that X-ray binaries are capable of accelerating particles to TeV energies. The HESS measurements have revealed that the flux and energy spectrum of the source are indeed modulated with the  $\sim 3.9$  d orbital period of the binary system (Aharonian et al. 2006). The VHE  $\gamma$ -ray emission is largely confined to half of the orbit and peaks around the inferior conjunction (INFC) epoch of the compact companion, where a hardening of the spectrum is observed. According to the authors these findings indicate that  $\gamma$ -ray absorption with pair production occurs in the system. Interestingly, orbital variability at TeV  $\gamma$ -ray energies has also been discovered in the similar system LS I+61°303 with MAGIC (Albert et al. 2006) and VERITAS (Acciari et al. 2008).

The nature of the compact object in the system RX J1826.2-1450/LS 5039 as well as the origin of the TeV emission is still under debate. A microquasar scenario (see e.g. Paredes et al. 2000; Bosch-Ramon & Paredes 2004; Torres & Sierpowska-Bartosik 2008) as well as a pulsar scenario (see e.g. Dubus 2006;

<sup>★</sup> Based on observations with *INTEGRAL*, an ESA project with instruments and science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland, and with the participation of Russia and the USA.



**Fig. 1.** A sketch of the system. The orbital values are listed in Casares et al. (2005) and the definition of phase intervals is from Aharonian et al. (2006).

Sierpowska-Bartosik & Torres 2007; Bosch-Ramon et al. 2008) have been discussed. In this Letter, based mainly on archival data from *INTEGRAL* which span 4.6 years between March 2003 and October 2007, we confirm the detection of the source reported by Bird et al. (2007) in the energy range from 25 keV up to 200 keV. We report the discovery of the modulation of the hard X-ray flux with orbital phase.

## 2. Observations and data analysis

### 2.1. Observations

The International Gamma-Ray Astrophysics Laboratory (*INTEGRAL*, Winkler et al. 2003) carries on board two main gamma-ray instruments, the *Spectrometer on INTEGRAL* (SPI, 20 keV–8 MeV, Vedrenne et al. 2003), and the *Imager on-Board INTEGRAL Satellite* (IBIS, 15 keV–10 MeV, Ubertini et al. 2003) as well as two monitoring instruments in the X-ray and optical range, the *Joint European X-Ray Monitor* (JEM-X, 3–35 keV, Lund et al. 2003), and the *Optical Monitoring Camera* (OMC, Mas-Hesse et al. 2003).

1648 science windows (ScWs) in the region that includes LS 5039 are found in the data archive available at the *INTEGRAL* Scientific Data Center (ISDC, <http://isdc.unige.ch/>), including public data as well as data from the Key Programme. These ScWs were preselected choosing an off-axis angle of the source smaller than 10 degrees and choosing only pointing observations that had a nominal observation time of more than 1000 s. 1627 ScWs were successfully processed through the analysis with OSA 7.0 up to the image step IMA.

In order to reduce statistical effects a further selection of “good science windows” was necessary. ScWs are included in the analysis when they fulfill the following criteria:

1. the effective exposure time after extraction is larger than 1000 s;
2. the number of good time intervals (GTI) does not exceed 10;
3. the mean background intensity level of a single ScW in the energy range 25–60 keV falls within  $3\sigma$  of the distribution of the mean background intensity levels of all other ScWs in the same energy range.

The resulting 1393 ScWs cover the time range from 03-11-2003 (ScW: 005000080010) to 10-25-2007 (ScW: 061400950010).

**Table 1.** Summary of selected and analyzed *INTEGRAL* data for LS 5039. Analyzed data were taken from 69 revolutions, i.e. from revolution 0050 (03-11-2003) to revolution 0614 (10-25-2007).

Number of all extracted ScWs	1648
Number of ScWs after selection	1393
Number of ScWs in phase SUPC	721
Number of ScWs in phase INFC	672
final total analyzed exposure time	$\approx 2.99$ Ms
splitted in SUPC phase	$\approx 1.56$ Ms
and in INFC phase	$\approx 1.43$ Ms

Firstly, the sample was split into two parts according to the orbital phase for imaging analysis, and secondly into 10 smaller parts according to the orbital phase to analyse the flux dependence on orbital phase (see Sect. 2.2). Table 1 summarizes the observational data of the source LS 5039 taken by the *INTEGRAL* satellite and the selection made for this analysis. Only in less than 10% of the selected ScWs was LS 5039 in the fully coded field of view of JEM-X. Based on the sensitivity of JEM-X<sup>1</sup> and the amount of available data, a JEM-X analysis for a faint source like LS 5039 is not suitable. In this paper we therefore focus our analysis on the data of the *Integral Soft Gamma-Ray Imager* (IBIS/ISGRI).

### 2.2. Analysis

For the analysis of ISGRI data the Off-line Science Analysis (OSA) software (version 7.0) provided by the ISDC was used. In addition, we cross checked the results and also generated the lightcurve in Sect. 2.2.2 with the analysis package developed by INAF-IFC Palermo<sup>2</sup> (Segreto & Ferrigno 2006).

A special handling of the OSA was used for this analysis. First, we used a user catalogue which includes 115 sources, i.e. all sources listed in the *INTEGRAL* General Reference Catalog<sup>3</sup>, which are in the field of view (FoV) and which turned out to be as bright as or brighter than RX J1826.2-1450 ( $\equiv$  LS 5039). This is required because the source of interest is quite faint and the crowded field produces too many ghost images if the known sources are not subtracted. Second, the spectrum was produced with the help of the OSA tool MOSAIC\_SPEC<sup>4</sup> from 13 mosaics, each covering one energy bin in the 13–250 keV range. The binning was chosen identical to the default energy binning OSA uses for the spectrum calculation step (SPE). This procedure is needed due to the fact that the source is faint and the FoV is too crowded, and provides more reliable results than the standard OSA pipeline does.

We also checked our results using the standard SPE and the lightcurve creation (LCR) steps of OSA. In this case we used a special catalogue including all sources with a detection significance greater than  $6\sigma$  in the energy band 20–60 keV. The catalogue includes 65 sources. According to the User Manual of the *INTEGRAL* IBIS/ISGRI data analysis this is beyond the standard method (see the IBIS Analysis Manual<sup>1</sup> for details).

<sup>1</sup> The User Manual to the OSA can be found at: <http://isdc.unige.ch/?Support+documents>. See p. 15 of JEM-X Analysis User Manual for a JEM-X sensitivity plot.

<sup>2</sup> <http://www.pa.iasf.cnr.it/~ferrigno/INTEGRALsoftware.html>

<sup>3</sup> The *INTEGRAL* General Reference Catalogue is available at ISDC: <http://isdc.unige.ch/index.cgi?Data+catalogs>

<sup>4</sup> [http://isdc.unige.ch/Soft/download/osa/osa\\_doc/osa\\_doc-5.0/osa\\_um\\_ibis-5.0/node108.html#mosaspec](http://isdc.unige.ch/Soft/download/osa/osa_doc/osa_doc-5.0/osa_um_ibis-5.0/node108.html#mosaspec)

However, our results show that a careful treatment of this standard method provides a comparable result in spite of the restrictions (see Fig. 4 and Sect. 2.2.2).

### 2.2.1. Imaging

The source is clearly seen at a significance level of  $7.7\sigma$  in the lower energy band of 25–60 keV in the significance map based on the sum of all ScWs fulfilling the selection criteria. In addition, we divided the data set into two phase intervals: the superior conjunction passage at orbital phase  $0.9 < \phi \leq 1.45$  (SUPC), and the inferior conjunction, at orbital phase  $0.45 < \phi \leq 0.9$  (INFC) (see Figs. 1 and 4 for details). Phase intervals are defined in Aharonian et al. (2006) for the HESS data. The two orbital resolved significance maps in the energy range 25–60 keV are shown in Fig. 2. Positions of sources (open boxes) are provided by the latest *INTEGRAL* General Reference Catalog<sup>3</sup>.

At the INFC phase interval, the source was detected with a significance  $\approx 7.8\sigma$ , for an exposure time of  $\approx 1.43$  Msec (Fig. 2, lower panel). At the SUPC phase interval the source was observed for a total exposure time of 1.56 Ms, and was only marginally detected: at the source position there is an excess at the level of  $2.8\sigma$  (Fig. 2, upper panel). For the total sample and the total energy band 25–200 keV the source is detected at the position fitted by OSA (RA = 276.517 deg, Dec =  $-14.812$  deg) with  $7.4\sigma$ . Within the errors, this is consistent with the source position determined by Bird et al. (2007).

### 2.2.2. Timing analysis


A lightcurve produced with the INAF-IFC data analysis software (Segreto & Ferrigno 2006) was used for the timing analysis. According to the authors this software is suitable to produce reliable results even in the case of the presence of a larger number of sources in the FoV.

Searching for an unambiguous period failed due to low statistics and large gaps in the sample of observations. But a periodogram produced by epoch folding provides a peak at a period of 3.903 days, which is within uncertainties identical to the orbital period determined by Casares et al. (2005) in the optical band. Figure 3 shows the periodogram.


In a next step the lightcurve was folded using the known orbital period of  $P_{\text{orb}} = 3.90603$  days with  $\text{HJD}(T_0) = 2451943.09 \pm 0.10$  ( $\equiv$  phase  $\phi = 0.0$  and periastron passage) determined by Casares et al. (2005), to produce a 10 bin orbital phase profile. In addition, 10 mosaics corresponding to each of the 10 orbital phase intervals were produced and the count rates at the catalogue<sup>3</sup> position were determined. The values of the fluxes were derived from a 2D-Gaussian fit with fixed width ( $HWHM = 6.0$  arcmin) and fixed source position. Both methods yield (within errors) comparable results: the hard X-ray emission is clearly modulated with the orbit, and appears to be in phase with the TeV emission (Fig. 4). The HESS data points in the figure are from Aharonian et al. (2006).

### 2.2.3. Spectral analysis

A statistically significant IBIS/ISGRI spectrum of LS 5039 was obtained for the orbital phase interval corresponding to INFC (not shown here) and was modeled with a power law with  $\Gamma = 2.0^{+0.2}_{-0.2}$  (90% confidence level). The reduced  $\chi^2$  for the fit is 1.4 (9 d.o.f.). Mean countrates for the two phase intervals allow us to estimate a flux for the INFC phase interval of



**Fig. 2.** *INTEGRAL* significance maps in the energy range 25–60 keV. The open boxes are known source positions. Comparable exposure times and a higher significance for LS 5039 in the INFC phase interval indicates variability of the hard X-ray emission with orbital phase (see also Fig. 4). The INFC phase interval includes the maximum flux state of the source. Details can be found in the text.




**Fig. 3.** An epoch folding on the *INTEGRAL* lightcurves shows a significant peak at a period that is nearly identical to the orbital period known and determined from optical observation of the system (Casares et al. 2005). All other peaks are likely beat frequencies due to the large gaps in the sample of observations.

$(3.54 \pm 2.30) \times 10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup> (90% conf. level). For the SUPC phase interval a flux upper limit of  $1.45 \times 10^{-12}$  erg cm<sup>-2</sup> s<sup>-1</sup> (90% conf. level) for the energy range 25–200 keV can be determined.

## 3. Summary and discussion

In this paper we report on a 3 Ms *INTEGRAL* monitoring of the HMXRB RX J1826-1450/LS 5039. The faintness of the source



**Fig. 4.** *Top:* orbital profile for LS 5039 in the energy range 25–200 keV observed with *INTEGRAL*. The lightcurve is folded with the orbital period of 3.9 days. *Bottom:* orbital profile for LS 5039 for the energy range above 1 TeV observed with HESS (Aharonian et al. 2006). The hard X-ray emission is in phase with the TeV  $\gamma$ -rays.

and the fact that it lies in a crowded field required a very careful, non-standard handling of the data.

Our results, obtained by applying different analysis methods, point out that the source significantly emits at hard X-rays (25–200 keV) and that this emission varies with the orbit in phase with the very high energy  $\gamma$ -rays detected with HESS (Fig. 4). The spectrum at the inferior conjunction is well described by a power law, while at the superior conjunction the hard X-ray emission, if any, is below the sensitivity of *INTEGRAL*. Our result indicates that accretion might not be the mechanism for the production of the hard emission, since, in this case, we would expect a rather sharp flux maximum near periastron.

Moreover, the observed close correlation of hard X-ray and VHE  $\gamma$ -ray emission in phase suggests that the emission would originate from the same source region, possibly from particles produced in the same acceleration process.

This casts doubt on the simple explanation that the TeV variability is mainly caused by  $\gamma$ - $\gamma$  absorption, as suggested in Aharonian et al. (2006).

We note that, from the correlation alone, more complicated scenarios involving  $\gamma$ - $\gamma$  absorption such as described by Dubus et al. (2008) cannot be ruled out. However the cascades initiated in the  $\gamma$ - $\gamma$  absorption process produce secondary electrons (Bednarek 1997) also emitting X-ray synchrotron emission. Therefore, soft and hard X-ray flux levels can be used to constrain the amount of absorption involved. The detection of TeV emission at SUPC together with the upper limit in the hard

X-ray domain derived here may pose problems for models invoking only one emission region close to the compact object, as recently discussed in Bosch-Ramon et al. (2008).

Radio observations interpreted as jets have led to a micro-quasar interpretation for this source. In this scenario, particles are accelerated to  $\approx$ TeV energies (Perucho & Bosch-Ramon 2008) due to the interaction between the jet and the stellar wind of the companion. Alternatively, Dubus (2006) has suggested the source to be a pulsar binary system, where VHE emission arises from the pulsar wind material shocked by the interaction with the stellar companion wind and afterwards flowing away in a comet-shape tail. Unfortunately, the observations presented here do not enable us to discriminate between the two scenarios, although the predicted X-ray variability by Paredes et al. (2006) seems to be not in accordance with our observations. Finally, we remark that a hadronic scenario (Romero et al. 2003; Romero & Vila 2008) is not ruled out by our findings. A longer and more detailed interpretation of our results is beyond the scope of this Letter, and will be the subject of further investigations.

*Acknowledgements.* We acknowledge the support of the Deutsches Zentrum für Luft- und Raumfahrt (DLR) under grant number 50OR0302. This work is based on observations with *INTEGRAL*, an European Space Agency (ESA) project with instruments and science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland, and with the participation of Russia and the USA. We also thank the anonymous referee for his/her valuable comments and suggestions.

## References

- Acciari, V. A., Beilicke, M., Blaylock, G., et al. 2008, *ApJ*, 679, 1427  
 Aharonian, F., Akhperjanian, A. G., Aye, K.-M., et al. 2005, *Science*, 309, 746  
 Aharonian, F., Akhperjanian, A. G., Bazer-Bachi, A. R., et al. 2006, *A&A*, 460, 743  
 Albert, J., Aliu, E., Anderhub, H., et al. 2006, *Science*, 312, 1771  
 Bednarek, W. 1997, *A&A*, 322, 523  
 Bird, A. J., Malizia, A., Bazzano, A., et al. 2007, *ApJS*, 170, 175  
 Bosch-Ramon, V., & Paredes, J. M. 2004, *A&A*, 417, 1075  
 Bosch-Ramon, V., Paredes, J. M., Ribó, M., et al. 2005, *ApJ*, 628, 388  
 Bosch-Ramon, V., Khangulyan, D., & Aharonian, F. 2008, [arXiv:0810.4494]  
 Casares, J., Ribó, M., Ribas, I., et al. 2005, *MNRAS*, 364, 899  
 Collmar, W. 2003, in Proc. of 4th Agile Science Workshop, Frascati (Rome), on 11–13 June 2003, 177  
 Dubus, G. 2006, *A&A*, 456, 801  
 Dubus, G., Cerutti, B., & Henri, G. 2008, *A&A*, 477, 691  
 Hartman, R. C., Bertsch, D. L., Bloom, S. D., et al. 1999, *ApJS*, 123, 79  
 Lund, N., Budtz-Jørgensen, C., Westergaard, N. J., et al. 2003, *A&A*, 411, L231  
 Martí, J., Paredes, J. M., & Ribo, M. 1998, *A&A*, 338, L71  
 Mas-Hesse, J. M., Giménez, A., Culhane, J. L., et al. 2003, *A&A*, 411, L261  
 Motch, C., Haberl, F., Dennerl, K., Pakull, M., & Janot-Pacheco, E. 1997, *A&A*, 323, 853  
 Paredes, J. M., Martí, J., Ribó, M., & Massi, M. 2000, *Science*, 288, 2340  
 Paredes, J. M., Ribó, M., Ros, E., Martí, J., & Massi, M. 2002, *A&A*, 393, L99  
 Paredes, J. M., Bosch-Ramon, V., & Romero, G. E. 2006, *A&A*, 451, 259  
 Perucho, M., & Bosch-Ramon, V. 2008, *A&A*, 482, 917  
 Ribó, M., Paredes, J. M., Moldón, J., Martí, J., & Massi, M. 2008, *A&A*, 481, 17  
 Romero, G. E., & Vila, G. S. 2008, *A&A*, 485, 623  
 Romero, G. E., Torres, D. F., Kaufman Bernadó, M. M., & Mirabel, I. F. 2003, *A&A*, 410, L1  
 Segreto, A., & Ferrigno 2006, in Proc. of the 6th *INTEGRAL* Workshop “The Obscured Universe”, Moscow, on 2–8 July 2006, ed. S. Grebenev, R. Sunyaev, & C. Winkler, 633  
 Sierpowska-Bartosik, A., & Torres, D. F. 2007, *ApJ*, 671, L145  
 Torres, D. F., & Sierpowska-Bartosik, A. 2008, [arXiv:0810.4435]  
 Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131  
 Vedrenne, G., Roques, J.-P., Schönfelder, V., et al. 2003, *A&A*, 411, L63  
 Winkler, C., Courvoisier, T. J.-L., Di Cocco, G., et al. 2003, *A&A*, 411, L1