

*Erratum*

**Circumstellar dust shells around long-period variables**

**IX. Molecular layers resulting from shocked carbon-rich atmospheres**

Ch. Helling<sup>1</sup> and J. M. Winters<sup>1,2,3</sup>

<sup>1</sup> Institut für Astronomie und Astrophysik, TU Berlin, Sekr. PN 8-1, Hardenbergstraße 36, 10623 Berlin, Germany

<sup>2</sup> Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany


<sup>3</sup> DEMIRM, UMR 8540, Observatoire de Paris, 61 Av. de l'Observatoire, 75014 Paris, France

A&A, 366, 229–240 (2001), DOI: 10.1051/0004-6361:20000206


By mistake, Fig. 3 has been plotted again in Fig. 4 (p. 233). The correct figures are given below.

---

*Send offprint requests to:* Ch. Helling  
e-mail: chris@astro.physik.tu-berlin.de



**Fig. 3.** Molecular number densities  $n$  for Model F (small constant gas opacity) in the hydrostatic case (l.h.s.) and for the hydrodynamic case (r.h.s.). The instant of time depicted is the same as in Fig. 1. (Upper panel: temperature (dashed black – static, full black – dynamic), gas density (dashed grey – static, full grey – dynamic))



**Fig. 4.** Same as Fig. 3 for Model B (Plank mean gas opacity). The instant of time depicted is the same as in Fig. 2