

## Full–Stokes spectropolarimetry of solar prominences<sup>\*</sup>

F. Paletou<sup>1</sup>, A. López Ariste<sup>2</sup>, V. Bommier<sup>3</sup>, and M. Semel<sup>4</sup>

<sup>1</sup> Observatoire de la Côte d’Azur, Département G. D. Cassini, BP 4229, 06304 Nice Cedex 4, France

<sup>2</sup> High Altitude Observatory, NCAR<sup>\*\*</sup>, PO Box 3000, Boulder CO 80303, USA  
e-mail: [alopez@hao.ucar.edu](mailto:alopez@hao.ucar.edu)

<sup>3</sup> Observatoire de Paris, DAMAp (CNRS/UMR 8588), 92195 Meudon, France  
e-mail: [v.bommier@obspm.fr](mailto:v.bommier@obspm.fr)

<sup>4</sup> Observatoire de Paris, DASOP–LPSH (CNRS/UMR 8645), 92195 Meudon, France  
e-mail: [meir.semel@obspm.fr](mailto:meir.semel@obspm.fr)

Received 18 June 2001 / Accepted 28 June 2001

**Abstract.** We report on first spectropolarimetric observations of solar prominences made at THEMIS. Hereafter we shall emphasize on the preliminary analysis of He I D<sub>3</sub> full-Stokes observations. Our measurements clearly show manifestations of both the Hanle and Zeeman effects. More generally, we demonstrate the very favourable performances of the THEMIS multi-line spectropolarimetric mode for further new investigations of the magnetic field in solar prominences.

**Key words.** Sun: prominences – Sun: magnetic fields – polarization

### 1. Introduction

Quoting Einar Tandberg-Hanssen (1974), “*the single, physically most important parameter to study in prominences may be the magnetic field. Shapes, motions, and in fact the very existence of prominences depend on the nature of the magnetic field threading the prominence plasma*”. Indeed, being low  $\beta$  structures, the channeling of plasma motions and of thermal conduction are controlled by the magnetic field, and it provides also support to the prominence material against gravity (Démoulin 1998). Details and further references can be found in the reviews of Leroy (1989) and Landi Degl’Innocenti (1990), as well as in Tandberg-Hanssen (1974, 1995).

Polarimetry constitutes a unique tool for the *direct* measurement of astrophysical magnetic fields. Up to now, the most comprehensive observational work on the determination of magnetic fields in solar prominences was led by J. L. Leroy using the facilities of the *Pic du Midi* observatory (France). First measurements, using the D<sub>3</sub> line of He I, were done in 1973–1976 with a photo-electric polarimeter attached to a 26 cm diameter coronagraph

(Leroy et al. 1977; Leroy 1977, 1978). This instrumentation offered a rather high polarimetric sensitivity (better than  $10^{-3}$ ) at the price of a poor spatial resolution ( $\sim 3''$ ) and, no spectral resolution. However, the measured linear polarization signals could be interpreted further in terms of magnetic field strength and orientation by use of the Hanle effect theory (Sahal-Bréchet et al. 1977; Bommier & Sahal-Bréchet 1978).

Shortly after, Bommier et al. (1981) pointed-out the possibility of a complete determination of the prominence magnetic field from two simultaneous (linear) polarization measurements made at two different wavelengths. One solution is indeed to observe two different spectral lines, such as D<sub>3</sub> and the H $\alpha$  or H $\beta$  Balmer lines of H I, for instance. This approach was successfully applied to diagnose new data taken during the 1974–1982 period (Leroy et al. 1983, 1984).

Another possibility comes from the use of the two components of D<sub>3</sub> itself, namely the  $3d^3D_{3,2,1} \rightarrow 2p^3P_{1,2}$  and  $3d^3D_1 \rightarrow 2p^3P_0$  groups of atomic transitions. However it requires to spectrally *resolve* those two components distant of some 34 pm. First profiles of the D<sub>3</sub> line observed in solar prominences were recorded in September 1977 with the HAO Stokes polarimeter available at the 40 cm coronagraph of the NSO Evans Solar Facility at Sacramento Peak (Baur et al. 1981). Those new measurements allowed for a complete determination of the magnetic fields in quiescent prominences (Landi Degl’Innocenti 1982; Athay et al. 1983; Querfeld et al. 1985).

Send offprint requests to: F. Paletou,  
e-mail: [paletou@obs-nice.fr](mailto:paletou@obs-nice.fr)

<sup>\*</sup> Based on observations made at THEMIS, operated on the Island of Tenerife by CNRS–CNR in the Spanish *Observatorio del Teide* of the *Instituto de Astrofísica de Canarias*.

<sup>\*\*</sup> The National Center for Atmospheric Research (NCAR) is sponsored by the National Science Foundation.

It was then well assessed from all these works that the mean magnetic field in quiescent prominences (i) has strength in the 3–15 G range, (ii) it is almost horizontal and (iii) it makes a small angle ( $\sim 25^\circ$ ) with the long axis of the structure. However, after the success of the 70's and 80's, and besides quite recent observations of *filaments* (i.e. a prominence as seen on projection against the solar disk) made in the 1083 nm line of He I by Lin et al. (1998), there has been up to now just very little “activity” in the field of the spectropolarimetry of solar prominences. On the other hand, nowadays instruments provide higher spatial and spectral resolution observations together with a  $\sim 10^{-3}$  (or better) polarimetric sensitivity.

THEMIS allows for multi-line spectropolarimetric observations at high resolution in the visible, which is definitely critical in order to perform a reliable diagnosis of prominences magnetic fields. The photon collector is large: the primary mirror's diameter is 90 cm, and it was super-polished at REOSC down to a 4 Å roughness (P. Assus, private communication). In combination with a unique multi-line capability, full-Stokes spectropolarimetry gives an opportunity to carry out new observations with the aim of improving our knowledge of the prominences magnetic field.


## 2. Observations and data reduction

A 7-days observing run dedicated to the full-Stokes spectropolarimetry of limb prominences in the He I D<sub>3</sub> and H I H $\alpha$  spectral lines simultaneously, was conducted in June 2000. We used the multi-line spectropolarimetric (MTR) observing mode available at THEMIS. The latter has been described in some details by Paletou & Molodij (2001).

### 2.1. Observational procedure

We shall focus on observations made on June 15th, 2000 between 07:18 and 09:40 UT. We selected a low-lying and rather bright prominence passing at the South-Western limb, at a polar angle of  $\sim 246^\circ$ . Synoptic observations made during the previous days showed that the filament long-axis was oriented  $\sim 45^\circ$  respective to the local meridian. Our observations were made in a sit-and-stare mode, with the slit oriented parallel to the nearest limb, and at some 30'' above the limb. We recorded a total of  $4 \times 50$  images in a row, according to the following temporal sequence for the polarization analysis:  $[(I \pm U), (I \mp Q), (I \mp U), (I \mp V)]$  (a calcite beam-splitter was used; therefore the 2 orthogonal states of polarization were measured simultaneously). This took us 31 min with an exposure time of 8 s per image. The scintillometer's real-time measurement available at THEMIS indicated a very stable and excellent seeing, of the order of 0.5'', during the whole observing sequence.

At the time of the observation, the sky's transparency was also quite good for the site. In order to evaluate the amount of diffused light contaminating the prominence



**Fig. 1.** The mean D<sub>3</sub> emission profile is displayed before having corrected it for diffused light. The thick line (slightly shifted in intensity for comparison) represents the spectral variations of the latter quantity as derived from our observations.

emission spectra, we recorded a set of images with the slit put off-limb at a neighbouring radial position, although at the same radial distance and with the same relative orientation vs. the nearest limb as in the prominence case. The level of diffused light as compared to the overall signal is shown in Fig. 1. Spectral variations are small but not negligible. A faint chromospheric D<sub>3</sub> emission component can be seen. At both ends of the spectral window, small absorption features can be clearly noticed, corresponding to photospheric Fe lines. No significant polarization signal was found from this set of images.


### 2.2. Data reduction

We adopted the numerical procedure first proposed by Trujillo Bueno et al. (2001). We could therefore retrieve a full Stokes vector with a temporal cadence of  $\sim 37$  s. Indeed, after each set of four successive independent measurements, a Stokes vector can be computed by inverting, for each beam, the “modulation” scheme which we may formally write as:

$$\begin{pmatrix} (I \pm U) \\ (I \mp Q) \\ (I \mp U) \\ (I \mp V) \end{pmatrix} = \begin{pmatrix} 1 & 0 & \pm 1 & 0 \\ 1 & \mp 1 & 0 & 0 \\ 1 & 0 & \mp 1 & 0 \\ 1 & 0 & 0 & \mp 1 \end{pmatrix} \begin{pmatrix} I^\pm \\ Q^\pm \\ U^\pm \\ V^\pm \end{pmatrix}, \quad (1)$$

and finally, recombine the two beams by a simple averaging ( $\sigma = \frac{1}{2}(\sigma^+ + \sigma^-)$ , with  $\sigma = I, Q, U, V$  respectively). Our naming convention for the linear polarization Stokes parameters  $Q$  and  $U$  agrees with the usual reference for which the direction of positive  $Q$  is parallel to the limb.

It is important to bear in mind that, as written above, the modulation scheme implicitly considers *perfect* retarders. On the contrary, the quarter-wave plates we used show some *phase chromatism* in the 400–700 nm range; they depart from being  $\lambda/4$  by roughly  $+3^\circ$  at 587.6 nm. However, one can easily extend the above-described reduction scheme in order to take into account for phase



**Fig. 2.** Maps of the time-averaged Stokes parameters  $I$ ,  $Q$ ,  $U$  and  $V$ .

chromatism, and therefore accurately consider crosstalks among the measured Stokes parameters.

Application of this procedure shows that  $U$  could be determined with a  $1.25 \times 10^{-3}$  polarimetric sensitivity per pixel, and twice as much for  $Q$  and  $V$ , in agreement with previous estimates of López Ariste et al. (2000). This can be improved by a factor of  $\sqrt{50}$  by averging the images of the full temporal sequence in time, while keeping intact the spatial-spectral multiplex on the CCD images.

### 3. First results and discussion

One can clearly see in Fig. 2 a very nice  $Q$  signal, persisting along most of the slit i.e. over a field of view of  $\sim 50''$ . The contribution of both “components” of the  $D_3$  line are detected, although the red component is indeed weaker and more affected by the noise level; typical values are of the order of  $Q/I_{\max} = 0.7\%$  for the blue component, while it goes down at the 0.2% level for the red one. Spatial variations along the slit are also noticeable.

Stokes  $U \sim 0$  at most of the upper part of the slit. This would agree with transverse magnetic field lines, as long as the field strength does not exceed a few tens of Gauss. However, examination of the lower half of the field of view reveals patches of small amplitude  $U$  signals at the  $-0.2\%$  level. These are signatures of a rotation of the plane of polarization such as predicted by the Hanle effect, i.e. a modification of non-magnetic resonance polarization

induced by the presence of magnetic field lines having line-of-sight components.

Our measurements of both  $Q$  and  $U$  Stokes parameters can be spectrally integrated giving mean values  $\bar{Q}$  and  $\bar{U}$  at different spatial positions along the slit. Then, following previous works done by Leroy’s group, a linear polarization rate can be derived:

$$p = \frac{\sqrt{\bar{Q}^2 + \bar{U}^2}}{\bar{I}}. \quad (2)$$


This obviously need a very careful correction of the dif-fused light in order to avoid spurious estimates of the integrated line intensity  $\bar{I}$ .

Afterwards, we can compute the linear polarization direction which is given by:

$$\theta = \frac{1}{2} \arctan \frac{\bar{U}}{\bar{Q}}. \quad (3)$$

Polarization diagrams of  $p$  vs.  $\theta$  can be built in order to inverse the data. More precisely, the rate of linear polarization was normalized to an upper limit value,  $p_{\max}$ , which corresponds to the maximum rate which would have been obtained in the *absence* of magnetic field. For this height above the limb,  $p_{\max} = 4.66\%$  (Leroy et al. 1977).

Points in Fig. 3 with the largest  $p/p_{\max}$  values denote magnetic field strengths typical for quiescent prominences (say, up to 15 G), according to polarization diagrams published earlier (e.g. Sahal-Bréchet et al. 1977). However,



**Fig. 3.** Polarization diagram using the spectrally integrated linear polarization measurements. Clear Hanle depolarizations are seen together with several points revealing a rotation of the plane of polarization.

lower values of  $p/p_{\max}$  are consistent with larger field strengths. From our polarization diagram, we can therefore conclude that we detected variations of the strength and orientation of  $\mathbf{B}$  within the  $\sim 50''$  of our field of view.

However, the most striking feature in these observations may very likely come from the Stokes  $V$  image in Fig. 2. Indeed, although very small signals appear along most of the upper half of the field of view, very conspicuous Zeeman-like signals are detected in the lower half of the field of view. This clearly proves the presence of rather large magnetic field values, and again its change in strength and orientation within the prominence body. Figure 4 shows a typical example of the agreement between Stokes  $V$  and  $\partial I/\partial\lambda$  at a given slit location. Indeed, according to the classical relationship for the Zeeman effect:

$$V \propto 4.67 \times 10^{-13} \lambda_0^2 (\text{\AA}) g_{\text{eff}} B_{\text{los}} (\text{G}) \frac{\partial I}{\partial \lambda}, \quad (4)$$


we can easily evaluate large magnetic field values in what seems to be an “active lobe” in the prominence (this is also suggested by the presence of large velocity fields). Values in the 30–45 G range are found.

#### 4. Conclusion

THEMIS proves to be a suitable instrument for the observation of cool coronal structures such as prominences. Our preliminary full-Stokes observations in the He I D<sub>3</sub> spectral line showed clear variations in strength and orientation of the prominence magnetic field.

New measurements are expected to bring new light on the study of the inhomogeneity of the magnetic field, and its relationship with the well-known fine structure of the prominences. Temporal variations will also be studied in more details.

*Acknowledgements.* Our warmest thanks go to J. L. Leroy for his advices and kind encouragements.



**Fig. 4.** Example of one of the Stokes  $V$  profiles as compared to the re-scaled derivative of  $I$  displayed as the thick curve. This corresponds to a line-of-sight magnetic field of 42 G.

#### References

- Athay, R. G., Querfeld, C. W., Smartt, R. N., Landi Degl’Innocenti, E., & Bommier, V. 1983, *Sol. Phys.*, 89, 3
- Baur, T. G., Elmore, D. E., Lee, R. H., Querfeld, C. W., & Rogers, S. R. 1981, *Sol. Phys.*, 70, 395
- Bommier, V., Leroy, J. L., & Sahal-Br  chot, S. 1981, *A&A*, 100, 231
- Bommier, V., & Sahal-Br  chot, S. 1978, *A&A*, 69, 57
- D  moulin, P. 1989, in *New Perspectives on Solar Prominences*, IAU Colloq. 167, ed. D. Webb, D. Rust, & B. Schmieder, ASP Conf. Ser., 150, 78
- Landi Degl’Innocenti, E. 1982, *Sol. Phys.*, 79, 291
- Landi Degl’Innocenti, E. 1990, in *Dynamics of Quiescent Prominences*, IAU Colloq. 117, ed. V. Ruzdjak, & E. Tandberg-Hanssen (Springer)
- Leroy, J. L. 1977, *A&A*, 60, 79
- Leroy, J. L. 1978, *A&A*, 64, 247
- Leroy, J. L. 1989, in *Dynamics and Structure of Quiescent Solar Prominences*, ed. E. R. Priest (Kluwer)
- Leroy, J. L., Bommier, V., & Sahal-Br  chot, S. 1983, *Sol. Phys.*, 83, 135
- Leroy, J. L., Bommier, V., & Sahal-Br  chot, S. 1984, *A&A*, 131, 33
- Leroy, J. L., Ratier, G., & Bommier, V. 1977, *A&A*, 54, 811
- Lin, H., Penn, M. J., & Kuhn, J. R. 1998, *ApJ*, 493, 978
- L  pez Ariste, A., Rayrole, J., & Semel, M. 2000, *A&AS*, 142, 137
- Paletou, F., & Molodij, G. 2001, in *Advanced Solar Polarimetry*, ed. M. Sigwarth, ASP Conf. Ser., 236, in press
- Querfeld, C. W., Smartt, R. N., Bommier, V., Landi Degl’Innocenti, E., & House, L. L. 1985, *Sol. Phys.*, 96, 277
- Sahal-Br  chot, S., Bommier, V., & Leroy, J. L. 1977, *A&A*, 59, 223
- Tandberg-Hanssen, E. 1974, *Solar Prominences* (Reidel)
- Tandberg-Hanssen, E. 1995, *The Nature of Solar Prominences* (Kluwer)
- Trujillo Bueno, J., Collados, M., Paletou, F., & Molodij, G. 2001, in *Advanced Solar Polarimetry*, ed. M. Sigwarth, ASP Conf. Ser., 236, in press