

Turbulent viscosity in clumpy accretion disks

Application to the Galaxy

B. Vollmer and T. Beckert

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany

Received 29 May 2001/ Accepted 16 November 2001

Abstract. The equilibrium state of a turbulent clumpy gas disk is analytically investigated. The disk consists of distinct self-gravitating clouds. Gravitational cloud–cloud interactions transfer energy over spatial scales and produce a viscosity, which allows mass accretion in the gas disk. Turbulence is assumed to be generated by instabilities involving self-gravitation and to be maintained by the energy input from differential rotation and mass transfer. Disk parameters, global filling factors, molecular fractions, and star formation rates are derived. The application of our model to the Galaxy shows good agreement with observations. They are consistent with the scenario where turbulence generated and maintained by gravitation can account for the viscosity in the gas disk of spiral galaxies. The rôle of the galaxy mass for the morphological classification of spiral galaxies is investigated.

Key words. ISM: clouds – ISM: structure – Galaxy: structure – galaxies: ISM

1. Introduction

Galactic gas disks are not continuous but clumpy. The structure of the interstellar medium (ISM) is usually hierarchical (Scalo 1985) over length scales of several magnitudes up to ~ 100 pc. The clouds are not uniform nor isolated and their boundaries are often of fractal nature (Elmegreen & Falgarone 1996). Whereas the atomic gas (HI) is mainly in the form of filaments, the molecular gas is highly clumped. The largest self-gravitating molecular clouds (giant molecular clouds = GMC) have sizes of ~ 30 pc and masses of $\sim 10^5 M_{\odot}$ (see e.g. Larson 1981). The GMCs have a volume filling factor $\phi_V \sim 10^{-4}$, with the ratio of the diameter of a typical GMC to the vertical scale height of the GMC distribution ($=130$ pc, Sanders et al. 1985) ~ 0.4 . In this respect, the GMC distribution resembles more a planetary ring (Goldreich & Tremaine 1987). The mass–size relation of the GMCs is $M \propto L^D$. The fractal dimension of a volume fractal is $D \sim 2.3$ (Elmegreen & Falgarone 1996). The origin of this dimension could be turbulent diffusion in an incompressible fluid with a Kolmogorov velocity spectrum ($D \sim 2 + \xi$ for $\Delta v \propto L^{\xi}$; see Meneveau & Sreenivasan 1990).

The Kolmogorov theory applies for fully developed subsonic incompressible fluids. However, the ISM is supersonic and compressible. Only recently, 3D numerical

studies of magneto-hydrodynamical and hydrodynamical turbulence in an isothermal, compressible, and self-gravitating fluid indicated that the energy spectrum of supersonic compressible turbulence follows a Kolmogorov law (Mac Low 1999; Klessen et al. 2000; Mac Low & Ossenkopf 2000). The molecular clouds themselves are stabilized against gravitational collapse by the turbulent velocity field within them (see e.g. Larson 1981).

Wada & Norman (1999) used high-resolution, 2D, hydrodynamical simulations to investigate the evolution of a self-gravitating multiphase interstellar medium in a galactic disk. They found that a gravitationally and thermally unstable disk evolves towards a globally quasi-stationary state where the disk is characterized by clumpy and filamentary structures. The energy source of the turbulence in this system originate in the shear driven by galactic rotation and self-gravitational energy of the gas. The effective Q parameter of the disk was found to have a value between 2 and 5. Without feedback the energy spectrum $E(k) \propto k^{-3}$ corresponds to a Kolmogorov law in two dimensions, but changes into $E(k) \propto k^{-2}$ if stellar energy feedback is included (Wada & Norman 2001). This power law is expected if shocks dominate the system (Passot et al. 1988). Furthermore, Wada & Norman (2001) derived a driving length scale of ~ 200 pc for the model without stellar energy feedback.

While the turbulent nature of the ISM is well established now, its origin and maintenance is still a matter

Send offprint requests to: B. Vollmer,
e-mail: bvollmer@mpifr-bonn.mpg.de

of debate. This is of great importance, because turbulence can provide angular momentum transport in the disk of spiral galaxies. The evolution and the structure of an accretion disk depends entirely on the effective viscosity caused by turbulence.

Two ingredients are necessary for a steady state turbulence: (i) a dynamical instability to generate and (ii) a steady energy input to maintain turbulence.

(i) Five possible instabilities were put forward by different authors:

- viscous instability (Lightman & Eardley 1974);
- thermal instability (Pringle et al. 1973);
- Parker instability (Parker 1966);
- magneto-hydrodynamic instability (Balbus & Hawley 1991);
- gravitational instability (Toomre 1964).

Because self-gravity is always present in the ISM, the gravitational instability can operate in conjunction with all other forms of instabilities. As soon as the density rises due to such a combined instability and increases over the value of the ambient medium by a factor 2, self-gravitation should be the dominant force (Elmegreen 1982). Moreover, the gravitational instability can produce a whole hierarchy of clumped structures inside the largest ones. Therefore, we will only discuss gravitational instability.

(ii) Two main mechanisms are proposed to provide the energy input in order to maintain the observed turbulence of $v_{\text{turb}} \sim 10 \text{ km s}^{-1}$:

- supernova explosions (see e.g. Spitzer 1968; McKee & Ostriker 1977);
- galactic differential rotation (Fleck 1981).

The supernova shocks, which can accelerate low mass clouds, are extremely ineffective in accelerating GMCs, because of the much larger mass to area ratio of GMCs. Jog & Ostriker (1988) applying directly the results of McKee & Ostriker (1977) found that supernova shocks can provide no more than 10% of the kinetic energy of the GMCs (however, Cui et al. (1996) suggested that uncertain background corrections could make this fraction larger). Therefore, we will only focus on energy input by galactic rotation.

The effective viscosity ν_{eff} due to turbulence can be expressed in general as

$$\nu_{\text{eff}} = \xi v_{\text{turb}} l_{\text{driv}}, \quad (1)$$

where v_{turb} is the turbulent velocity on large scales, l_{driv} the driving wavelength, and ξ a constant. For continuous disks the most widely used form is the so-called α -ansatz introduced by Shakura (1972) and Shakura & Sunyaev (1973). They assumed that the turbulent velocity is limited by the sound velocity ($v_{\text{turb}} = c_s$), the turbulent length scale equals the disk scale height $l_{\text{driv}} = H$, and $\xi = \alpha < 1$. Lin & Pringle (1987a) derived a prescription

for the effective viscosity due to gravitational instability: $l_{\text{driv}} = H$ and $\xi = Q^{-2}$, where Q is the Toomre parameter (Toomre 1964). Recently, Duschl et al. (2000) suggested a prescription for the viscosity due to hydrodynamically driven turbulence at the critical effective Reynolds number Re_{crit} , where $\xi = 1$, $v_{\text{turb}} = Re_{\text{crit}}^{-\frac{1}{2}} v_{\text{rot}}$, and the driving wavelength is a well defined fraction of the local radius in the disk $l_{\text{driv}} = Re_{\text{crit}}^{-\frac{1}{2}} R$.

For clumpy accretion disks, Goldreich & Tremaine (1978) and Stewart & Kaula (1980) elaborated models where the shear viscosity in a rotating disk is due to cloud–cloud interactions. Their prescription has the form:

$$\nu_{\text{eff}} = \frac{v_{\text{turb}}^2}{\Omega} \frac{\tau}{\tau^2 + 1} \simeq \tau^{-1} \frac{v_{\text{turb}}^2}{\Omega} \quad \text{for } \tau \gg 1, \quad (2)$$

where Ω is the rotational angular velocity and $\tau = \Omega t_{\text{int}}$ is the number of cloud interactions per rotation period with the interaction time scale t_{int} between the clouds. Vertical hydrostatic equilibrium $R/H \sim v_{\text{rot}}/v_{\text{turb}}$ gives

$$\nu_{\text{eff}} = \tau^{-1} v_{\text{turb}} H, \quad (3)$$

where $v_{\text{rot}} = \Omega R$ is the rotation velocity. In this article we derive a prescription for the turbulent viscosity of comparable form with $\xi = Re^{-1}$ in terms of the turbulent Reynolds number Re . All disk parameters are given as functions of Re , Q , and the mass accretion rate \dot{M} . We apply our model to the Galaxy and derive a Galactic mass accretion rate.

2. The basic picture

We consider a gaseous accretion disk in a given gravitational potential Φ which gives rise to an angular velocity $\Omega = \sqrt{R^{-1} \frac{d\Phi}{dR}}$. The Toomre parameter (Toomre 1964) is treated as a free parameter

$$Q = \frac{v_{\text{turb}} \kappa}{\pi G \Sigma}, \quad (4)$$

with the restriction $Q \geq 1$, where G is the gravitational constant, Σ the average gas column density, and κ the local epicyclic frequency. The disk consists of distinct self-gravitating clouds, which are orbiting in an external gravitational field and have a velocity dispersion Δv . These clouds might be embedded in a low density medium as long as $\Delta v/c_s^{\text{ext}} \leq 1$, where c_s^{ext} is the sound speed of the external medium. This is the case for the warm atomic or ionized phase ($T \sim 8000 \text{ K}$) and the hot ionized phase ($T \sim 10^6 \text{ K}$). The disk scale height is determined uniquely by the turbulent pressure $p_{\text{turb}} = \rho v_{\text{turb}}^2$, where ρ is the averaged density in the disk. The clouds have a small volume filling factor $\phi_V \ll 1$. The transport of angular momentum is due to cloud–cloud interactions. These can be gravitational (elastic) encounters or direct (inelastic) collisions. Thus, gravitational scattering of the massive clouds off each other in a differentially rotating galactic disk gives rise to an effective “gravitational” viscosity. Jog & Ostriker (1988) determined analytically the one-dimensional velocity dispersion of such a system to

be $\Delta v_{1D} = 5\text{--}7 \text{ km s}^{-1}$ which is in good agreement with observations. We assume that energy is dissipated in the disk when clouds become self-gravitating. Turbulence is generated by instabilities involving self-gravitation and is maintained by the energy input, which is provided by differential rotation via cloud–cloud interactions. Thus, the dissipative scale length l_{diss} is equivalent to the size of the largest self-gravitating clouds. Since we assume clouds to form from local instabilities due to self-gravitation, which do not depend on the distance to the galaxy center, $Q = \text{const.}$ Furthermore, we assume that the turbulent Reynolds number Re is also independent of the galactic radius.

3. The equations

3.1. The viscosity prescription

In the inertial range of a turbulent medium kinetic energy is transferred from large scale structures to small scale structures practically without losing energy. So there is a constant energy flux from large scales to small scales where the energy is finally dissipated. Since the velocity dispersion ($\Delta v \sim 10 \text{ km s}^{-1}$) within the disk is more important than the shear, there is no preferred transfer direction. The turbulence is therefore assumed to be isotropic. In this case the similarity theory of Kolmogorov applies (see e.g. Landau & Lifschitz 1959). The assumption of a universal Kolmogorov equilibrium implies that the kinetic energy spectrum of the turbulence depends only on the energy dissipation rate per unit mass ϵ and the characteristic size of the turbulent eddy $l \simeq \frac{1}{k}$, where k is the wave number. The kinetic energy $E(k)$ is related to the mean kinetic energy in the following way:

$$\frac{1}{2} \langle \mathbf{u}(x)^2 \rangle = \int_0^{+\infty} E(k) dk, \quad (5)$$

where \mathbf{u} is the velocity of the medium. Kolmogorov's theory yields

$$E(k) = C \epsilon^{\frac{2}{3}} k^{-\frac{5}{3}}, \quad (6)$$

where C is a constant of order unity. Considering a schematic energy spectrum given by $E(k) = 0$, for $k < k_{\text{driv}}$ and for $k > k_{\text{diss}}$ and by Eq. (6) for $k_{\text{driv}} < k < k_{\text{diss}}$, one derives an expression for the dissipative length scale $l_{\text{diss}} \simeq k_{\text{diss}}^{-1}$ and the large scale velocity v_{turb} :

$$l_{\text{diss}} = (\nu^3 / \epsilon)^{\frac{1}{4}}, \quad (7)$$

where ν is the large scale viscosity due to turbulence;

$$v_{\text{turb}}^2 = \langle \mathbf{u}^2 \rangle \simeq \epsilon^{\frac{2}{3}} k_{\text{driv}}^{-\frac{2}{3}}. \quad (8)$$

This leads to a relation between the two length scales and the turbulent Reynolds number, which is defined by $Re \equiv v_{\text{turb}} \cdot l_{\text{driv}} / \nu$:

$$l_{\text{driv}} \simeq Re^{\frac{3}{4}} l_{\text{diss}}. \quad (9)$$

This turbulent Reynolds number is not equivalent to the macroscopic Reynolds number defined by e.g. Frank et al. (1992)

$$Re_{\text{macro}} = \frac{R v_{\text{rot}}}{\nu}. \quad (10)$$

We will show in Sect. 7 that $Re_{\text{macro}} \gg Re$. In this work we will use the following viscosity prescription:

$$\nu = \frac{1}{Re} v_{\text{turb}} l_{\text{driv}}. \quad (11)$$

In our model the Reynolds number Re defined by Eq. (9) is a free parameter with the restriction $Re \geq 1$. This viscosity prescription will be compared to others in Sect. 5.

It is formally equivalent to the expression for a clumpy accretion disk (Ozernoy et al. 1998) where the viscosity is due to cloud–cloud interactions (Eq. (2)).

We interpret this equivalence as two different pictures for a turbulent self-gravitating medium. We prefer the point of view where the whole ISM (all phases) is taken as one turbulent gas which change phases (i) on turbulent time scales $t_{\text{turb}} = l_{\text{turb}} / v_{\text{turb}}$ and (ii) due to external processes (energy output by stars, i.e. supernovae, UV radiation by O/B stars). Only near the midplane of the disk can the gas become molecular and self-gravitating which leads to a maximum GMC lifetime of $\Delta t \sim H_{\text{mol}} / v_{\text{turb}} \sim 10^7 \text{ yr}$, where ($H_{\text{mol}} \sim 100 \text{ pc}$ Sanders et al. 1985). On the other hand the lifetime of a molecular cloud is approximately given by the crossing time $\Delta t = d_{\text{cl}} / v_{\text{turb}}$, where d_{cl} is the cloud size. With a cloud size of 30 pc this results in $\Delta t \sim 5 \times 10^6 \text{ yr}$. These values are comparable to the lifetimes of star-forming molecular clouds given by Blitz & Shu (1980). Since these lifetimes are about 10 times smaller than the expected mean collision time ($\Delta t_{\text{coll}} \sim 2 \times 10^8 \text{ yr}$ Elmegreen 1987), direct cloud–cloud collisions are very rare and are not important for the effective viscosity.

Even with the limited lifetime of molecular clouds, gravitational interactions always take place between the continuously appearing and disappearing clouds. These interactions give rise to a collision term in the Boltzmann equation with an average collision time for gravitational encounters

$$t_{\text{G}} = \left\langle \frac{3v_{\text{turb}}^3}{4\sqrt{\pi}G^2 m_{\text{cl}}^2 n_{\text{cl}}} \right\rangle, \quad (12)$$

(Braginskii 1965) where m_{cl} is the cloud mass and n_{cl} is the spatial number density of the clouds. For molecular clouds the turbulent velocity is approximately independent of the cloud mass ($v_{\text{turb}} \propto m_{\text{cl}}^{0.2}$ Larson 1981), thus

$$t_{\text{G}} = \frac{3v_{\text{turb}}^3}{4\sqrt{\pi}G^2 \langle m_{\text{cl}}^2 n_{\text{cl}} \rangle}, \quad \text{with} \quad (13)$$

$$\langle m_{\text{cl}}^2 n_{\text{cl}} \rangle = \int m_{\text{cl}}^2 n(m_{\text{cl}}) dm_{\text{cl}} =$$

$$6.6 \times 10^{-6} M_{\odot}^{0.5} \text{ pc}^{-3} \int m_{\text{cl}}^{0.5} dm_{\text{cl}}, \quad (14)$$

where we have used the differential number density of molecular clouds given by Elmegreen (1987). Using integration boundaries $m_1 = 10^4 M_\odot$, $m_2 = 10^6 M_\odot$, and $v_{\text{turb}} = 10 \text{ km s}^{-1}$ one obtains $t_G \sim 4 \times 10^9 \text{ yr}$. Assuming a galactic radius $R = 10 \text{ kpc}$ and a rotation velocity of $v_{\text{rot}} = 200 \text{ km s}^{-1}$ gives $\tau = Re \sim 80$.

Since the dissipative scale length is of the order of the largest self-gravitating clouds $l_{\text{diss}} \sim d_{\text{cl}} \sim 30 \text{ pc}$, we can calculate the value of the driving wavelength $l_{\text{driv}} = Re^{\frac{3}{4}} l_{\text{diss}} \simeq 800 \text{ pc}$. This is of the same order as the thickness of the distribution of the atomic gas (Kulkarni & Heiles 1987; Dickey 1993). We will show in Sect. 4 that in our model $l_{\text{driv}} \sim H$. The scale height of the molecular disk H_{mol} is smaller, because of the limited lifetime of molecular clouds.

3.2. Non Kolomogorov energy spectra

Wada & Norman (2001) found an energy spectrum of the form $E(k) \propto k^{-2}$ in their 2D hydrodynamical simulations of a supersonic, compressible turbulence including stellar energy feedback. This power law corresponds to that of a shock dominated medium. In contrast to a compression dominated medium, the exponent of the power law for an incompressible medium is the same in two and three dimensions (Passot et al. 1988). In this section we assume that $E(k) \propto k^{-2}$ in three dimensions. In this case, an alternative explanation for this exponent comes from the theory of intermittency (Frisch 1995). The open parameter of this model is the dimension D . In the case of intermittent turbulence the energy flux from scales $\sim l$ to smaller scales is

$$\epsilon = \frac{v_l^3}{l} \left(\frac{l}{l_{\text{driv}}} \right)^{3-D} = \text{const.} \quad (15)$$

The local spatial average of the energy dissipation can then be expressed as

$$\epsilon_l = \frac{v_l^3}{l} = \epsilon \left(\frac{l}{l_{\text{driv}}} \right)^{D-3}. \quad (16)$$

Within this framework $E(k) \propto k^{-2}$ implies $D = 2$ and consequently $\epsilon_l = \epsilon \left(\frac{l}{l_{\text{driv}}} \right)^{-1}$, i.e. the local spatial average of the energy dissipation decreases with decreasing length scale l . The ratio between the driving and the dissipation length scale is then

$$\frac{l_{\text{driv}}}{l_{\text{diss}}} = Re, \quad (17)$$

instead of $Re^{\frac{3}{4}}$ in the case of Kolmogorov turbulence. For a Reynolds number $Re \sim 50$ the difference is about a factor 2.5. Since the power law exponent of the ISM energy spectrum is not yet well established, both possibilities must be considered.

3.3. Angular momentum equation

In a steady state accretion disk, the mass accretion rate is $\dot{M} = 2\pi R \Sigma (-v_{\text{rad}})$,

where v_{rad} is the radial velocity. The angular momentum equation can be integrated giving

$$\nu \Sigma = -\frac{\dot{M}}{2\pi R} \Omega \left(\frac{\partial \Omega}{\partial R} \right)^{-1}. \quad (19)$$

Furthermore, we use

$$\Sigma = \rho H \quad (20)$$

for the surface density of the disk, where ρ is the mass density at the midplane.

3.4. Energy flux conservation

The ISM turbulence in the disk is initiated by an instability involving self-gravitation. The transport of angular momentum is due to cloud–cloud interactions giving rise to radial mass accretion. We assume that the necessary energy input to maintain the turbulence is the gravitational energy, which is gained when the ISM is accreted to smaller Galactic radii, i.e. the energy input is supplied by the Galactic differential rotation. Kinetic energy of the Galactic rotation is transferred to the turbulent cascade at the driving wavelength l_{driv} and reaches the dissipative length scale l_{diss} without losing energy. The energy per unit time which is transferred by turbulence is

$$\dot{E} = \rho \nu \int \nabla (v_{\text{turb}}^2) d\mathbf{f} \simeq \rho \nu \frac{v_{\text{turb}}^2}{l_{\text{driv}}} A, \quad (21)$$

where the integration is over the area $\int d\mathbf{f} = A$. Thus, the energy flux per unit time and unit area is

$$\frac{\Delta E}{\Delta t \Delta A} = \rho \nu \frac{v_{\text{turb}}^2}{l_{\text{driv}}}. \quad (22)$$

From standard fluid dynamics, the viscosity ν generates dissipation in the disk at a rate

$$\frac{\Delta E}{\Delta t \Delta A} = \nu \Sigma \left(R \frac{\partial \Omega}{\partial R} \right)^2 = -\frac{1}{2\pi} \dot{M} v_{\text{rot}} \frac{\partial \Omega}{\partial R} \quad (23)$$

(see e.g. Pringle 1981), where $v_{\text{rot}} = \Omega R$. This is valid for $R \gg R_*$, where $(\partial \Omega / \partial R)_{R=R_*} = 0$. Conservation of energy flux leads thus to

$$\rho \nu \frac{v_{\text{turb}}^2}{l_{\text{driv}}} = -\frac{1}{2\pi} \dot{M} v_{\text{rot}} \frac{\partial \Omega}{\partial R} \quad (24)$$

and relates the mass accretion rate to the turbulence in the disk.

3.5. The vertical pressure equilibrium

Since we assume that the sound velocity of the ambient medium is smaller than the turbulent velocity dispersion of the clouds, the only pressure which counterbalances gravitation in the vertical direction is the turbulent

pressure $p_{\text{turb}} = \rho v_{\text{turb}}^2$. We distinguish three cases for the gravitational force in the vertical z direction:

1. $M_{\text{d}}(R) \leq 0.5 (H/R) M$, where M_{d} is the disk mass and M is the central mass (dominating central mass);
2. $\rho \ll \rho_*$ and $M_{\text{d}}(R) \ll M_*(R)$, where $\rho_*/M_*(R)$ is the stellar central density/disk mass within a radius R (dominating stellar disk mass);
3. $\rho \geq \rho_*$ and $0.5 (H/R) M(R) < M_{\text{d}}(R) < M(R)$, where $M(R)$ is the total mass enclosed within a radius R (self-gravitating gas disk in z direction).

For these cases the gravitational pressure p_{grav} has the following forms:

1. $p_{\text{grav}} = \rho \Omega^2 H^2$ (see e.g. Pringle 1981);
2. $p_{\text{grav}} = \pi G \Sigma_* \Sigma$, where Σ_* is the stellar surface density of the disk (see e.g. Binney & Tremaine 1987);
3. $p_{\text{grav}} = \pi G \Sigma^2$ (Paczynski 1978).

The equilibrium in the vertical direction is given by

$$p_{\text{grav}} = p_{\text{turb}}. \quad (25)$$

3.6. Global gravitational stability in z direction

The basic principles underlying the gravitational instability of a thin rotating disk can be found in Toomre (1964). A gaseous disk is locally stable to axisymmetric perturbations, if

$$Q = \frac{v_{\text{turb}} \kappa}{\pi G \Sigma} > 1, \quad (26)$$

where $\kappa = \sqrt{R \frac{d\Omega^2}{dR} + 4\Omega^2}$. Since in general $\Omega \leq \kappa \leq 2\Omega$, we will use the following equation:

$$Q = \frac{v_{\text{turb}} \Omega}{\pi G \Sigma}. \quad (27)$$

Multiplying the numerator and the denominator of the right hand side by R^2 gives

$$Q = \frac{v_{\text{turb}} M_{\text{tot}}}{v_{\text{rot}} M_{\text{gas}}}, \quad (28)$$

where $M_{\text{tot/gas}}$ is the total enclosed mass and the total enclosed gas mass at radius R . Thus for a given velocity dispersion Q^{-1} is proportional to the ratio of gas mass to total mass.

4. Results

We have solved the set of equations Eq. (11), Eq. (19), Eq. (20), Eq. (24), Eq. (25), and Eq. (27), for the three cases:

1. dominating central mass;
2. dominating stellar disk mass;
3. self-gravitating gas disk in z direction.

We will use $\Omega' = \partial\Omega/\partial R$ with $\Omega' < 0$ in the region of interest.

4.1. Dominating central mass

Pressure equilibrium (Eq. (25)) leads to

$$H = v_{\text{turb}} \Omega^{-1}, \quad (29)$$

which is equivalent to the hydrostatic equilibrium where the sound speed is replaced by the turbulent velocity dispersion.

From energy flux conservation (Eq. (24)) and the angular momentum equation (Eq. (19)) it follows

$$H = \left(R \frac{\Omega'}{\Omega} \right)^2 l_{\text{driv}}. \quad (30)$$

Thus, the driving wavelength is proportional to the disk height $l_{\text{driv}} \propto H$ and the viscosity prescription reads $\nu \sim Re^{-1} v_{\text{turb}} H$, which is formally equivalent to the viscosity prescription for clumpy disks of Goldreich & Tremaine (1978), Stewart & Kaula (1980) (see Eq. (3)), and the α -viscosity (Shakura & Sunyaev 1973) with $v_{\text{turb}} = c_s$ and $Re^{-1} = \alpha$.

Inserting Eqs. (29) in (27) and using $\rho = \Sigma/H$ gives

$$\rho = \frac{1}{\pi Q} \frac{\Omega^2}{G}. \quad (31)$$

If we define the critical cloud density for stability against tidal shear $\rho_{\text{crit}} = \pi^{-1} \Omega^2 G^{-1}$, the Toomre parameter Q represents the ratio between the critical density and the volume averaged density in the disk $Q = \rho_{\text{crit}}/\rho$.

Inserting the viscosity prescription (Eq. (11)) into the angular momentum equation (Eq. (19)) leads to

$$H = \left(\frac{1}{2} G Re Q \dot{M} R (-\Omega') \Omega^{-4} \right)^{\frac{1}{3}}. \quad (32)$$

With $\Sigma = \rho/H$ we obtain

$$\Sigma = \rho H = \frac{1}{2^{\frac{2}{3}} \pi} G^{-\frac{2}{3}} Re^{\frac{1}{3}} Q^{-\frac{2}{3}} \dot{M}^{\frac{1}{3}} R^{\frac{1}{3}} (-\Omega')^{\frac{1}{3}} \Omega^{\frac{2}{3}}. \quad (33)$$

Inserting Eqs. (33) into (19) gives

$$\nu = \frac{1}{2^{\frac{2}{3}}} G^{\frac{2}{3}} Re^{-\frac{1}{3}} Q^{\frac{2}{3}} \dot{M}^{\frac{2}{3}} R^{-\frac{4}{3}} (-\Omega')^{-\frac{4}{3}} \Omega^{\frac{1}{3}}. \quad (34)$$

The turbulent velocity can be calculated using the pressure equilibrium (Eq. (25)):

$$v_{\text{turb}} = H \Omega = \left(\frac{1}{2} G Re Q \dot{M} R (-\Omega') \Omega^{-1} \right)^{\frac{1}{3}}. \quad (35)$$

If $\Omega' \simeq -\Omega/R$, the viscosity is inversely proportional to the angular velocity $\nu \propto \Omega^{-1}$ and the turbulent velocity is constant. In the case of a Keplerian velocity field due to a point mass $\nu \propto R^{3/2}$. The behaviour of the disk parameters (v_{turb} , l_{driv} , Σ , H , ρ , and ν) are shown for a constant rotation curve (solid line) and a rising rotation curve $v_{\text{rot}} \propto \sqrt{R}$ (dashed line) in Fig. 1. The free parameters $Q = 1$, $Re = 50$, and $\dot{M} = 10^{-2} M_{\odot} \text{ yr}^{-1}$ were chosen such that the disk parameters fit the observed values for the Galaxy (see Sect. 7).




Fig. 1. Disk parameters for the case of a dominating central mass. Solid lines: constant rotation curve. Dashed line: rising rotation curve $v_{\text{rot}} \propto \sqrt{R}$. If $Q = 1$ the behaviour of these parameters is the same in the case of a selfgravitating gas disk in z direction.

4.2. Dominating stellar disk

Pressure equilibrium (Eq. (25)) together with $v_{\text{rot}}^2 = \pi G \Sigma_* R$ leads to

$$\frac{H}{R} = \left(\frac{v_{\text{turb}}}{v_{\text{rot}}} \right)^2. \quad (36)$$

Inserting Eq. (36) in the equation for the Toomre parameter (Eq. (27)) and using the viscosity prescription (Eq. (11)), the angular momentum equation (Eq. (19)) and energy flux conservation (Eq. (24)) gives

$$v_{\text{turb}} = \left(\frac{1}{2} G Q Re \dot{M} (-\Omega') \Omega^{-2} \right)^{\frac{1}{2}}. \quad (37)$$

In the case of $\Omega' \simeq -\Omega/R$, the turbulent velocity is inversely proportional to the square root of the rotation velocity $v_{\text{turb}} \propto v_{\text{rot}}^{-1/2}$.

Inserting Eqs. (37) into (36) gives the disk scale height

$$H = \frac{1}{2} G Re Q \dot{M} R^{-1} (-\Omega') \Omega^{-4}. \quad (38)$$

Equations (11), (19), and (24) together with Eq. (38) lead to the following expression for the driving wavelength

$$l_{\text{driv}} = (-\Omega')^{-2} \Omega^2 R^{-1}. \quad (39)$$

From the same equations the density, column density, and viscosity read

$$\rho = \frac{\sqrt{2}}{\pi} G^{-\frac{3}{2}} Q^{-\frac{3}{2}} Re^{-\frac{1}{2}} \dot{M}^{-\frac{1}{2}} R (-\Omega')^{-\frac{1}{2}} \Omega^4, \quad (40)$$

$$\Sigma = \frac{1}{\sqrt{2}\pi} G^{-\frac{1}{2}} Q^{-\frac{1}{2}} Re^{\frac{1}{2}} \dot{M}^{\frac{1}{2}} (-\Omega')^{\frac{1}{2}}, \quad (41)$$

and

$$\nu = \sqrt{2} G^{\frac{1}{2}} Q^{\frac{1}{2}} Re^{-\frac{1}{2}} \dot{M}^{\frac{1}{2}} R^{-1} (-\Omega')^{-\frac{3}{2}} \Omega. \quad (42)$$

If $\Omega' \simeq -\Omega/R$, the driving wavelength is proportional to the galactic radius $l_{\text{driv}} \propto R$. Therefore, in the case of a dominating stellar disk, one obtains for the viscosity prescription (Eq. (11)) $\nu \sim Re^{-1} v_{\text{turb}} R$. In this case, the viscosity is proportional to the galactic radius divided by the




Fig. 2. Disk parameters for the case of a dominating stellar disk. Solid lines: constant rotation curve. Dashed line: rising rotation curve $v_{\text{rot}} \propto \sqrt{R}$.

square of the rotation velocity $\nu \propto R v_{\text{rot}}^{-1/2}$. Furthermore, if the rotation curve is flat, i.e. $v_{\text{rot}} = \text{const.}$, $\nu \propto R$. This behaviour can be observed in Fig. 2 where the disk parameters (v_{turb} , l_{driv} , Σ , H , ρ , and ν) are shown for a constant rotation curve (solid line) and a rising rotation curve $v_{\text{rot}} \propto \sqrt{R}$ (dashed line) for $Q = 1000$, $Re = 50$, and $\dot{M} = 10^{-2} M_{\odot} \text{ yr}^{-1}$. Re , and \dot{M} are the same as for the Galaxy (see Sect. 7) and the value of Q is chosen such that the turbulent velocity dispersion corresponds to that measured of an S0 galaxy ($\sim 50 \text{ km s}^{-1}$ D’Onofrio et al. 1999).

4.3. Self-gravitating gas in z direction

The pressure equilibrium (Eq. (25)) together with the equation for the Toomre parameter (Eq. (27)) leads to

$$\rho = \frac{1}{\pi Q^2} \frac{\Omega^2}{G}. \quad (43)$$

The square of the Toomre parameter thus represents the fraction between the critical density with respect to tidal

shear and the volume averaged density in the disk $Q^2 = \rho_{\text{crit}}/\rho$. We want to recall here that for a dominating point mass $Q = \rho_{\text{crit}}/\rho$.

Inserting Eq. (43) into the pressure equilibrium equation (Eq. (25)) gives

$$H = Q \Omega^{-1} v_{\text{turb}}. \quad (44)$$

Then, using the angular momentum equation (Eq. (19)), it follows

$$l_{\text{driv}} = \frac{1}{2} Re G Q \dot{M} v_{\text{turb}}^{-2} R^{-1} (-\Omega')^{-1}. \quad (45)$$

From the energy flux conservation (Eq. (24)) one obtains

$$v_{\text{turb}} = \left(\frac{1}{2} G Q^2 Re \dot{M} R (-\Omega') \Omega^{-1} \right)^{\frac{1}{3}}. \quad (46)$$

For $\Omega' \simeq -\Omega/R$, the turbulent velocity is constant. This is what is observed in the nearby galaxy NGC 3938 (van der Kruit & Shostak 1982).

Inserting this expression in Eq. (45) leads to

$$\frac{H}{l_{\text{driv}}} = Q^2 R^2 (-\Omega')^2 \Omega^{-2}. \quad (47)$$

For $\Omega' \simeq -\Omega/R$, $H/l_{\text{driv}} \simeq Q^2$. Thus, the viscosity prescription yields $\nu = Re^{-1} Q^{-2} v_{\text{turb}} H$. Here, Re is not the macroscopic Reynolds number $Re_{\text{macro}} = v_{\text{rot}} R/\nu$ (see Sect. 7). In the case of $Re = 1$ this is the prescription given by Lin & Pringle (1987a).

Inserting Eqs. (46) into (44) leads to the disk height

$$H = \frac{1}{2^{\frac{1}{3}}} Q^{\frac{5}{3}} Re^{\frac{1}{3}} G^{\frac{1}{3}} \dot{M}^{\frac{1}{3}} R^{\frac{1}{3}} (-\Omega')^{\frac{1}{3}} \Omega^{-\frac{4}{3}}. \quad (48)$$

With $\Sigma = \rho/H$ one obtains the following expression for the column density:

$$\Sigma = \frac{1}{2^{\frac{1}{3}} \pi} Q^{-\frac{1}{3}} Re^{\frac{1}{3}} G^{-\frac{2}{3}} \dot{M}^{\frac{1}{3}} R^{\frac{1}{3}} (-\Omega')^{\frac{1}{3}} \Omega^{\frac{2}{3}}. \quad (49)$$

Equation (11), Eqs. (46)–(48) lead to

$$\nu = \frac{1}{2^{\frac{1}{3}}} Q^{\frac{1}{3}} Re^{-\frac{1}{3}} G^{\frac{2}{3}} \dot{M}^{\frac{2}{3}} R^{-\frac{4}{3}} (-\Omega')^{-\frac{4}{3}} \Omega^{\frac{1}{3}}. \quad (50)$$

In the case of $\Omega' \simeq -\Omega/R$, the column density is proportional to and the viscosity is inversely proportional to the angular velocity: $\Sigma \propto \Omega$, $\nu \propto \Omega^{-1}$. If in addition the rotation velocity is constant, then $\nu \propto R$.

Comparing this result with Eq. (42) shows that both models with a gravitational potential Φ due to an extended mass distribution give the same viscosity prescription if the rotation velocity $v_{\text{rot}} = \sqrt{R d\Phi/dR}$ is constant.

On the other hand, the dominating central mass model and the self-gravitating gas disk model have the same analytical solutions for $Q = 1$ (see Fig. 1).

5. Discussion of the results

5.1. Comparison with previous models of turbulent, selfgravitating gas disks

Models of turbulent, self-gravitating gas disks can be divided into two different approaches: (i) The disk is assumed to be quasi continuous and at the edge of fragmentation ($Q \sim 1$); (ii) The disk is already clumpy and the viscosity is due to cloud–cloud interactions ($Q \geq 1$).

We will first discuss the quasi continuous approach. Paczyński (1978) investigated a self-gravitating disk with a polytropic index of $\gamma = \frac{4}{3}$ which corresponds to a radiation pressure dominated disk. Furthermore, he assumed the disk luminosity to be at the Eddington limit. With the acceleration due to a central mass M and the disk gas surface density Σ : $g = GMR^{-3}z + 2\pi G\Sigma$, he obtained $\Sigma \propto R^{-3}$ and thus $\nu \propto R^3$ for a constant rotation curve. Lin & Pringle (1987a) proposed a viscosity prescription based on the Toomre instability criterion (Toomre 1964): $\nu = Q^{-2} H^2 R$. They showed that under certain conditions this prescription allows a similarity solution $\Sigma \propto R^{-\frac{3}{2}}$. Their viscosity prescription can be generalized $\nu \propto \Sigma^\alpha R^{-\beta}$ with $\alpha, \beta \geq 0$ or $\nu \propto \Sigma^\alpha \Omega^{-\beta}$ for

self-gravitating disk in z and R direction (Saio & Yoshii 1990). Shlosman & Begelman (1989) also considered a disk at the edge of selfgravitation, i.e. $Q \sim 1$. They obtained $\Sigma \propto v_{\text{turb}} v_{\text{rot}} R^{-1} \propto R^{-1}$ for constant turbulent and rotation velocities. On the other hand, Duschl et al. (2000) made a completely different ansatz: $\nu \propto \beta v_{\text{rot}} R$, with $\beta \ll 1$. They found $\Sigma \propto R^{-1}$ and $\nu \propto R$ for a disk with $v_{\text{rot}} = \text{const.}$

On the other hand, using a clumpy disk model Silk & Norman (1981) derived a viscosity prescription in assuming that the cooling time for cloud–cloud collisions $t_{\text{cool}} = l_0(v_{\text{turb}}\eta)^{-1}$ equals the viscous time scale $t_\nu = R^2\nu^{-1}$ for all radii R , where η is the fraction of cloud kinetic energy radiated in a collision and l_0 is the cloud mean free path. This leads to $\nu = \nu_0(R/R_0)$, where $\nu_0 = \sqrt{\eta} R_0 v_{\text{turb}}$ and R_0 is a characteristic length scale. Lynden Bell & Pringle (1974) showed that for this viscosity prescription $\Sigma \propto (Rt)^{-1} \exp(-\xi R/t)$, where ξ is a constant and t is time. Later Shlosman & Begelman (1987) also used $t_{\text{coll}} \sim t_{\text{visc}}$. Ozernoy (1998) and Kumar (1999) suggested a viscosity prescription based on the collisional Boltzmann equation (see Sect. 3.1) $\nu = \tau v_{\text{turb}}^2 \Omega$, where $\tau = (t_{\text{coll}} \Omega)^{-1}$.

In this section we will only discuss the self-gravitating gas disk in z direction, because only for this set of equation $Q \sim 1$, i.e. the gravitational instability is active. This point will be further discussed in Sect. 5.2.

Whereas the above cited viscosity prescriptions (except Duschl et al. 2000) use $l_{\text{driv}} = H$, the driving wavelength l_{driv} is a priori a free parameter in our set of equations. It is then defined by the energy flux conservation (Eq. (24)). In terms of the standard accretion disk equations (see e.g. Pringle 1981), the “thermostat” equation, i.e. the viscous heating is radiated by the disk, is replaced by the turbulent energy flux equation (Eq. (24)), where the energy gained through accretion and differential rotation is transported by turbulence to smaller scales where it is dissipated. The addition of this energy flux conservation to the “traditional” set of disk equations leads to $l_{\text{driv}} = H$. Once this relation is established, the viscosity prescription has the form of that derived from the collisional Boltzmann equation with $Re = t_{\text{coll}} \Omega$ (see Sect. 3.1). On the other hand, our viscosity is a factor Re^{-1} smaller than that of Lin & Pringle (1987a) for $Q \sim 1$.

Thus, a big difference to previous models is that we do not use a disk “thermostat”, i.e. that the viscous heating is radiated by the disk. Radiation is not included in our model. Duschl et al. (2000) noticed that the α viscosity together with a pressure term due to self-gravitation leads to a constant sound velocity. They declared this unphysical within the framework of a disk “thermostat”. On the other hand, if one replaces the sound velocity by the turbulent velocity this is exactly what is observed in spiral galaxies. Since $v_{\text{turb}} = \text{const}$ we have $H/R = \text{const.}$, and consequently $\nu \propto R$ and $\Sigma \propto R^{-1}$.

5.2. Why can the Toomre parameter be greater than 1?

Whereas the two approaches of a continuous and a clumpy disk can be unified, this can not be done for a disk with $Q > 1$. In this case the disk is not globally gravitational unstable (Toomre 1964). The turbulent velocity is so high, or the surface density is so low that the disk is globally gravitationally stable. However, we can imagine two possibilities for the formation of $Q > 1$ disks:

(i) In Sect. 7 we will show that the mass accretion rate within the disk is much smaller than the star formation rate. Thus, starting from a $Q \sim 1$ disk with star formation, Q will increase with time (if there is no external mass infall). This could be the case for S0 galaxies.

(ii) The gas which falls into the given gravitational potential is already clumpy. This could be the case for the Circumnuclear Disk in the Galactic Center (Vollmer & Duschl 2000).

For these scenarios the above models for a dominating central mass and a dominating stellar disk are valid. We are the first to solve these sets of equations.

We will now show that $Q = \text{const} > 1$ can be understood in terms of star formation. With the critical density for tidal disruption $\rho_{\text{crit}} = \Omega^2/(\pi G)$ and using $\dot{\Sigma}_* = Re^{-1}\Sigma\Omega$ (Eq. (71)) one obtains:

$$\dot{\rho}_* = \frac{1}{ReQ} \frac{\rho_{\text{crit}}}{t_{\text{H}}}, \quad (51)$$

where $t_{\text{H}} = H/v_{\text{turb}}$ is the vertical crossing time. Thus, $Q = \text{const} > 1$ together with $Re = \text{const} > 1$ means that the star formation rate is proportional to the critical density with respect to tidal shear divided by the vertical crossing time.

The detailed comparison between observations and our model will show if $Q > 1$ models are useful to describe clumpy disks with a low gas mass or high velocity dispersion.

5.3. Energy dissipation due to selfgravitation

If turbulence is dissipated by self-gravitation, the energy dissipation rate due to self-gravitation of the disk in z direction ϵ_{sg} must equal the constant, turbulent energy dissipation rate per unit mass ϵ_{turb} .

The energy due to self-gravitation of a gaseous disk is

$$\Delta E = p \Delta A H = \pi G \Sigma^2 \Delta A H = \pi G \Sigma M_{\text{gas}} H, \quad (52)$$

where ΔA is the disk surface. The growth rate of the gravitational instability in z direction is approximately the free fall time $\Delta t \sim \sqrt{H/(G\Sigma)} = \sqrt{(G\rho)^{-1}}$. Thus, one obtains for the energy dissipation rate per unit mass due to self-gravitation

$$\epsilon_{\text{sg}} = \frac{\Delta E}{\Delta M \Delta t} = \pi G \Sigma H \sqrt{G\rho}. \quad (53)$$

Using Eqs. (43) and (44) it can be shown that

$$\epsilon_{\text{sg}} \sim \epsilon_{\text{turb}} = \frac{v_{\text{diss}}^3}{l_{\text{diss}}} = \frac{v_{\text{turb}}^3}{l_{\text{driv}}} = \frac{v_{\text{turb}}^3}{H}. \quad (54)$$

In order to estimate the mass of a self-gravitating cloud, we use $\Delta E \sim N_{\text{cl}} \frac{3}{5} M_{\text{cl}}^2 G / l_{\text{diss}}$, where N_{cl} is the number of selfgravitating clouds and M_{cl} is the cloud mass. With the free fall time of the clouds $\Delta t \sim \sqrt{(G\rho)^{-1} \Phi_{\text{V}}}$ this leads to

$$\frac{\Delta E}{\Delta M \Delta t} = \frac{3}{5} Re^{-\frac{3}{4}} H^{-1} \sqrt{G\rho \Phi_{\text{V}}^{-1}} = \epsilon_{\text{turb}}, \quad (55)$$

where Φ_{V} is the volume filling factor (see Sect. 6.1) and $\Delta M = N_{\text{cl}} M_{\text{cl}}$. Using the expression for Φ_{V} derived in Sect. 6.1 and Eq. (43) gives

$$M_{\text{cl}} \sim \frac{5\sqrt{\pi}}{3} G^{-1} Re^{-\frac{5}{4}} Q^{-1} \Omega^{-1} v_{\text{turb}}^3. \quad (56)$$

With $Q = 1$, $Re = 50$, $\Omega = 2 \times 10^{-8} \text{ yr}^{-1}$, and $v_{\text{turb}} = 10 \text{ km s}^{-1}$ one obtains $M_{\text{cl}} \sim 2 \times 10^5 M_{\odot}$. This is consistent with the mass of GMCs.

In the present paper we have assumed that the potential energy gained through accretion is dissipated locally. This is true only in the absence of radial energy transport, which can be checked a posteriori. The energy dissipation rate due to self-gravitation of the disk in z direction is

$$\delta_{\text{sg}} = \frac{\Delta E}{\Delta A \Delta t} \sim \frac{\pi G \Sigma^2 H}{t_{\text{H}}}. \quad (57)$$

The radial energy flux is given by

$$\delta_{\text{r}} = \Sigma v_{\text{r}} \frac{\partial e}{\partial R} - \Sigma v_{\text{turb}}^2 R^{-2} \frac{\partial}{\partial R} (R^2 v_{\text{r}}), \quad (58)$$

where $v_{\text{r}} = -\dot{M}/(2\pi R \Sigma)$ is the radial velocity and e the specific internal energy. For supersonic turbulence in clumpy disks e is dominated by the specific kinetic energy v_{turb}^2 . Since $v_{\text{turb}} = \text{const}$, $\partial e / \partial R \sim 0$. With $v_{\text{r}} = \text{const}$ one obtains

$$\delta_{\text{r}} = -2\Sigma v_{\text{turb}}^2 v_{\text{r}} R^{-1} = -2\rho v_{\text{turb}}^2 v_{\text{r}} H R^{-1}. \quad (59)$$

Using $t_{\text{H}} = H/v_{\text{turb}}$ and Eqs. (43), (46), (48), (49) one obtains

$$\frac{\delta_{\text{r}}}{\delta_{\text{sg}}} = \sqrt{\frac{3}{32}} 2^{-\frac{2}{3}} \pi Q^{\frac{4}{3}} Re^{-\frac{1}{3}} G^{\frac{2}{3}} \dot{M}^{\frac{2}{3}} v_{\text{rot}}^{-2}. \quad (60)$$

With $Q = 1$, $Re = 50$, and $\dot{M} = 10^{-2} M_{\odot} \text{ yr}^{-1}$ (see Sect. 7) this leads to

$$\frac{\delta_{\text{r}}}{\delta_{\text{sg}}} \sim \frac{2}{(v_{\text{rot}}[\text{km s}^{-1}])^2} \ll 1. \quad (61)$$

The radial advective energy flux is much smaller than the energy gain by mass accretion and can therefore be neglected as long as Eq. (61) holds.

We can conclude here that the energy flux transported to small scales by turbulence has the same order of magnitude as the energy dissipation rate due to selfgravitation of the gaseous disk. Moreover, the potential energy gained through accretion is dissipated locally at small scales. Since our model fits observations for a reasonable choice of parameters, we conclude that, based on energy flux conservation, it is possible that turbulence is generated and dissipated through gravitational instabilities, and maintained by the energy input due to mass inflow and differential rotation (Sect. 3.4).

6. Implications

6.1. Volume filling factors

In this section we compare the crossing time of a turbulent cloud to the gravitational free fall time in order to derive an expression for the volume filling factor as a function of the Reynolds number Re and the Toomre parameter Q . We assume turbulence with a Kolmogorov spectrum for $l_{\text{diss}} \leq l \leq l_{\text{driv}}$. This implies: $l_{\text{diss}} = Re^{-\frac{3}{4}} l_{\text{driv}}$ and $v_{\text{cl}} = Re^{-\frac{1}{3}} v_{\text{turb}}$, where v_{cl} is the turbulent velocity of a cloud of size l_{diss} .

The characteristic turbulent time scale of clouds whose size is comparable to the dissipative length scale is

$$t_1 = l_{\text{diss}}/v_{\text{cl}}. \quad (62)$$

The gravitational free fall time is given by

$$t_{\text{ff}} = \sqrt{\frac{3\pi}{32G\rho_{\text{cl}}}}, \quad (63)$$

where ρ_{cl} is the density of a single cloud, which is related to the overall disk density ρ by the volume filling factor ϕ_V : $\rho_{\text{cl}} = \phi_V^{-1}\rho$.

The clouds are self gravitating for $t_1 = t_{\text{ff}}$. Inserting the expressions given in Sect. 4 leads to:

1. dominating central mass:

$$\phi_V = \frac{32}{3\pi^2} Q^{-1} Re^{-1} R^{-4} (-\Omega')^{-4} \Omega^4;$$

2. dominating stellar disk mass:

$$\begin{aligned} \phi_V &= \frac{64\sqrt{2}}{3\pi^2} G^{-\frac{3}{2}} Re^{-\frac{5}{2}} Q^{-\frac{5}{2}} \dot{M}^{-\frac{3}{2}} R^{-1} (-\Omega')^{-\frac{11}{2}} \Omega^{10} \\ &= \frac{32}{3\pi^2} Re^{-1} Q^{-1} R^{-4} (-\Omega')^{-4} \Omega^4 \left(\frac{v_{\text{rot}}}{v_{\text{turb}}} \right)^3; \end{aligned}$$

3. self-gravitating gas disk in z direction:

$$\phi_V = \frac{32}{3\pi^2} Q^{-4} Re^{-1} R^{-4} (-\Omega')^{-4} \Omega^4.$$

For a non Kolmogorov spectrum $E(k) \propto k^{-2}$, the volume filling factors have to be multiplied by Re^{-1} : $\Phi_V^{\text{nonK.}} = Re^{-1} \Phi_V^{\text{K.}}$.

6.2. Bondi–Hoyle accretion limit

We will now consider the limit $\phi_V = Q = Re \sim 1$, which means $M_{\text{gas}} \sim M_{\text{tot}}$ and $v_{\text{turb}} \sim v_{\text{rot}}$. This implies $H \sim R$, i.e. this is the limit for a spherical configuration. It is assumed that $\partial\Omega/\partial R \simeq -\Omega/R$. The viscosity prescription (Eq. (11)) together with energy flux conservation equation (Eq. (24)) gives

$$\frac{1}{Re} \rho v_{\text{turb}}^3 = \frac{\dot{M}}{2\pi} v_{\text{rot}} \frac{\Omega}{R}. \quad (64)$$

Setting $v_{\text{turb}} = v_{\text{rot}}$, $Re = 1$ and using $v_{\text{rot}}^2 = MG/R$ leads to

$$\dot{M} = \frac{2\pi G^2 M^2 \rho}{v_{\text{rot}}^3}. \quad (65)$$

This is equivalent to the Bondi–Hoyle accretion rate (Bondi & Hoyle 1944), which is given by

$$\dot{M}_{\text{B-H}} = \frac{2.5\pi G^2 M^2 \rho}{v_{\text{turb}}^3} \quad (66)$$

for $v_{\text{turb}} = v_{\text{rot}}$.

6.3. The molecular fraction

In order to derive an expression for the molecular fraction of the gas in the disk, we compare the crossing time of the turbulent layer t_{turb} and the H–H₂ transition time scale $t_{\text{H}_2} = \alpha/\rho$ (Hollenbach & Tielens 1997). We define the molecular fraction here as $f_{\text{mol}} = t_{\text{turb}}/t_{\text{H}_2}$.

For the three different cases we obtain using a Kolmogorov spectrum:

1. dominating central mass:

$$f_{\text{mol}} = 1/(\pi\alpha) Q^{-1} Re^{-\frac{1}{2}} G^{-1} R^{-2} (-\Omega')^{-2} \Omega^3;$$

2. dominating stellar disk mass:

$$\begin{aligned} f_{\text{mol}} &= 2/(\pi\alpha) G^{-2} Q^{-2} Re^{-\frac{3}{2}} \dot{M}^{-1} (-\Omega')^{-3} \Omega^7 \\ &= 2/(\pi\alpha) G^{-2} Q^{-2} Re^{-\frac{3}{2}} \dot{M}^{-1} R^{-3} (-\Omega')^{-3} v_{\text{rot}}^3 \Omega^4; \end{aligned}$$

3. self-gravitating gas disk in z direction:

$$f_{\text{mol}} = 1/(\pi\alpha) Q^{-3} Re^{-\frac{1}{2}} G^{-1} R^{-2} (-\Omega')^{-2} \Omega^3.$$

For a non Kolmogorov spectrum $E(k) \propto k^{-2}$, the molecular fractions have to be multiplied by $Re^{-\frac{1}{2}}$: $f_{\text{mol}}^{\text{nonK.}} = Re^{-\frac{1}{2}} f_{\text{mol}}^{\text{K.}}$.

In the following we will only treat the case of a self-gravitating gas disk in z direction, because it represents the most realistic description of normal field spiral galaxies. We will motivate this choice in Sect. 6.4.

For a galaxy with a constant rotation velocity the molecular gas surface density is given by

$$\Sigma_{\text{H}_2} = \frac{1}{2^{\frac{1}{3}} \pi^2 \alpha} Q^{-\frac{10}{3}} Re^{-\frac{1}{6}} G^{-\frac{5}{3}} \dot{M}^{\frac{1}{3}} \Omega^2. \quad (67)$$

The total integrated molecular gas mass is then

$$M_{\text{H}_2} = \pi \Sigma_{\text{H}_2} R^2 = \frac{1}{2^{\frac{1}{3}} \pi \alpha} Q^{-\frac{10}{3}} Re^{-\frac{1}{6}} G^{-\frac{5}{3}} \dot{M}^{\frac{1}{3}} v_{\text{rot}}^2, \quad (68)$$

which does not depend on the galactic radius.

With these results we can now calculate the ratio of molecular to atomic gas in spiral galaxies

$$M_{\text{H}_2}/M_{\text{HI}} = M_{\text{H}_2}/(M_{\text{gas}}^{\text{tot}} - M_{\text{H}_2}), \quad (69)$$

where $M_{\text{gas}}^{\text{tot}} = \pi \Sigma R^2$ is the total gas mass.

6.4. Star formation and the Tully–Fisher relation

The star formation rate per unit area $\dot{\Sigma}_*$ is usually described by a Schmidt law (Schmidt 1959) of the form

$$\dot{\Sigma}_* \propto \Sigma_{\text{gas}}^N, \quad (70)$$

where $N \sim 1.4$ (Kennicutt 1998). An alternative description of the star formation rate was proposed by Elmegreen (1997) and Silk (1997). They suggested that it scales with

the ratio of the gas density to the average orbital time scale

$$\dot{\Sigma}_* = 0.017 \Sigma_{\text{gas}} \Omega. \quad (71)$$

This parameterization provides a fit that is nearly as good as the Schmidt law.

On the basis of our model, we suggest that the star formation time scale is $\tau_{\text{SF}} = Re \Omega^{-1}$, which is the interaction time scale between the clouds. This corresponds to the time scale of gravitational encounters between the clouds. We thus find $Re \sim 60$, which is only a factor 2 higher than our estimate based on the disk height and the cloud size. In the case of a constant rotation velocity our star formation prescription yields in the case of self-gravitation in z direction:

$$\dot{\Sigma}_* = \frac{1}{2^{\frac{1}{3}} \pi} Q^{-\frac{1}{3}} Re^{-\frac{2}{3}} G^{-\frac{2}{3}} \dot{M}^{\frac{1}{3}} \Omega^2. \quad (72)$$

Thus both, the star formation rate per unit area and the molecular gas density, are proportional to the square of the angular velocity $\Sigma_{\text{H}_2} \propto \Omega^2$, $\dot{\Sigma}_* \propto \Omega^2$. Both profiles have thus the same dependence on the galactic radius. This is what is found when comparing CO and H α , radio continuum, B band luminosity profiles of most of the Virgo cluster galaxies observed by Kenney & Young (1989). More recently, Rownd & Young (1999) combined H α imaging and CO-line observations to derive radial star formation efficiencies for a sample of cluster and field spiral galaxies. They found that $\dot{\Sigma}_*/\Sigma_{\text{H}_2} = \text{const}$, which is consistent with our findings. It is worth noting, that in the framework of our model this proportionality only exists between the star formation rate and the molecular gas mass and *not* the total gas mass.

Whereas the star formation laws of Kennicutt (1998) or Silk (1997) depend explicitly on the gas surface density, our prescription only depends on the angular velocity Ω . The proportionality factor depends on \dot{M} and the free parameters Re , Q . The results of Rownd & Young (1999) imply that the combination of these parameters or all parameters individually are the same for all spiral galaxies. The prescription of Silk (1997) is equivalent to ours $\dot{\Sigma}_* = \xi \Sigma \Omega$. We identify $\xi = Re^{-1}$. The observationally derived ξ is consistent with the definition of $Re = (l_{\text{driv}}/l_{\text{diss}})^{\frac{3}{4}}$. For the comparison between the Kennicutt and Silk law we refer to Kennicutt (1998).

The integrated star formation rate is

$$\dot{M}_* = \frac{Q}{2} \left(\frac{v_{\text{rot}}}{v_{\text{turb}}} \right)^2 \dot{M} = \frac{1}{2^{\frac{1}{3}}} Q^{-\frac{1}{3}} Re^{-\frac{2}{3}} G^{-\frac{2}{3}} \dot{M}^{\frac{1}{3}} v_{\text{rot}}^2. \quad (73)$$

Thus the integrated star formation rate is proportional to the square of the rotation velocity as the total amount of molecular gas. Observationally, Devereux & Hameed (1997) found that the ratio of star formation rate to molecular gas ($L_{\text{FIR}}/M(\text{H}_2)$) is independent of the morphological type of a spiral galaxy, thus $\dot{M}_* \propto M_{\text{H}_2}$.

The blue Tully–Fisher relation is dominated by light associated with current star formation. In general, the

galaxy luminosity is proportional to the rotation velocity at a certain power $L \propto v_{\text{rot}}^\gamma$. Compilations of Tully–Fisher data for available samples find that $\gamma = 2.1\text{--}2.2$ for the B band with a systematic increase with increasing wavelength (Silk 1997). Our model would ideally predict $\dot{M}_* \propto v_{\text{rot}}^2$, i.e. $\gamma = 2$. Since the B band luminosity is provided by a mixture of stars of different ages, we would expect that γ is slightly higher than predicted by our model.

We will now calculate the integrated star formation rate for the case of a galaxy with dominating stellar disk. Inserting the expression for the volume filling factor for a constant rotation velocity disk $\phi_V \simeq \frac{4}{3\pi} Q^{-1} Re^{-1} (v_{\text{rot}}/v_{\text{turb}})^3$ and Eq. (41) into the expression for the integrated star formation rate $\dot{M}_* = \pi \Sigma R^2 Re^{-1} \Omega$ yields

$$\dot{M}_* = \frac{v_{\text{rot}}}{v_{\text{turb}}} \dot{M} = \left(\frac{3\pi}{4} Re Q \Phi_V \right)^{\frac{1}{3}} \dot{M}. \quad (74)$$

Taking $\phi_V = 10^{-2}$, $Re = 50$, $\dot{M} = 10^{-2} M_\odot \text{ yr}^{-1}$, and $v_{\text{rot}} = 200 \text{ km s}^{-1}$ gives $v_{\text{turb}} \sim 50 \text{ km s}^{-1}$ and $\dot{M}_* \sim 4 \times 10^{-2} M_\odot \text{ yr}^{-1}$. Thus, the integrated star formation rate is comparable to the mass accretion rate. Such low integrated star formation rates are only observed for S0 galaxies (Kennicutt 1998). The Toomre parameter for such a disk is $Q \sim 7 \times 10^3$. For $\phi_V = 1$, $Q Re^{\frac{2}{3}} = 4(v_{\text{rot}}/v_{\text{turb}})^3 \geq 10^3$. This is not the case for normal field spirals. Thus, for most of the field spiral galaxies, the case of a self-gravitating disk in z direction applies.

6.5. Low surface brightness galaxies

Low surface brightness (LSB) galaxies have low density/column density gas but a high gas fraction $M_{\text{gas}}/M_{\text{tot}}$ (de Blok 1999). These LSB galaxies can be either dwarf galaxies with small rotation velocities (see e.g. Walter & Brinks 1999) or large galaxies with high rotation velocities (see e.g. Pickering et al. 1997). Both have small angular velocities $\Omega = v_{\text{rot}}/R$. Since our model yields $\Sigma_{\text{gas}} \propto \Omega$ and $\rho \propto \Omega^2$, a small angular velocity implies a low gas column density. Thus, the observed small gas density/column density of LSB galaxies can be partly due to their small angular velocity.

7. Application to the Galaxy

As motivated in Sect. 6.4 the model of a vertically self-gravitating gas disk describes the gas disks of spiral galaxies. We will use the local parameter of the ISM in the solar neighbourhood given by Binney & Tremaine (1987): $\rho = 0.042 M_\odot \text{ pc}^{-3}$, $\Sigma = 5.3 M_\odot \text{ pc}^{-2}$, $\Omega = 2.62 \times 10^{-8} \text{ yr}^{-1}$ and $R = 10 \text{ kpc}$. Furthermore, we have shown in Sects. 3.1 and 6.4 that $Re \sim 60\text{--}80$. The rotation curve is assumed to be constant. Equation (43) leads to $Q \sim 1$. On the other hand, the definition of the Q parameter (Eq. (27)) would yield $v_{\text{turb}} \sim 3 \text{ km s}^{-1}$. This equation would thus require $Q \sim 3$ in order to match the gas




Fig. 3. Radial profiles of the rotation velocity, angular velocity, turbulent velocity, driving wavelength, gas column density, disk scale height, gas density, viscosity, molecular fraction, and volume filling factor for three different rotation velocities (solid line: $v_{\text{rot}} = 250 \text{ km s}^{-1}$, dashed line: $v_{\text{rot}} = 175 \text{ km s}^{-1}$, dotted line: $v_{\text{rot}} = 100 \text{ km s}^{-1}$).

velocity dispersion of $v_{\text{turb}} = 10 \text{ km s}^{-1}$ found by van der Kruit & Shostak (1982). We will here adopt $Q = 1$, $Re = 50$, $\alpha = 10^7 \text{ yr } M_{\odot} \text{ pc}^{-3}$ (this corresponds to a molecular

transition time scale of $t_{\text{mol}} \sim 5 \times 10^8/n \text{ yr}$, where n is the particle density), $R = 10 \text{ kpc}$, and $v_{\text{turb}} = 10 \text{ km s}^{-1}$. Equation (46) then leads to $\dot{M} \sim 10^{-2} M_{\odot} \text{ yr}^{-1}$.




Fig. 4. Radial profiles of the total gas (dotted line), the molecular (solid line), and the atomic (dashed line) gas surface density.

Thus, we suggest that the Galactic mass accretion rate is $\dot{M} \sim 10^{-2} M_{\odot} \text{ yr}^{-1}$. Figure 3 shows the derived turbulent velocity, driving wavelength, gas column density, disk scale height, gas density, viscosity, molecular fraction, and volume filling factor for three different rotation velocities. For the model of the Galaxy, we have adopted $v_{\text{rot}} = 250 \text{ km s}^{-1}$.

This results in a total gas mass of $M_{\text{gas}}^{\text{tot}} = 5.4 \times 10^9 M_{\odot}$, a total molecular gas mass $M_{\text{H}_2} = 1.7 \times 10^9 M_{\odot}$, a total atomic gas mass $M_{\text{HI}} = 3.7 \times 10^9 M_{\odot}$, and $M_{\text{H}_2}/M_{\text{HI}} = 0.3$. Scoville & Sanders (1987) give $M_{\text{HI}} \sim 2.5 \times 10^9 M_{\odot}$, $M_{\text{H}_2} \sim 0.7 \times 10^9 M_{\odot}$ (we have used the conversion factor $X = 10^{20} \text{ cm}^{-2} (\text{K km s}^{-1})^{-1}$ of Digel et al. 1996) and thus $M_{\text{H}_2}/M_{\text{HI}} \sim 0.3$. Our model calculations are consistent within a factor of 2 with the absolute values of the total gas mass and the observed ratio of molecular to atomic gas in the Galaxy. Figure 4 shows the radial profiles of the total gas (dotted line), the molecular (solid line), and the atomic (dashed line) gas surface density for $v_{\text{rot}} = 250 \text{ km s}^{-1}$ and $\alpha = 7 \times 10^6 \text{ yr } M_{\odot} \text{ pc}^3$. These profiles well resemble that found by Kenney & Young (1989) (their Fig. 12). Honma et al. (1995) examined the HI and H₂ radial distributions of four face-on spiral galaxies. They found that the gas phase transition occurs suddenly within a narrow range of radius. This transition is explained using a theory whose main parameters are the ISM pressure, the UV radiation field, and the metallicity. It was found that the metallicity gradient is most crucial for the formation of a narrow range transition. Our model (Fig. 4) also shows a sharp phase transition. For a given Re , Q , and \dot{M} , the location of this transition region depends mainly on the formation time scale α of H₂ from HI. If α depends on metallicity (and that is what one would expect), the two models are in good agreement, because radiation, which is not included in our model, does not play a major rôle.

The disk height equals the driving wavelength $H \sim l_{\text{driv}} \sim 400 \text{ pc}$, and the viscosity is $\nu \sim 8 \times 10^{-5} \text{ pc}^2 \text{ yr}^{-1} = 2 \times 10^{25} \text{ cm}^2 \text{ s}^{-1}$. Thus, the macroscopic Reynolds number is $Re_{\text{macro}} = v_{\text{rot}} R / \nu \sim 3 \times 10^4 \gg Re$. This value of the driving wavelength is only a factor of 2 larger than that found by Wada & Norman (2001) in their 2D hydrodynamical simulations.

The derived mass accretion rate is much smaller than the star formation rate (Eq. (73) yields $\dot{M}_{*} \sim 3 M_{\odot} \text{ yr}^{-1}$ compared to $\dot{M}_{*} \sim 3 - 5 M_{\odot} \text{ yr}^{-1}$ as suggested by the observations of other Sbc spirals; Tamman et al. 1994; Cappellaro et al. 1997). The current viscous time scale $t_{\nu} \sim R^2 / \nu \sim 10^{12} \text{ yr}$ is larger than a Hubble time. Since the viscous time scale is given by

$$t_{\nu} \sim \frac{R^2}{\nu} \propto \dot{M}^{-\frac{2}{3}} R^2 \Omega \sim \dot{M}^{-\frac{2}{3}} v_{\text{rot}} R, \quad (75)$$

viscous evolution has taken place at a time when the mass accretion rate was higher and/or the disk was smaller and/or rotation velocity was smaller. After this evolution the gas distribution reached an equilibrium state that is described by our model. Lin & Pringle (1987b) and later Saio & Yoshii (1990) have shown that one obtains an exponential stellar disk if $t_{\nu} \sim t_{*}$. In our model this is only the case for a dominating stellar disk, where $\dot{M}_{*} \sim \dot{M}$. However, we have to stress here that \dot{M} is the mass accretion rate in the disk. If there is also external accretion present, this formally increases t_{*} , which can lead to $t_{\nu} \sim t_{*}$.

8. Morphological type versus mass

In this section we investigate the link between the morphological classification of galaxies and their mass on the basis of our model for vertically selfgravitating gas disks. In general, disk galaxies of later types (Sc–Scd) are smaller and less massive than galaxies of earlier type (Sa–Sab) (Roberts & Haynes 1994). Some of the properties which lead to the morphological classification could thus be due to the galaxies' gravitational potential as proposed by Gavazzi et al. (1996).

8.1. The velocity dispersion and the fraction of gas to total mass

It is a surprising fact that galactic gas disks all have dispersion velocities between 8 and 10 km s^{-1} (see e.g. Freeman 1999). Assuming a constant rotation velocity Eq. (46) implies that $Q^2 Re \dot{M}$ is approximately constant for spiral galaxies of all types and all luminosities. Furthermore, if the galaxy forms still actively stars Q should be ~ 1 . Thus $Re \dot{M}$ can only vary by a factor ~ 10 .

From Eq. (28) we find $M_{\text{gas}}/M_{\text{tot}} = Q^{-1} v_{\text{turb}}/v_{\text{rot}}$. Since the gas velocity dispersion is almost constant, $M_{\text{gas}}/M_{\text{tot}} \propto v_{\text{rot}}^{-1}$. Galaxies with high rotation velocities have thus smaller gas to total mass ratios. This trend is also observed in the Hubble sequence: early type spirals




Fig. 5. Ratio of molecular to atomic gas mass as a function of the galaxy rotation velocity.

have larger rotation velocities and smaller gas to total mass ratios (Roberts & Haynes 1994).

8.2. The fraction of molecular to atomic gas

Young & Knezek (1989) have shown that the ratio of molecular to atomic gas mass in spiral galaxies increases from early type galaxies to late type galaxies by a factor ~ 10 . We have calculated the ratio

$$\frac{M_{\text{H}_2}}{M_{\text{HI}}} = \frac{\pi f_{\text{mol}} \Sigma R^2}{\pi \Sigma R^2 - \pi f_{\text{mol}} \Sigma R^2} = \frac{f_{\text{mol}}}{1 - f_{\text{mol}}}. \quad (76)$$

We used the following parameters: $Q = 1$, $Re = 50$, $\alpha = 10^7 \text{ yr } M_{\odot} \text{ pc}^{-3}$, and $R = 10 \text{ kpc}$. The result can be seen in Fig. 5. There is almost a factor 10 between the molecular to atomic gas ratio of a galaxy with $v_{\text{rot}} = 50 \text{ km s}^{-1}$ and a galaxy with $v_{\text{rot}} = 300 \text{ km s}^{-1}$. Of course, galaxies with small rotation velocities tend to be smaller, but their HI diameter should be larger, justifying $R = \text{const.}$ as a first approach. Furthermore, we expect that there is an influence of the molecule formation time scale α on the ratio between molecular and atomic gas mass. This time scale should depend on the galaxy’s metallicity. Thus the observed trend might be a mixture of (i) higher metallicities (and therefore smaller α) of early type spiral galaxies and (ii) higher rotation velocities of early type spiral galaxies compared to late type spirals, i.e. that early type galaxies are more massive.

8.3. Star formation rate per unit mass

The integrated star formation rate times the rotation period divided by the total mass should be comparable to the ratio of FIR luminosity to H band luminosity L_{FIR}/L_H . Using Eq. (73) and $v_{\text{rot}}^2 \sim M_{\text{tot}}G/R$, we obtain for our model

$$\frac{\dot{M}_* \Omega^{-1}}{M_{\text{tot}}} \propto v_{\text{rot}}^{-1}. \quad (77)$$

This means that rapidly rotating galaxies have smaller L_{FIR}/L_H . Devereux & Hameed (1997) found indeed that galaxies of later type have smaller L_{FIR}/L_H . Since the FIR luminosity is assumed to be a measure of recent star formation and the H band luminosity a measure of the total mass, $(\dot{M}_* \Omega^{-1})/M_{\text{tot}}$ decreases with increasing mass. Moreover, Gavazzi et al. (1996) found a trend of smaller $H\alpha$ equivalent width for more luminous (H band) galaxies. This also translates into a decreasing $(\dot{M}_* \Omega^{-1})/M_{\text{tot}}$ with increasing galaxy mass. Thus, again, a part of this observed trend might be due to the fact that early spirals have higher rotation velocities than late type spirals.

9. Summary

We have analytically investigated the equilibrium state of a turbulent clumpy gas disk. The disk consists of distinct self-gravitating clouds, which are embedded in a low density medium, and evolve in the fixed gravitational potential of the galaxy. The disk scale height results from the balance between the gravitational acceleration and the pressure due to the turbulent velocity dispersion of the clouds. Gravitational cloud–cloud interactions in the disk give rise to an effective viscosity and allows the transport of angular momentum and mass in the gas disk. In our scenario turbulence is assumed to be generated by instabilities involving self-gravitation and to be maintained by the energy input, which is provided by differential rotation of the disk and mass transfer to smaller galactic radii via cloud–cloud interactions.

A description of the turbulent viscosity is given, which is formally equivalent to a prescription derived from the collisional Boltzmann equation. The vertical gravitational acceleration in the gas disk is either due to a (i) dominating central mass, (ii) dominating stellar disk mass, or (iii) the vertically self-gravitating gas disk itself. For all three cases we derive analytical expressions for disk parameters (density, surface density, velocity dispersion, disk height, driving wavelength, and viscosity) as functions of the Toomre parameter Q , the turbulent Reynolds number Re , the mass accretion rate \dot{M} , the galactic radius R , and the angular velocity Ω of the gas.

The model does not include radiation. The “thermostat” equation of the standard accretion disk model is replaced by an equation which assumes that the energy flux generated by mass inflow and differential rotation is entirely transported by turbulence to smaller scales where it is dissipated. Whereas case (iii) corresponds to the $Q \sim 1$ disk, case (i) and (ii) require $Q > 1$.

The structure of the resulting clumpy gas disks allows us to derive global volume filling factors of self-gravitating clouds as a function of Q and Re . Along the same line analytical expressions for the molecular fraction and the star formation rate are given.

On the basis of our analytical model we conclude that

1. constant velocity dispersions in the gas as observed for spiral galaxies can be reproduced by models with a (i)

dominating central mass and (iii) a vertically selfgravitating gas disk.

2. In models (ii) with a dominating stellar disk the velocity dispersion depends on the rotation velocity.
3. The driving wavelength of the turbulence equals approximately the disk scale height if the vertical acceleration is not dominated by the stellar disk.
4. The effective turbulent viscosity ν depends on the galactic radius. For a disk in Keplerian rotation with a (i) dominating central mass $\nu \propto R^{\frac{3}{2}}$. The other two model classes (ii) and (iii) give $\nu \propto R$ for a constant velocity rotation curves.
5. The molecular gas surface density and the star formation rate are proportional to the square of the angular velocity $\Sigma_{\text{H}_2}, \dot{\Sigma}_* \propto \Omega^2$ in self-gravitating gas disks.

The model of a selfgravitating disk in vertical direction is applied to the Galaxy and gives a good description of its gas disk. Furthermore, we investigate the rôle of the galaxy mass for the morphological classification of spiral galaxies.

A comparison of our analytical model to the Galaxy shows that

1. The observed global star formation time scale implies a Reynolds number $Re \sim 50$.
2. Our model is consistent with observations. That is the model reproduces the physical parameters of the Galaxy as derived from observations.
3. The derived mass accretion rate of the galaxy is $\dot{M} \sim 10^{-2} M_{\odot} \text{yr}^{-1}$. It is thus much smaller than the star formation rate $\dot{M}_*/\dot{M} \sim 100$.
4. The dependence of several physical parameters of spiral galaxies on morphological type might be at least partly due to the mass–morphology relation, i.e. galaxies of later types are more massive.

In our model we assume that turbulence is generated by local instabilities due to selfgravitation. The energy input from accretion and differential rotation is high enough to maintain the ISM turbulence. Moreover, the energy dissipation rate per unit mass due to the self-gravitation of the gaseous disk in z direction has the same order of magnitude as the energy per unit time and unit mass transported to smaller scales by turbulence. Thus, in terms of energy conservation, turbulence generated and maintained by gravitation can account for the viscosity of spiral galaxies. This does not exclude energy input in the turbulence through supernovae. Nevertheless, our model where accretion in the galactic potential is sufficient to maintain ISM turbulence is fully consistent with observations.

Acknowledgements. We would like to thank the referee, K. Wada, for helping us to improve this article considerably and W. J. Duschl for fruitful discussions.

References

- Balbus, S. A., & Hawley, J. F. 1991, *ApJ*, 376, 214
 Binney, J., & Tremaine, S. 1987, in *Galactic Dynamics* (Princeton University Press)
 Blitz, L., & Shu, F. H. 1980, *ApJ*, 238, 148
 Bondi, H., & Hoyle, F. 1944, *MNRAS*, 104, 273
 Braginskii, S. I. 1965, in *Review of Plasma Physics*, ed. M. A., Leontovich, Consultants Bureau, N. Y., 1, 205
 Cappellaro, E., Turatto, M., Tsvetkov, D., et al. 1997, *A&A*, 322, 43
 Cui, W., Sanders, W. T., McCammon, D., Snowden, S. L., & Womble, D. S. 1996, *ApJ*, 468, 102
 de Blok, W. J. G. 1999, in *Low Surface Brightness Universe*, ASP Conf. Ser., 170, ed. J. I. Davies, C. Impey, & S. Phillipps, Astronomical Society of the Pacific (San Francisco), 161
 Devereux, N. A., & Hameed, S. 1997, *AJ*, 113, 599
 Dickey, J. M. 1993, in *The Minnesota lectures on the structure and dynamics of the Milky Way*, ed. R. M. Humphreys, ASP Conf. Ser., 39, 93
 Digel, S. W., Grenier, I. A., Heithausen, A., Hunter, S. D., & Thaddeus, P. 1996, *ApJ*, 463, 609
 D’Onofrio, M., Capaccioli, M., Merluzzi, P., Zaggia, S., & Boulesteix, J. 1999, *A&AS*, 134, 437
 Duschl, W. J., Strittmatter, P. A., & Biermann, P. L. 2000, *A&A*, 357, 1123
 Elmegreen, B. G. 1982, *ApJ*, 253, 655
 Elmegreen, B. G. 1987, in *Interstellar processes*, ed. D. J. Hollenbach, & H. A. Thronson Jr. (Dordrecht: Kluwer), 259
 Elmegreen, B. G. 1997, in *Starburst Activity in Galaxies*, ed. J. Franco, R. Terlevich, & A. Serano, Rev. Mex. Conf. Ser., 6, 165
 Elmegreen, B. G., & Falgarone, E. 1996, *ApJ*, 471, 816
 Fleck Jr., R. C. 1981, *ApJ*, 246, L151
 Frank, J., King, A., & Raine, D. 1992, *Accretion Power in Astrophysics* (Cambridge University Press)
 Frisch, U. 1995, *Turbulence – The Legacy of A.N. Kolmogorov* (Cambridge University press)
 Freeman, K. C. 1999, *Ap&SS*, 269, 119
 Gavazzi, G., Pierini, D., & Boselli, A. 1996, *A&A*, 312, 397
 Goldreich, P., & Tremaine, S. 1978, *Icarus*, 34, 227
 Hollenbach, D. J., & Tielens, A. G. G. M. 1997, *ARA&A*, 35, 179
 Honma, M., Sofue, Y., & Arimoto, N. 1995, *A&A*, 304, 1
 Jog, C. J., & Ostriker, J. P. 1988, *ApJ*, 328, 404
 Kenney, J. D. P., & Young, J. S. 1989, *ApJ*, 344, 171
 Kennicutt Jr., R. C. 1998, *ARA&A*, 36, 189
 Klessen, R. S., Heitsch, F., & Mac Low, M.-M. 2000, *ApJ*, 535, 887
 Kulkarni, S. R., & Heiles, C. 1988, in *Galactic and Extragalactic Radio Astronomy*, ed. G. L. Verschuur, & K. I. Kellermann (New York: Springer Verlag), 95
 Kumar, P. 1999, *ApJ*, 519, 599
 Landau, L. D., & Lifschitz, E. M. 1959, *Fluid Mechanics*
 Larson, R. B. 1981, *MNRAS*, 194, 809
 Lightman, A. P., & Eardley, D. M. 1974, *ApJ*, 187, L1
 Lin, D. N. C., & Pringle, J. E. 1987a, *MNRAS*, 225, 607
 Lin, D. N. C., & Pringle, J. E. 1987b, *ApJ*, 320, L87
 Lynden-Bell, D., & Pringle, J. E. 1974, *MNRAS*, 168, 603
 Mac Low, M.-M. 1999, *ApJ*, 524, 169
 Mac Low, M.-M., & Ossenkopf, V. 2000, *A&A*, 353, 339

- McKee, C. F., & Ostriker, J. P. 1977, *ApJ*, 218, 148
- Meneveau, C., & Sreenivasan, K. R. 1990, *Phys. Rev. A*, 41, 2246
- Ozernoy, L. M., Fridman, A. M., & Biermann, P. L. 1998, *A&A*, 337, 105
- Paczynski, B. 1978, *Acta Astron.*, 28, 91
- Parker, E. N. 1966, *ApJ*, 145, 811
- Passot, T., Pouquet, A., & Woodward, P. 1988, *A&A*, 197, 228
- Pickering, E., Impey, C. D., van Gorkom, J. H., & Bothun, G. D. 1997, *AJ*, 114, 1858
- Pringle, J. E. 1981, *ARA&A*, 19, 137
- Pringle, J. E., Rees, M. J., & Pacholczyk, A. G. 1973, *A&A*, 29, 179
- Roberts, M. S., & Haynes, M. P. 1994, *ARA&A*, 32, 115
- Rownd, B. K., & Young, J. 1999, *ApJ*, 118, 670
- Saio, H., & Yoshii, Y. 1990, *ApJ*, 363, 40
- Sanders, D. B., Scoville, P. M., & Solomon, P. M. 1985, *ApJ*, 289, 373
- Scalo, J. 1985, in *Protostars and Planets II*, ed. D. C. Black, & M. S. Matthews (Tucson: Univ. Arizona), 201
- Scoville, N. Z., & Sanders, D. B. 1987, in *Interstellar processes*, ed. D. J. Hollenbach, & H. A. Thronson Jr. (Dordrecht: Kluwer), 21
- Schmidt, M. 1959, *ApJ*, 129, 243
- Shlosman, I., & Begelman, M. C. 1987, *Nature*, 329, 810
- Shlosman, I., & Begelman, M. C. 1989, *ApJ*, 341, 685
- Shakura, N. I. 1972, *Astron. Zhur.* 49, 921 (translated in: *SvA AJ* 16, 921 (1973))
- Shakura, N. I., & Sunyaev, R. A. 1973, *A&A*, 24, 337
- Silk, J. 1997, *ApJ*, 481, 703
- Silk, J., & Norman, C. 1981, *ApJ*, 247, 59
- Spitzer, L. 1968, *Diffuse Matter in Space* (New York: Wiley Interscience)
- Stewart, G. L., & Kaula, W. 1980, *Icarus*, 44, 154
- Tammann, G., Loeffler, W., & Schroder, A. 1994, *ApJS*, 92, 487
- Toomre, A. 1964, *ApJ*, 139, 1217
- van der Kruit, P. C., & Shostak, G. S. 1982, *A&A*, 105, 351
- Vollmer, B., & Duschl, W. J. 2000, *A&A*, 367, 72
- Wada, K., & Norman, C. A. 1999, *ApJ*, 516, L13
- Wada, K., & Norman, C. A. 2001, *ApJ*, 547, 172
- Walter, F., & Brinks, E. 1999, in *Low Surface Brightness Universe*, ASP Conf. Ser., 170, ed. J. I. Davies, C. Impey, & S. Phillipps, Astronomical Society of the Pacific (San Francisco), 224
- Young, J. S., & Knezek, P. M. 1989, *ApJ*, 347, L55