

On the effective temperatures and surface gravities of superficially normal main sequence band B and A stars

Saul J. Adelman^{1,2}, O. I. Pintado^{3,4}, F. Nieva³, K. E. Rayle¹, and S. E. Sanders, Jr.¹

¹ Department of Physics, The Citadel, 171 Moultrie Street, Charleston, SC 29409, USA

² Guest Investigator, Dominion Astrophysical Observatory, Herzberg Institute of Astrophysics, National Research Council of Canada, 5071 W. Saanich Road, Victoria V9E 2E7 Canada

³ Departamento de Física, Facultad de Ciencias Exactas y Tecnología, Universidad Nacional de Tucumán, Av. Independencia 1800, 4000 San Miguel de Tucumán, Argentina

⁴ Member of the Carrera del Investigador Científico, CONICET, Consejo Nacional de Investigaciones Científicas y Técnicas de la República Argentina

Received 26 February 2002 / Accepted 12 June 2002

Abstract. Effective temperatures and surface gravities for 48 main sequence band B and A stars were found by matching optical region spectrophotometry and H γ profiles with the predictions of ATLAS9 solar composition model atmospheres. When these values were compared with those found using Strömberg *uvby* photometry based on ATLAS6 model atmospheres, we found a difference (photometry-spectrophotometry) of 25 ± 118 K for 29 stars with $8000 \text{ K} \leq T_{\text{eff}} \leq 10\,050 \text{ K}$ compared to 76 ± 105 K for 14 stars with $10\,050 \text{ K} \leq T_{\text{eff}} \leq 17\,000 \text{ K}$. The surface gravity scales are in agreement. These stars are sufficiently hot that their effective temperatures and surface gravity determinations are unaffected by discrepancies due to the choice of Mixing-Length or Canuto-Mazzitelli convection theories.

Key words. stars: fundamental parameters – stars: early-type

1. Introduction

Astronomers can determine the effective temperatures and surface gravities of single main sequence band B and A stars by matching their optical region flux distributions as measured spectrophotometrically and their observed H γ profiles with the predictions of the best model atmospheres. When these observations are not available, investigators have used photometric indices calibrated by this method, e.g. for Strömberg *uvby* photometry the work of Moon & Dworetzky (1985), which has been updated by Napiwotzki et al. (1993). The errors are of order ± 200 K and ± 0.2 dex in the middle B to early F star regime (Lemke 1989) (see Ribas et al. 1997 concerning which values are better). Our current ability to calculate physically realistic model atmospheres is the greatest in this region of the HR Diagram. These calibrations are based mainly on model atmospheres used prior to 1992. In the optical region, there can be small differences in the predicted fluxes between ATLAS9 (Kurucz 1993) and the older ATLAS6 (Kurucz 1979) models due in part to an improved treatment of line blanketing and to the use of opacity distribution functions with different micro-turbulences. This paper to a large extent investigates the effects of these differences.

In determining stellar effective temperatures and surface gravities, two basic concerns are the agreement of the predictions of the models with the observations as this is what many abundance workers use to select appropriate models and whether these results show systematic errors when one uses flux observations covering a much wider range of wavelengths. In this paper we are concerned mainly with the former. One can obtain infrared and perhaps ultraviolet fluxes and use the results of this study as a basis for determining effective temperatures via the infrared flux method. But for single and binary stars with sufficiently faint companions if there are no systematic errors in the model fluxes, our approach should yield as good results without the complications of additional sources of error.

The ATLAS9 code produces fully line blanketed LTE plane parallel model atmospheres. From converged model atmospheres, one can calculate the fluxes using ATLAS9 and the H γ region using the synthetic spectrum code SYNTHE (Kurucz & Avrett 1981). For comparison with observations the synthesized spectra were convolved with the measured stellar rotational velocity and the instrumental profile of the short camera of the coude spectrograph of the Dominion Astrophysical Observatory (DAO) 1.22-m telescope. Trends in recent elemental abundance studies indicate for main sequence band stars with temperatures greater than 10 250 K their

Send offprint requests to: Saul J. Adelman,
e-mail: adelmans@citadel.edu

microturbulence is 0 km s^{-1} , for those with temperatures between 10 250 and 9500 K 1 km s^{-1} , and for those with temperatures less than 9500 K 2 km s^{-1} (Adelman 1999). In this investigation, grids of ATLAS9 models with effective temperature and surface gravity spacings of 500 K and 0.25 dex, respectively, were employed. Additional models were calculated to confirm the interpolated values. In general the greater the number of spectrophotometric flux values, the better the resultant temperature determination.

2. Observational data

To assess the discrepancies of the effective temperatures and surface gravities found with the algorithm of Napiwotzki et al. (1993) and those found by comparison of fluxes and H γ profiles with ATLAS9 model predictions, 48 stars with fluxes based on the Hayes-Latham (1975) calibration of Vega mainly from the catalogs of Breger (1976) and Adelman et al. (1989) were studied. The homogeneous $uvby\beta$ values of Hauck & Mermilliod (1980) were employed. The H γ profiles were extracted from 20 \AA mm^{-1} spectrograms obtained with either Reticon or CCD detectors at the Dominion Astrophysical Observatory (DAO) and had a resolution of 0.6 \AA (2 pixels). The exposures were flat fielded with those of an incandescent lamp placed in the Coudé mirror train as viewed through a filter to eliminate first order light. A central stop removed light from the beam in the same manner as the secondary mirror of the telescope. The spectra were rectified with the interactive computer graphics program REDUCE (Hill et al. 1982). A 3.5% correction (Gulliver et al. 1996) was applied for scattered light along the dispersion direction.

The normalization of the H γ observations is not without problems. The continuum can be difficult to locate for late A and cooler stars and while the sensitivity of the spectrograph at the DAO changes over the 200 \AA interval centered on this Balmer line due to the high reflective coatings used in the Coudé mirror train. In retrospect one might be able to improve somewhat the normalization by the use of all of the observations from a single night, especially if observations were made of a range of spectral types, rather than processing each spectrum independently. This would necessitate some changes in reduction software. Alternatively if such data becomes available one might be able to use spectrophotometry having continuous wavelength coverage and a resolution of a few \AA to help the normalization process.

3. B and A star temperatures and surface gravities

As most spectrophotometers capable of determining high quality stellar fluxes in the optical region were retired from service about twenty years ago, the candidate stars for H γ observation were those whose values were cataloged and could be observed from the DAO in Victoria. Additional selection criteria were that the spectrophotometric wavelength range must span the Balmer jump and that Strömgren indices derived from the spectrophotometry were in good accord with those values directly observed. The reddening, if any, was deduced using the computer program of Napiwotzki et al. (1993) and is uncertain

to ~ 0.005 mag. If positive and would result in an observable change in the flux distribution, the spectrophotometric values were dereddened using those of Schild (1977). The information content of the spectrophotometric values is greater than that of the photometric values.

In this study we also wanted to avoid studying stars for which uncertainties in convection theory have observable effects on the fluxes and H γ profiles. Smalley & Kupka (1997) argued that the turbulent convection theory of Canuto & Mazzitelli (1991, 1992) should be more realistic than Mixing-Length theory (Castelli et al. 1997). Kupka (private communication) supplied the necessary subroutine for Canuto-Mazzitelli convection in ATLAS9. Smalley & Kupka (1997) indicate the discrepancies in effective temperatures and surface gravities between these theories.

As it would be useful to know where these two theories predict observable discrepancies, we found the last stellar models which exhibit minimal differences (less than 1%) in the optical region fluxes and H γ profiles as one proceeds towards cooler temperatures. These models had the fine solar abundance opacity distribution function, a microturbulence of 2 km s^{-1} , and 64 depths converged usually to better than 1% in flux and in flux derivative. The convergence properties of the models with Canuto & Mazzitelli convection are often nicer than those for Mixing-Length theory. The synthetic spectra whose resolution is 500 000 were broadened by the instrumental profile of the DAO long camera (Gulliver & Hill 1990). Our comparisons were made by making plots of the H γ profiles and flux distributions that utilized most of the dimensions of letter sized pages. Initially we used a grid of models with a 500 K spacing in effective temperature. As the comparisons proceeded we reduced the spacing to find the onset of differences to $\pm 25 \text{ K}$. Thus the coolest models with no significant differences occur at 7725 K for $\log g = 3.00$, 7850 K for $\log g = 3.25$, 8000 K for $\log g = 3.50$, 8150 K for $\log g = 3.75$, 8300 K for $\log g = 4.00$, and 8475 K for $\log g = 4.25$. These boundary values are likely to depend on microturbulence and metallicity, variables not investigated at this time. The stars studied in this paper are on the hot side of this boundary.

Table 1 contains the determinations of the temperatures and surface gravities along with the HD number and another identification, the metallicities (with $[m/H] = 0$ being solar), the sources of the spectrophotometry, and the reddenings for 48 stars whose effective temperatures are $\geq 8000 \text{ K}$. The values longward of H α were given lower weights than other values owing to the authors' previous difficulties in obtaining simultaneous fits to them and other spectrophotometric values (see, e.g., Adelman et al. 1999). Figure 1 shows as an example the best fit for ν Cap. For most stars the discrepancies are less between the observations and the best theoretical fits. But in the Paschen continuum the spectrophotometric values of β Lib, τ Her, and ϵ Cas produced only a fair fit to the predicted fluxes of the models. As the photometric values of T_{eff} and $\log g$ for θ Vir were determined without the use of a measured β value, they are not used for comparison with the spectrophotometrically derived values. For 22 Dra which has spectrophotometry by Schild et al. (1971), we found $T_{\text{eff}} = 12 500 \text{ K}$ and $\log g = 3.55$, but this star lacks published $uvby\beta$ values.

Table 1. Effective temperature and surface gravity determinations.

Star Name	HD number	$uvby\beta$ T_{eff}	$\log g$	spectro. + T_{eff}	Hy $\log g$	$[m/H]$	spectro. source	reddening $E(b-y)$
τ Vir	122 408	8085	3.84	8150	3.75	0.0	APW	0.00
95 Leo	103 578	8331	4.14	8300	3.65	0.0	APW	0.01
δ Her	154 494	8361	3.95	8500	4.00	0.0	DP	0.01
HR 6410	156 164	8615	3.70	8500	4.00	0.0	APW	0.00
δ UMa	106 591	8664	3.87	8650	4.10	0.0	AP	0.00
λ Boo	125 162	8925	4.11	8750	4.00	0.0	OKE	0.02
				8700	4.15	-1.5		
δ Aqr	216 677	8657	3.56	8750	3.50	0.0	MOR	0.00
68 Tau	27 962	9025	3.95	9000	4.00	0.0	LL	0.00
				8900	4.00	0.2		
λ UMa	89 021	8790	3.75	9000	3.75	0.0	SPO	0.00
				9000	3.75	0.2		
θ And	1280	8968	3.87	9000	4.00	0.0	APW	0.00
γ Oph	161 868	9388	4.09	9100	4.00	0.0	APW	0.02
γ Gem	47 105	9277	3.51	9150	3.60	0.0	TAY	0.00
ϵ Aqr	198 001	9229	3.56	9200	3.50	0.0	SPO	0.00
ν Tau	25 490	9226	3.93	9250	4.00	0.0	MOR	0.00
θ Leo	97 633	9289	3.62	9250	3.60	0.0	APW	0.00
27 Lyn	67 006	9320	3.77	9250	3.75	0.0	A	0.01
60 Leo	95 608	9053	4.22	9250	4.25	0.0	BVJ	0.00
				9250	4.25	0.5		
γ UMa	103 287	9361	3.79	9350	3.75	0.0	APW	0.00
θ Vir	114 330	9407*	3.87*	9350	3.50	0.0	SPO	0.00
21 Lyn	58 142	9601	3.74	9425	3.75	0.0	AP	0.00
HR 6127	148 330	9670	4.00	9500	3.80	0.0	AP93	0.01
ζ Aql	177 724	9542	3.84	9500	3.90	0.0	SPO	0.01
o Peg	214 994	9591	3.64	9525	3.70	0.0	A	0.00
γ Lyr	176 437	9674	2.67	9550	2.75	0.0	SPO	0.01
β UMa	95 418	9601	3.85	9600	3.80	0.0	AP	0.00
				9600	3.80	0.2		
HR 7086	174 262	9507	4.12	9600	4.20	0.0	APW	0.00
HR 5169	119 763	9639	4.12	9700	4.15	0.0	APW	0.00
α Sex	87 887	9862	3.54	9875	3.55	0.0	MOR	0.01
α Dra	123 299	9975	3.63	10 000	3.50	0.0	A	0.00
σ Aqr	213 320	10 188	3.94	10 100	3.85	0.0	A	0.00
29 Vul	196 724	10 397	4.14	10 200	4.00	0.0	WKH	0.00
κ Cep	192 907	10 341	3.64	10 250	3.75	0.0	APW	0.00
ν Cap	193 432	10 311	3.86	10 250	4.00	0.0	A	0.00
21 Peg	209 459	10 375	3.47	10 350	3.55	0.0	AP	0.01
14 Cyg	185 872	10 953	4.11	10 750	3.55	0.0	APW	0.00
134 Tau	38 899	10 928	4.01	10 750	4.10	0.0	A	0.00
5 Aqr	198 667	11 288	3.37	11 125	3.55	0.0	A	0.01
β Lib	135 742	12 036	3.26	12 125	3.50	0.0	KUB	0.01
21 Aql	179 761	13 029	3.44	13 000	3.60	0.0	A	0.05
π Cet	17 081	13 174	3.70	13 100	3.85	0.0	A	0.01
HR 2154	41 692	14 330	3.21	14 500	3.50	0.0	A	0.02
τ Her	147 394	15 022	3.93	15 000	4.10	0.0	A	0.02
ϵ Cas	11 415	15 290	3.85	15 125	3.55	0.0	SPO	0.02
η Aur	32 630	16 887	4.08	16 375	4.10	0.0	SPO	0.00
η UMa	120 315	17 319	4.31	16 900	4.30	0.0	AP	0.00
γ Peg	886	21 482	3.92	21 250	4.00	0.0	A	0.00
1 Cas	218 376	27 334	4.10	29 000	4.00	0.0	A	0.17
10 Lac	214 680	32 845	4.61	33 000	4.40	0.0	KUB	0.06

Notes: Breger (1976) converted data from DP, KUB, MOR, OKE, SPO, and WKH to the Hayes & Latham (1975) calibration of Vega. The spectrophotometric sources are: A = Adelman (1978), AP = Adelman & Pyper (1983), AP93 = Adelman & Pyper (1993), APW = Adelman et al. (1980), BVJ = Böhm-Vitense & Johnson (1978), DP = Dickens & Penny (1971), KUB = Kubiak (1973), LL = Lane & Lester (1980), MOR = Gutierrez-Moreno et al. (1968), OKE = Oke (1964), SPO = Schild et al. (1971), TAY = Taylor (1984), and WKH = Wolff et al. (1968).

* = estimated with predicted β value.




Fig. 1. The spectrophotometric fluxes for ν Cap (open squares) compared with the predictions of a solar composition $T_{\text{eff}} = 10\,250$ K, $\log g = 4.00$ ATLAS9 model atmosphere with a microturbulence of 0 km s^{-1} (thin line connecting solid triangles). The axes are inverse microns and the magnitude of the flux in frequency units in $\text{ergs cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1}$. The constant was chosen to make the flux at 5000 \AA be 0.0 for the observations. The theoretical fluxes were shifted for the best match over as much of the observed wavelength range as possible.




Fig. 2. The synthesized spectrum of γ Peg (thicker line) centered at $H\gamma$ calculated with a solar composition $T_{\text{eff}} = 21\,250$ K, $\log g = 4.00$ ATLAS9 model atmosphere with a microturbulence of 0 km s^{-1} (thinner line).

For most stars the synthesized $H\gamma$ regions match the observations quite well, especially the Balmer line, but not perfectly. Three representative examples are shown in Figs. 2–4 with the thicker lines being the synthesized spectra and the thinner lines

the observations. For γ Peg, the He I line $\lambda 4388$ is not satisfactorily synthesized and for γ Gem the metal lines in the synthesized spectrum are slightly too deep. For 10 Lac, the worst case in fitting, the Balmer line profile fit was to the $H\gamma$ line wings




Fig. 3. The synthesized spectrum of σ Aqr (thicker line) centered at Hy calculated with a solar composition $T_{\text{eff}} = 10\,100$ K, $\log g = 3.85$ ATLAS9 model atmosphere with a microturbulence of 1 km s^{-1} (thinner line).

with the observed core being deeper, especially in the central 5 \AA , by about 5%, which may represent non-LTE effects while for 60 Leo models with a 4 km s^{-1} microturbulence were used consistent with the analysis by Adelman et al. (1999).

For 48 stars the average difference in T_{eff} (photometric minus spectrophotometric-Hy value) is 29 ± 293 K. For those stars with $8000\text{ K} \leq T_{\text{eff}} \leq 10\,050\text{ K}$, we find a difference of 25 ± 118 K and for stars with $10\,050\text{ K} \leq T_{\text{eff}} \leq 17\,000\text{ K}$, 76 ± 105 K. This suggests a slight decrease of the derived temperature scale which is not unexpected as it reflects an increase in the metal line opacity for ATLAS9 relative to ATLAS6 models. Figure 5 shows the difference (ΔT_{eff} (K)) between the effective temperatures as derived using photometry and spectrophotometry as a function of the photometric temperature ($T_{\text{eff}}(uvby\beta)$ (K)).

The mean difference in $\log g$ (photometric minus spectrophotometric-Hy value) for 47 stars is -0.007 ± 0.169 dex. The scatter is similar to the uncertainty found by Lemke (1989) 0.20 dex. But two stars, 95 Leo and 14 Cyg, have $\log g$ values with rather large discrepancies, 0.49 dex and 0.56 dex, respectively. Together they have a 0.02 dex effect on the average. Figure 6 shows the difference ($\Delta \log g$ (dex)) between the surface gravity as derived using photometry and from spectrophotometry and the Hy profile as function of the photometric temperature $T_{\text{eff}}(uvby\beta)$ (K). It is best described as a scatter diagram. Since the discrepancies in the temperatures are small relative to the derived temperatures, the basic agreement of the surface gravity scales is not unexpected.

In Table 2 we compare the calibration or “known” temperatures that Napiwotzki et al. (1993) used from

Table 2. Comparison of effective temperature and surface gravity determinations.

Star	“known”	T_{eff}	
		$uvby\beta$	spectro.+Hy
γ Gem	9240	9277	9150
β UMa	9170	9601	9600
ν Cap	9950	10 311	10 250
134 Tau	10 790	10 928	10 750
π Cet	12 820	13 174	13 100
η Aur	17 580	16 887	16 375

Code et al. (1976), Beeckmans (1977), and Malagini et al. (1986) with those from their $uvby\beta$ calibration and our values. Smalley & Dworetsky (1995) re-evaluated the fundamental values of Code et al. (1976) using more recent flux measurements and found no significant changes. In general our results agreed better with the photometric values than the calibrator values and exhibit agreement with the calibrator similar to that of the photometric values. However this comparison can be done only for a few calibrators. How the nonuniformity of the spectrophotometric data affects these results is unclear and is a possible source of some differences as the data quality is not uniform among the spectrophotometric sources.

4. Conclusions

To substantially decrease the errors from this type of study we need a set of uniform high quality optical flux measurements with near continuous coverage for many more stars. Small systematic differences in effective temperature, but not in surface




Fig. 4. The synthesized spectrum of γ Gem (thicker line) centered at H γ calculated with a solar composition $T_{\text{eff}} = 9150$ K, $\log g = 3.60$ ATLAS9 model atmosphere with a microturbulence of 2 km s^{-1} (thinner line).




Fig. 5. The differences (ΔT_{eff} (K)) (photometric minus spectrophotometric-H γ values) of the effective temperatures as a function of the photometric temperature T_{eff} ($uvby\beta$) (K).




Fig. 6. The differences ($\Delta \log g$ (dex)) (photometric minus spectrophotometric-H γ values) of the surface gravities as a function of the photometric temperature T_{eff} ($uvby\beta$) (K).

gravity have been found between matching optical spectrophotometry and H γ profiles with the predictions of ATLAS9 solar composition model atmospheres and those derived from Strömgren $uvby\beta$ photometry based on ATLAS6 model atmospheres. These discrepancies have been attributed to small differences in the line opacities. In performing elemental abundance analyses, their effects will be small. But as one must do many steps as well as possible, the total effect of many such differences might be substantial.

Acknowledgements. SJA's contribution to this paper was support in part by grants from The Citadel Foundation as was that of KER.

Some H γ observations at the DAO were made by Frank Younger and Charles Perry. We thank F. Kupka for his implementation of Canuto-Mazzitelli convection in ATLAS9.

References

- Adelman, S. J. 1978, *ApJ*, 222, 547
- Adelman, S. J. 1999, *MNRAS*, 310, 146
- Adelman, S. J., & Pyper, D. M. 1983, *ApJ*, 266, 732
- Adelman, S. J., & Pyper, D. M. 1993, *A&AS*, 101, 393

- Adelman, S. J., Pyper, D. M., & White, R. E. 1980, *ApJS*, 43, 491
- Adelman, S. J., Pyper, D. M., Shore, S. N., White, R. E., & Warren, W. H. Jr 1989, *A&AS*, 81, 221
- Adelman, S. J., Caliskan, H., Cay, T., Kocer, D., & Gokmen Tektunali, H. 1999, *MNRAS*, 305, 591
- Beeckmans, F. 1977, *A&A*, 60, 1
- Böhm-Vitense, E., & Johnson, P. 1978, *ApJ*, 225, 514
- Breger, M. 1976, *ApJS*, 32, 7
- Canuto, V. M., & Mazzitelli, I. 1991, *ApJ*, 370, 295
- Canuto, V. M., & Mazzitelli, I. 1992, *ApJ*, 389, 724
- Castelli, F., Gratton, R. G., & Kurucz, R. L. 1997, *A&A*, 318, 841
- Code, A. D., Davis, J., Bless, R. C., & Hanbury Brown, R. 1976, *ApJ*, 203, 417
- Dickens, R. J., & Penny, A. J. 1971, *MNRAS*, 153, 287
- Gulliver, A. F., & Hill, G. 1990, *PASP*, 102, 1200
- Gulliver, A. F., Hill, G., & Adelman, S. J. 1996, in 5th Vienna Symp. on Stellar Atmospheres and Spectrum Synthesis, ed. S. J. Adelman, F. Kupka, & W. W. Weiss (San Francisco, Astron. Soc. Pacific), ASP Conf. Ser., 108, 232
- Gutierrez-Moreno, A., Moreno, H., & Stock, J. 1968, Univ. Chile Dept. Astron. Pub., 8, 127
- Hauck, B., & Mermilliod, M. 1980, *A&AS*, 40, 1
- Hayes, D. S., & Latham, D. W. 1975, *ApJ*, 197, 593
- Hill, G., Fisher, W. A., & Poeckert, R. 1982, *Publ. Dom. Astrophys. Obs. Victoria*, 16, 27
- Lane, M. C., & Lester, J. B. 1980, *ApJ*, 228, 210
- Lemke, M. 1989, *A&A*, 225, 125
- Kubiak, M. 1973, *Acta Astr.*, 23, 23
- Kurucz, R. L. 1979, *ApJS*, 40, 1
- Kurucz, R. L. 1993, Atlas9 Stellar Atmosphere Programs and 2 km/s grid, Kurucz CD-Rom No. 13 (Smithsonian Astrophysical Observatory, Cambridge)
- Kurucz, R. L., & Avrett, E. H. 1981, Smithsonian Astrophysical Observatory Report, 391
- Malagnini, M. L., Morossi, C., Rossi, L., & Kurucz, R. L. 1986, *A&A*, 162, 140
- Moon, T. T., & Dworetsky, M. M. 1985, *MNRAS*, 217, 305
- Napiwotzki, R., Schönberner, D., & Wenske, V. 1993, *A&A*, 268, 653
- Oke, J. B. 1967, *ApJ*, 150, 513
- Ribas, I., Jordi, C., Torra, J., & Gimenez, A. 1997, *A&A*, 327, 207
- Schild, R. E. 1977, *AJ*, 82, 337
- Schild, R., Peterson, D. M., & Oke, J. B. 1971, *ApJ*, 166, 95
- Smalley, B., & Dworetsky, M. M. 1995, *A&A*, 293, 446
- Smalley, B., & Kupka, F. 1997, *A&A*, 328, 349
- Taylor, B. J. 1984, *ApJS*, 54, 256
- Wolff, S. C., Kuhl, L. V., & Hayes, D. S. 1968, *ApJ*, 152, 871