

Erratum

Asteroid (216) Kleopatra

Tests of the radar-derived shape model

D. Hestroffer¹, J. Berthier¹, P. Descamps¹, P. Tanga^{2,*}, A. Cellino², M. Lattanzi²,
M. Di Martino², and V. Zappalà²

¹ IMCCE, UMR CNRS 8028, Observatoire de Paris, 77 Av. Denfert Rochereau, 75014 Paris, France

² Osservatorio Astronomico di Torino (OATo), Strada Osservatorio 20, 10025 Pino Torinese (TO), Italy

A&A, 392, 729–733 (2002), DOI: 10.1051/0004-6361:20021006

Figure 1 has been misprinted so we print it again.




Fig. 1. Comparison between the observed occultation data and the radar model. Left panel: occultation of Oct. 10, 1980 at 7.00 h UTC. Right panel: occultation of Jan. 1, 1991 at 5.28 h UTC. The dashed lines correspond to negative observations. Error bars are given by crosses and are negligible for the photoelectric observations.

Send offprint requests to: D. Hestroffer, e-mail: hestro@bd1.fr

* Associate researcher at the IMCCE. Present address: Laboratoire Cassini, UMR 6529, Observatoire de la Côte d'Azur, BP 4229, 06304 Nice Cedex 4, France.