

*Erratum*

**The absence of the 10  $\mu\text{m}$  silicate feature in the isolated  
Herbig Ae star HD 100453**

G. Meeus<sup>1</sup>, J. Bouwman<sup>2</sup>, C. Dominik<sup>3</sup>, L. B. F. M. Waters<sup>3</sup>, and A. de Koter<sup>3</sup>

<sup>1</sup> Astrophysikalisches Institut Potsdam (AIP), An der Sternwarte 16, 14482 Potsdam, Germany

<sup>2</sup> Service d'Astrophysique, CEA Saclay, 91191 Gif-sur-Yvette, France

<sup>3</sup> Astronomical Institute Anton Pannekoek, University of Amsterdam, Kruislaan 403,  
1098 SJ Amsterdam, The Netherlands

A&A, 392, 1039 (2002), DOI: 10.1051/0004-6361:20020952

In this paper we refer to a 1.2 mm measurement of HD 100453 and quote a flux of 30 mJy. This flux is incorrect. The correct value obtained in the measurement is  $265 \pm 21$  mJy (S. Wolf and M. Albrecht, private communication). When considering this value, the long wavelength slope  $\alpha$  of the SED

is now  $-2.8$  instead of  $-3.7$ . The new value is consistent with large grains. Therefore, the general lack of small silicate grains is the most likely explanation for the weakness of the 10  $\mu\text{m}$  silicate feature in this source. We are grateful to M. Albrecht for reducing and discussing the SIMBA data.