

First NH₃ detection of the Orion Bar[★]

B. Larsson¹, R. Liseau¹, P. Bergman², P. Bernath³, J. H. Black², R. S. Booth², V. Buat⁴, C. L. Curry³, P. Encrenaz⁵, E. Falgarone⁶, P. Feldman⁷, M. Fich³, H. G. Florén¹, U. Frisk⁸, M. Gerin⁶, E. M. Gregersen⁹, J. Harju¹⁰, T. Hasegawa¹¹, L. E. B. Johansson², S. Kwok¹¹, A. Lecacheux¹², T. Liljeström¹³, K. Mattila¹⁰, G. F. Mitchell¹⁴, L. H. Nordh¹⁵, M. Olberg², G. Olofsson¹, L. Pagani⁵, R. Plume¹¹, I. Ristorcelli¹⁶, Aa. Sandqvist¹, F. v. Schéele⁸, N. F. H. Tothill¹⁴, K. Volk⁹, C. D. Wilson⁹, and Å. Hjalmarson²

¹ Stockholm Observatory, SCFAB, Roslagstullsbacken 21, 106 91 Stockholm, Sweden

² Onsala Space Observatory, 439 92 Onsala, Sweden

³ Department of Physics, University of Waterloo, Waterloo, ON N2L 3G1, Canada

⁴ Laboratoire d'Astronomie Spatiale, BP 8, 13376 Marseille Cedex 12, France

⁵ LERMA & FRE 2460 du CNRS, Observatoire de Paris, 61 Av. de l'Observatoire, 75014 Paris, France

⁶ LERMA & FRE 2460 du CNRS, École Normale Supérieure, 24 rue Lhomond, 75005 Paris, France

⁷ Herzberg Institute of Astrophysics, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

⁸ Swedish Space Corporation, PO Box 4207, 171 04 Solna, Sweden

⁹ Department of Physics and Astronomy, McMaster University, Hamilton, ON L8S 4M1, Canada

¹⁰ Observatory, PO Box 14, University of Helsinki, 00014 Helsinki, Finland

¹¹ Department of Physics and Astronomy, University of Calgary, Calgary, ABT 2N 1N4, Canada

¹² LESIA, Observatoire de Paris, Section de Meudon, 5 place Jules Janssen, 92195 Meudon Cedex, France

¹³ Metsähovi Radio Observatory, Helsinki University of Technology, Otakaari 5A, 02150 Espoo, Finland

¹⁴ Department of Astronomy and Physics, Saint Mary's University, Halifax, NS B3H 3C3, Canada

¹⁵ Swedish National Space Board, Box 4006, 171 04 Solna, Sweden

¹⁶ CESR, 9 avenue du Colonel Roche, BP 4346, 31029 Toulouse, France

Received 29 November 2002 / Accepted 1 February 2003

Abstract. Odin has successfully observed three regions in the Orion A cloud, i.e. Ori KL, Ori S and the Orion Bar, in the 572.5 GHz rotational ground state line of ammonia, ortho-NH₃ (J, K) = (1, 0) → (0, 0), and the result for the Orion Bar represents the first detection in an ammonia line. Several velocity components are present in the data. Specifically, the observed line profile from the Orion Bar can be decomposed into two components, which are in agreement with observations in high- J CO lines by Wilson et al. (2001). Using the source model for the Orion Bar by these authors, our Odin observation implies a total ammonia abundance of NH₃/H₂ = 5 × 10⁻⁹.

Key words. ISM: individual objects: Orion A – clouds – molecules – abundances – stars: formation

1. Introduction

The central parts of the Orion A cloud contain several different source regions, among which one can identify Orion KL, Orion S and the Orion Bar (see, e.g., Fig. 3 of Wilson et al. 2001). On angular scales of two arcminutes (the Odin beam size, Sect. 2), the Ori KL region, in particular, harbours a highly complex source structure, apparently with several components along the line of sight. The region has been reviewed by Genzel & Stutski (1989) and more recent results and references can be

found in the papers of Wilson et al. (2001) and of Wiseman & Ho (1998). The latter authors presented VLA-maps in two inversion lines of para-ammonia, showing the emission to arise from finger-like structures. However, as in previous observations (e.g. Ho et al. 1979), the Orion Bar was not detected.

The structure of the NH₃ molecule and its versatility as an astrophysical tool have been discussed by Ho & Townes (1983). These authors also provide an energy level diagram. The rotational ground state line of ammonia, NH₃(J, K) = (1, 0) → (0, 0), has a wavelength of 524 μm and an upper state energy of 28 K above ground. For the (1, 1) and (2, 2) inversion lines at 1.3 cm, the corresponding values are 24 K and 65 K, respectively. The radiative lifetime of the (1, 0) → (0, 0) transition is shorter by orders of magnitude than those of the inversion lines and the line profile could therefore potentially

Send offprint requests to: B. Larsson, e-mail: bem@astro.su.se

[★] Based on observations with Odin, a Swedish-led satellite project funded jointly by the Swedish National Space Board (SNSB), the Canadian Space Agency (CSA), the National Technology Agency of Finland (Tekes) and Centre National d'Études Spatiales (CNES). The Swedish Space Corporation has been the industrial prime contractor.

be probing regions of very different excitation conditions (cf. Table 2 of Liseau et al., this issue). In particular for dynamical studies, the NH₃ line is expected to complement or, in regions of undetectably low water vapour abundance, to substitute for the ground state line of ortho-H₂O ($1_{10} - 1_{01}$), since these resonance lines share the property of being “effectively transparent” up to very high optical depths, as they exhibit an essentially linear growth of the intensity (for τ of several hundred; see, e.g., Snell et al. 2000 and Liseau 2001).

Using the Kuiper Airborne Observatory (KAO), Keene et al. (1983) observed the NH₃ ($1,0 \rightarrow 0,0$) line toward Ori KL and made a five point cross with 1' spacing. Their observations will be compared to those presented in this letter, which have been obtained with Odin (Sect. 2), a spaceborne submillimeter telescope (Frisk et al., Hjalmarsen et al. and Nordh et al., this issue). The reductions of these data, which required some special and careful treatment, are described in detail in Sect. 3 and the results are presented in Sect. 4. The concluding discussion (Sect. 5) of this Letter will focus on the NH₃ emission from the Orion Bar, as this provides an entirely new piece of information for the complex Orion A region.

2. The Odin observations

The NH₃ observations of Orion A were made on September 27 to 29, 2001, simultaneously with observations of the H₂¹⁸O ($1_{10} - 1_{01}$) and O₂ ($1_1 - 1_0$) lines (Olofsson et al. and Pagani et al., this issue). Aiming at Ori KL, the coordinates of the (0,0) position were RA = 5^h35^m15^s.1 and Dec = -5°22'12" (J2000). The absolute pointing of the Odin telescope is presently known to within 30". At 572.5 GHz the half power beam width is 2'. In addition, two positions with the nominal relative offsets in arcmin (0, -2; Ori S) and (+2, -2; Orion Bar) were observed. During the observations, the relative pointing accuracy (rms) was (Δ RA, Δ Dec) \lesssim (7", 14"). The data were obtained in a sky-switching mode (Frisk et al. and Hjalmarsen et al., this issue) by observing a total of 4.5 hr each on-source and on blank sky, with 10 s integrations per individual scan. As back-end, a digital hybrid autocorrelator (AC) was used, with an 800 MHz bandwidth at the resolution of 1 MHz (0.5 km s⁻¹).

3. Data reductions

The phase-lock system for the 572 GHz receiver is not working properly. However, it is reasonably well locked at the tuning frequency and the telluric ozone line O₃ ($J_{K^-,K^+} = 30_{4,26} \rightarrow 30_{3,27}$) 572.9 GHz is frequently observed along parts of the Odin orbit, drifting across the receiver band. The line is sufficiently close in frequency to the ammonia line NH₃ ($J, K = 1,0 \rightarrow 0,0$) 572.5 GHz that it can be used to establish the observing frequency scale, and hence allows us to restore the NH₃ data.

The drift of the O₃ line can be expected to correlate with the equipment temperature and, in particular, inversely with the temperature measured at the local oscillator (LO). That this is indeed the case is demonstrated in the upper panel of

Table 1. Gaussian fit parameters for the observed line profiles.

Offset (arcmin)	Component 1			Component 2		
	$T_{0,1}$ (K)	$v_{\text{LSR},1}$ (km s ⁻¹)	Δv_1 (km s ⁻¹)	$T_{0,2}$ (K)	$v_{\text{LSR},2}$ (km s ⁻¹)	Δv_2 (km s ⁻¹)
(0, 0)	2.0	9.1	4.9	0.9	9.9	16.0
(0, -2)	1.5	8.3	5.5
(+2, -2)	0.35	8.1	5.4	0.35	10.5	3.3

Fig. 1, where the center channel for the O₃ line, C_{O₃} and identified by plus-signs, is shown as a function of the time, expressed as satellite revolution numbers. The solid curve in that panel depicts the relation for the reciprocal LO-temperature, $f(T_{\text{LO}}) \propto 1/T_{\text{LO}}$. Subtraction of the T_{LO} relation from the O₃ channel data points results in the swarm of plus signs in the middle panel of Fig. 1. A polynomial fit to these points is shown by the full drawn curve [= $g(\Delta)$], depicting the temperature behaviour measured at the telescope (main reflector). In the lower panel, the residual displays the final scatter about the fixed central channel for the O₃ line (rms = 0.7 channels = 0.36 km s⁻¹).

The drift corrections to the Orion observations were done by applying the empirically determined relation

$$C_i = C_{i0} - [f(T_{\text{LO}}) + g(\Delta) + C_{\text{O}_3}], \quad (1)$$

where the frequencies are expressed in channel units and where the constant, C_{O₃}, is derived from the ozone line observations. The final spectra are given in the main beam temperature scale, defined as

$$T_{\text{mb}} = \frac{T_A}{\eta_{\text{mb}}} = \frac{1}{\eta_{\text{mb}}} \frac{T_{\text{on}} - T_{\text{sky}}}{T_{\text{sky}}} T_{\text{sys}}, \quad (2)$$

with obvious notations (average $T_{\text{sys}} = 3750$ K). In the upper panel of Fig. 2, the averaged spectral scans (T_A vs. C) toward the three observed positions in Orion are shown, together with polynomial fits to their baselines. Finally, the baseline subtracted spectra (T_{mb} vs. v_{LSR}) are shown in the lower panel. The choice of polynomials is conservative, as a sinusoidal baseline fit to the remaining standing wave pattern would result in more extended wings in especially the (0,0) spectrum.

4. Results

The NH₃ ($J, K = 1,0 \rightarrow 0,0$) spectra observed toward the three positions in Ori A are shown in Fig. 2 (lower panel). The agreement with the KAO data toward Ori KL, also shown in that figure for comparison, is striking, although the signal-to-noise ratio of the Odin data is far better. Keene et al. (1983) speculated whether there existed some low intensity broad component in their data. That this is indeed the case has now been confirmed by Odin. In our observations, the line intensity peaks at the KL position and decreases by about a factor of two at Ori S and by a factor of five toward the Orion Bar. Changes in radial velocity of the line centers are also evident. In the figure, the results of Gaussian fits to the line profiles are also




Fig. 1. Correction for frequency drift during the observation. **Upper:** the plus-signs denote the channel number of the center of the telluric O₃ line during the Orion observations, expressed as Odin revolutions. The upper curve (solid line) depicts the behaviour of the reciprocal LO-temperature. **Middle:** the plus-signs designate the difference between the values shown above. The full-drawn curve is a polynomial fit to these points. This curve describes the average behaviour of the temperature measured at the main reflector. **Lower:** the rms-spread of the residuals to the polynomial fit corresponds to 0.7 of the AC-channel width ($=0.36 \text{ km s}^{-1}$).

shown and their parameters are reported in Table 1. Four distinct velocity systems can be identified: (1) a $v_{\text{LSR}} = 9 \text{ km s}^{-1}$ component of width 5 km s^{-1} (*FWHM*) is present toward Ori KL, (2) a very broad feature ($\Delta v = 16 \text{ km s}^{-1}$), but centered at $v_{\text{LSR}} = 10 \text{ km s}^{-1}$, is also present in Ori KL, (3) a blueshifted line ($v_{\text{LSR}} = 8 \text{ km s}^{-1}$) of width 5.5 km s^{-1} toward the outflow source Ori S and the Orion Bar and (4) a component at $v_{\text{LSR}} = 10.5 \text{ km s}^{-1}$ of width 3.3 km s^{-1} from the Orion Bar, identical to CO ($J = 4 - 3$) (Wilson et al. 2001). The result




Fig. 2. NH₃ (J, K) = (1, 0) \rightarrow (0, 0) spectra toward Orion A. **Upper:** the reduced spectra for the three positions discussed in the text (antenna temperature vs. correlator channel) are shown together with the adopted polynomial fits to their baselines. **Lower:** same as above, but with the baselines subtracted (T_{mb} vs. v_{LSR}). The data are displayed as histograms. Uppermost, the KAO observation ($2'$ *FWHM*) by Keene et al. (1983) (thin line) is shown together with the Odin spectrum toward (0, 0). The much higher S/N of the Odin data aside, the agreement between these two observations is strikingly good. Following below, the spectra toward the three positions observed with Odin are displayed, together with Gaussian fits to the line profiles as thin lines (cf. Table 1). For reference, a vertical dashed line is drawn at $v_{\text{LSR}} = 9.1 \text{ km s}^{-1}$.

for the Orion Bar represents the first detection in an ammonia line.

The radial velocities found in the NH₃ (1, 0) \rightarrow (0, 0) spectra are different from those quoted for various inversion lines toward Ori KL by Hermsen et al. (1988a), but are in good agreement with those observed toward all positions in the ortho-H₂O ($1_{10} - 1_{01}$) 557 GHz line during an independent

observing run (Olofsson et al., this issue). The line shapes of these transitions do agree in the direction of Ori S and the Bar, but at the resolution of 1 MHz, the NH₃ 572.5 GHz line core toward Ori KL is highly symmetric and does not display any self-absorption like feature as does the H₂O 557 GHz line.

5. Discussion and conclusions

For the Orion Bar, no previous NH₃ line information, other than upper limit results, is in existence. For the interpretation of this observation, we will have to rely on source models based primarily on other data. The observed width of the Bar ranges from about 10'' to 60'' for a variety of molecules (e.g., Tielens et al. 1993; Wilson et al. 2001). Based on observations in high- J CO lines, estimates of the physical parameters of the Orion Bar have been given by Wilson et al. (2001). These authors argue that the geometry is that of a 30'' wide rod in the plane of the sky and we assume that their model is applicable also to NH₃. The observed width of the NH₃ and CO lines, $\Delta v = 3.3 \text{ km s}^{-1}$, can be interpreted as a velocity gradient of $45 \text{ km s}^{-1} \text{ pc}^{-1}$. The hyperfine structure of the NH₃ transition (Townes & Schawlow 1955; cf. Liseau et al., this issue) does not significantly contribute to the observed width of the line. The column density of H₂ is given by Wilson et al. (2001) as $N(\text{H}_2) = 2.2 \times 10^{22} \text{ cm}^{-2}$. As kinetic gas temperature we adopt the peak value of their observed CO (7–6) line, $T_k = 145 \text{ K}$.

We use a large velocity gradient code to estimate the statistical equilibrium level populations and to solve the equation of radiative transfer along the line of sight. We have calculated the level energies and Einstein- A values according to Poynter & Kakar (1975), using the dipole moment of Cohen & Poynter (1974). For the computation of the collision rates, we adopted the values provided by Danby et al. (1988) for collisions with para-H₂ ($J = 0$), in conjunction with the assumption of detailed balance for the inverse rates.

For the source model of Wilson et al. (2001), we find the abundance of ortho-ammonia relative to H₂ in the Orion Bar to be $X_o(\text{NH}_3) = 2.5 \times 10^{-9}$, where the NH₃ (1, 0) \rightarrow (0, 0) line radiation temperature is $T_R = 1.7 \text{ K}$, corrected for partial beam filling (~ 0.2). This result for X_o remains unaltered, if the kinetic temperature were higher, e.g., $T_k = 200 \text{ K}$. The model is also in agreement with the upper limits on the inversion lines of Ho et al. (1979) and Wiseman & Ho (1998), provided the ammonia ortho-to-para ratio is not largely different from unity. This would be expected for a gas at such elevated temperatures, since for ammonia the ortho-to-para ratio $\rightarrow g_o/2g_p = 1$ as $T_k \rightarrow \infty$. The high temperature ortho-to-para ratio of 1.0 has

previously been determined for Ori KL by Morris et al. (1973). For the Orion Bar, such ratio would thus imply a total ammonia abundance $X(\text{NH}_3) = 5 \times 10^{-9}$. The ammonia emission region is probably not much narrower than 10'' and average densities do probably not exceed 10^6 cm^{-3} by large, so that it is likely that $X(\text{NH}_3) < 2 \times 10^{-8}$. These estimates can be compared to the values deduced for the Ori KL components “hot core” (10^{-7} – 10^{-6} , Dishoeck 1993; Hermsen et al. 1988b) and “plateau” ($< 10^{-8}$, van Dishoeck et al. 1993). The density of the applied model is lower than the critical density of the (1, 0) \rightarrow (0, 0) transition by more than one order of magnitude and, hence, the line is very subthermally excited ($T_{\text{ex}} = 11 \text{ K}$). For the inferred abundance, the line is moderately optically thick ($\tau = 1$).

The validity of the presented results rests on the assumption that the CO model by Wilson et al. (2001) is applicable to NH₃. The observation of the Orion Bar in the (3, 3) inversion line could provide a test, as this line is predicted to be the strongest NH₃ transition.

References

- Cohen, E. A., & Poynter, R. L. 1974, *J. Mol. Spec.*, 53, 131
 Danby, G., Flower, D. R., Valiron, P., Schilke, P., & Walmsley, C. M. 1988, *MNRAS*, 235, 229
 Genzel, R., & Stutski, J. 1989, *ARA&A*, 27, 41
 Hermsen, W., Wilson, T. L., Walmsley, C. M., & Henkel, C. 1988a, *A&A*, 201, 285
 Hermsen, W., Wilson, T. L., & Bieging, J. H. 1988b, *A&A*, 201, 276
 Ho, P. T. P., Barrett, A. H., Myers, P. C., et al. 1979, *ApJ*, 234, 912
 Ho, P. T. P., & Townes, C. H. 1983, *ARA&A*, 21, 239
 Keene, J., Blake, G. A., & Phillips, T. G. 1983, *ApJ*, 271, L 27
 Liseau, R. 2001, in *The Promise of the Herschel Space Observatory*, ed. G. L. Pilbratt, J. Cernicharo, A. M. Heras, T. Prusti, & R. Harris, ESA SP-460, 313
 Lovas, F. J. 1992, *J. Phys. Chem. Ref. Data*, 21, 181
 Morris, M., Zuckerman, B., Palmer, P., & Turner, B. E. 1973, *ApJ*, 186, 501
 Poynter, R. L., & Kakar, R. K. 1975, *ApJS*, 29, 87
 Snell, R. L., Howe, J. E., Ashby, M. L. N., et al. 2000, *ApJ*, 539, L 93
 Tielens, A. G. G. M., Meixner, M. M., van der Werf, P. P., et al. 1993, *Science*, 262, 86
 Townes, C. H., & Schawlow, A. L. 1955, *Microwave Spectroscopy* (Dover Publications, Inc.)
 Wilson, T. L., Muders, D., Kramer, C., & Henkel, C. 2001, *ApJ*, 557, 240
 Wiseman, J. J., & Ho, P. T. P. 1998, *ApJ*, 502, 676
 van Dishoeck, E. F., Blake, G. A., Draine, B. T., & Lunine, J. I. 1993, in *Protostars and Planets III*, ed. E. H. Levy, & J. I. Lunine, (University of Arizona Press), 163