

Refraction in a pulsar magnetosphere – the effect of a variable emission height on pulse morphology

P. Weltevrede¹, B. W. Stappers^{2,1}, L. J. van den Horn^{1,3}, and R. T. Edwards¹

¹ Astronomical Institute “Anton Pannekoek”, University of Amsterdam, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

² Stichting ASTRON, Postbus 2, 7990 AA Dwingeloo, The Netherlands

³ Institute for Theoretical Physics, University of Amsterdam, Valckenierstraat 65, 1018 XE Amsterdam, The Netherlands

Received 19 May 2003 / Accepted 15 September 2003

Abstract. The Petrova (2000) model to calculate pulse profiles is extended to a variable emission height model to make it physically self-consistent. In this context variable means that the emission height is no longer considered to be the same for different magnetic field lines. The pulse profiles calculated using this new model seem to be less realistic due to a focusing effect and cannot be used to fit (typical) multifrequency pulsar observations. Apart from the focusing effect the general morphology of pulse profiles is not greatly affected by introducing a variable emission height. Additional extensions of the model will be needed to be able to fit observations, and several suggestions are made.

Key words. plasmas – waves – stars: pulsars general

1. Introduction

Arons & Barnard (1986) have derived the dispersion relation for three wave modes which can propagate through the plasma of a pulsar magnetosphere: the ordinary subluminescent mode (subluminescent O-mode), the ordinary superluminescent mode (superluminescent O-mode) and the extraordinary mode (X-mode). The X-mode does not suffer refraction, but refraction of the subluminescent O-mode can be considerable in pulsar magnetospheres (Barnard & Arons 1986). The subluminescent O-mode cannot escape the pulsar magnetosphere due to Landau damping, so it does not contribute directly to the observed emission. Lyubarskii (1996) has shown that the subluminescent O-mode can be converted into the superluminescent O-mode – which can escape the magnetosphere – by induced scattering off plasma particles. As pointed out by Barnard & Arons (1986) refraction of the superluminescent O-mode is less severe than for the subluminescent O-mode. It can, however, be important in the presence of a transverse plasma density gradient.

For the superluminescent O-mode Petrova (2000) (hereafter P2000) shows how pulse profiles can be calculated taking into account the transverse plasma density gradient. This model demonstrated that complex profiles can be produced by a “simple” ring-shaped emission region (as predicted by Ruderman & Sutherland 1975), and thus that the wealth of observed pulse profile shapes may be due to different magnetospheric conditions rather than more complex emission region-shapes.

Furthermore it was shown that the observed phenomenon of high frequency core splitting could be an effect of refraction.

The emission height is an important ingredient in calculating pulse profiles. The emission height is frequency dependent; i.e. there is radius-to-frequency mapping (Cordes 1978). Plasma waves with higher frequencies are excited closer to the star. The observed frequency dependence of pulse profiles is often very complex, perhaps more complex than can be expected from just radius-to-frequency mapping. Because refraction itself is a frequency dependent phenomenon, a more complex frequency dependence of pulse profiles can be expected if refraction is important in pulsar magnetospheres. Other effects that can be understood by taking into account refraction are the occurrence of orthogonal polarization modes (Petrova 2001) and the spectral breaks of pulsars (Petrova 2002).

To link the observed pulse profiles to the shape of the emission region, so as to be able to check emission theories, one must know the refractive properties of pulsar magnetospheres. This calls for the development of improved refraction models. As noted by P2000, the emission surface at one observing frequency should be, strictly speaking, an isodensity surface of the plasma distribution. Yet, for simplicity, a constant emission height (CEH) was assumed in P2000, in the expectation that the qualitative features of profile formation would not be sensitive to that assumption. In the present paper we do adopt a surface of constant density as required for self-consistency of the refractive model, and we investigate the effects on the pulse morphology. This “variable” emission height (VEH) appears to introduce a focusing effect which causes the profiles to have unrealistically sharp edges. As a consequence, the VEH

Send offprint requests to: P. Weltevrede,
e-mail: wltvrede@science.uva.nl




Fig. 1. The ray at position (r, χ) is propagating in the direction θ . This ray was emitted at (r_0, χ_0) and the field line through this point is indicated by χ_s . The plasma density peaks at the characteristic field lines indicated by χ_c . The angles χ_c and χ_s are defined at $r = 1$ and the angles χ_0 and θ_0 at the emission height r_0 .

model cannot be used to fit multifrequency pulsar observations without relaxing additional restrictive assumptions, a number of which are discussed at the end of the paper.

2. Refraction model

2.1. The ray equations

The refraction model below is essentially that of P2000, and we refer to that paper for details. The plasma distribution and the magnetic field are assumed to be axisymmetric around the magnetic pole, so the refraction model can be described in two dimensions. A position on a ray trajectory is indicated by the polar coordinates r and χ , and the direction along the trajectory by θ (see Fig. 1).

The geometrical optics description applies and the time evolution of these quantities is given by the Hamilton equations. For a highly magnetized ultrarelativistic electron-positron plasma, which is cold in the proper restframe, the dispersion relation has been derived by Arons & Barnard (1986) and the associated Hamilton equations by Barnard & Arons (1986). On the condition that the plasma flows with the same velocity for all field lines and when rays are emitted parallel to the local (dipolar) magnetic field, the dispersion law describing the two ordinary wave modes can be written as (P2000)

$$\eta \left(1 - \frac{4N}{f_0^2 (1 + \eta)^2} \right) - \frac{9}{4} \chi_0^2 \gamma^2 (\theta_n - \chi_n)^2 = 0, \quad (1)$$

while from the Hamilton equations one finds

$$\begin{aligned} r \frac{d\chi_n}{dr} &= \frac{\chi_n}{2} + \frac{3(1 + \eta)^3 (\theta_n - \chi_n)}{A_1} \\ r \frac{d\theta_n}{dr} &= \frac{N}{A_1 f_0^2} \left[6(1 - \eta) (\theta_n - \chi_n) - A_2 \frac{\partial \ln N}{\partial \chi_n} \right], \end{aligned} \quad (2)$$

where

$$\begin{aligned} A_1 &= (1 + \eta)^3 - \frac{4(1 - \eta)N}{f_0^2} \\ A_2 &= \frac{4\eta(1 + \eta)}{3\chi_0^2 \gamma^2}. \end{aligned}$$

The radial plasma density derivative has been omitted, because it can be neglected for the plasma density we will adopt (P2000). The parameter η is related to the component of the wave vector \mathbf{k} in the direction of the local magnetic field and is defined as

$$\eta = 2\gamma^2 (1 - n_{\parallel}) \quad (3)$$

with $\gamma \gg 1$ the Lorentz factor of the outflowing plasma, and $n_{\parallel} = ck_{\parallel}/\omega$ where ω is the frequency of the plasma wave. The refractive index is $n = (n_{\parallel}^2 + n_{\perp}^2)^{1/2}$, where parallel/perpendicular is with respect to the local magnetic field. It is assumed that n_{\parallel} is such that $\eta \ll 2\gamma^2$. The plasma waves are assumed to be generated close to the local Lorentz-shifted plasma frequency $\omega_p \sqrt{\gamma}$,

$$\omega_p = \sqrt{\frac{4\pi N_p e^2}{m}}, \quad (4)$$

with e the electron charge, m the electron rest mass and N_p the particle number density (electrons plus positrons) of the plasma. The ratio

$$f = \frac{\omega}{\omega_p \sqrt{\gamma}} \quad (5)$$

should then be close to unity.

Equations (1) and (2) are *normalized*, i.e. the coordinates r , χ_n and θ_n , as well as the plasma number density distribution N , are normalized to their values at the emission height (so $\chi = \chi_0 \chi_n$ and $\theta = \theta_0 \theta_n$). The emission height can be different for different rays as will be discussed in Sect. 2.3, so in this context *the* emission height is the emission height of the particular ray that is being considered. The values of f and χ at the emission height are denoted as f_0 and χ_0 respectively. All angles are assumed to be small compared to 1 throughout this paper, so the propagation direction of the plasma waves should always be nearly parallel to the magnetic axis.

For the superluminal O-mode which is considered here, n_{\parallel} cannot be larger than 1, therefore η is required to be positive (or zero). At the emission height (where $\theta_n = \chi_n = N = 1$) the solution of the dispersion relation follows immediately and we have for the superluminal O-mode

$$\eta_0 = \begin{cases} 2/f_0 - 1 & \text{for } 0 < f_0 < 2 \\ 0 & \text{for } f_0 \geq 2 \end{cases} \quad (6)$$




Fig. 2. Rays (dashed) with the observing frequency are generated at the emission height and are refracted toward lower plasma densities. The scale is in units emission height and the gray scale indicates the hollow cone plasma density. The plasma density is proportional to the numbers of the gray scale bar. The plots on the left are for the CEH model, the plots on the right are for the VEH model, the top plots are calculated with $\chi_c = 0.03$ and the bottom plots with $\chi_c = 0.01$. The other input parameters are: $\gamma = 30$, $f_0 = 0.5$, $\varepsilon_1 = 3$, $\varepsilon_2 = 4$.

Note that η_0 is continuous at $f_0 = 2$. The solution of the dispersion relation applicable above the emission height is given by the general solution of the cubic (A.1).

Equations (1) and (2) describe, to first order in χ_n and θ_n , the refraction of an ordinary (both the sub- and superluminal) plasma wave. The two differential equations for $\chi_n(r)$ and $\theta_n(r)$ can be solved numerically if $\eta(r)$ is known. As noted above, η can be calculated analytically from the dispersion equation. We use a fourth order Runge-Kutta method with adaptive stepsize control (Press et al. 1986) to solve the set of differential equations. For the plasma density distribution we adopt the hollow cone model (P2000)

$$N_p = \frac{N_\star}{r^3} \exp\left(-\varepsilon \left(\frac{|\chi| - \chi_c \sqrt{r}}{\chi_c \sqrt{r}}\right)^2\right), \quad (7)$$

where N_\star is the particle number density at ($r = 1, \chi = \chi_c$). This plasma density is Gaussian shaped around a “characteristic field line” indicated by χ_c . The decrease of the plasma density may be different for the inner and outer regions, so we set

$$\varepsilon = \begin{cases} \varepsilon_1 & \text{for } |\chi| \leq \chi_c \sqrt{r} \\ \varepsilon_2 & \text{for } |\chi| \geq \chi_c \sqrt{r} \end{cases}. \quad (8)$$

Equations (2) do not contain r_0 , so the whole problem is independent of the scaling of r . However, in Eq. (7) χ_c is defined at $r = 1$. The normalized plasma density is given by

$$N(\chi, r) = \frac{N_p(\chi, r)}{N_p(\chi_0, r_0)} \quad (9)$$

and its derivative with respect to χ_n is

$$\frac{\partial \ln N}{\partial \chi_n} = -2\varepsilon \chi_0 \frac{\chi - \chi_c \sqrt{r} \operatorname{sign}(\chi)}{(\chi_c \sqrt{r})^2}. \quad (10)$$

The parameters required to calculate a single ray trajectory are those of the plasma density distribution (χ_c , ε_1 and ε_2), the plasma outflow Lorentz factor γ , the frequency of the plasma wave (expressed in f_0) and the start position χ_0 of the ray. Solving Eqs. (1) and (2) with the start condition $\chi_n = \theta_n = 1$ will give the ray trajectory.

Plasma waves are refracted toward lower plasma densities in the magnetosphere until refraction becomes inefficient due to the decreasing plasma density along its trajectory. As can be seen in Fig. 2 refraction results in a redistribution of rays; i.e. the rays are no longer equi-spaced above the emission height and two “conal components” of outer rays and a “core component” of inner rays are formed.

2.2. Calculation of the pulse profiles

The effect of refraction is quantified in a plot of the final ray direction (Fig. 3). Here the propagation direction θ_f at a height where refraction has become inefficient is plotted versus χ_s . The initial ray position (r_0, χ_0) corresponds to a value of χ_s by tracing back the field line from the emission height to $r = 1$ (see Fig. 1).

Both inside and outside the characteristic field line cone refraction is toward lower plasma densities, which results in a




Fig. 3. The final propagation direction θ_f versus χ_s for both the VEH and the CEH model, for two values of χ_c . In these plots χ_s ranges from $\chi_c/5$ to $5\chi_c/2$ and the other parameters are the same as for Fig. 2. The final propagation direction θ_f is in radians and χ_s in units χ_c .

steepening of the final ray direction plot near $\chi_s = \chi_c$. Inside the plasma cone the rays are refracted toward the magnetic axis and the innermost rays may even intersect the magnetic axis; in that case the final ray direction plot crosses the line $\theta_f = 0$.

For small values of θ_f , rays originating from several discrete values of χ_s leave the magnetosphere in the same direction and at the corresponding pulse longitude different parts of the emission ring can be observed simultaneously. Note that the final ray direction curve for the opposite half of the emission ring is found by mirroring the curve with respect to the line $\theta_f = 0$. If the final ray direction plot crosses this line, some parts of both sides of the emission ring have the same θ_f . In that case both sides of the emission ring can be observed simultaneously, if the impact angle β is small enough.

If the curve in Fig. 3 is horizontal at the θ_f value corresponding to the line of sight (θ_{LOS}), a large part of the emission surface is observed simultaneously while if the curve is steep only a small part is observed. This means that the observed intensity in the pulse profile is proportional to the value of $d\chi_s/d\theta_f$ at $\theta_f = \theta_{\text{LOS}}$ which is just an energy conservation argument (P2000).

Apart from refraction effects the pulse profile will depend on the intensity distribution at the emission height. If the pair production is somehow related to the observed coherent microwave radiation (Ruderman & Sutherland 1975), then similar distributions for the plasma density and the intensity at the emission height can be expected such as (P2000)

$$W_{r_0} = \exp\left(-\varepsilon\Upsilon\left(\frac{|\chi_s| - \chi_c}{\chi_c}\right)^2\right). \quad (11)$$

This corresponds to an emission ring which peaks at the characteristic field lines and its thickness is set by Υ . For $\Upsilon = 1$ the intensity distribution follows exactly the plasma density distribution. The shape is Gaussian as a function of the field line parameter χ_s , but the choice of another parameter (such as the length along the emission surface) is also conceivable. However for simplicity the parameter χ_s is used. As will be discussed later on, the conclusions do not depend on this choice.

The refraction model is axisymmetric around the magnetic axis, so the beam-pattern of the pulsar is also axisymmetric around the same axis. The shape of observed pulse profiles depends on how the line of sight cuts the magnetic pole of the star. We only consider the most simple geometry; i.e. the magnetic axis is orthogonal to the rotation axis ($\alpha = 90^\circ$) and the line of sight cuts the magnetic pole centrally (impact angle $\beta = 90^\circ$).

For this geometry the pulse longitude ϕ is equal to the final ray direction θ_f . Because of this choice of geometry and the axisymmetry, all the information of the beam-pattern is in the calculated pulse profiles. The model itself is independent of α and β , only the mapping between ϕ and θ_f changes.

2.3. Variable emission height model

The model described above may be applied with both a constant (CEH) and a variable emission height (VEH), but (as we will argue) a VEH is needed to make the model self-consistent. The requirement of a VEH was not met in P2000; it is the basic conceptual difference between the model presented here and the P2000 model. Its effect on the pulse profiles turns out to be appreciable, as discussed below.

The emission height can be derived when a plasma density distribution has been specified. The plasma density decreases as r^{-3} , so the local plasma frequency decreases away from the star resulting in the excitation of plasma waves with higher frequencies closer to the star. This results in rays propagating in a direction which is more aligned with the magnetic axis at the emission height. But there is another effect involved in the frequency dependence of the pulse profile morphology: refraction becomes more prominent.

The assumption that both f_0 and γ are constant implies that plasma waves of one particular frequency are generated at one particular equi-plasma density surface (Eqs. (4) and (5)). Because a transverse plasma density gradient is needed for refraction, the emission height of a given frequency varies with polar angle χ_0 . If the magnetic field is dipolar, we have

$$\chi_0 = \chi_s \sqrt{r_0}, \quad (12)$$

where χ_s is shown in Fig. 1. Combining the plasma density (Eq. (7)) with Eqs. (4) and (5) at the emission height, and using Eq. (12) leads to the following expression for the emission height

$$r_0^3 = R^3 \exp\left(-\varepsilon\left(\frac{|\chi_s| - \chi_c}{\chi_c}\right)^2\right) \\ R^3 = \frac{4\pi\gamma e^2 f_0^2 N_\star}{m\omega^2}. \quad (13)$$

Because γ and f_0 are assumed to be constant, R should be constant and r_0 is not constant. As noted earlier, the whole problem is independent of the scale of r , so R can be set equal to 1.




Fig. 4. Pulse profiles for different χ_c (a small χ_c corresponds to a high observing frequency) for both the CEH and the VEH model. For the top row χ_s ranges from $\chi_c/2$ to $3\chi_c/2$ with $\Upsilon = 0$ (all field lines having equal intensity) and for the bottom row χ_s ranges from $\chi_c/5$ to $5\chi_c/2$ with $\Upsilon = 1$ (intensity coupled to the plasma density). The other parameters are the same as in Figs. 2 and 3. The scale is such that the integrated intensity is the same for all profiles.

The frequency dependence of the pulse profiles is then in the parameter χ_c (P2000)

$$\chi_c \propto \omega^{-1/3}. \quad (14)$$

The ray trajectory is solved as a function of the distance to the star, expressed in units of the emission height and different χ_c correspond to relative observing frequencies. The physical emission height can be calculated from Eq. (13) when N_\star and ω are specified.

The emission surface specified by Eq. (13) corresponds to an isodensity surface, so the plasma density distribution has a more prominent role in this VEH model than in the CEH model. Apart from causing refraction, it also determines the shape of the emission surface.

Refraction becomes more severe for the inner and outer rays in the VEH model, because the plasma gradients are larger at lower emission heights. Moreover a lower emission height implies that the rays are emitted closer to, and are initially propagating more aligned with the magnetic axis. For the inner rays this means that the rays can intersect the magnetic axis more easily. For the outer rays there are two counteracting effects. A lower emission height implies that the rays are refracted in a more outward direction, but at the same time the rays are also emitted more aligned to the magnetic axis.

3. Results

Model calculations of pulse profiles for both a VEH and a CEH are presented in Fig. 4 for the most simple geometry ($\beta = 0^\circ$ and $\alpha = 90^\circ$). For this geometry the pulse longitude ϕ is equal to the final ray direction θ_f .

Observationally the core component behaves differently from conal components, both in the frequency dependence of its morphology and in its polarization properties. This is what can be expected from refraction (P2000), because the core

component consists of “mixed” rays; i.e. the order of the beams changes. This is true for both a VEH and a CEH.

In Fig. 2 one can see that refraction becomes more prominent for higher frequencies (lower χ_c). This can also be seen in Fig. 3 where the final propagation direction of rays versus χ_s is plotted. The curve becomes more complex for lower χ_c . Besides this refractive effect, a lower emission height implies smaller propagation angles θ at the emission height, resulting in narrower pulse profiles with increasing frequency (decreasing χ_c) in Fig. 4. This is again true for both a VEH and a CEH.

The pulse profiles in Fig. 4 for a CEH are more spiky than the pulse profiles presented in P2000. The main reason for this is the higher resolution of the calculations presented here.

There are three reasons why the profiles, for both a VEH and a CEH, are spiky. First of all, if the intensity distribution at the emission height is flat, the emission ring has sharp edges resulting in sharp edges in the pulse profile. This effect can be reduced by making Υ larger (see Eq. (11)), as is seen in the bottom row of Fig. 4. Making the parameter Υ larger results in the edge of the intensity distribution becoming Gaussian blurred.

The second effect is caused by rays crossing the magnetic axis, so this applies especially to the core component. This means that at certain pulse longitude both sides of the emission ring can be seen simultaneously. When the number of sides visible changes at a particular pulse longitude, a step in intensity appears in the pulse profile. This effect can again be reduced by increasing Υ as can be seen in the bottom row of Fig. 4.

The last effect contributing to the spikiness of the profiles is a focusing effect. If a large patch of the emission ring is focused at one pulse longitude, a peak is observed. This focusing effect corresponds to a horizontal part in the final ray direction curve; rays emitted at a range of χ_s are focused to a single θ_f . At high frequencies this can be seen for the innermost rays (Fig. 3), causing peaks in the core component.




Fig. 5. Pulse profiles calculated for the case of no refraction. For the top row χ_s ranges from $\chi_c/2$ to $3\chi_c/2$ with $\Upsilon = 0$ and for the bottom row χ_s ranges from $\chi_c/5$ to $5\chi_c/2$ with $\Upsilon = 1$. These profiles have been calculated for $\chi_c = 0.02$. The other parameters and the normalization of the profiles are the same as in Fig. 4.

By comparing the pulse profiles for the VEH and the CEH model in Fig. 4, the most striking difference is the conal components. The edges of the VEH profiles are very sharp, and introducing a large Υ will not reduce their sharpness. The reason can be found in Fig. 3. The curves for the VEH model show a global maximum at $\chi_s \approx 1.5$. Because it is a maximum, there is focusing and because the maximum is global, the peaks occur at the edges of the profile.

As discussed in Sect. 2.3, there are two counteracting effects for the outer rays in the VEH model. A lower emission height makes the outward directed refraction stronger, but the propagation angle is more aligned with the magnetic axis at the emission height. The reason for the global maximum is that the latter effect dominates for the outermost rays.

This focusing effect due to the variable emission height is also visible in the pulse profiles of Fig. 5, which were calculated without using refraction. The focusing is caused by the geometry of the emission surface, not by the intensity distribution at the emission height (Eq. (11)). This means that this focusing is independent of the precise form of Eq. (11), and therefore also of the choice to use the field line parameter χ_s instead of for example the length along the emission surface in this equation.

The edges of the profiles are produced by rays emitted from $\chi_s \approx 1.5$ and the rays are focused, so there should be only very little radiation produced at $\chi_s \approx 1.5$ to avoid the sharp edge. This means that the emission ring should be very thin, so Υ

should be large. At high frequencies Υ should be at least ≈ 5 and at the lowest frequencies ($\chi_c = 0.03$) Υ should be at least ≈ 15 . A large Υ physically means that only the middle part of the emission ring is producing coherent microwave radiation, although the whole ring is producing streams of particles. Such a scenario is in conflict with the expectation that pair production and coherent emission are related.

A VEH leads to stronger refraction. Besides introducing a VEH refraction can also be increased by changing the values of ε_1 , ε_2 , f_0 or γ in the CEH model. Experimenting with a range of values of these parameters did not lead to the formation of the sharp edge of the profiles with a CEH. Therefore the focusing effect is a typical property of the VEH model.

4. Discussion

Contrary to the expectation expressed in P2000, the qualitative features of profile formation turn out to be different for the VEH and the CEH refraction models. Although the VEH is a physical improvement in the sense that it makes the emission model self-consistent, the profiles obtained are less realistic. The model, therefore, needs further improvements before it can serve as a tool to fit (typical) multifrequency pulsar observations.

The most pronounced difference between the CEH and the VEH model is that for the VEH model the rays emitted at the outside of the emission surface do not form the edges of the pulse profile. The edges of the pulse profiles in the VEH model are generated by a focusing effect causing the edges to be sharp. If the thickness of the emission ring at the emission height were much thinner than the thickness of the plasma cone at the emission height the sharp edges would disappear, but this seems physically unrealistic.

It must be noted that the results depend strongly on the plasma distribution adopted. The density profile not only causes refraction, but it also determines the shape of the emission surface. If the plasma density falls off more slowly than the Gaussian distribution assumed here, the results may be more realistic although in that case refraction will be less prominent.

Several other effects could contribute to smoother pulse profiles. There are probably more frequencies generated at one point in the magnetosphere, so there would be a f_0 range rather than a fixed f_0 value. Also the rays are not emitted strictly aligned with the magnetic field lines, rather there will be an elementary beam pattern of finite angular width. A beam pattern with a width of γ^{-1} can be considerable compared with the pulse width (for a plasma outflow Lorentz factor $\gamma \approx 30$ the beam is about 2° wide). If the outflow Lorentz factor is different for different field lines, the shape of the emission surface is changed. Moreover refraction becomes more complex, because it depends on gradients of the γ factor as well as gradients in the plasma density (Barnard & Arons 1986).

Appendix A: Analytical solution of the normalized dispersion relation

The normalized dispersion relation (1) is of the third degree in η , so the analytical solution of η is given by the cubic.

The solution for the superluminal O-mode (the solution with positive η at the emission height) is

$$\eta = s_+ + s_- - \frac{a_2}{3}, \quad (\text{A.1})$$

where

$$\begin{aligned} s_{\pm} &= \left(r \pm \sqrt{q^3 + r^2} \right)^{1/3} \\ q &= \frac{a_1}{3} - \frac{a_2^2}{9} \\ r &= \frac{1}{6}(a_1 a_2 - 3a_0) - \frac{a_2^3}{27} \end{aligned} \quad (\text{A.2})$$

and

$$\begin{aligned} a_2 &= 2 + a_0 \\ a_1 &= 1 - 4 \frac{N}{f_0^2} + 2a_0 \\ a_0 &= -\frac{9}{4} \chi_0^2 \gamma^2 (\theta_n - \chi_n)^2. \end{aligned} \quad (\text{A.3})$$

Acknowledgements. We thank Svetlana Petrova for making available her code and for her valuable comments as referee of this article. We also thank her as well as Joeri van Leeuwen, Ramachandran and John Barnard for constructive discussions.

References

- Arons, J., & Barnard, J. J. 1986, ApJ, 302, 120
 Barnard, J. J., & Arons, J. 1986, ApJ, 302, 138
 Cordes, J. M. 1978, ApJ, 222, 1006
 Lyubarskii, Y. E. 1996, A&A, 308, 809
 Petrova, S. A. 2000, A&A, 360, 592
 Petrova, S. A. 2001, A&A, 378, 883
 Petrova, S. A. 2002, A&A, 383, 1067
 Press, W. H., Flannery, B. P., Teukolsky, S. A., & Vetterling, W. T. 1986, Numerical Recipes: The Art of Scientific Computing (Cambridge: Cambridge University Press)
 Ruderman, M. A., & Sutherland, P. G. 1975, ApJ, 196, 51