

## Serpens X-1 observed by *INTEGRAL* ★

N. Masetti<sup>1</sup>, L. Foschini<sup>1</sup>, E. Palazzi<sup>1</sup>, V. Beckmann<sup>2,3</sup>, N. Lund<sup>4</sup>, S. Brandt<sup>4</sup>, N. J. Westergaard<sup>4</sup>,  
L. Amati<sup>1</sup>, E. Caroli<sup>1</sup>, S. Del Sordo<sup>5</sup>, G. Di Cocco<sup>1</sup>, P. Durouchoux<sup>6</sup>, R. Farinelli<sup>7</sup>, F. Frontera<sup>1,7</sup>,  
M. Orlandini<sup>1</sup>, and A. Zdziarski<sup>8</sup>

<sup>1</sup> Istituto di Astrofisica Spaziale e Fisica Cosmica - Sezione di Bologna, CNR, via Gobetti 101, 40129 Bologna, Italy  
e-mail: masetti@bo.iasf.cnr.it

<sup>2</sup> NASA Goddard Space Flight Center, Code 661, Greenbelt, MD 20771, USA

<sup>3</sup> Joint Center for Astrophysics, Department of Physics, University of Maryland, Baltimore County, MD 21250, USA

<sup>4</sup> Danish Space Research Institute, 2100 Copenhagen 0, Denmark

<sup>5</sup> Istituto di Astrofisica Spaziale e Fisica Cosmica, Sezione di Palermo, CNR, via La Malfa 153, 90146 Palermo, Italy

<sup>6</sup> CEA Saclay, DSM, DAPNIA, Service d'Astrophysique, 91191 Gif-sur-Yvette Cedex, France

<sup>7</sup> Dipartimento di Fisica, Università di Ferrara, via Paradiso 12, 44100 Ferrara, Italy

<sup>8</sup> N. Copernicus Astronomical Center, Bartycka 18, 00-716 Warsaw, Poland

Received 5 February 2004 / Accepted 11 May 2004

**Abstract.** Here we report results of an *INTEGRAL*-AO1 observation of the X-ray burst and atoll source Ser X-1 performed in May 2003. The object was observed for a total on-source time of 400 ks but nearly 8° off-axis due to its amalgamation with an observation of SS 433, the pointing target source. Ser X-1 has been clearly detected up to 30 keV with unprecedented positional accuracy for high-energy emission. The 20–30 keV light curve showed substantial variability during the observation. Comparison with previous observations indicates that the source was in its high (“banana”) state and displayed a soft spectrum during the *INTEGRAL* pointing. A (non simultaneous) radio-to- $\gamma$ -rays broad-band spectral energy distribution is also presented for the first time and discussed.

**Key words.** X-rays: binaries – X-rays: individuals: Ser X-1 – stars: neutron

### 1. Introduction

Low Mass X-ray Binaries (LMXBs) form the most populated class of Galactic X-ray binaries. They are X-ray binary systems in which the mass donor star is less than about  $1 M_{\odot}$  and loses matter via Roche-lobe overflow to a very compact accretor. Matter forms an accretion disk as it falls towards the compact object. Many of these systems contain a weakly magnetized neutron star (NS) and are persistently bright in X-rays.

The study of spectral and temporal properties of NS LMXBs in the X-ray energy band below 20 keV has shown the existence of two subclasses of sources. These were identified by their difference in the hardness ratio from two adjacent X-ray bands as a function of the source intensity in the total energy band (hardness-intensity diagram) and/or their track on an X-ray “colour–colour” diagram (CD) constructed by using two hardness ratios using different energy bands. These two subclasses were labeled as Z sources, if they track a Z-shape,

or part of it, in the CD and atoll sources if their position in the CD draws an atoll-shaped figure. Indeed, Z sources are in general permanently bright, while atoll sources show flux variations by factors ranging from about 5 to about 1000 or more and, in parallel, substantial spectral variations (e.g., Piraino et al. 1999). Also, the timing behaviour is different between the two subclasses; see e.g. van der Klis (2000) for details on this classification and on the properties of these subclasses. However, recent results indicate that atoll sources actually track a Z shape on year-long timescales (Gierliński & Done 2002; Munro et al. 2002).

Ser X-1, or 4U 1837+04, has been classified as an atoll source (e.g., Liu et al. 2001). Archival *EXOSAT* data (Seon & Min 2002) showed this object in the “banana” (i.e. high luminosity) state during the observations. More recent *BeppoSAX* and *RXTE* pointings (Oosterbroek et al. 2001) caught the source while it was again in a high activity state, with an unabsorbed flux (1–200 keV) of  $8.0 \times 10^{-9}$  erg cm<sup>-2</sup> s<sup>-1</sup>. We refer the reader to the work of these authors for the most recent description of the broadband X-ray spectral and timing properties of this source. Ser X-1 has never been seen in the hard (“island”) state that is observed in atoll sources at low luminosities (Hasinger & van der Klis 1989).

\* Based on observations with *INTEGRAL*, an ESA project with instruments and science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland, and with the participation of Russia and the USA.

This source also exhibits Type I X-ray bursts (e.g., Lewin et al. 1995); moreover, during 2001, *BeppoSAX* pinpointed a very long ( $\sim 4$  h) X-ray burst (Cornelisse et al. 2002), making this source join the group of “superbursters” (see Kuulkers 2003, for a review). By studying X-ray bursts observed with *Einstein*, Christian & Swank (1997) deduced a distance to the source of 8.4 kpc. This implies a 1–200 keV luminosity of  $6.7 \times 10^{37}$  erg s $^{-1}$  during the *BeppoSAX* observation of Oosterbroek et al. (2001), which means roughly one third of the Eddington luminosity for a NS.

The optical counterpart to Ser X-1, located in a crowded stellar field (Thorstensen et al. 1980), was correctly identified by Wachter (1997) and, subsequently, spectroscopically confirmed and studied by Hynes et al. (2004). Very recently, the radio counterpart was discovered with the VLA (Migliari et al. 2004).

In this paper we report and discuss the results of an observation of Ser X-1 performed with the INTERnational GAMMA-RAY Laboratory (*INTEGRAL*; Winkler et al. 2003) during AO1 on May 2003, i.e. less than 7 months after the launch of this spacecraft. This paper is thus structured as follows: Sect. 2 describes the observations and the data analysis, Sect. 3 reports the results of the *INTEGRAL* pointing as well as the construction of a broadband spectral energy distribution (SED) for Ser X-1 using literature data along with the high-energy ones presented here; then, in Sect. 4 a discussion will be given.

## 2. Observations and data analysis

The observations presented here were acquired during revolutions 67 to 69, i.e. between 12:00 UT of 3 May 2003 and 09:26 UT of 9 May 2003, for a total on-source time of 400 ks, with the IBIS (Ubertini et al. 2003) and SPI (Vedrenne et al. 2003) instruments onboard *INTEGRAL*. These detectors allow an actual 20 keV–10 MeV spectral coverage altogether. In particular, the IBIS telescope is composed of two detector layers: ISGRI (Lebrun et al. 2003), optimized for the energy range 20–200 keV, and PICsIT (Di Cocco et al. 2003; Labanti et al. 2003), covering the range from 175 keV to 10 MeV. Data were acquired with a spacecraft rectangular  $5 \times 5$  dithering pattern mode.

The field of Ser X-1 was also intermittently observed with Unit 2 of the coded-mask X-ray monitor JEM-X on board *INTEGRAL* (Lund et al. 2003). During the period in which the observation reported here was performed, Unit 1 of JEM-X was not available, being turned off to avoid fast degradation of the detector. The present *INTEGRAL* observation of Ser X-1 was amalgamated with one targeting SS433 (Cherepashchuk et al. 2003), thus leaving Ser X-1 about 7:8 off axis; hence, JEM-X could only observe this object for a small fraction of the total 400 ks exposure time. All JEM-X observations of Ser X-1 were at an off-axis angle larger than 3:6; we only considered JEM-X pointings in which the source was less than 4:5 off axis, because of uncertainties in the knowledge of the instrument response at larger offset angles. Thus, JEM-X data from 22 pointings, corresponding to about 40 ks of observation time, have been used in the present analysis.

**Table 1.** Fluxes and luminosities for Ser X-1 measured during the present *INTEGRAL* observation. For each measurement, the instrument and the energy band are reported. Quoted errors are at  $1\sigma$  confidence level, while upper limits are at  $3\sigma$  level. Luminosities are computed assuming a distance  $d = 8.4$  kpc.

Energy range (keV)	<i>INTEGRAL</i> instrument	Flux (erg cm $^{-2}$ s $^{-1}$ )	Luminosity (erg s $^{-1}$ )
3–5	JEM-X	$(2.2 \pm 0.2) \times 10^{-9}$	$1.9 \times 10^{37}$
5–10	JEM-X	$(1.76 \pm 0.15) \times 10^{-9}$	$1.5 \times 10^{37}$
10–15	JEM-X	$(0.44 \pm 0.10) \times 10^{-9}$	$3.7 \times 10^{36}$
15–35	JEM-X	$< 1 \times 10^{-9}$	$< 8.4 \times 10^{36}$
20–30	ISGRI	$(6.2 \pm 0.3) \times 10^{-11}$	$5.2 \times 10^{35}$
20–30	SPI	$< 5.0 \times 10^{-11}$	$< 4.2 \times 10^{35}$
30–40	ISGRI	$< 6 \times 10^{-12}$	$< 5 \times 10^{34}$
30–40	SPI	$< 2.6 \times 10^{-11}$	$< 2.2 \times 10^{35}$
220–280	PICsIT	$< 1 \times 10^{-10}$	$< 8 \times 10^{35}$

Ser X-1 was outside the field of view of the fourth instrument carried by *INTEGRAL*, i.e. the optical camera OMC (Mas-Hesse et al. 2003), and therefore no data collected with OMC were available for this source during the present *INTEGRAL* pointing.


The data reduction and analysis was performed with the standard Offline Scientific Analysis (OSA v3.0<sup>1</sup>), whose algorithms are described in Goldwurm et al. (2003a) for IBIS and in Diehl et al. (2003) for SPI. The count rates extracted from the standard pipeline were normalized to those measured during off-axis observations of the Crab, in order to obtain fluxes in terms of physical units. Given the present uncertainties in the flux evaluation for off-axis sources (see Goldwurm et al. 2003b), we adopted a procedure similar to that explained by Goldoni et al. (2003) to analyse the IBIS data for off-axis sources. For JEM-X the analysis was performed using software still under development at DSRI, Copenhagen (Denmark). The count rates extracted for Ser X-1 were compared to results from off-axis Crab observations analyzed with the same software. For details concerning the JEM-X analysis software see Westergaard et al. (2003).

## 3. Results

Measurements and upper limits for the X-ray emission in various energy bands from Ser X-1 during the present *INTEGRAL* pointing are listed in Table 1. Below we report the findings obtained with each instrument onboard *INTEGRAL*.

Ser X-1 was clearly detected with ISGRI in the 20–30 keV band (Fig. 1) with a significance of  $\sim 25\sigma$  over the whole 400 ks observation, with a count rate of  $0.84 \pm 0.03$  counts s $^{-1}$ . Nothing was detected above 30 keV at the position of this source: the upper limit to the emission in the 30–40 keV range is reported in Table 1. Our ISGRI results agree with those

<sup>1</sup> Available through the *INTEGRAL* Science Data Centre (ISDC, Geneva, Switzerland) at the web address: <http://isdc.unige.ch/index.cgi?Soft+download>



**Fig. 1.** Part of the mosaic image of the Ser X-1 field as imaged by ISGRI in the 20–30 keV band and accumulated over the whole observation (400 ks on-source time). The field is about  $30^\circ \times 25^\circ$ . North is up, East is left. Ser X-1 is detected at  $\sim 25\sigma$  significance. The image greyscale is histogram-equalization coded and scaled to the intensity, with the white color corresponding to zero counts. The large unidentified features northwest of Ser X-1 and west of J1855–026 are artifacts (ghost residuals) produced by the bright source GRS 1915+105.

obtained by Molkov et al. (2004) using *INTEGRAL* observations acquired before and after ours.

The object was also not detected in the whole spectral range covered by PICsIT. A  $3\sigma$  upper limit of  $4 \times 10^{-6}$  photons  $\text{cm}^{-2} \text{s}^{-1}$  in the energy band centered at 250 keV and  $\Delta E = 60$  keV wide was obtained from the PICsIT data (see Table 1).

SPI also did not reveal any significant emission from Ser X-1. From Table 1 one can note an apparent discrepancy between the flux obtained from the ISGRI detection and the SPI upper limit: this can be explained in terms of a not yet complete correction for systematic errors.


JEM-X detected the source in the 3–15 keV band with significance  $\sim 30\sigma$  in 2200 s when the source was  $3.6$  off axis. Using off-axis observations of the Crab (as mentioned in the previous section), we measured, in the three bands 3–5, 5–10 and 10–15 keV, the fluxes shown in Table 1. A  $3\sigma$  upper limit was instead obtained from the 15–35 keV band data; its value is in agreement with the ISGRI 20–30 keV detection.

The ISGRI source detection was at coordinates  $\alpha = 18^{\text{h}}40^{\text{m}}00^{\text{s}}.0$ ,  $\delta = +05^\circ 02' 06''$  (J2000). The significance of the detection implies a 90% confidence positional accuracy of  $1'$

(Gros et al. 2003). As regards JEM-X, it detected Ser X-1 at  $\alpha = 18^{\text{h}}39^{\text{m}}56^{\text{s}}.2$ ,  $\delta = +05^\circ 01' 57''$  (J2000) with a 99% confidence uncertainty of  $40''$  in both coordinates. The two positions are in good agreement with each other, given the off-axis location of the source within the fields of view of ISGRI and JEM-X. The object localization afforded by these two instruments onboard *INTEGRAL* is moreover fully consistent with that of the optical counterpart MM Ser (Thorstensen et al. 1980; Wachter 1997) as well as with the radio position (Migliari et al. 2004).

The 20–30 keV light curve of Ser X-1, rebinned to  $\sim 40$  ks, is shown in Fig. 2. We chose this temporal resolution as it turned out to be the best tradeoff between the time sampling and the S/N ratio of each bin. Substantial fluctuations are apparent in this band; so, in order to test their reality, we performed a fit to the data points on the hypothesis that the emission is constant. By means of a standard  $\chi^2$  test it was found that the constancy of the 20–30 keV emission from this source is rejected at the 99.99% confidence level.

Figure 2 also shows the Ser X-1 light curves in the 3–5, 5–10 and 10–15 keV ranges as obtained with JEM-X. Each point corresponds to a single JEM-X pointing ( $\sim 2$  ks long).




**Fig. 2.** Light curves of Ser X-1 as observed by ISGRI (20–30 keV) and JEM-X (3–5 keV, 5–10 keV and 10–15 keV) onboard *INTEGRAL* during the pointing presented in this paper. ISGRI data are rebinned to  $\sim 40$  ks, while JEM-X data are accumulated over each single pointing (with duration  $\sim 2$  ks).

We also explored the JEM-X light curves using higher time resolutions down to 500 s in the parts of the observation covered by the instrument (slots lasting  $\sim 5$  ks on average and separated by off-source intervals of about 20 ks). We did not find any hint of X-ray bursts or superbursts in these light curves. We of course cannot exclude the possibility that X-ray (super)bursts from Ser X-1 may have occurred in the 6 days covered by our *INTEGRAL* observation during the time intervals in which the source was outside the JEM-X field of view.

Observations with the All-Sky Monitor (ASM<sup>2</sup>; Levine et al. 1996) onboard the satellite *RXTE* (Bradt et al. 1993) showed that Ser X-1 had an average flux of  $208 \pm 8$  mCrab in the 1.5–12 keV band at the time of the *INTEGRAL* pointing, taking into account that 75 ASM counts  $s^{-1}$  correspond to 1 Crab in this band. For comparison, during the *BeppoSAX* observation, the ASM data indicate that the source was at  $239 \pm 3$  mCrab in the same energy band. This  $\sim 15\%$  difference in brightness apparently was not reflected in the overall spectral shape in the 1.5–12 keV range: indeed, the ASM hardness ratios HR1 and HR2 (defined as the ratio between the ASM counts in the 3–5 keV band and those in the 1.5–3 keV band, and as the ratio of the counts in the 5–12 keV band to

<sup>2</sup> ASM light curves are available at [http://xte.mit.edu/ASM\\_lc.html](http://xte.mit.edu/ASM_lc.html)



**Fig. 3.** Broadband SED of Ser X-1 constructed with simultaneous data and upper limits from ISGRI (filled diamonds), PICsIT (filled pentagon), JEM-X (filled squares) and ASM data (open square) plus non-simultaneous optical measurements (filled dots), NIR and far-infrared upper limits (open circles and open triangles, respectively) and radio data (filled triangles). Error bars of data points are not reported as their size is comparable to, or smaller than, that of the symbols used in the figure.

those in the 3–5 keV band, respectively), were  $1.01 \pm 0.11$  and  $1.20 \pm 0.10$  during the *INTEGRAL* pointing and  $1.02 \pm 0.04$  and  $1.20 \pm 0.03$  at the time of the *BeppoSAX* observation, respectively.

In addition, the JEM-X fluxes measured from Ser X-1 during the present *INTEGRAL* observation were found to be in general agreement with those obtained with the ASM in the spectral range over which the sensitivities of these two instruments overlap.

Feeling justified by these results, we expanded the spectral coverage of Ser X-1 down to 1.5 keV by adding to the *INTEGRAL* data set simultaneous 1.5–3 keV data acquired by the ASM. These ASM counts in the 1.5–3 keV were thus averaged over the duration of the *INTEGRAL* pointing; next, they were converted into fluxes assuming the spectral description for Ser X-1 provided by Oosterbroek et al. (2001), care being taken to rescale it with the use of the archival ASM data obtained simultaneously with the *BeppoSAX* observation.

We then collected published optical data (Wachter 1997) and near-infrared (NIR) 2MASS data (Skrutskie et al. 1997), as well as far-infrared upper limits (Beichman et al. 1988) and radio data (Migliari et al. 2004; Wendker 1995, and references therein). This allowed us to construct, for the first time for Ser X-1, a broadband SED (Fig. 3) spanning radio to  $\gamma$ -ray frequencies. Although non-simultaneous with the high-energy part of the spectrum, these lower frequency data and upper limits can give us a general description of the overall SED of this source thanks to the relative long-term stability of its emission with time.

Optical and NIR data were corrected assuming a color excess  $E(B - V) = 0.8$  (e.g., Hynes et al. 2004, and references therein) and converted into fluxes using the normalizations by Fukugita et al. (1995) for the optical and the ones referring to the 2MASS<sup>3</sup> for the NIR. Given the resolution of the 2MASS survey (pixel size: 2'') and the crowding of the Ser X-1 field (e.g., Wachter 1997), we considered the 2MASS NIR detections as conservative upper limits to the actual source fluxes in the *JHK* bands.

#### 4. Discussion

We observed Ser X-1 with *INTEGRAL*; we detected the source in the 3–15 keV band with the JEM-X instrument, and in the 20–30 keV range with the ISGRI detector of the IBIS instrument. Upper limits for the source flux were instead obtained in the 30–40 keV band and around 250 keV with the ISGRI and PICsIT detectors, respectively. The positional accuracy in the 20–30 keV band (1') is unprecedented, and the observation indicates that no significant nearby hard source is present within a radius of few degrees from Ser X-1. Thus we can exclude any contamination from persistent field sources in past observations of Ser X-1 made with previous non-imaging hard X-ray detectors (e.g., the HPGSPC and the PDS onboard *BeppoSAX*).

The source appears to be variable by a factor two in intensity in the 20–30 keV band on timescales of ~100 ks (Fig. 2). A similar behaviour was present in the 2–10 keV band as observed by the MECS instrument onboard *BeppoSAX* (Oosterbroek et al. 2001), although in this case the variability amplitude was about 25%, thus not as strong as in the *INTEGRAL* observations. This however is most likely a sampling effect due to the shorter duration (~60 ks) of the *BeppoSAX* pointing: indeed, during the first 80 ks of ISGRI data the flux variation in the 20–30 keV band is comparable to that seen with *BeppoSAX*.

The difference between the ISGRI and MECS light curves might however be due to a further reason: as the 20–30 keV flux is likely dominated by the Comptonization component, while that between 2 and 10 keV by the blackbody (or disk-blackbody) originating close to the NS surface, the two components might display variability with different timescales which in turn may be reflected in the light curve shape of the two above mentioned bands. Unfortunately, again the ASM and JEM-X light curves do not carry enough information to confirm or disprove this hypothesis.

The light curve sampling of JEM-X and ISGRI did not allow us to look for X-ray bursts from the surface of the NS harboured in this system throughout the entire observation: only in the ~40 ks covered by JEM-X can we exclude their presence. We therefore cannot rule out that some of the variability we see in the 20–30 keV range is caused by these phenomena occurring when JEM-X was pointing off-source; however, for typical values of their spectral shape, rate and overall energy output (see, e.g., Sztajno et al. 1983), we do not expect that X-ray bursts can contribute substantially to the total energy

radiated in 40 ks by Ser X-1 in the 20–30 keV band, which amounts to  $\sim 2 \times 10^{40}$  erg.

The observed ~100 ks timescale variability also cannot be explained assuming the occurrence of a ‘‘superburst’’ like the one reported by Cornelisse et al. (2002), which lasted ~4 h and emitted  $\approx 10^{40}$  erg in the 20–30 keV range. Indeed such a phenomenon, if present, can be contained in a single 40-ks bin of the 20–30 keV light curve reported in Fig. 2. The variations detected between adjacent bins in this spectral range might instead have been produced by a superburst but, again, the sparse ASM dwell-by-dwell and JEM-X data coverage during the *INTEGRAL* pointing does not allow us to check this hypothesis.

It should moreover be noted that the trend seen in the 20–30 keV light curve suggests that the source seems to vary periodically on a ~3 days timescale: this roughly matches a possible (superorbital?) X-ray periodicity of 3.4 days already reported for Ser X-1 (Ponman 1981; Ritter & Kolb 2003).

The 20–30 keV flux measurement obtained with *INTEGRAL* is about 20% lower than that obtained from the model by Oosterbroek et al. (2001) in the same band ( $8 \times 10^{-11}$  erg cm<sup>-2</sup> s<sup>-1</sup>); this roughly agrees with the difference (~15%) in the ASM flux as seen between the epochs of the *BeppoSAX* and *INTEGRAL* pointings. The 30–40 keV to 20–30 keV flux ratio in the *INTEGRAL* data is <0.1 which is consistent with the value of 0.054 as measured by *BeppoSAX*. The lack of sufficient coverage in the soft X-rays did not allow us to give a better description of the spectrum of this object; however, from the above results we can confidently say that we observed the atoll source Ser X-1 in its bright (banana) state, albeit at a lower intensity with respect to the time in which the *BeppoSAX* pointing was done, i.e., September 1999.

The broadband SED of Ser X-1 clearly shows a cut-off in the energy distribution above  $3 \times 10^{18}$  Hz ( $\approx 10$  keV). This can be reasonably taken to be due to the drop of the Comptonization tail induced by the electron temperature,  $kT_{e-} \sim 3$  keV as determined by Oosterbroek et al. (2001), and to the lack of further components in the  $\gamma$ -ray domain. No simple model can fit the overall SED: while the optical data fall above the extrapolation of the *BeppoSAX* X-ray model by a factor ~100, a smooth connection between optical and radio data may be present. Indeed, radio and optical fluxes are marginally consistent with lying on the same power law,  $F_{\nu} \propto \nu^{\beta}$ , with  $\beta \sim 0.2$ . This might indicate that the mechanism responsible for the radio emission (possibly a jet) can play a substantial role in the observed optical light from this source also (see e.g., Fender 2003, and Migliari et al. 2004).

*Acknowledgements.* We thank Simone Migliari for having communicated to us the Ser X-1 radio detection result prior to publication. Pavel Binko is acknowledged for the help in the *INTEGRAL* data retrieval from the ISDC archive. This work has made use of the NASA Astrophysics Data System Abstract Service, of the SIMBAD database, operated at CDS, Strasbourg, France, and of data products from the 2MASS. ASM data were provided by the *RXTE* ASM teams at MIT and at the *RXTE* SOF and GOF at NASA’s GSFC. This research was partially funded by ASI. We also thank the referee for several comments which helped us to improve this paper.

<sup>3</sup> These normalizations are available at <http://www.ipac.caltech.edu/2mass/releases/allsky/faq.html>

## References

- Beichman, C., Neugebauer, G., Habing, H. J., Clegg, P. E., & Chester, T. J. 1988, IRAS catalogs and atlases explanatory supplement (Washington, DC: GPO), NASA RP-1190, 1
- Bradt, H. V., Rothschild, R. E., & Swank, J. H. 1993, *A&AS*, 97, 355
- Cherepashchuk, A. M., Sunyaev, R. A., Seifina, E. V., et al. 2003, *A&A*, 411, L441
- Christian, D. J., & Swank, J. H. 1997, *ApJS*, 109, 177
- Cornelisse, R., Kuulkers, E., in 't Zand, J. J. M., Verbunt, F., & Heise, J. 2002, *A&A*, 382, 174
- Di Cocco, G., Caroli, E., Celesti, E., et al. 2003, *A&A*, 411, L189
- Diehl, R., Baby, N., Beckmann, V., et al. 2003, *A&A*, 411, L117
- Fender, R. P. 2003, in Compact stellar X-ray sources, ed. W. H. G. Lewin, & M. van der Klis (Cambridge: Cambridge Univ. Press), in press [arXiv:astro-ph/0303339]
- Fukugita, M., Shimasaku, K., & Ichikawa, T. 1995, *PASP*, 107, 945
- Gierliński, M., & Done, C. 2002, *MNRAS*, 331, L47
- Goldoni, P., Bonnet-Bidaud, J. M., Falanga, M., & Goldwurm A., 2003, *A&A*, 411, L399
- Goldwurm, A., David, P., Foschini, L., et al. 2003a, *A&A*, 411, L223
- Goldwurm, A., Gros, A., Goldoni, P., et al. 2003b, IBIS/ISGRI instrument specific software scientific validation report, v1.0
- Gros, A., Goldwurm, A., Cadolle-Bel, M., et al. 2003, *A&A*, 411, L179
- Hasinger, G., & van der Klis, M. 1989, *A&A*, 225, 79
- Hynes, R. I., Charles, P. A., van Zyl, L., et al. 2004, *MNRAS*, 348, 100
- Kuulkers, E. 2003, The observers' view of (very) long X-ray bursts: they are super!, in Proc. of the 2nd BeppoSAX Meet. The rest-less high-energy Universe, ed. E. P. J. van den Heuvel, J. J. M. in 't Zand & R. A. M. J. Wijers, Nucl. Phys. B Suppl. Ser., in press [arXiv:astro-ph/0310402]
- Labanti, C., Di Cocco, G., Ferro, G., et al. 2003, *A&A*, 411, L149
- Lebrun, F., Relay, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141
- Levine, A. M., Bradt, H. V., Cui, W., et al. 1996, *ApJ*, 469, L33
- Lewin, W. H. G., van Paradijs, J., & Taam, R. E. 1995, in X-ray Binaries, ed. W. H. G. Lewin, J. van Paradijs, & E. P. J. van den Heuvel (Cambridge: Cambridge Univ. Press), 175
- Liu, Q. Z., van Paradijs, J., & van den Heuvel, E. P. J. 2001, *A&A*, 368, 1021
- Lund, N., Budtz-Jørgensen, C., Westergaard, N. J., et al. 2003, *A&A*, 411, L231
- Mas-Hesse, J. M., Giménez, A., Culhane, J. L., et al. 2003, *A&A*, 411, L261
- Migliari, S., Fender, R. P., Rupen., M., et al. 2004, *MNRAS*, 351, 186
- Molkov, S. V., Cherepashchuk, A. M., Lutovinov, A. A., et al. 2004, *Astron. Lett.*, in press [arXiv:astro-ph/0402416]
- Muno, M. P., Remillard, R. A., & Chakrabarty, D. 2002, *ApJ*, 568, L35
- Oosterbroek, T., Barret, D., Guainazzi M., & Ford, E. C. 2001, *A&A*, 366, 138
- Piraino, S., Santangelo, A., Ford, E. C., Kaaret, P. 1999, *A&A*, 349, L77
- Ponman, T. J. 1981, *Space Sci. Rev.*, 30, 353
- Ritter, H., & Kolb, U. 2003, *A&A*, 404, 301
- Seon, K.-I., & Min, K. W. 2002, *A&A*, 395, 141
- Skrutskie, M. F., Schneider, S. E., Stiening, R., et al. 1997, The Two Micron All Sky Survey (2MASS): Overview and Status, in The Impact of Large Scale Near-IR Sky Surveys, ed. F. Garzon, N. Epchtein, A. Omont, B. Burton, & P. Persi (Dordrecht: Kluwer Acad. Pub.), 25
- Sztajno, M., Basinska, E. M., Cominsky, L. R., et al. 1983, *ApJ*, 267, 713
- Thorstensen, J. R., Charles, P. A., & Bowyer, S. 1980, *ApJ*, 238, 964
- Ubertini, G., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131
- van der Klis, M. 2000, *ARA&A*, 38, 717
- Vedrenne, G., Roques, J.-P., Schönfelder, V., et al. 2003, *A&A*, 411, L63
- Wachter, S. 1997, *ApJ*, 490, 401
- Wendker, H. J. 1995, *A&AS*, 109, 177
- Westergaard, N. J., Kretschmar, P., Oxborrow, C. A., et al. 2003, *A&A*, 411, L257
- Winkler, C., Courvoisier, T. J.-L., Di Cocco, G., et al. 2003, *A&A*, 411, L1