

## The broad-band spectrum of the persistent emission from SGR 1806-20<sup>\*</sup>

S. Molkov<sup>1,3</sup>, K. Hurley<sup>2</sup>, R. Sunyaev<sup>1,3</sup>, P. Shtykovsky<sup>1,3</sup>, M. Revnivtsev<sup>1,3</sup>, and C. Kouveliotou<sup>4</sup>

<sup>1</sup> Space Research Institute, Russian Academy of Sciences, Profsovnaya 84/32, 117997 Moscow, Russia  
e-mail: molkov@hea.iki.rssi.ru

<sup>2</sup> University of California at Berkeley, Space Sciences Laboratory, 7 Gauss Way, Berkeley, CA 94720-7450, USA

<sup>3</sup> Max-Planck-Institute für Astrophysik, Karl-Schwarzschild-Str. 1, 85740 Garching-bei-München, Germany

<sup>4</sup> National Space Science Technology Center, NASA/MSFC, XD-12, Huntsville, AL 35805, USA

Received 26 November 2004 / Accepted 16 February 2005

**Abstract.** We present the results of an analysis of the quiescent X-ray emission from the Soft Gamma-Ray Repeater SGR 1806-20, taken during an INTEGRAL ultra-deep survey of the Galactic center region in Aug.–Sep. 2003. The total effective exposure time spent on the source by the IBIS telescope during these observations was 1.6 million seconds. Combining the INTEGRAL results with results from the XMM-Newton observatory, we present the broad band (1–200 keV) spectrum of the quiescent emission from this source. This is the first spectrum of the persistent emission from an SGR in the broad energy range up to 200 keV. The spectrum of the source is very hard and has a power law shape ( $\Gamma = 1.6 \pm 0.1$ ) without any trace of a high energy cutoff up to  $\sim 160$  keV. No strong cyclotron line was detected in the persistent spectrum. The luminosity of SGR 1806-20 in this range (1–200 keV) was  $\sim 3.6 \times 10^{36}$  erg s<sup>-1</sup> for an assumed distance of 15 kpc. We show that weak undetected bursts should not contribute significantly to the quiescent emission. During our next observations in August 2004 the source went into an active phase and its average flux between powerful bursts was 2–3 times higher than in 2003. During Aug.–Sep. 2003, two other SGR candidates, SGR 1801-23 and SGR 1808-20, were in the field of view. Neither persistent hard X-ray emission nor bursts were detected from them. The upper limit on the persistent flux from each of them in the energy band 18–100 keV is about  $4 \times 10^{-11}$  erg s<sup>-1</sup> cm<sup>-2</sup>.

**Key words.** gamma rays: bursts – pulsars: general – stars: individual: SGR 1806-20

### 1. Introduction

Soft gamma repeaters (SGRs) are rare objects which have been discovered by their sporadic bursting activity. The bursts typically last  $\sim 10$ – $100$  ms, and their spectra may extend up to  $\sim 100$  keV or beyond. These properties make them easily observable for small, isotropic hard X-ray and soft gamma ray experiments (see e.g. Mazets & Golenetskii 1981a, Hurley 2000, Aptekar et al. 2001 and Woods & Thompson 2004 for reviews). In particular, SGR 1806-20 was discovered and localized on January 7, 1979 during a single  $< 0.25$  s long burst in soft gamma rays (Mazets & Golenetskii 1981b; Laros et al. 1986). Numerous bursts have been detected from this source since then (Atteia et al. 1987; Kouveliotou et al. 1987; Kuznetsov et al. 1987; Laros et al. 1987) and intermittent bursting activity has continued up to the present day.


The energy emitted by SGRs is in many cases significantly larger than that available from the rotational spin-down.

Therefore an additional energy reservoir is needed to power the observed emission. The most widely accepted model for SGRs suggests that this power comes not from accretion but rather from the decay of an ultra-strong magnetic field of an isolated neutron star (the “magnetar” model, Duncan & Thompson 1992; Paczyński 1992; Thompson & Duncan 1995).

Studies of the four known SGRs in 2–10 keV X-rays have revealed that they emit a persistent X-ray flux (Hurley et al. 1996; Hurley et al. 1999; Mereghetti et al. 2000; Woods et al. 2001; Kouveliotou et al. 2003; Kulkarni et al. 2003). The study of their persistent hard X-ray emission is more difficult, because of the weak fluxes and the locations of the sources in crowded fields. The INTEGRAL mission, however, with its imaging detectors operating in the hard X-ray and gamma-ray ranges (Winkler et al. 2003), has made the study of these sources possible up to energies which have not been explored before.

In this letter we present the broad-band spectrum (1–200 keV) of the persistent emission of SGR 1806-20 obtained with long exposures by the INTEGRAL observatory in August–September 2003. The source flux was  $\sim 1.3 \times 10^{-10}$  erg s<sup>-1</sup> cm<sup>-2</sup>, which corresponds to a luminosity  $\sim 3.6 \times 10^{36}$  erg s<sup>-1</sup> in this energy band (assuming a distance of

<sup>\*</sup> Based on observations with INTEGRAL, an ESA project with instruments and science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland, and with the participation of Russia and the USA.



**Fig. 1.** The differential distribution of IBIS exposure time on SGR 1806-20 as a function of the angle between the optical axis of the instrument and the direction to the source. The dashed line denotes the half angle of the IBIS Field of View (FOV). The total exposure time is 1.6 Ms.

15 kpc). In a companion paper, Mereghetti et al. (2005) also present evidence, from a separate observation, of hard, persistent X-ray emission from this source. Note that the persistent emission of SGR 1806-20 varies with time. During the 2004 burst activation of SGR 1806-20, the persistent flux was 2–3 times higher in soft X-rays (Woods et al. 2004) and hard X-rays (Molkov et al. 2004b) than during observations prior to 2004.

## 2. Observations and data analysis

SGR 1806-20 is located  $\sim 10^\circ$  from the Galactic center and it was therefore observed by the wide field telescopes aboard INTEGRAL (IBIS and SPI), but was outside the field of view of JEM-X during the ultra-deep survey of the Galactic center field performed during the Russian share of the INTEGRAL open time AO-1 observing program (ID0120213; August 23–September 24, 2003). The total exposure time of these observations is  $\sim 2$  Ms. In order to distinguish the flux of SGR 1806-20 from that of nearby bright X-ray sources we used only data from the IBIS/ISGRI telescope (Ubertini et al. 2003) which has moderate angular resolution ( $\sim 12'$ ). The SPI telescope has angular resolution  $\sim 2$ – $2.5'$  and cannot work in such a crowded field. IBIS/ISGRI effectively operates in the energy range 18–400 keV with typical sensitivity  $\sim 1.5$  mCrab for one orbit ( $\sim 3$  days) of observations. The fully coded field of view of this telescope is  $10^\circ \times 10^\circ$ , and the zero response field of view is  $29^\circ \times 29^\circ$ . Due to the fact that the source was practically always outside of the fully coded field of view (the region of maximum effective area of the telescope) the effective exposure is lower than the total exposure time, namely  $\sim 1.6$  Ms. The distribution of exposure time as a function of offset angle is presented in Fig. 1

The IBIS/ISGRI data analysis was carried out with the software developed by Eugene Churazov at the Space Research Institute, Moscow and described in Revnivtsev et al. (2004). For the spectral deconvolution we used the ratio of the fluxes measured in various energy channels to the fluxes measured

by the ISGRI detector from the Crab Nebula in the same energy bands; wherever possible the ratios were calculated for the same positions in the telescope field of view.

In order to extend the spectrum of SGR 1806-20 to lower energies we have used the data from an XMM-Newton observation performed on October 7, 2003, just two weeks after our one month long observing run that included SGR 1806-20. In the analysis we have used the data of the PN and MOS1 cameras, which were processed with the Science Analysis System (SAS) software v6.0.0.

## 3. Results

### 3.1. SGR 1806-20


The first deep IBIS/ISGRI map of the Galactic center region was presented in Revnivtsev et al. (2004). SGR 1806-20 was detected in the 18–60 keV energy range, along with sixty other sources. This was the first report of the detection of this source in hard X-rays.

During the period of our INTEGRAL observations the source was in a bursting phase (Sunyaev et al. 2003; Götz et al. 2004). The burst and quiescent emission of SGR 1806-20 are believed to originate from different regions on and around the neutron star, and are thought to be due to different mechanisms. Therefore, to obtain the spectrum of the quiescent emission, it is necessary to eliminate any possible contribution from bursts.

To do this, we constructed the statistical distribution of the detector count rates with time resolution 0.01 s. We identified candidate bursts as time intervals where the count rate during at least four consecutive bins exceeded the mean value of the count rate by more than four times the RMS value. Only those candidate bursts which demonstrated the clear presence of a point-like source on the image constructed during the burst intervals were considered as “real”. All the real bursts detected during our observations originated from SGR 1806-20. Their typical durations were 50–200 ms. The minimum number of counts from the source during these bursts was 40–50, which corresponds to a minimum fluence  $\sim 2$ – $3 \times 10^{-8}$  erg  $\text{cm}^{-2}$ . The majority of the remaining candidate events are probably caused by the anomalous short-term activation of one of the detector pixels (the so-called “short-lived hot pixels”). During these intervals more than 50% of all detected counts come from only one pixel, while for real bursts this fraction is typically less than 10%. Also, these bursts did not produce a point source image.

In order to demonstrate that the origin of these bursts was SGR 1806-20 and not other possible candidates in the wide field of view of IBIS/ISGRI (SGR 1808-20, Lamb et al. 2003; or SGR 1801-23, Cline et al. 2000) we present in the lower panel of Fig. 2 the  $\sim 3^\circ \times 2^\circ$  image around SGR 1806-20, collected during all the bursts which were detected during our observations. Detailed analysis of these bursts and others detected by INTEGRAL will be presented in a separate paper. In the top panel of Fig. 2 we present the time-averaged map for the same region. Note that SGR 1806-20 is clearly detected.


The summed spectrum of all the real bursts is presented in Fig. 3. The best fit spectrum using an optically thin



**Fig. 2.** *Top panel:* time averaged image of the region around SGR 1806-20 constructed during the ultra-deep Aug.–Sep. 2003 observations of the Galactic center region. *Bottom panel:* image of the same region constructed during the bursts detected in this observation. Both images are reconstructed in the 18–60 keV energy band. The entire error boxes of the sources SGR 1808-20 and SGR 1801-23 were in the field of view, but neither persistent emission nor bursts from these sources was detected during our observations.

thermal plasma fit (*brems* model in XSPEC fitting package) has  $kT \sim 47 \pm 5$ , which is consistent with that presented by Götz et al. (2004) for one of the brightest bursts detected in a different set of observations.

We have filtered out both the real and the hot pixel bursts from our data and have constructed the time-averaged spectrum of the total (i.e. pulsed plus unpulsed) persistent emission from SGR 1806-20. In order to complement this hard X-ray spectrum with the spectrum at lower, X-ray energies we have used the publicly available data from the XMM-Newton (EPIC-pn and EPIC-MOS1) observations of the source carried out on October 7, 2003. The broad-band spectrum of SGR 1806-20 is presented in Fig. 3. In this figure we have rescaled the XMM spectrum by 35% in order to match the INTEGRAL/IBIS normalization. This discrepancy in the normalization factor could be partially caused by the intrinsic source variability, but it is likely that most of the difference originates in the uncertain cross-calibration of the INTEGRAL and XMM instruments (see e.g. Courvoisier et al. 2003). The spectrum can be well described by a single power law with photoabsorption (presumably interstellar). We did not detect statistically significant temporal variations of the source spectral shape in our observations. The best fit photon index for the time-averaged spectrum between 1 and 200 keV is  $\Gamma = 1.6 \pm 0.1$  and the absorption column  $N_{\text{H}} = (6.6 \pm 0.2) \times 10^{22} \text{ cm}^{-2}$ .



**Fig. 3.** The quiescent energy spectrum of SGR 1806-20 (lower spectrum) and the summed spectrum of all detected bursts rescaled down by a factor of 1000 (upper spectrum).

A lower limit on a possible exponential cutoff in the spectrum of the form  $dN/dE \propto E^{-\Gamma} \exp(-E/E_{\text{cut}})$  is  $E_{\text{cut}} > 160 \text{ keV}$  ( $2\sigma$  lower limit). The reduced  $\chi^2_r$  for the best fit model is  $\chi^2_r = 1.15$  for 294 degrees of freedom. The total observed flux in the 1–200 keV energy band is  $\sim 1.3 \times 10^{-10} \text{ erg s}^{-1} \text{ cm}^{-2}$  (not corrected for absorption). This is approximately 8 times higher than the source emissivity in X-rays  $< 10 \text{ keV}$ . For an assumed source distance of 15 kpc (Corbel & Eikenberry 2004) the observed broad-band source flux corresponds to a luminosity  $\sim 3.6 \times 10^{36} \text{ erg s}^{-1}$ . The energy loss due to the spin-down of the neutron star in SGR 1806-20 is approximately three orders of magnitude smaller, namely  $\dot{E}_{\text{rot}} \sim 10^{33} \text{ erg s}^{-1}$  (Kouveliotou et al. 1998). Obviously the rotational energy loss cannot power either the observed X-ray emission or the hard X-ray (20–200 keV) emission.

The RXTE/PCA has detected a relatively strong narrow line at 5 keV during some bright bursts, interpreted as cyclotron absorption (Ibrahim et al. 2002, 2003). The high quality 1–10 keV XMM spectrum of SGR 1806-20 allows us to put constraints on the possible presence of such a feature in the persistent spectrum of the source during this observation. Fixing the energy of the line to the 4.0–6.0 keV range, we can place an upper limit on a possible absorption line component in the spectrum. The  $2\sigma$  upper limit on the line equivalent width varies between  $EW \lesssim 50 \text{ eV}$  for 0.1 keV linewidth and  $EW \lesssim 160 \text{ eV}$  for 0.5 keV linewidth. This is significantly smaller than the equivalent width of the line detected by the RXTE/PCA during the bursts,  $EW \sim 500\text{--}800 \text{ eV}$ . The large difference between the two equivalent widths may be connected with differences in the physical conditions leading to the formation of burst and persistent spectra.

### 3.2. SGR 1801-23 and SGR 1808-20

During our INTEGRAL observations two SGR candidates were in the field of view. Both sources have rather large

location uncertainties. SGR 1808-20 is localized to a  $\sim 6$  arcmin radius circle and SGR 1801-23 to a  $\sim 200 \times 0.5$  arcmin error box. The error boxes of both sources were entirely in the IBIS field of view (Fig. 2). Neither persistent nor bursting emission was detected from these sources. The upper limit on the persistent flux from these sources is about  $4 \times 10^{-11}$  erg s $^{-1}$  cm $^{-2}$ .

#### 4. Discussion

Although we do not know the exact mechanism which produces the persistent emission of SGR 1806-20, we can nevertheless exclude a significant contribution to it from weak, undetected bursts by the following argument. We can estimate the contribution using the number-intensity distribution of the bursts presented in Göğüş et al. (2000) and Götz et al. (2004). The peak number of detected bursts with fluence  $\geq 2-3 \times 10^{-8}$  erg cm $^{-2}$  is approximately 20–30 bursts per 10 days. The total flux that bursts with fluences lower than this can produce during the exposure time of our observations is not more than a few  $\times 10^{-13}$  erg s $^{-1}$  cm $^{-2}$  if we assume that the number-fluence distribution of bursts from SGR 1806-20 can be described by a function  $N(>F) \propto F^{-0.7}$  (Göğüş et al. 2000; Götz et al. 2004). This flux does not contribute even one percent to the detected persistent flux of SGR 1806-20. Therefore we conclude that the detected quiescent spectrum of SGR 1806-20 is unlikely to be strongly affected by weak undetected bursts. Additional indirect support for this conclusion comes from the completely different spectral shapes of the bright bursts and the quiescent emission detected during our observations (Fig. 3).

*Acknowledgements.* The authors thank E. Churazov for the development of the analysis methods for the IBIS data and software. This research has made use of data obtained through the INTEGRAL Science Data Center (ISDC), Versoix, Switzerland, and the Russian INTEGRAL Science Data Center (RSDC), Moscow, Russia. K.H. is grateful for support under the INTEGRAL US guest investigator program, NASA grant NAG5-13738.

#### References

- Aptekar, R., Frederiks, S., Golenetskii, V., et al. 2001, *ApJS*, 137, 227  
 Atteia, J.-L., Boer, M., Hurley, K., et al. 1987, *ApJ*, 320, L105  
 Cline, T., Frederiks, D., Golenetskii, S., et al. 2000, *ApJ*, 531, 407  
 Corbel, S., & Eikenberry, S. S. 2004, *A&A*, 419, 191  
 Courvoisier, T., Beckmann, V., Bourban, G., et al. 2003, *A&A*, 411, L343  
 Duncan, R., & Thompson, C. 1992, *ApJ*, 392, L9  
 Göğüş, E., Woods, P., Kouveliotou, C., et al. 2000, *ApJ*, 532, L121  
 Götz, D., Mereghetti, S., Mirabel, F., & Hurley, K. 2004, *A&A*, 417, L45  
 Hurley, K., Li, P., Vrba, F., et al. 1996, *ApJ*, 463, L13  
 Hurley, K., Li, P., Kouveliotou, C., et al. 1999, *ApJ*, 510, L111  
 Hurley, K. 2000, in *Gamma-ray Bursts, 5th Huntsville Symposium*, ed. R. M. Kippen, R. S. Mallozzi, & G. J. Fishman, AIP Conf. Proc., 526, 763  
 Ibrahim, A., Safi-Harb, S., Swank, J., et al. 2002, *ApJ*, 574, L51  
 Ibrahim, A., Swank, J., & Parke, W. 2003 *ApJ*, 584, L17  
 Kouveliotou, C., Norris, J., Cline, T., et al. 1987, *ApJ*, 322, L21  
 Kouveliotou, C., Dieters, S., Strohmayer, T., et al. 1998, *Nature*, 393, 235  
 Kouveliotou, C., Eichler, D., Woods, P., et al. 2003, *ApJ*, 596, L79  
 Kulkarni, S., Kaplan, D., Marshall, H., et al. 2003, *ApJ*, 585, 948  
 Kuznetsov, A. V., Sunyaev, R. A., Terekhov, O. V., et al. 1987, *SvA Lett.*, 13, 419  
 Lamb, D., Graziani, C., Shirasaki, Y., et al. 2003, *GCN Circ.*, 2351  
 Laros, J., Fenimore, E., Fikani, M., et al. 1986, *Nature*, 322, 152  
 Laros, J., Fenimore, E., Klebesadel, R., et al. 1987, *ApJ*, 320, L111  
 Mazets, E. P., & Golenetskii, S. V. 1981a, *Astrophys. Space Phys. Rev.*, 1, 205  
 Mazets, E. P., & Golenetskii, S. V. 1981b, *Ap&SS*, 75, 47  
 Mereghetti, S., Cremonesi, D., Feroci, M., & Tavani, M. 2000, *A&A*, 361, 240  
 Mereghetti, S., Götz, D., Mirabel, I. F., & Hurley, K. 2005, *A&A*, 433, L9  
 Molkov, S., Ubertini, P., Jorgensen, C., et al. 2004, *The Astronomer's Telegram*, 324  
 Molkov, S., Hurley, K., Lutovinov, A., & Sunyaev, R. 2004, *The Astronomer's Telegram*, 325  
 Paczyński, B. 1992, *Acta Astron.*, 42, 145  
 Revnivtsev, M., Sunyaev, R., Varshalovich, D., et al. 2004, *Astr. Lett.*, 30, 382  
 Sunyaev, R., Zheleznyakov, V., Cherepashchuk, A., et al. 2003, *The Astronomer's Telegram*, 183  
 Thompson, C., & Duncan, R. 1995, *MNRAS*, 275, 255  
 Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131  
 Winkler, C., Gehrels, N., Schönfelder, V., et al. 2003, *A&A*, 411, L349  
 Woods, P., Kouveliotou, C., Göğüş, E., et al. 2001, *ApJ*, 552, 748  
 Woods, P., & Thompson, C. 2004 [[arXiv:astro-ph/0406133](https://arxiv.org/abs/astro-ph/0406133)]  
 Woods, P., Kouveliotou, C., Göğüş, E., et al. 2004, *The Astronomer's Telegram*, 313