

On metal-deficient barium stars and their link with yellow symbiotic stars^{★,★★}

A. Jorissen^{1,★★★}, L. Začs², S. Udry³, H. Lindgren⁴, and F. A. Musae^{5,6,7}

¹ Institut d'Astronomie et d'Astrophysique, Université Libre de Bruxelles, CP 226, Boulevard du Triomphe, 1050 Bruxelles, Belgium
e-mail: ajorisse@astro.ulb.ac.be

² Institute of Atomic Physics and Spectroscopy, University of Latvia, Raiņa bulvāris 19, Rīga, LV 1586 Latvia

³ Observatoire de Genève, 1290 Sauverny, Suisse

⁴ Lund Observatory, Box 43, 221 00 Lund, Sweden

⁵ Special Astrophysical Observatory and Isaac Newton Institute of Chile, SAO Branch, Nizhnij Arkhyz 369167, Russia

⁶ ICAMER, National Academy of Sciences of Ukraine, 361605 Peak Terskol, Kabardino-Balkaria, Russia

⁷ Shamakhy Astrophysical Observatory, National Academy of Sciences of Azerbaijan, Azerbaijan

Received 25 April 2005 / Accepted 20 May 2005

ABSTRACT

This paper addresses the question of why metal-deficient barium stars are not yellow symbiotic stars (YSyS). Samples of (suspected) metal-deficient barium (mdBa) stars and YSyS have been collected from the literature, and their properties reviewed. It appears in particular that the barium nature of the suspected mdBa stars needs to be ascertained by detailed abundance analyses. Abundances are therefore derived for two of them, HD 139409 and HD 148897, which reveal that HD 148897 should not be considered a barium star. HD 139409 is a mild barium star, with overabundances observed only for elements belonging to the first s-process peak (Y and Zr). It is only moderately metal-poor ($[Fe/H] = -0.4$). The evidence for binarity among mdBa stars is then reviewed, using three different methods: (i) radial-velocity variations (from CORAVEL observations), (ii) Hipparcos astrometric data, and (iii) a method based on the comparison between the Hipparcos and Tycho-2 proper motions. An orbit is obtained for HIP 55852, whereas evidence for the (so far unknown) binary nature of HIP 34795, HIP 76605, HIP 97874 and HIP 107478 is presented. No conclusion regarding the binary nature of HIP 11595, HIP 25161 could be reached. Two stars with no evidence for binarity whatsoever (HIP 58596 and BD +3°2688) are candidates low-metallicity thermally-pulsing asymptotic giant branch stars, as inferred from their large luminosities. The reason why mdBa stars are not YSyS is suggested to lie in their different orbital period distributions: mdBa stars have on average longer orbital periods than YSyS, and hence their companion accretes matter at a lower rate, for a given mass loss rate of the giant star. The definite validation of this explanation should nevertheless await the determination of the orbital periods for the many mdBa stars still lacking periods, in order to make the comparison more significant.

Key words. binaries: symbiotic – stars: abundances – stars: AGB and post-AGB – binaries: spectroscopic

1. The problem

Our understanding of the link between chemically-peculiar red giants like barium stars or CH stars, and yellow symbiotic stars (YSyS) has made substantial progress in the last decade (see the reviews by Jorissen 2003a,b), mainly with the realisation that likely all yellow symbiotics (i.e., involving a giant of spectral type G or K as primary component) involve barium stars (Smith et al. 1996, 1997;

Pereira & Porto de Mello 1997; Pereira et al. 1998). The metal-deficient nature of the giant is a key factor, because it implies a rather large luminosity for the giant. Since evolutionary tracks of metal-deficient stars are shifted towards the blue, metal-deficient giants of spectral type K must lie on the upper part of the (asymptotic) giant branch (see Fig. 11 of Smith et al. 1996), where they suffer strong mass loss. If such metal-deficient K giants are in binary systems, their strong wind will interact with the companion and trigger symbiotic activity.

The other facet of this problem, namely whether all metal-deficient barium stars are symbiotic stars, is not yet fully answered. The present paper offers a first step in that direction.

We have collected a list of candidate metal-deficient barium stars and have assembled new observations to check (i) whether

* Based on observations carried out at the European Southern Observatory (ESO, La Silla, Chile) and with the 1-m Swiss telescope at the Haute-Provence Observatory.

** Table 7 is only available in electronic form at <http://www.edpsciences.org>

*** Senior Research Associate, FNRS, Belgium.

Table 1. The barium syndrome among YSyS. The spectral type of the cool component is taken from Mürset & Schmid (1999), or references therein. In column labelled “nebula”, “y” means that an optical nebula has been detected, and “PN” that, based on its emission line spectrum, the star has traditionally been included in planetary nebulae catalogues, even though no optical nebula may be visible. The column labelled “*P*” lists the orbital period, from Mürset & Schmid (1999).

Name	Sp. typ.	[Fe/H]	V_r (km s ⁻¹)	b (°)	[Ba/Fe]	$V \sin i$ (km s ⁻¹)	Nebula	P (d)	Ref.
d'-type									
V417 Cen	G8-K2	~0.0		-1	0.5	70	y	247	(5, 11)
HDE 330036	G5	0.02	-14	+4	0.88	100	PN	-	(14, 17)
=Cn 1-1									
AS 201	G5	0.07		+7	0.63	25	y	-	(12, 17)
V471/V741 Per	G5	?	-12	-9	>0		PN	-	(2)
=M 1-2									
St H _α 190	G5	0.0	~10	-35	~0.5	100	bip. outf.	-	(10, 13)
Wray 157	G5	?							
Hen 1591	<K4	?							
s-type									
UKS Ce-1	C4, 5Jch	?	+20	+20	>0			-	(6)
S 32	C1, 1CH	?	+325	-30	>0			612	(6, 14)
Hen 2-467	K0	-1.1	-109	-12	+0.8		n	478	(4, 16)
BD-21°3873	K2	-1.1	+204	+37	+0.5		n	282	(3, 15, 16)
		-1.3			+0.3				(9)
AG Dra	K2	-1.3	-148	+41	+0.5		n	554	(8, 16)
CD -43°14304	K7	-1.4	+27	-41	?			1448	(7, 18)
Chemical evolution of the Galaxy									
		-1.0			<0.2				(1)

References: (1) Edvardsson et al. 1993, A&A, 275, 101; (2) Grauer & Bond 1981, PASP, 93, 630; (3) Pereira et al. 1997, AJ, 114, 2128; (4) Pereira et al. 1998, AJ, 116, 1977; (5) Pereira et al. 2003, in Symbiotic stars probing stellar evolution, ed. R. L. M. Corradi, J. Mikołajewska, & T. J. Mahoney, Astron. Soc. Pacific Conf. Ser. (San Francisco), 85; (6) Schmid 1994, A&A, 284, 156; (7) Schmid et al. 1998, A&A, 329, 986; (8) Smith et al. 1996, A&A, 315, 179; (9) Smith et al. 1997, A&A, 324, 97; (10) Smith et al. 2001, ApJ, 556, L55; (11) Van Winckel et al. 1994, A&A, 285, 241; (12) Schwarz 1991, A&A, 243, 469; (13) Munari et al. 2001, A&A, 369, L1; (14) Schmid & Nussbaumer 1993, A&A, 268, 159; (15) Munari & Patat 1993, A&A, 277, 195; (16) Corradi et al. 1999, A&A, 343, 841; (17) Pereira et al. 2005, A&A, 429, 993; (18) The metallicity is from Pereira, priv. comm.

these stars are binaries, (ii) whether they are barium stars and (iii) whether they exhibit symbiotic activity.

2. The samples

Before discussing metal-deficient barium stars, it is useful to first summarize the properties of YSyS, to which metal-deficient barium stars may be compared.

2.1. Yellow symbiotic stars

All known YSyS are listed in Table 1, which shows that all the stars studied so far exhibit the barium syndrome. YSyS with a *stellar* infrared continuum (s-type, as opposed to the dusty d'-type; see below) are clearly metal-deficient objects, as revealed by their low metallicities and high space velocities (CD -43°14304 may be an exception; however, it is of spectral type K7, and should perhaps not be included in the family of YSyS). The presence of the barium syndrome among

a family of binary metal-deficient stars fully supports the commonly accepted hypothesis that the s-process is more efficient at low metallicities (Clayton 1988; Jorissen 2003a). s-Type YSyS, with their metallicities lower than classical barium stars, may be expected to be, on average, more luminous than the latter (see Fig. 11 of Smith et al. 1996, comparing the luminosity function of Pop.I and Pop.II K giants). This is a direct consequence of the fact that evolutionary tracks shift towards the blue in the Hertzsprung-Russell (HR) diagram as metallicity decreases, as shown in Fig. 1b. Figure 1a confirms that the YSyS AG Dra and BD -21°3873 are indeed more luminous than classical barium stars. This difference in the average luminosity – and hence mass-loss rate – of the two populations thus explains why YSyS, despite hosting a K giant, exhibit symbiotic activity whereas barium stars do not. The larger mass-loss rates for the cool components of s-type YSyS – as compared to Ba stars – may be inferred from the comparison of their IRAS [12]–[25] color indices, which reflect the amount of dust present in the system: $([12]–[25])_{\text{Ba}} < 0.1$, as compared to 0.45




Fig. 1. *Left panel a):* evolutionary tracks of Schaller et al. (1992) compared with the locations of classical barium stars (stars labelled “G” in Mennessier et al. (1997); open dots) and the yellow SyS AG Dra (filled circle), BD $-21^{\circ}3873$ (filled triangle) and Hen 2-467 (filled square). The bolometric magnitudes were taken from the references listed in Table 1. These bolometric magnitudes were combined with bolometric corrections from Bessell et al. (1998) and $B - V$ indices from Munari et al. (1992) and Munari & Buson (1992) to yield the absolute visual magnitudes. *Right panel b):* same as a) but for YSyS (filled symbols as in the left panel) and metal-deficient barium stars [open triangles: stars flagged as “H” by Mennessier et al. (1997); crosses: CH stars also flagged as “H” by Mennessier et al. (1997); small open squares: additional metal-deficient, s-process-rich stars from Table 2; open circle: HD 104340, large open square: HD 206983 from Junqueira & Pereira (2001)]. The dashed horizontal line represents the luminosity ($M_{\text{bol}} = -3$, corresponding to $M_V \sim -2$) at the first thermal pulse in a $1 M_{\odot}$ AGB star of metallicity $[\text{Fe}/\text{H}] = -1.8$ according to Lattanzio (1991).

for AG Dra (Smith et al. 1996). Mürset et al. (1991) and Drake et al. (1987) provide direct measurements (or upper limits) for the mass loss rates of AG Dra and of Ba stars, respectively, which confirm the above conclusion.

YSyS with a *dusty* infrared continuum (d' -type; Allen 1982; Schmid & Nussbaumer 1993) differ from their s-type counterparts in several respects (Table 1): they host a complex circumstellar environment (including cool dust, bipolar outflows, extended optical nebulae or emission-line spectra closely resembling those of planetary nebulae), the cool components have early spectral types (F to early K), they are often fast rotators (with the possible exception of M 1–2 = V471 Per; Grauer & Bond 1981) and, finally, they belong to the galactic disk unlike s-type YSyS which belong to the halo.

All these arguments suggest that the hot component in d' -type SyS has just evolved from the AGB to the WD stage. The rather cool dust (Schmid & Nussbaumer 1993) is a relic from the mass lost by the AGB star. The optical nebulae observed in d' -type SyS are most likely genuine planetary nebulae rather than the nebulae associated with the ionized wind of the cool component (Corradi et al. 1999). This is especially clear for AS 201 which actually hosts *two* nebulae (Schwarz 1991): a large fossil planetary nebula detected by direct imaging, and a small nebula formed in the wind of the current cool component. Finally, the rapid rotation of the cool component has likely been caused by spin accretion from the former AGB wind like in WIRRING systems (Jeffries & Stevens 1996; Jorissen 2003b). The fact that the cool star has not yet been slowed down by magnetic braking is another indication

that the mass transfer occurred fairly recently (Theuns et al. 1996). Corradi & Schwarz (1997) obtained 4000 y for the age of the nebula around AS 201, and 40 000 y for V417 Cen.

2.2. Metal-deficient barium stars

Metal-deficient barium stars (with metallicities in the range -1.1 to -1.8 , comparable to that of YSyS) were identified by Luck & Bond (1991), Mennessier et al. (1997) and Začs et al. (2000), and occupy the same region of the HR diagram as YSyS (Fig. 1b). The question thus arises why metal-deficient barium stars are not YSyS. Different answers must be sought, depending upon their absolute visual magnitudes M_V . The most luminous systems, with $M_V < -2$, are likely located on the thermally-pulsing AGB, so that their Ba syndrome may be explained by internal nucleosynthesis. They thus should not be binaries, and therefore cannot be YSyS! HD 104340 (open circle in Fig. 1b), a metal-deficient Ba star studied by Junqueira & Pereira (2001), and BD $+03^{\circ}2688$ (Table 2) provide good illustrations of this situation, since they both lie above the TP-AGB threshold and CORAVEL radial-velocity measurements spanning several years do not reveal any clear orbital motion (Figs. 9 and 10, as well as Sect. 4).

The less luminous and warmest among metal-deficient Ba stars, clumping around $M_V \sim +1$ in the HR diagram, are also sometimes classified as CH stars (crosses in Fig. 1b). They are not losing mass at a large enough rate to trigger any symbiotic activity, as revealed by their small $[12]-[25]$ color indices (<0.3 ; Smith et al. 1996).

Table 2. Stars classified as metal-deficient barium stars by Mennessier et al. (1997). The absolute magnitude M_V is a maximum likelihood estimate obtained by Mennessier et al. (1997), except for the additional stars where it is derived from a straight inversion of the Hipparcos parallax (BD +75°348), from a spectroscopic estimate of gravity and an educated guess for the mass (BD +3°2688) or from a fit to the M 92 isochrone (CS stars). The column labelled “Ba” indicates whether detailed chemical analyses have confirmed the Ba nature of the star.

HIP	HD/DM	M_V	$B - V$	[Fe/H]	Ref.	Ba	Ref.	Rem.
4347	5424	-0.6	1.14					
11595	15589	-0.5	1.15	-0.7	9			
25161	-27°2233	-1.1	1.13					
29740	43389	-3.5	1.48					
34795	55496	-1.7	0.96	-1.55	5	y	5	
58596	104340	-1.9	1.36	-1.72	3	y	3	
69834	123396	-0.9	1.19					
76605	139409	-0.5	1.04	-0.42	2	mild	2	$M_V = 1.5$ is derived from spectroscopic $\log g$
80843	148897	-1.8	1.27	-1.0	2	n	2	
				-0.62	7			
				-1.16	8			
97874	187762	-1.2	1.10					
107478	206983	0.2	1.64	-1.43	3	y	3	
Additional stars (from refs. (1), (5) and (10))								
43042	+75°348	-0.4	1.17	-0.8	1	y	1	
55852	+4°2466	?	0.60	-1.85	5,11	y	4, 5	
	+3°2688	-5.0	1.22	-1.42	5,11	y	5	
	CS 22942-019	0	0.74	-2.67	10	y	10	
	CS 22948-027	-1	0.93	-2.60	10	y	10	

References to the table: (1) Zacs et al. (2000); (2) this work; (3) Junqueira & Pereira (2001); (4) Burris et al. (2000); (5) Luck & Bond (1991); (6) Jorissen et al. (1998); (7) Kyröläinen et al. (1986); (8) Luck (1991); (9) Barbuy et al. (1992); (10) Preston & Sneden (2001); (11) $B - V$ from the Tycho-2 catalog (Høg et al. 2000b).

Finally, at intermediate luminosities ($-2 \leq M_V \leq +1$) where YSyS are located, metal-deficient Ba stars are not luminous enough to be TP-AGB (hence they should be binaries), but yet their mass loss rates must be large enough to trigger symbiotic activity (provided that the orbital separation is not too large, since it is the mass *accretion* rate by the compact companion which is in fact the key parameter; see Sect. 5 and Jorissen 2003a). It is thus of great interest to check (i) the Ba nature of those metal-deficient stars with intermediate luminosities, (ii) their binary nature, and (iii) their suspected symbiotic activity. The first two issues are addressed in Sects. 3 and 4, respectively.

As far as a possible symbiotic activity is concerned, there is no indication from their photometric $U - B$ and $B - V$ indices that the metal-deficient stars in Table 2 have a strong blue continuum which could betray their symbiotic nature. It is thus very likely that none among these stars is a full-fledged symbiotic star. No signature of weak symbiotic activity (of the kind exhibited by some binary S stars; see Fig. 14 and Van Eck & Jorissen 2002) is observed in the H_α line profile either (Fig. 2).

3. Abundances

The classification of the stars in Table 2 as metal-deficient Ba stars is subject to caution, as it does not rely on spectroscopic abundance analyses, but rather on a maximum-likelihood assignment based on kinematic, spatial and

luminosity properties (Mennessier et al. 1997) for barium stars from the list of Lü (1991). Nevertheless, when a metallicity determination is available, it confirms the metal-deficient nature of the object (see Table 2). HD 139409 is an exception, though, since detailed spectral analyses reveal that it is neither metal-poor nor strongly enriched in s-process elements (see below). It may nevertheless be hoped that the metal-deficient assignment made by Mennessier et al. (1997) is valid in all the other cases. Regarding the Ba nature of these stars, it is known that the Lü (1991) catalogue of barium stars, from which the sample of barium stars used by Mennessier et al. (1997) was drawn, is contaminated by many non-barium stars (Griffin & Keenan 1992; Jorissen et al. 1996), especially among those stars having a Ba index smaller than 1.

It would therefore in principle be necessary to re-evaluate the Ba nature of all the stars listed in Table 2. So far, spectra could be obtained for two of them, HD 139409 and HD 148897, which are discussed in detail in the present section. Of these, only HD 139409 appears to be a mild barium star, thus confirming the suspicion about the Lü (1991) catalogue expressed above.

3.1. The case of HD 148897 and HD 139409

HD 148897 (=HR 6152) has been tagged as a “likely marginal barium star” by Boyle & McClure (1975) and as a marginal CH star by Vilhu et al. (1977). It therefore found its way into




Fig. 2. H_{α} profiles for two candidate metal-deficient Ba stars (HD 139409 and HD 148897; see however the discussion in Sect. 3), compared to that of the K giant Arcturus. The wavelength scale has been corrected for the stellar radial velocity. The profiles show no indication of symbiotic activity.

the barium-star catalogs of Lü et al. (1983) and Lü (1991), as well as the catalog of CH and metal-deficient Ba stars of Šleivyte & Bartkevičius (1990). The star was classified as G8.5III CN-2 Fe-1 CH-1 by Keenan & McNeil (1989), and this classification as CN- and CH-weak contradicts the earlier assignments.

A detailed abundance analysis has thus been performed to clarify the situation, and its results are compared with previous analyses by Kyröläinen et al. (1986) and Luck (1991) in Sect. 3.1.2.

HD 139409 (=HIP 76605) has been classified as a marginal barium star by MacConnell et al. (1972), as G5 III Ba1 by Yamashita & Norimoto (1981) and as K0 III/II Ba 0.5 by Lü (1991).

3.1.1. Observations

A high-resolution spectrum of HD 148897 was obtained using the Coudé Matrix Echelle Spectrometer (MAESTRO; Musaev et al. 1999) delivering a resolving power of 45 000 and installed on the 2-m Zeiss telescope of the Terskol Observatory (located in Northern Caucasus at an altitude of 3100 m). The spectrometer is equipped with a Wright Instruments CCD detector with 1242×1152 pixels ($22.5 \times 22.5 \mu\text{m}$). A total exposure of 1800 s was taken on February 18, 2003. The spectrum covers the range 365 to 1020 nm spread over 85 spectral orders.




Fig. 3. Comparison of equivalent widths measured on the HARPS spectrum of Arcturus with those of the Arcturus spectral atlas (Hinkle et al. 2000), for the same set of lines as that measured in HD 139409.

A high-resolution spectrum of HD 139409 was obtained on the HARPS spectrograph (Mayor et al. 2003), delivering a resolution of 115 000 and installed on the ESO La Silla 3.6 m telescope. A total exposure of 200 s was obtained by Xavier Bonfils on May 31, 2004. In order to check equivalent widths delivered by HARPS, a spectrum of the standard star Arcturus was obtained as well by Fabien Carrier. The HARPS spectra were reduced by the observers using standard pipeline processing. Equivalent widths for the same set of lines as those studied in HD 139409 have been measured by one of us (L.Z.) in the HARPS spectrum of Arcturus and compared to those from the Arcturus spectral atlas (Hinkle et al. 2000). The agreement between the two sets of equivalent widths is excellent (Fig. 3), thus qualifying HARPS for abundance analyses. Spectra around the $\lambda 614.172$ nm Ba II line are shown in Fig. 4 for the two analyzed stars and for Arcturus (Hinkle et al. 2000).

3.1.2. Reduction and analysis

HD 148897

The reduction of the CCD frames (subtraction of bias, dark and scattered light, flat fielding, extraction of echelle orders and wavelength calibration) was performed with the DECH20T software (Galazutdinov 1992). More than 500 weak to medium-strong atomic lines, free of blends, were identified and their equivalent widths were measured with the DECH routines. The radial velocity was measured using a large number of symmetric absorption lines. The observed velocity has been brought to the heliocentric system by adding $+22.7 \text{ km s}^{-1}$. The mean heliocentric radial velocity for HD 144897 was found to be $+16.7 \text{ km s}^{-1}$. The atmospheric parameters for that star cannot be derived from photometry, since the standard temperature calibrations refer to stars of normal chemical composition. To obtain a colour-independent estimate of the temperature, a spectroscopic temperature has been derived from the




Fig. 4. The spectra of the target stars in the region of the Ba II line at 614.172 nm. Measured lines are marked. Also shown is the spectrum of the standard star Arcturus (Hinkle et al. 2000).

excitation equilibrium of Fe I lines. The surface gravity $\log g$ was determined using the Fe I/Fe II ionization balance, whereas the microturbulent velocity v_t was derived by forcing the abundances of individual Fe I lines to be independent of the reduced equivalent width. The resulting atmospheric parameters for HD 148897 are as follows: $T_{\text{eff}} = 4350$ K, $\log g = 1.0$ (cgs), and $v_t = 2.0$ km s $^{-1}$. An independent determination of the surface gravity, using the above T_{eff} value, Mennessier et al. (1997) absolute visual magnitude ($M_V = -1.8$), a bolometric correction of -0.5 and a mass of $1 M_{\odot}$, yields $\log g = 1.1$, in agreement with the adopted value (see Luck 1991, for a discussion of the discrepancy usually observed between the photometric and spectroscopic gravities). Comparison with atmospheric parameters from the literature is presented in Table 3.

The abundance analysis has been performed with the standard LTE line analysis program WIDTH9 developed by Kurucz. The model atmospheres were taken from Gustafsson et al. (1975). The synthetic spectra were generated using the spectral synthesis code STARSP (Tsymbol 1996). Oscillator strengths have been taken from the VALD database (Piskunov et al. 1995). The resulting abundances (normalized by the solar-system abundances of Grevesse & Sauval 1998) are listed in Table 3, from which it may be concluded that HD 148897 appears to be a rather typical metal-deficient star (McWilliam 1997), and should certainly *not be considered as a (metal-deficient) barium star*.

Interestingly enough, there are no indications whatsoever from the Hipparcos and Tycho data (applying the methods




Fig. 5. The iron abundances derived from Fe I (open circles) and Fe II (filled circles) lines are displayed as a function of the excitation potential for the lower energy level of the line. The absence of a trend in these data is used to derive the spectroscopic temperature of HD 139409.




Fig. 6. The abundances derived from the Fe I (open circles) and Fe II (filled circles) lines are displayed as a function of the line equivalent widths. The absence of a trend in these data is used to derive the microturbulent velocity of HD 139409.

described in Sect. 4) that this star is binary, in agreement with the fact that it is not a barium star.

HD 139409

The effective temperature of HD 139409 has been derived from the excitation equilibrium of Fe I, Ti I and Cr I lines (see Fig. 5, for Fe). The surface gravity $\log g$ was determined from the Fe I/Fe II ionization balance, and the microturbulent velocity by forcing the abundances of individual Fe I, Ti I and Cr I lines to be independent of equivalent width (see Fig. 6, for Fe). Although spectroscopic gravity and temperature determinations in late-type, metal-deficient stars are probably affected by non-LTE effects, these effects remain small when $[\text{Fe}/\text{H}] \geq -1.0$ (see, for example, Allende Prieto et al. 1999). The stellar parameters of Arcturus ($[\text{Fe}/\text{H}] = -0.6$, $\log g = 1.3$) derived by the spectroscopic method (applied on the HARPS spectrum) are in good agreement indeed with those derived by other (non-spectroscopic) methods.

The resulting atmospheric parameters for HD 139409 are as follows: $T_{\text{eff}} = 5000$ K, $\log g = 2.8$ (cgs), and $v_t = 2.0$ km s $^{-1}$. The spectroscopic gravity, combined with a mass of $1 M_{\odot}$, leads to $M_V \simeq +1.5$. Thus our calculations indicate

Table 4. Abundances for HD 139409, in the scale where $\log \epsilon(\text{H}) = 12$, and normalized with respect to the solar abundances (Grevesse & Sauval 1998). The standard deviations σ and the number N of lines used in the analysis are also given.

Element(X)	Z	$\log \epsilon(X)$	σ	N	[X/Fe]
C I	6	8.46	0.10	4	+0.36
Na I	11	6.02	0.05	5	+0.11
Mg I	12	7.31	0.12	6	+0.15
Si I	14	7.33	0.06	20	+0.20
Ca I	20	6.20	0.07	6	+0.26
Sc II	21	2.79	0.06	10	+0.04
Ti I	22	4.77	0.09	51	+0.17
Ti II	22	4.72	0.11	4	+0.12
V I	23	3.76	0.11	23	+0.18
Cr I	24	5.21	0.11	11	-0.04
Cr II	24	5.36	0.09	3	+0.11
Mn I	25	5.13	0.11	9	+0.16
Fe I	26	7.08	0.08	136	-
Fe II	26	7.08	0.12	14	-
Ni I	28	5.86	0.07	44	+0.03
Y I	39	2.22	0.10	2	+0.40
Y II	39	2.41	0.07	6	+0.59
Zr I	40	2.58	0.09	3	+0.40
Zr II	40	2.78	0.12	3	+0.60
Ba II	56	1.89	0.12	3	+0.18
La II	57	1.05	0.12	3	+0.30
Ce II	58	1.46	0.14	5	+0.30
Nd II	60	1.38	0.09	6	+0.30




Fig. 7. Abundance pattern in HD 139409 (abundances derived from ionized species are represented by black dots, and from neutral species by crosses).

- checking for astrometric orbital motion, indirectly from a comparison of the Hipparcos and Tycho-2 proper motions.

These methods are now described in turn.

4.1. Radial-velocity variations

Several stars from Table 5 (namely HIP 4347, HIP 29740, HIP 34795, HIP 55852, HIP 58956, HIP 76605 and BD +3°2688) have been monitored for many years using the CORAVEL spectrovelocimeter (Baranne et al. 1979), as part of a larger program aiming at finding the frequency of spectroscopic binaries among s-process-rich late-type giants (see Jorissen & Mayor 1988, 1992; Jorissen et al. 1998, for details and other results from this CORAVEL monitoring). Individual radial-velocity measurements for those stars (in the CORAVEL-ELODIE system as defined in Udry et al. 1999) are given in Table 7 or in Udry et al. (1998). For a few other stars (HIP 43042, CS 22942-019 and CS 22948-027), radial velocities were monitored using other instruments, and their results were taken from the literature (Preston & Sneden 2001; Začs et al. 2005).

Orbits were already available for HIP 4347 and HIP 29740 (Udry et al. 1998), HIP 43042 (Začs et al. 2005), as well as for CS 22942-019 and CS 22948-027 (Preston & Sneden 2001). A new orbital solution has been derived for BD +04°2466 (Table 6 and Fig. 8). The binary nature of those stars is therefore beyond doubt.

The radial-velocity standard deviation of HIP 58596 is larger than expected based on the uncertainty on one measurement (Fig. 9), but no satisfactory orbital solution could be found. A 3-d orbit (with an eccentricity of 0.30) is possible, but this short orbital period is not consistent with the giant nature of HIP 58596, which imposes orbital periods of at least 20 d (see Fig. 4 of Pourbaix et al. 2004). The large standard deviation exhibited by HIP 58596 is therefore very likely another example of the large intrinsic jitter often observed for metal-deficient stars, as discussed by McClure (1984) and Carney et al. (2003).

A 14 y radial-velocity monitoring for HIP 34795 with the northern and southern CORAVELs is not very conclusive either, mostly because there are difficulties in finding the zero-point offset between the two instruments for such large radial velocities (Udry et al. 1999). When a -1 km s^{-1} offset is applied to the northern velocities (with respect to the values listed in Table 7), a long-term trend seems to be present, albeit with some superimposed jitter (Fig. 11). The analysis of the Hipparcos astrometric data presented in Sect. 4.2 suggests that the star might be binary, although the evidence is not very conclusive.

Finally, there is no sign of radial-velocity variations for BD +3°2688 (Fig. 10).

4.2. Orbital motion from Hipparcos astrometric data

A tailored reprocessing of the Hipparcos *Intermediate Astrometric Data* (hereafter IAD; van Leeuwen & Evans 1998) makes it possible to look for a possible orbital signature in the astrometric motion, following the method outlined by Pourbaix & Jorissen (2000), Pourbaix & Boffin (2003), Pourbaix (2004) and applied to barium stars by Jorissen et al. (2004b). We give here only a brief summary of the method.

The basic idea is to quantify the likelihood of the fit of the Hipparcos astrometric data with an orbital model. For that

Table 5. Binary properties of confirmed or suspected metal-deficient barium stars. The column labeled “Ba” indicates whether detailed chemical analyses have confirmed the Ba nature of the star. The columns labeled $\chi^2_{\mu_{\text{HIP}} - \mu_{\text{Tyco}}}$ and $\text{Prob}(\chi^2 > \chi^2_{\mu_{\text{HIP}} - \mu_{\text{Tyco}}})$ provide the χ^2 (and its associated probability) involved in the comparison of Hipparcos and Tycho-2 proper motions (see text for details). The column “Bin.” has been set to “y” if the first kind risk of rejecting the null hypothesis that the Tycho and Hipparcos proper motions are equal is smaller than 10%. The column labeled IAD indicates whether the signature of an orbital motion is present in the Hipparcos Intermediate Astrometric Data, according to the various tests described in the text. The column labeled $\sigma(Vr)$ provides the radial-velocity standard deviation, Δt and N correspond to the time span and number of observations, respectively. The columns labeled P and “Ref” list the orbital period (when available) and the reference for the radial velocity and/or orbital data. The column “Binary” gives the final binary diagnostics.

HIP	HD/DM	Ba	Ref.	ϖ (mas)	$\mu_{\text{HIP}} - \mu_{\text{Tyco}}$			IAD	$\sigma(Vr)$ (km s ⁻¹)	Δt (d)	N	P (d)	Ref.	Binary
					χ^2_{obs}	$\text{Prob}(\chi^2 > \chi^2_{\text{obs}})$	Bin.							
4347	5424			0.22 ± 1.42	1.05	0.59	n	n	2.32	3306	13	1881	6	y
11595	15589			2.03 ± 1.21	1.63	0.44	n	n	–	–	–	–	–	?
25161	-27°2233			0.89 ± 1.35	0.23	0.89	n	n	–	–	–	–	–	?
29740	43 389			-1.25 ± 1.00	4.54	0.10	y	y	3.85	3350	24	1689	6	y
34795	55 496	y	5	2.44 ± 1.04	0.02	0.99	n	y	0.73	5121	24	–	2	y?
58596	104 340	y	3	-0.97 ± 1.09	3.39	0.18	n	n	1.64	2587	16	–	2	n
69834	123 396			1.73 ± 0.86	1.98	0.27	n	n	–	–	–	–	–	?
76605	139 409	y	2	5.51 ± 1.14	4.98	0.08	y	n	0.66	1478	2	–	2	y?
97874	187 762			2.07 ± 1.53	9.00	0.01	y	y	–	–	–	–	–	y
107478	206 983	y	3	3.75 ± 1.86	2.94	0.23	n	y	–	–	–	–	–	y?
Additional stars (from refs. (1), (5) and (10))														
43042	+75°348	y	1	1.02 ± 1.32	0.12	0.94	n	y	4.64	1436	33	1042	11	y
55852	+4°2466	y	4,5	0.96 ± 1.83	0.43	0.81	n	n	3.38	6191	41	4592	2	y
	+3°2688	y	5	-	-	-	-	-	0.45	986	7	-	2	n?
	CS 22942-019	y	10	-	-	-	-	-	3.42	3274	15	2800	10	y
	CS 22948-027	y	10	-	-	-	-	-	1.87	2560	13	505	10	y

References to the table: (1) Začs et al. (2000); (2) this work (3) Junqueira & Pereira (2001); (4) Burris et al. (2000); (5) Luck & Bond (1991); (6) Udry et al. (1998); (10) Preston & Sneden (2001); (11) Začs et al. (2005).

Table 6. Orbital elements for BD +04°2466.

P (d)	4592.7 ± 51.1
e	0.286 ± 0.02
ω (deg)	266.8 ± 5.3
V_γ (km s ⁻¹)	$+39.29 \pm 0.11$
K (km s ⁻¹)	5.67 ± 0.12
T (JD)	$2\,445\,076.6 \pm 73.0$
$f(M)$ (M_\odot)	0.076
$a_1 \sin i$ (Gm)	343.18
N	41

purpose, Pourbaix & Arenou (2001) (see also Jancart et al. 2005) introduced several statistical indicators which allow us to decide whether to keep or to discard an orbital solution. Those indicators relevant to our purpose are the following:

- The addition of 4 supplementary parameters (the four Thiele-Innes orbital constants) describing the orbital motion should result in a statistically significant decrease of the χ^2 for the fit of the N IAD with an orbital model with 9 free parameters (χ^2_T), as compared to a fit with a single-star solution with 5 free parameters (χ^2_S). This criterion is expressed by an F -test:

$$Pr_2 = Pr[\hat{F} < F(4, N - 9)], \quad (1)$$




Fig. 8. Radial velocities as a function of heliocentric Julian Day for BD +4°2466, superimposed on the orbital solution corresponding to the orbital elements listed in Table 6.

where

$$\hat{F} = \frac{N - 9}{4} \frac{\chi^2_S - \chi^2_T}{\chi^2_T}. \quad (2)$$




Fig. 9. Radial velocities as a function of heliocentric Julian Day for HIP 58596 = HD 104340 (open circles correspond to measurements obtained with the southern CORAVEL, and filled circles with the northern one).




Fig. 10. Radial velocities as a function of heliocentric Julian Day for BD +3°2688.

Pr_2 is thus the first kind risk associated with the rejection of the null hypothesis: “*there is no orbital wobble present in the data*”.

- Getting a substantial reduction of the χ^2 with the Thiele-Innes model does not necessarily imply that the four Thiele-Innes constants A, B, F, G are significantly different from 0. The first kind risk associated with the rejection of the null hypothesis “*the orbital semi-major axis is equal to zero*” may be expressed as

$$Pr_3 = Pr[\chi_{ABFG}^2 < \chi^2(4)], \quad (3)$$

where

$$\chi_{ABFG}^2 = \mathbf{X}^t \mathbf{V}^{-1} \mathbf{X}, \quad (4)$$




Fig. 11. Radial velocities as a function of heliocentric Julian Day for HIP 34795 = HD 55496. The symbols are as in Fig. 9. With respect to the data listed in Table 7, an offset of -1 km s^{-1} has been applied to the northern measurements to make them consistent with the southern ones.

and \mathbf{X} is the vector of components A, B, F, G and \mathbf{V} is its covariance matrix².

- For the orbital solution to be a significant one, its parameters should not be strongly correlated with the other astrometric parameters (e.g., the proper motion). In other words, the covariance matrix of the astrometric solution should be dominated by its diagonal terms, as measured by the *efficiency* ϵ of the matrix being close to 1 (Eichhorn 1989). The efficiency is simply expressed by

$$\epsilon = \sqrt{\frac{\prod_{k=1}^m \lambda_k}{\prod_{k=1}^m V_{kk}}}, \quad (5)$$

where λ_k and V_{kk} are respectively the eigenvalues and the diagonal terms of the covariance matrix \mathbf{V} .

With the above notations, the requirements for a star to qualify as a binary is then

$$\alpha \equiv (Pr_2 + Pr_3) / \epsilon \leq 0.02, \quad (6)$$

where the threshold value of 0.02 has been chosen to minimize false detections (Jorissen et al. 2004b).

Hipparcos data are, however, seldom precise enough to derive the orbital elements from scratch. Therefore, when a spectroscopic orbit is available beforehand, it is advantageous to import e, P, T from the spectroscopic orbit and to derive the remaining astrometric elements (as done by Pourbaix & Jorissen 2000; Pourbaix & Boffin 2003). If a spectroscopic orbit is not

² Since it may be shown that $\chi_S^2 - \chi_T^2 = \chi_{ABFG}^2$, the Pr_2 and Pr_3 tests are in fact equivalent provided that $\chi_T^2 \sim N - 9$. Thus, if Pr_2 and Pr_3 are significantly different, it means either that the Thiele-Innes orbital model does not fit the data very well ($\chi_T^2 \gg N - 9$), or that it fits much better than could be expected ($\chi_T^2 \ll N - 9$). We are indebted to L. Lindgren for this clarification (see also Jancart et al. 2005).




Fig. 12. The α statistics (Eq. (6)) as a function of the trial orbital period (assuming $e = 0$) for metal-deficient barium stars. The period from the spectroscopic orbit (represented by an arrow) of HIP 29740 (upper left panel) indeed lies within the range of minimum α values. By comparison, the suspected binaries HIP 97874 and HIP 107478 are likely to have periods $P \sim 800$ d and ~ 250 – 450 d, respectively. For comparison, stars non-flagged as binaries have no (e, P, T) grid points with $\alpha < 0.02$ (see Fig. 13). The vertical dashed lines represent multiple, or integer fractions, of 1 y. At those periods, there is a strong correlation between the parallactic and orbital signals, which degrades the α statistics and makes binaries difficult to find at those 1-y alias periods.

available, trial (e, P, T) triplets scanning a regular grid (with $10 \leq P(\text{d}) \leq 5000$ imposed by the Hipparcos scanning law and the mission duration) may be used. The quality factor α is then computed for each trial (e, P, T) triplet, and if there exist triplets yielding $\alpha < 0.02$, the star is flagged as a binary. To test its success rate, this method has been applied by Jorissen et al. (2004b) on a sample of barium stars. These authors show that, when $\varpi > 5$ mas and $100 < P(\text{d}) < 4000$, the (astrometric) binary detection rate is close to 100%, i.e., the astrometric method recovers all known spectroscopic binaries (see also Jancart et al. 2005). When the orbit is not known beforehand,

the method makes it even possible to find a good estimate for the orbital period, provided, however, that the true period is not an integer fraction, or a multiple, of one year. Here the method is applied to the sample of metal-deficient barium stars listed in Table 5.

The method flags as definite binaries the stars HIP 29740, 34795, 43042, 97874 and 107478 (Figs. 12 and 13). In two cases (HIP 29740 and 43042), the IAD method thus confirms the conclusion from the radial-velocity monitoring, but yields as well three new binaries (HIP 34795, 97874 and 107478). Two spectroscopic binaries (HIP 4347 and




Fig. 13. Same as Fig. 12 for HIP 34795 (*left panel*, flagged as binary) and HIP 58596 (*right panel*, non-binary).

HIP 55852) are not detected by the IAD method because of their small parallax or long orbital period. The non-binary nature of HIP 58596, already suspected from the radial-velocity data, is confirmed by the analysis of the IAD (Fig. 13).

4.3. Orbital motion from a comparison of Hipparcos and Tycho-2 proper motions

Kaplan & Makarov (2003) suggested that the comparison of Hipparcos and Tycho-2 (Høg et al. 2000b) proper motions offers a way to detect binaries with long periods (typically from 2000 to 4000 d). The Hipparcos proper motion, being based on observations spanning only 3 y, may be altered by the orbital motion, especially for systems with periods in the range of 2000 to 4000 d whose orbital motion was not recognized by Hipparcos. On the other hand, this effect should average out in the Tycho-2 proper motion, which is derived from observations covering a much longer time span. This method, already used by Makarov (2004), Pourbaix (2004) and Jancart et al. (2006), works best when applied to stars with parallaxes in excess of about 5 mas.

The method evaluates the quantity

$$\chi_{\text{obs}}^2 = (\boldsymbol{\mu}_{\text{HIP}} - \boldsymbol{\mu}_{\text{Tyc}})^t \mathbf{W}^{-1} (\boldsymbol{\mu}_{\text{HIP}} - \boldsymbol{\mu}_{\text{Tyc}}), \quad (7)$$

where $\boldsymbol{\mu}_{\text{HIP}}$ and $\boldsymbol{\mu}_{\text{Tyc}}$ are the vectors of α and δ components of the Hipparcos and Tycho-2 proper motions, respectively, and \mathbf{W} is the associated 2×2 variance-covariance matrix. The covariance between $\mu_{\alpha, \text{HIP}}$ and $\mu_{\delta, \text{HIP}}$, as provided by field H28 of the Hipparcos catalogue (ESA 1997) and the correlation between Tycho-2 and Hipparcos proper motions, as encapsulated in the quantity R of Table 1 of Høg et al. (2000a), have both been considered (see Jancart et al. 2006, for details).

Since the above quantity follows a χ^2 probability distribution function with 2 degrees of freedom, it is then possible to compute the probability $\text{Prob}(\chi^2 > \chi_{\text{obs}}^2)$, giving the first kind

risk of rejecting the null hypothesis $\boldsymbol{\mu}_{\text{Tycho}} = \boldsymbol{\mu}_{\text{HIP}}$ while it is actually true. This probability is listed in Table 5, along with χ_{obs}^2 , and the star is flagged as binary if $\text{Prob} < 0.1$.

Only HIP 29740, HIP 76605 and HIP 97874 satisfy the test at the 10% threshold. Note, however, that all the other stars have parallaxes smaller than 5 mas, which make the test less efficient.

5. Summary of the binary criteria and discussion

The situation may be summarized as follows (see also last column of Table 5):

- Definite binaries with known orbits: HIP 4347, HIP 29740 (passes all three binarity tests), HIP 43042, HIP 55852, CS 22942-019, CS 22948-027;
- Suspected binaries from astrometric data (either IAD or proper motions; no or inconclusive radial-velocity data): HIP 34795, HIP 76605, HIP 97874 (both astrometric tests yield positive results), HIP 107478;
- Data inconclusive (mainly because of too small a parallax): HIP 11595, HIP 25161, HIP 69834;
- Non-binary stars (mainly from radial-velocity data, not contradicted by astrometry): HIP 58596, BD +3°2688.

The latter two non-binary stars are in fact good candidate thermally-pulsing AGB stars, as revealed by their location in the HR diagram (Table 2 and right panel of Fig. 1). Hence, they must not be binaries.

Finally, we come to the central question of this paper: why do the metal-deficient barium stars, despite being binaries and occupying the same location of the HR diagram as YSyS, do not exhibit symbiotic activity? Three possible answers have been suggested in this paper:

- (i) Some among the stars listed in Table 2 and displayed in Fig. 1 are in fact not barium stars (especially HIP 80843 = HD 148897).




Fig. 14. Comparison of the period distributions for samples of binary systems with different kinds of red giant primaries: S stars (Jorissen et al. 1998), red SyS (excluding symbiotic novae and symbiotic Miras; Mürset & Schmid 1999), M giants (Jorissen et al. 2004a) and K giants (Mermilliod 1996). The lower two panels present the orbital-period distribution for barium stars (Jorissen et al. 1998) and yellow SyS (Mürset & Schmid 1999). In the S star panel, the arrow marked H_{α} indicates the period range where binary S stars exhibit H_{α} emission as a signature of weak symbiotic activity. The shaded area marks stars with only a lower limit available on their orbital period. The numbers in the upper right corner of each panel correspond to the sample size.

- (ii) Among those which are barium stars, some seem to lie on the TP-AGB, and thus need not be binaries (HIP 58596 = HD 104340, BD +3°2688). They therefore cannot exhibit symbiotic activity.
- (iii) Finally, there remain a few genuine metal-deficient barium stars in the sample. Why are they not symbiotic stars? It seems that the answer to that question lies in the different period distributions for YSyS and metal-deficient barium stars: YSyS have shorter orbital period than metal-deficient barium stars, as seen in Fig. 14. This argument seems to apply especially to HIP 29740 (=HD 43389), which has been assigned a very bright absolute visual magnitude of -3.5 by the maximum likelihood method of Mennessier et al. (1997). It is therefore expected to have a rather strong mass loss rate, and be a good candidate YSyS. However, with its orbital period of 1689 d, it lies at the long-period edge of the period distribution of YSyS (Fig. 14). The same difference seems to exist between the period distributions of red symbiotics and binary S stars (Van Eck & Jorissen 2002, and Fig. 14).

However, a firm conclusion on this issue should await the determination of the orbital periods for the metal-deficient barium stars flagged as binaries by the IAD method (especially HIP 11595, HIP 25161, HIP 69834, HIP 97874, HIP 107478),

so as to make the comparison between the orbital period distributions of metal-deficient barium stars and YSyS more meaningful.

Acknowledgements. This work was performed in the framework of the NATO Collaborative Linkage Grant SA (PST.CLG.979128)6774/FP. We thank D. Pourbaix for the processing of the Hipparcos data of the metal-deficient stars. F. Carrier and X. Bonfils are thanked for obtaining the Arcturus and HD 139409 spectra. L.Z. thanks I. Platais for valuable discussions and support. The *Fonds National Suisse de la Recherche Scientifique* has funded the operations of the CORAVEL spectrometer and of the Swiss 1-m telescope installed at the Haute-Provence Observatory.

References

- Allen, D. A. 1982, in *The Nature of Symbiotic Stars*, ed. M. Friedjung, & R. Viotti (Dordrecht: Reidel), IAU Coll., 70, 27
- Allende Prieto, C., García López, R. J., Lambert, D. L., & Gustafsson, B. 1999, *ApJ*, 527, 879
- Baranne, A., Mayor, M., & Poncet, J. L. 1979, *Vistas Astron.*, 23, 279
- Barbuy, B., Jorissen, A., Rossi, S. C. F., & Arnould M. 1992, *A&A*, 262, 216
- Bessell, M. S., Castelli, F., & Plez, B. 1998, *A&A*, 333, 231
- Boyle, R. J., & McClure, R. D. 1975, *PASP*, 87, 17
- Burris, D. L., Pilachowski, C. A., Armandroff, T. E., et al. 2000, *ApJ*, 544, 302
- Carney, B. W., Latham, D. W., Stefanik, R. P., Laird, J. B., & Morse, J. A. 2003, *AJ*, 125, 293
- Clayton, D. D. 1988, *MNRAS*, 234, 1
- Corradi, R., & Schwarz, H. E. 1997, in *Physical Processes in Symbiotic Binaries and Related Systems* (Warsaw: Copernicus Foundation for Polish Astronomy), 147
- Corradi, R. L. M., Brandi, E., Ferrer, O. E., & Schwarz, H. E. 1999, *A&A*, 343, 841
- Drake, S. A., Simon, T., & Linsky, J. L. 1987, *AJ*, 93, 163
- Eichhorn, H. 1989, *Bull. Astron. Inst. Czechosl.*, 40, 394
- ESA 1997, *The Hipparcos and Tycho Catalogues*, ESA
- Galazutdinov, G. 1992, *Special Astrophys. Obs. Prepr.*, 92, 52
- Grauer, A. D., & Bond, H. E. 1981, *PASP*, 93, 630
- Grevesse, N., & Sauval, A. J. 1998, *Space Sci. Rev.*, 85, 161
- Griffin, R. F., & Keenan, P. C. 1992, *The Observatory*, 112, 168
- Gustafsson, B., Bell, R. A., Eriksson, K., & Nordlund, A. 1975, *A&A*, 42, 407
- Hinkle, K., Wallace, L., Valenti, J., & Harmer, D. 2000, *Visible and Near Infrared Atlas of the Arcturus Spectrum 3727–9300 Å* (San Francisco: ASP)
- Høg, E., Fabricius, C., Makarov, V. V., et al. 2000a, *A&A*, 357, 367
- Høg, E., Fabricius, C., Makarov, V. V., et al. 2000b, *A&A*, 355, L27
- Jancart, S., Jorissen, A., Babusiaux, C., & Pourbaix, D. 2005, *A&A*, in press
- Jancart, S., Jorissen, A., Babusiaux, C., & Pourbaix, D. 2006, *A&A*, in preparation
- Jeffries, R. D., & Stevens, I. R. 1996, *MNRAS*, 279, 180
- Jorissen, A. 2003a, in *Symbiotic stars probing stellar evolution*, ed. R. L. M. Corradi, J. Mikołajewska, & T. J. Mahoney, (San Francisco), ASP Conf. Ser., 303, 25
- Jorissen, A. 2003b, in *Asymptotic Giant Branch Stars*, ed. H. Habing, & H. Olofsson (New York: Springer Verlag), 461
- Jorissen, A., Famaey, B., Dedecker, M., et al. 2004a, *Rev. Mex. Astron. Astrof. Conf. Ser.*, 21, 71

- Jorissen, A., Jancart, S., & Pourbaix, D. 2004b, in *Spectroscopically and Spatially Resolving the Components of Close Binary Stars*, ed. R. W. Hilditch, H. Hensberge, & K. Pavlovski (San Francisco), ASP Conf. Ser., 318, 141
- Jorissen, A., & Mayor, M. 1988, *A&A*, 198, 187
- Jorissen, A., & Mayor, M. 1992, *A&A*, 260, 115
- Jorissen, A., Schmitt, J. H. M. M., Carquillat, J. M., Ginestet, N., & Bickert, K. F. 1996, *A&A*, 306, 467
- Jorissen, A., Van Eck, S., Mayor, M., & Udry, S. 1998, *A&A*, 332, 877
- Junqueira, S., & Pereira, C. B. 2001, *AJ*, 122, 360
- Kaplan, G. H., & Makarov, V. V. 2003, *Astron. Nachrichten*, 324, 419
- Keenan, P. C., & McNeil, R. C. 1989, *ApJS*, 71, 245
- Kyröläinen, J., Tuominen, I., Vilhu, O., & Virtanen, H. 1986, *A&AS*, 65, 11
- Lattanzio, J. C. 1991, *ApJS*, 76, 215
- Lü, P. K. 1991, *AJ*, 101, 2229
- Lü, P. K., Dawson, D. W., Upgren, A. R., & Weis, E. W. 1983, *ApJS*, 52, 169
- Luck, R. E. 1991, *ApJS*, 75, 579
- Luck, R. E., & Bond, H. E. 1991, *ApJS*, 77, 515
- MacConnell, D. J., Frye, R. L., & Upgren, A. R. 1972, *AJ*, 77, 384
- Makarov, V. V. 2004, *ApJ*, 600, L71
- Mayor, M., Pepe, F., Queloz, D., et al. 2003, *The Messenger*, 114, 20
- McClure, R. D. 1984, *ApJ*, 280, L31
- McWilliam, A. 1997, *ARA&A*, 35, 503
- Mennessier, M. O., Luri, X., Figueras, F., et al. 1997, *A&A*, 326, 722
- Mermilliod, J.-C. 1996, in *The Origins, Evolution and Destinies of Binary Stars in Clusters*, ed. E. F. Milone, & J.-C. Mermilliod (San Francisco: ASP), ASP Conf. Ser., 90, 95
- Munari, U., & Buson, L. M. 1992, *A&A*, 255, 158
- Munari, U., Yudin, B. F., Taranova, O. G., et al. 1992, *A&AS*, 93, 383
- Mürset, U., Nussbaumer, H., Schmid, H. M., & Vogel, M. 1991, *A&A*, 248, 458
- Mürset, U., & Schmid, H. M. 1999, *A&AS*, 137, 473
- Musaev, F. A., Galazutdinov, G. A., Sergeev, A. V., Karpov, N. V., & Podyachaev, Y. V. 1999, *Kinematika I Fizika Nebesnyh Tel*, 15, 282
- Pereira, C. B., & Porto de Mello, G. F. 1997, *AJ*, 114, 2128
- Pereira, C. B., Smith, V. V., & Cunha, K. 1998, *AJ*, 116, 1977
- Pinsonneault, M. H., Sneden, C., & Smith, V. V. 1984, *PASP*, 96, 239
- Piskunov, N. E., Kupka, F., Ryabchikova, T. A., Weiss, W. W., & Jeffery, C. S. 1995, *A&AS*, 112, 525
- Pourbaix, D. 2004, in *Spectroscopically and Spatially Resolving the Components of Close Binary Stars*, ed. R. W. Hilditch, H. Hensberge, & K. Pavlovski (San Francisco), ASP Conf. Ser., 318, 132
- Pourbaix, D., & Arenou, F. 2001, *A&A*, 372, 935
- Pourbaix, D., & Boffin, H. M. J. 2003, *A&A*, 398, 1163
- Pourbaix, D., & Jorissen, A. 2000, *A&AS*, 145, 161
- Pourbaix, D., Tokovinin, A. A., Batten, A. H., et al. 2004, *A&A*, 424, 727
- Preston, G. W., & Sneden, C. 2001, *AJ*, 122, 1545
- Schaller, G., Schaerer, D., Meynet, G., & Maeder, A. 1992, *A&AS*, 96, 269
- Schmid, H. M., & Nussbaumer, H. 1993, *A&A*, 268, 159
- Schwarz, H. E. 1991, *A&A*, 243, 469
- Šleivyte, J., & Bartkevičius, A. 1990, *Vilnius Astronomijos Observatorijos Biuletenis*, 85, 3
- Smith, V. V., Cunha, K., Jorissen, A., & Boffin, H. M. J. 1996, *A&A*, 315, 179
- Smith, V. V., Cunha, K., Jorissen, A., & Boffin, H. M. J. 1997, *A&A*, 324, 97
- Theuns, T., Boffin, H. M. J., & Jorissen, A. 1996, *MNRAS*, 280, 1264
- Tsymbal, V. 1996, in *Model Atmospheres and Spectrum Synthesis*, ed. S. J. Adelman, F. Kupka, & W. W. Weiss (San Francisco: ASP), ASP Conf. Ser., 108, 198
- Udry, S., Jorissen, A., Mayor, M., & Van Eck, S. 1998, *A&AS*, 131, 25
- Udry, S., Mayor, M., & Queloz, D. 1999, in *Precise Stellar Radial Velocities*, ed. J. B. Hearnshaw, & C. D. Scarfe (San Francisco), ASP Conf. Ser., 170, 367
- Van Eck, S., & Jorissen, A. 2002, *A&A*, 396, 599
- van Leeuwen, F., & Evans, D. W. 1998, *A&AS*, 130, 157
- Vilhu, O., Tuominen, I., Kyröläinen, J., & Holsti, N. 1977, *Obs. Astrophys. Lab. Univ. Helsinki Rep. No. 3*
- Yamashita, Y., & Norimoto, Y. 1981, *Annals of the Tokyo Astronomical Observatory*, 18, 125
- Začs, L., Schmidt, M. R., Musaev, F. A., Galazutdinov, G. A., & Sperauskas, J. 2005, *A&A*, 441, 303
- Začs, L., Schmidt, M. R., & Schuster, W. J. 2000, *A&A*, 358, 1022

Online Material

Table 7. Individual CORAVEL radial velocities and associated errors for HIP 34795, HIP 58596, HIP 76605, BD +3°2688 and BD +04°2466. The last column indicates which one of the two CORAVEL spectrovelocimeters has been used (NO = CORAVEL north at the *Observatoire de Haute Provence*; SO = CORAVEL south on the Danish 1.54-m telescope at ESO).

HIP 34795 = HD 55496				
DDMMYY	HJD	RV (km s ⁻¹)	ε (km s ⁻¹)	
050183	45 340.695	315.30	0.33	SO
151283	45 684.797	314.79	0.33	SO
120684	45 864.434	315.71	0.46	SO
310185	46 097.658	314.78	0.34	SO
271185	46 397.826	314.95	0.30	SO
111186	46 746.858	314.56	0.31	SO
101286	46 775.595	316.07	0.56	NO
150187	46 811.682	314.98	0.31	SO
160288	47 208.394	316.01	0.49	NO
231088	47 458.704	315.83	0.70	NO
151288	47 511.857	315.11	0.33	SO
090289	47 567.393	315.39	0.56	NO
241189	47 855.642	315.44	0.57	NO
070390	47 958.339	317.36	0.66	NO
050490	47 987.497	314.78	0.31	SO
061190	48 202.697	315.63	0.45	NO
031290	48 229.645	316.10	0.56	NO
180391	48 334.380	314.95	0.62	NO
110393	49 058.361	315.00	0.67	NO
130393	49 060.338	314.32	0.80	NO
180194	49 371.471	315.37	0.58	NO
301295	50 082.752	313.99	0.34	SO
231196	50 411.600	316.66	0.56	NO
120197	50 461.480	317.72	0.63	NO
HIP 58596 = HD 104340				
DDMMYY	HJD	RV (km s ⁻¹)	ε (km s ⁻¹)	
141289	47 875.832	265.29	0.45	SO
140392	48 696.805	264.18	0.30	SO
120393	49 059.755	266.40	0.29	SO
190393	49 066.707	264.89	0.33	SO
050493	49 083.708	265.41	0.28	SO
200493	49 098.453	263.93	0.46	NO
080394	49 420.525	263.34	0.47	NO
030494	49 446.435	266.38	0.41	NO
140594	49 487.360	265.37	0.46	NO
260195	49 744.638	264.59	0.56	NO
300395	49 807.475	263.16	0.52	NO
210795	49 920.516	264.34	0.30	SO
191295	50 071.721	265.60	0.37	NO
150296	50 129.556	263.70	0.43	NO
130197	50 462.684	265.61	0.45	NO
HIP 76605 = HD 139409				
DDMMYY	HJD	RV (km s ⁻¹)	ε (km s ⁻¹)	
090790	48 082.629	62.43	0.30	SO
260794	49 560.513	63.36	0.27	SO

Table 7. continued.

HIP 55852 = BD +04°2466				
DDMMYY	HJD	RV (km s ⁻¹)	ε (km s ⁻¹)	
070282	45 008.814	38.11	0.69	SO
130684	45 865.491	44.73	0.74	SO
030285	46 100.832	45.14	0.68	SO
290385	46 154.675	44.78	0.58	SO
200186	46 451.831	43.54	0.53	SO
140286	46 476.783	43.32	0.49	SO
291186	46 764.706	42.05	0.96	NO
090187	46 805.864	41.36	0.49	SO
080287	46 835.808	41.99	0.41	SO
280188	47 189.812	40.50	0.43	SO
040388	47 225.726	39.83	0.41	SO
250488	47 277.657	40.17	0.48	SO
051288	47 501.707	37.84	0.69	NO
060389	47 592.565	38.14	0.64	NO
220589	47 669.526	37.64	0.49	SO
241289	47 885.857	37.48	0.45	SO
050290	47 928.800	36.83	0.48	SO
160290	47 939.743	37.05	0.43	SO
260290	47 949.595	37.00	0.69	NO
040490	47 986.631	37.01	0.42	SO
110490	47 993.594	35.99	0.49	SO
050590	48 017.647	36.35	0.61	SO
020690	48 045.501	36.82	0.46	SO
260291	48 314.487	34.56	0.73	NO
260391	48 342.724	34.92	0.47	SO
150392	48 697.747	34.26	0.42	SO
120393	49 059.726	33.77	0.39	SO
070493	49 085.670	33.15	0.40	SO
080493	49 086.448	32.60	0.70	NO
100294	49 394.609	34.54	0.77	NO
020394	49 414.713	35.84	0.43	SO
070394	49 419.515	34.92	0.52	NO
080494	49 451.424	37.51	0.88	NO
140594	49 487.351	36.78	0.64	NO
310195	49 749.592	39.24	0.74	NO
150395	49 792.681	41.24	0.44	SO
020495	49 810.460	40.55	0.44	NO
190795	49 918.506	41.94	0.53	SO
191295	50 071.715	43.32	0.60	NO
130197	50 462.674	44.33	0.90	NO
200199	51 199.634	41.42	0.80	NO
BD +03°2688				
DDMMYY	HJD	RV (km s ⁻¹)	ε (km s ⁻¹)	
080493	49 086.481	33.16	0.43	NO
050394	49 417.506	32.35	0.47	NO
030494	49 446.510	32.33	0.44	NO
140594	49 487.388	33.00	0.47	NO
030295	49 752.658	33.52	0.52	NO
300395	49 807.506	32.64	0.71	NO
201295	50 072.729	33.07	0.46	NO