

Proper identification of RR Lyrae stars brighter than 12.5 mag^{*}

G. Maintz

Sternwarte der Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany
e-mail: gmaintz@astro.uni-bonn.de

Received 12 April 2005 / Accepted 13 July 2005

ABSTRACT

RR Lyrae stars are of great importance for investigations of Galactic structure. However, a complete compendium of all RR-Lyraes in the solar neighbourhood with accurate classifications and coordinates does not exist to this day. Here we present a catalogue of 561 local RR-Lyrae stars ($V_{\max} \leq 12.5$ mag) according to the magnitudes given in the Combined General Catalogue of Variable Stars (GCVS) and 16 fainter ones. The Tycho2 catalogue contains ≈ 100 RR Lyr stars. However, many objects have inaccurate coordinates in the GCVS, the primary source of variable star information, so that a reliable cross-identification is difficult. We identified RR Lyrae from both catalogues based on an intensive literature search. In dubious cases we carried out photometry of fields to identify the variable. Mennessier & Colome (2002, A&A, 390, 173) have published a paper with Tyco2-GCVS identifications, but we found that many of their identifications are wrong.

Key words. astrometry – stars: variables: RR Lyr – catalogs

1. Introduction

RR Lyrae stars (RR Lyrs) are classical variables of the instability strip. They are one of the primary distance indicators, which is why they have often been investigated to determine their absolute magnitude and their distances (see, e.g., references in Fernley et al. 1998). In addition, RR Lyrs provide an excellent sample of stars to investigate kinematics in the galaxy (see, e.g., Beers et al. 2000; Layden 1994; Layden et al. 1996; Dambis & Rastorguev 2001; Martin & Morrison 1998; Altmann & de Boer 2000).

Several studies have recently been carried out on other horizontal-branch (HB) stars. Altmann et al. (2004) published a study of the orbits and halo distribution of sdB stars, extending earlier work by de Boer et al. (1997). A similar study on RHB stars has been completed, too (Kaempfer et al. 2005). It thus appeared worthwhile to also investigate the orbits and the ensuing z -distribution of RR Lyrs (Maintz & de Boer 2005).

For such purposes the distance of the stars as well as accurate data of proper motion and radial velocity are required. A good source for astrometric data is the Tycho-2 catalogue (Høg et al. 2000, Tyco2) since it provides astrometric data for more than 2.5 million stars. However, one first has to identify the RR Lyrs in that catalogue.

We took the Combined General Catalogue of Variable Stars (Kholopov et al. 1998, GCVS) as a basic source for compiling an unbiased sample of RR Lyrs. The GCVS is, as the authors

write, “the only reference source on all known variable stars ... The total number of designated variable stars has now reached 31 918”. However, the coordinates given in GCVS are not as accurate as one would like for, specifically, cross-correlations with other catalogues. The positions in GCVS are said to be accurate “to 1 s of arc”. But we found that this accuracy is too optimistic. In some cases the coordinates denoted not the variable but another star. In even more cases, at the location given by the coordinates there was no star at all. According to the authors of GCVS, “the Sternberg Institute has started preparation of an electronic release GCVS 4.2 which will contain improved light elements, classifications etc., along with sufficiently accurate positional information”.

In this paper we present a list of RR Lyrs, with $V_{\max} \leq 12.5$ mag, all present in the GCVS, but now with absolutely reliable positions. It includes 286 Hipparcos and Tycho stars (ESA 1997; HIP), of which 104 only in Tyco2, as well as 273 in the GSC (Lasker et al. 1996), and a few USNO-A2 stars (the USNO-A2 is published by Monet et al. 1998). Our list contains the names used in each of these catalogues.

2. Comparing catalogues

As a first step we began with an identification of RR Lyr stars in the GCVS with a brightness at maximum of $V_{\max} \leq 12.5$ mag. We then proceeded to inspect their positions by looking up the original finding charts. It emerged from that already that several RR Lyrs were not identified properly. Thus a full inspection of all the basic data was required.

^{*} Full Table 1 is only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/442/381>




Fig. 1. Light-curve of PW Lac (*left panel*) and FN Lyr (*right panel*) observed at the observatory of Maintz. Such data was used to identify the RR Lyrae stars in their fields.

Because it was impossible to observe all RR Lyrs ourselves, and especially stars from the southern hemisphere could not be observed by us, we made a literature search to find the missing ones. We followed very strict principles. First, papers which give a finding chart or image identifying the variable by observing its light-curve were regarded as a reliable reference. Second, we used papers giving the Tyc2 or GSC number of the star, by the same criteria. Beside this we tried to find the original record of discovery and the finding chart given there. It takes considerable effort to do so, especially if only very old finding notes exist, which are not available electronically.

All images, finding charts and records of the discoverer were examined in the same way as our own observations. We compared them with the DSS and overlaid them with Tyc2 and GSC from the Vizier catalogue service using the Aladin interactive sky atlas. However, for some GCVS stars the identification was still elusive and it was decided to go and observe the fields anew to properly identify the particular RR Lyrs.

3. Observation and identification of RR Lyrae stars

The best way to identify RR Lyrs in the field of stars is to observe the fields and look for the star with RR Lyr like variability.

For 34 fields time series were obtained in Bonn, using a 8" Schmidt-cassegrain Meade telescope and an OES Alpha-Mini CCD-camera with an IRR-UV-Cut filter (B&W 486). This equipment is fully sufficient for taking series of images for several hours allowing to define the light-curve. Usually about every minute an image was taken. For an example of light-curves see Fig. 1.

The data reduction was made with the OES-Fleischmann software provided with the camera. We identified the RR Lyr easily by its characteristic variability. We then compared our image with the DSS as described in Sect. 2. With this procedure we extracted name and coordinates of the star in question from the Tyc2 and GSC catalogue, respectively.

For 22 stars time series were observed at the Observatory Hoher List with the 1 m Cassegrain-Telescope using the Cassegrain focus with focal reducer ($f = 368$ cm) and the HoLiCam 2048×2048 pix CCD camera. For the reduction of these CCD-images we used standard CCD reduction routines




Fig. 2. Light-curve of CE Vul observed at the observatory “Hoher List” of the Sternwarte of the Univ. of Bonn. Such data was required to identify RR Lyrae stars in their fields.

(DAOPHOT under IRAF) to determine positions and magnitudes of all stars on the frames. The RR Lyr in question was identified by its light-curve (for an example see Fig. 2) and its identification number and coordinates were obtained by comparing with the DSS overlaying with the Tyc2 or GSC, too.

4. The final catalogue

We give all RR Lyrs with $V_{\max} \leq 12.5$ mag (according to GCVS) in one list, but we mention that the 186 RR Lyrs from the HIP which are included in our catalogue, are taken from HIP directly needing no further identification.

Our final catalogue contains 577 RR Lyrs with the following information in the columns

- 1: name of the variable (GCVS);
- 2+3: catalogue (H = Hip, T = Tyc2, G = GSC, UA2 = USNO – A2) + number; an asterisk points at an individual note to the star¹;
- 4+5: α, δ (2000) given in the particular catalogue;
- 6: V magnitude in maximum light;
- 7: class of RR Lyr based on light-curve;
- 8+9: first epoch and period;
- 10: date of last observed maximum light known by us;
- 11: source of first epoch and period if not taken from GCVS or HIP.

If the star is taken from HIP, the magnitude in maximum light, type of variability, first epoch and period are from the HIP Variability Annex. If this information comes from another catalogue, it is from the GCVS, except when new information for first epoch and period are available in the literature.

The date of the last observation of maximum light mostly comes from amateur observers. They regularly observe variables of all kinds and they obtain light-curves of RR Lyrs as well, achieving correct maximum epochs. This column is, although not complete, very useful to make predictions of maximum light.

¹ The source of identification is available from gmaintz@astro.uni-bonn.de

Table 1. 561 RR-Lyrae Stars with maximum light ≤ 12.5 mag according to GCVS plus 16 fainter ones.

Name of star	Cat	Nr	α_{2000}	δ_{2000}	V_{\max} (mag)	Type	First ep. (JD)	Per. (d)	Last observ. (JD)	Source of period
SW And	H	1878	00 23 43.090	29 24 03.62	9.22	RRab	2448500.0384	0.4422620000	53260.1300	
XX And	H	6029	01 17 27.415	38 57 02.03	10.20	RRab	2448500.6580	0.7227550000	53045.3410	
AT And	H	116958	23 42 30.832	43 00 51.66	10.51	RRab	2448500.2310	0.6169170000	53232.5870	
BK And	G	323500127	23 35 06.022	41 06 10.51	12.50	RRab	2429146.4500	0.4215985000	51321.8076	
CI And	H	8939	01 55 08.294	43 45 56.47	11.78	RRab	2448500.0022	0.4847280000	52695.3060	
DK And	G*	364500701	23 28 45.910	50 34 29.35	12.50	RR:	2429130.4070	0.2436553000		
DM And	T	277300253	23 32 00.707	35 11 48.87	12.40	RRab	2435717.4310	0.6303890000	51335.8300	
DR And	T	228600352	01 05 10.707	34 13 06.24	12.00	RRab	2437220.3190	0.5631180000	52620.4385	
DU And	G	283600362	02 30 31.337	40 50 33.11	12.50	RR	2436051.4500	0.6067160000	51469.4641	
OV And	T	278701874	00 20 44.206	40 49 40.80	10.40	RRab	2439026.4780	0.4705810000	52983.4265	1
V395 And	H	117111	23 44 32.143	46 22 48.59	7.57	RRc:	2448500.2660	0.3423280000		
WY Ant	H	50289	10 16 04.946	-29 43 42.41	10.40	RRab	2448500.5339	0.5743410000	51869.6630	
SY Aps	G*	926500960	14 39 24.130	-72 49 35.62	12.20	RR:	2451904.0650	0.2789100000		2
TY Aps	H	72444	14 48 50.012	-71 19 41.88	11.27	RRab	2448500.4217	0.5016950000	52823.5250	
TY Aps	H	72444	14 48 50.012	-71 19 41.88	11.27	RRab	2448500.4217	0.5016950000	52823.5250	
UY Aps	UA2	0150 14253525	14 59 34.440	-71 47 53.77	12.00	RRab	2425326.5400	0.4825200000	51921.9800	
VX Aps	G	942900076	15 59 56.690	-75 13 21.00	11.50	RRab	2434239.3610	0.4845780000	51927.5010	
...	
...	

The sources of first epoch and period if not from HIP or GCVS:

- 1 Rossiger, S., & Busch, H. 1988, Mitt. Verand. Sterne 11, 133.
- 2 A. Paschke, priv. communication.

... ..
... ..

Individual notes:

- DK And type EW is also possible. A. Paschke, priv. communication.
 SY Aps type EW is also possible. A. Paschke, priv. communication.

If a star could not be identified in its field, because no second source for identification besides the GCVS was available (and no variable was found in our observation of the field) our catalogue has 999999999 as number. This happened in eight cases. The position given for these stars is taken from the GCVS. If there are no data for period or first epoch in the GCVS and no information is found otherwise, its first epoch and period is given as 9999999.9999 and 9.99999999, respectively.

For 6 stars without Tyc2 or GSC number we give identifications and coordinates (2000) from the USNO-A2.0. Magnitudes and elements are taken from the GCVS.

For three stars (DM And, CQ Lac and UZ Eri) being members of Tyc2, there are no proper motions available in the Tyc2. Therefore those stars are not useful for further investigations based on Tyc2.

X CMi is identified as GSC 168-406 while Schmidt et al. (1995) identified it as GSC 168-562. However, comparing their finding chart with the chart from Tsessevich & Kazanasmas (1971), we found that in both charts there is only one star at the position of X CMi while the POSS gives a close pair. GSC 168-562 is the preceding star of the pair and is very faint

(15.25 mag according to GSC III aJ). That is why we adopt that X CMi is the following star (168-406) with a magnitude of 12.34 mag (GSC III aJ).

Mennessier & Colome (2002, hereafter MC) gave results for 172 RR Lyrae stars identified to be Tyc2 stars. They did the identification blindly, using the GCVS as source of the positions and magnitudes of the RR Lyrs. They were successful in finding many RR Lyrs at Tyc2 star positions. But they did not find all and, as it turned out during our search, their list is wanting since they misidentified a large fraction of their stars.

For 78 stars of their 172 we agree with the identifications of MC. In 93 cases we found their stars being misidentified. We found 27 Tyc2 stars which are not given by MC. From the misidentified ones we found 84 in the GSC and 3 in the USNO-A2, while for 3 Tyc2 stars (NO Cas, V672 Aql, V1823 Cyg²) we found another identification number than given by MC. The misidentification mostly happens, when the star was fainter than the magnitudes given in the GCVS. Of the MC paper stars, 3 are not mentioned in our list: V1823 Cyg (see IBVS 4997), UUCam (Schmidt et al. 1995), CE Aqr

² According to IBVS Nr 4997, V1823 Cyg is a W-UMa star and not a RR Lyrae star. It is of course not included in our catalogue.

(A. Paschke, priv. commun.), *because they are not* RR Lyrs. 14 stars of their list are fainter than our limiting magnitude, but we added these stars to our list for completeness.

5. Summary

We present a catalogue listing 561 RR Lyrs with $V_{\max} \leq 12.5$ mag (plus 16 fainter ones) with exact coordinates. Most of them come from the GCVS. 21 have recently been recovered (of the new found RR Lyrs 2 have a V_{\max} of 12.52 and 12.53 mag respectively). 27 RR Lyrs are previously unknown as members of Tyc2. Data of first epoch and period are given if available. The dates of the last maxima observed come mostly from amateur observers and (even if this data is incomplete) allow to calculate trust-able predictions.

As an example, the beginning of the catalogue is given in Table 1.

Acknowledgements. This research has made use of the Simbad database, operated at CDS, Strasbourg, France. We thank Karl Wälke and Wolfgang Grimm who donated the Atlas of Finding Charts for Variable Stars by Tsevisch and Kazanasmas. We thank Franz Agerer, Wolfgang Moschner, Konstantin von Poschinger and K. Bernhard for sending several identifications of stars from their observations. We thank Anton Paschke for sending maximum light times of recent observations.

References

- Altmann, M., & de Boer, K. S. 2000, A&A, 353, 135
 Altmann, M., Edelmann, H., & de Boer, K. S. 2004, A&A, 414, 181
 Beers, T. C., Chiba, M., Joshii, Y., et al. 2000, AJ, 119, 2688
 de Boer, K. S., Aguilar Sanchez, Y., Altmann, M., et al. 1997, A&A, 327, 577
 Dambis, A. K., & Rastorguev, A. S. 2001, AstL, 27, 108
 ESA, 1997, The Hipparcos and Tycho Catalogues, ESA SP1200, and its Variability Annex: Periodic variables
 Fernley, J., Barnes, T. G., Skillen, I., et al. 1998, A&A, 330, 515
 Høg, E., Fabricius, C., Makarov, V. V., et al. 2000, Tycho-2 catalogue (Tyc2)
 Kaempf, K. S., de Boer, K. S., & Altmann, M. 2005, A&A, 432, 879
 Kholopov, P. N., Samus, N. N., Durlevich, O. V., et al. 1990, General Catalogue of Variable Stars, 4rd edn., Vol. IV (Moscow: Nauka) (GCVS)
 Lasker, B. M., Russel, J. N., & Jenkner, H. 1996, The HST Guide Star Catalogue, Version 1.2
 Layden, A. C. 1994, AJ, 108, 1016
 Layden, A. C., Hanson, R. B., Hawley, S. L., et al. 1996, AJ, 112, 2110
 Maintz, G., & de Boer, K. S. 2005, A&A, 442, 229
 Martin, J. C., & Morrison, H. L. 1998, AJ, 116, 1724
 Mennessier, M. O., & Colomé, J. 2002, A&A, 390, 173
 Schmidt, E. G., Chab, J. R., & Reiswig, D. E. 1995, AJ, 109, 1239
 Monet, D., Bird, A., Canzian, B., et al. 1998, A Catalogue of Astrometric Standards (USNO-A V2.0)
 Tsevisch, V. P., & Kazanasmas, M. S. 1971, Atlas of Finding Charts for Variable Stars, Moscow