Identification of a new short-period comet near the sun[★]

S. F. Hönig

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany
 e-mail: shoenig@mpifr-bonn.mpg.de

Received 5 August 2005 / Accepted 6 September 2005

ABSTRACT

We present the identification of comet C/1999 R1 (SOHO) with comet C/2003 R5 (SOHO). Both apparitions were only observed with the Solar and Heliospheric Observatory (SOHO) at distances smaller than 0.1 AU from the sun with the LASCO coronagraphs onboard the spacecraft. Although SOHO comets usually have poor orbital coverage, the 1999 and 2003 arcs are sufficient to generate a link that seems to satisfy all observations. We also analyze comet C/2002 R5 (SOHO) which has similar orbital elements. A fragmentation scenario is proposed and discussed which would support the linkage of C/1999 R1 and C/2003 R5 and thus its short periodic nature.

Key words. comets: general – comets: individual: SOHO (C/1999 R1) – comets: individual: SOHO (C/2003 R5) – comets: individual: SOHO (C/2002 R5)

1. Introduction

Since 1996 the *Solar and Heliospheric Observatory (SOHO)* has regularly observed comets close to the sun with its Large Angle Spectrometric coronagraph (LASCO) C2 and C3 cameras. C2 has a circular field of view with a diameter of around 1.5° while C3 reaches out to almost 8° solar elongation. Most of the observed comets belong to the well-known Kreutz comet group (Biesecker et al. 2002). A number of non-Kreutz comets with perihelion distances $0.01 \text{ AU} < q < 0.1 \text{ AU}$ have appeared. Similarities in orbital elements of a number of objects led to the establishment of three other groups of near-sun comets (Green 2002; Meyer 2003).

While the Meyer group has a mean inclination $i = 72.44^\circ$, comets of the Marsden and Kracht groups have rather small inclinations (see Table 1). Shortly after the discovery of these new groups Marsden (2002) remarked on similarities in the orbital elements of the Marsden group comets and comet 96P/Machholz, which has already been associated with various meteor streams. Discussions on the long-term evolution of the Quadrantids (Williams et al. 1979) and 96P (Rickman & Froeschlé 1988) have been brought together by McIntosh (1990), who included the Daytime Arietids as well as the Southern and perhaps Northern δ Aquarids into this complex of meteor streams and a comet. More recently it was suspected that the near-earth asteroid 2003 EH1 is the parent body of the Quadrantids (Jenniskens 2003) and that C/1490 Y1, the Ursids, Carinids and κ Velids might also fit into the scheme

Table 1. Mean orbital elements for new comet groups.

	Meyer group	Marsden group	Kracht group
q [AU]	0.0359	0.0481	0.0449
ω [deg]	57.26	22.65	59.38
Ω [deg]	72.98	81.92	42.78
i [deg]	72.44	27.14	13.25

q ...Perihelion distance, ω ...Argument of perihelion,
 Ω ...Ascending node, i ...Inclination.

(McIntosh 1990; Babadzhanov & Obrubov 1992; Ohtsuka et al. 2003; Sekanina & Chodas 2005). Given that the Kracht and Marsden groups fit into this complex with short periodic components in the range of 4 to 6 years and their low inclination, attempts have been made to find identifications of objects with two perihelion passages within the comet groups. Two pairs of Marsden group comets could be found where formal links have been presented by Marsden (2004b, 2005). On the other hand, two suspect Kracht group pairs could not be unequivocally linked due to poor measurements and ambiguities.

There exist another three SOHO comets with similar low- i orbital elements which do not fit into the Machholz interplanetary complex. These are C/1999 R1 (SOHO), C/2002 R5 (SOHO) and C/2003 R5 (SOHO). Remarks on the similarity of the apparent C2 tracks of C/1999 R1 and C/2002 R5 came from Kracht shortly after discovery. He also noted the possible relation of C/2003 R5 to this pair.

[★] Table of residuals is only available in electronic form at <http://www.edpsciences.org>

Table 2. Parabolic orbital elements for the individual apparitions from C2 observations only that have been used as a starting point for the identification, in comparison to the actual C2 link. For the elements two data points from 2003 with overall residuals larger than 50 arcsec were excluded.

	C/1999 R1 (SOHO)	C/2003 R5 (SOHO)	C/1999 R1 = C/2003 R5 (SOHO)
Epoch	1999 Sep. 5.0	2003 Sep. 8.0	2003 Sep. 8.0
T	1999 Sep. 5.53	2003 Sep. 8.82	2003 Sep. 8.82
q [AU]	0.0556	0.0569	0.0570
e	1.0	1.0	0.97757
ω [deg]	43.88	44.04	43.58
Ω [deg]	5.65	4.81	5.04
i [deg]	13.60	14.14	13.67
P [yr]	n/a	n/a	4.01
rms ["]	5.4	13.7	13.6

T ...Time of perihelion, q ...Perihelion distance, e ...Eccentricity, ω ...Argument of perihelion, Ω ...Ascending node, i ...Inclination, P ...Orbital period, rms...Root mean square of residuals.

2. Observations and initial orbits

Astrometric positions and initial parabolic orbital elements for C/1999 R1 (SOHO), C/2002 R5 (SOHO) and C/2003 R5 (SOHO) have been published in *MPEC 1999–R19*, *MPEC 2002–S35* and *MPEC 2004–J59*, respectively. First indications about similarities of C/1999 R1 and C/2003 R5 come from their appearance in SOHO images: both cover the same orbital arc and have same apparent tracks, a fact noted by Kracht shortly after discovery. In addition, they showed a very similar brightness evolution during their passages through C2 and C3, without any sign of degrading or break-up. On the other hand, C/2002 R5 was less bright which results in a shorter orbital arc. Its apparent track, however, was quite similar to the other comets and led Kracht to the conclusion that they may form the core of another new comet group (Marsden 2004a).

3. Identification C/1999 R1 (SOHO) = C/2003 R5 (SOHO)

Orbital calculations were performed using the *Find_Orb* software¹. It includes perturbations by all major planets plus the Earth's moon and the three largest minor planets (1) Ceres, (2) Pallas and (4) Vesta. Additionally, relativistic effects due to the comet's small perihelion distance are considered in the software.

The first step is to choose the most validated observations. C3 has a scale of 55"/pixel, while C2 has 12"/pixel. For the initial orbital calculation we thus chose only C2 observations. It has to be noted that the 2003 data show higher individual residuals than the 1999 observations, regardless of whether they are taken as the starting point or not. For the calculations, initial parabolic elements for one apparition were taken and the eccentricity e was forced also to match the other apparition. After a rough fit this constraint was removed and further integration steps led to convergence of the final link. Table 2 presents elements for C/1999 R1 (SOHO) and C/2003 R5 (SOHO) from C2 observations only. They were used as a

Table 3. Orbital elements of the C2 double weighted link (see text) for the 2003 apparition, as well as a prediction for the next perihelion passage in 2007.

	C/1999 R1 = C/2003 R5 (SOHO)	
Epoch	2003 Sep. 9.0	2007 Sep. 11.0
T	2003 Sep. 8.819	2007 Sep. 11.263
q [AU]	0.0570	0.0538
e	0.97741	0.97860
ω [deg]	43.713	48.612
Ω [deg]	5.022	0.019
i [deg]	13.687	12.727
P [yr]	4.01	3.99
rms ["]	28.213	

starting point for the linking. Additionally, a link with both C2 and C3 data was calculated (Table 3) where individual C2 observations are weighted 6 times more than individual C3 observations. Considering that there are around 3 times as many C3 positions as C2, the latter are overweighted by roughly a factor of two – although C3 still covers a much longer orbital arc. The residuals are presented in Table A.1. Table 3 shows predictions for the next perihelion passage calculated from the C2 double weighted elements. The whole solution appears to be stable in the sense that use of shorter integration steps does not lead to other elements with smaller residuals or residuals with similar quality. These can be taken as an indication that the identification may be correct. Actually a comet with a period of around 4 years is somewhat unusual since there is only comet 2P/Encke with a shorter period. Further arguments for the periodicity of the object, however, will be given in Sect. 4 and the nature of the object – comet or asteroid – discussed in Sect. 5.

4. The role of C/2002 R5 (SOHO)

Since C/2002 R5 (SOHO) has similar orbital elements to C/1999 R1 (SOHO) = C/2003 R5 (SOHO) it is worth considering whether there may be some physical association.

¹ See http://www.projectpluto.com/find_orb.htm

Fig. 1. Backwards integration of various possible orbital elements for C/2002 R5 (SOHO). Each line represents one set of elements with a certain $3.40 \text{ yr} < P < 4.65 \text{ yr}$, according to the results for C/1999 R1 = C/2003 R5 (SOHO). The black line indicates the orbital evolution of C/1999 R1 = C/2003 R5 (SOHO), green and blue lines represent elements with $P < 3.90 \text{ yr}$ and $P > 3.95 \text{ yr}$, respectively. The red lines mark the elements for $P = 3.90 \text{ yr}$ and $P = 3.95 \text{ yr}$ which show best overall agreement with C/1999 R1 = C/2003 R5 (SOHO) (see text). The x -axis of each panel represents Julian Date (JD), while the y -axis represents the Ascending Node Ω , Inclination i , Perihelion Distance q and Eccentricity e , respectively.

Calculations show that a direct linkage with a periodicity of $P \sim 3.0 \text{ yr}$ to C/1999 R1 is not possible due to large overall residuals and systematic trends therein. The same applies to the 2003 apparition and C/2002 R5. Together with its difference in brightness compared to C/1999 R1 and C/2003 R5 we can rule out that it is identical to one of the latter two objects. In view of the fragmentation history of some members of the other low-inclination comet groups (Marsden 2005) C/1999 R1 = C/2003 R5 (SOHO) and C/2002 R5 (SOHO) may have a common origin. Thus, C/2002 R5 (SOHO) may have separated from the parent body C/1999 R1 = C/2003 R5 (SOHO) sometime in the past – which seems plausible due to the potential parent body’s recurring close approaches to the sun. Therefore, it is interesting to perform backwards integration of both objects to see if the orbital elements were more identical at a certain time in the past. For this purpose we have to assume that C/2002 R5 is also of short periodic nature. If it is possible to find hints for a common origin within this scenario, it would be a strong argument for the periodicity of C/1999 R1 = C/2003 R5 (SOHO) aside from the linkage of the orbital arcs.

As a starting point we assume that C/2002 R5 has similar periodicity to C/1999 R1 = C/2003 R5. This can be satisfied by the assumption of fragmentation. We did, however, take a broad range of orbital elements with $3.40 \text{ yr} < P < 4.65 \text{ yr}$ in steps of

0.05 yr for the backwards integration. A set of orbital elements for each step was calculated and put into the freely available orbital integrator *Solex* 8.5², developed by Vitagliano (1997). All 26 solutions for C/2002 R5, together with the orbital elements for C/1999 R1 = C/2003 R5 have been integrated backwards simultaneously. In general, orbits for SOHO comets show large rms residuals due to non-perfect astrometrical circumstances. Thus backwards integration is a problematic issue, especially in the case of C/2002 R5 where there is only a small orbital arc available and the data quality is poorer. Since we have the P -constraint from C/1999 R1’s orbit we think that integrating the abovementioned set of elements will give us a rough idea of the object’s past 200 000 days ($\sim 544 \text{ yr}$), although no final statement will be possible.

Our calculations illustrate that the orbital developments of C/1999 R1 = C/2003 R5 and C/2002 R5 are quite different within the simulated time span (see Fig. 1). The inclination evolution is quite extreme for most of the elements of C/2002 R5 in comparison to C/1999 R1 = C/2003 R5. There is, however, a certain time span around 2 410 000 JD where the elements were quite similar for both objects for periodic solutions around the interval $3.90 \text{ yr} < P < 3.95 \text{ yr}$. There is no other

² See <http://chemistry.unina.it/~alvitagl/solex/>

Table 4. Comparison of the orbital elements of C/2002 R5 (SOHO) with a initial periodicity of $P = 3.90$ yr to C/1999 R1 = C/2003 R5 (SOHO) at around the assumed separation time in the late 19th century.

	C/1999 R1 = C/2003 R5	C/2002 R5 (SOHO)
q [AU]	0.060	0.060 ± 0.006
e	0.976	0.976 ± 0.002
a [AU]	2.52	2.51 ± 0.25
ω [deg]	30.3	34.6 ± 3.0
Ω [deg]	18.3	24.2 ± 3.2
i [deg]	18.8	17.4 ± 1.1

time within the last 550 yr where such a good agreement in orbital elements can be achieved. Around 2 410 000 JD a good match of the Argument of Perihelion ω , the inclination i , the eccentricity e , the perihelion distance q , and thus also the semi-major axis a to C/1999 R1 = C/2003 R5 is found. Furthermore the difference in Longitude of Ascending Node Ω is smaller than 6° (see Table 4). Although the orbits of C/2002 R5 and C/1999 R1 = C/2003 R5 seem to be close together in the late 19th century, the difference in the angular elements hardly improved with respect to the present elements. Additionally, the longitude of perihelion of C/2002 R5 ($L = 58^\circ$) is still far from the value for C/1999 R1 = C/2003 R5 ($L = 48^\circ$), while the latitude of perihelion matches quite well ($B = 9^\circ$ for both objects). This may be an effect of the unknown scatter in initial conditions for C/2002 R5. As a measure for this scatter we illustrated the uncertainties of the elements, based on the variations for one periodicity interval $dP = 0.05$ yr, in Table 4. The resulting deviation may partly explain the large difference in L while the agreement is still not striking. It is necessary to improve the quality of the current orbit. If C/2002 R5 is periodic and separated from the parent body in the late 19th century (probably around 1890 ± 10), we may soon reobserve it.

The first step to test this scenario would be a search for C/2002 R5 in LASCO data at the previous return. The problem is that it should have appeared in September or early October, 1998. At around this time there are no LASCO images available since SOHO experienced major problems. Therefore we encourage observations at the possible next return in 2006 (see Table 5). On the other hand, periodicities that placed the object around perihelion later than the end of October 1998 or earlier than July 1998 can almost certainly be ruled out since there was nothing found in the LASCO archives, although thoroughly inspected by several people. We want to emphasize that the actual time of perihelion for the next return is much more uncertain than the difference in the orbits may suggest.

5. Comet or minor planet?

It is widely known that almost no SOHO comet displays a tail, except for a number of Kreutz group members and very few other bright comets. Nevertheless, all objects are designated and referred to as comets, mostly because of the parabolic elements that fit into this scheme, and the origin

Table 5. Predicted orbital elements for a possible next return of C/2002 R5 (SOHO), based on $P = 3.90$ yr and $P = 3.95$ yr for the 2002 apparition, respectively.

	C/2002 R5 (SOHO)	
Epoch	2006 Jul. 30.0	2006 Aug. 17
T	2006 Jul. 30.4	2006 Aug. 17.7
q [AU]	0.0470	0.0470
e	0.981	0.981
ω [deg]	45.92	45.94
Ω [deg]	13.15	13.13
i [deg]	14.10	14.09
P [yr]	3.90	3.95

of the Kreutz group where small tail-less members are fragments of well-observed cometary progenitors. The suggested connection between comet 96P/Machholz and the Kracht and Marsden groups supports this procedure. Additionally there are hints that the identified recurring objects of these groups show a trend to have fainter second perihelion passages as a result of their losing a lot of material and thus desintegrate over time (Sekanina & Chodas 2005).

Given the semi-major axis $a \sim 2.52$ AU of C/1999 R1 = C/2003 R5, one may ask whether this object is really a comet or an Alinda-type asteroid close to the 3:1 resonance with Jupiter. No tail was visible, in either C3 or in C2 images. It is also not clear if the object displayed a coma. We accomplished an analysis of photometric data acquired during the 1999 perihelion passage (Biesecker 1999) and searched for possible phase effects which would be typical of asteroid behaviour. Photometric measurements from LASCO images are highly affected by a number of disturbing factors, such as severe radial vignetting or obscuration by the pylon (Biesecker et al. 2002). An analysis of the absolute magnitude of this object with standard minor planet ($G = 0.15$) and cometary ($n = 2.0$) slope parameters give no conclusive answer (Table 6). Both possibilities satisfy the photometric data and allow for similar standard deviation in their absolute magnitudes. The absolute value for H_{MP} is, however, not consistent with the standard translation to the object's size (~ 10 km). From estimations for other SOHO comets its size should be of the order of 10s of meters up to perhaps 100 m, depending on its actual nature.

6. Discussion

We present the identification of a new short periodic comet observed with the SOHO spacecraft. We linked the objects C/1999 R1 and C/2003 R5 as two consecutive perihelion passages of the same object. Orbital elements for the next return in 2007 were calculated. Additionally, we show that the third object, C/2002 R5, with similar orbital elements may be a fragment of the periodic comet that separated from it some time in the late 19th century. Aside from the orbital link this supports the idea of the short periodicity of C/1999 R1 = C/2003 R5. If the scenario is correct, C/2002 R5 may be observed in

Table 6. Determination of absolute magnitude from photometric data of C/1999 R1 (SOHO).

Magnitude m [mag]	Distance		Phase angle	Absolute magnitude	
	r (helioc.) [AU]	d (geoc.) [AU]	ϕ [deg]	cometary H_C [mag]	asteroidal H_{MP} [mag]
6.6	0.0577	0.9978	116.9	12.84	12.84
6.5	0.0573	0.9833	110.9	12.75	12.75
6.4	0.0572	0.9870	106.6	12.64	12.64
6.3	0.0573	0.9907	102.3	12.53	12.53
6.2	0.0575	0.9945	98.1	12.41	12.41
6.2	0.0580	0.9993	93.3	12.38	12.38
6.4	0.0592	1.0071	85.9	12.52	12.52
6.4	0.0600	1.0109	82.1	12.49	12.49
6.7	0.0609	1.0148	78.4	12.75	12.75
6.9	0.0619	1.0186	74.9	12.90	12.90
6.9	0.0645	1.0272	68.2	12.79	12.80
7.4	0.0676	1.0355	62.2	13.17	13.20
7.4	0.0706	1.0427	67.7	13.07	13.11

2006 with the SOHO satellite. In the last section we discussed the nature of C/1999 R1 = C/2003 R5 and concluded that from currently available photometric measurements it is impossible to tell if it is a comet or a minor planet. With this identification the object should also become accessible to ground-based observations, so that the situation may soon improve. Therefore, it is strongly encouraged to attempt ground-based observations of the object within the observable window just before the perihelion approach in 2007.

References

- Babadzhanov, P. B., & Obrubov, Y. V. 1992, *Astronomicheskii Vestnik*, 26, 70
- Biesecker, D. A. 1999, *IAU Circ.*, 7251
- Biesecker, D. A., Lamy, P., St. Cyr, O. C., Llebaria, A., & Howard, R. A. 2002, *Icarus*, 137, 323
- Green, D. W. E. 2002, *IAU Circ.*, 7832
- Jenniskens, P. 2003, *IAU Circ.*, 8252
- Marsden, B. G. 2002, *MPEC 2002-E25*
- Marsden, B. G. 2004a, *MPEC 2004-J59*
- Marsden, B. G. 2004b, *MPEC 2004-X73*
- Marsden, B. G. 2005, *MPEC 2005-E87*
- McIntosh, B. A. 1990, *Icarus*, 86, 299
- Meyer, M. 2003, *Int. Comet Quart.*, 25, 115
- Ohtsuka, K., Nakano, S., & Yoshikawa, M. 2003, *PASJ*, 55, 321
- Rickman, H., & Froeschlé, C. 1988, *Celest. Mech.*, 43, 243
- Sekanina, Z., & Chodas, P. W. 2005, *ApJ*, submitted
- Vitagliano, A. 1997, *Celest. Mech. Dyn. Astron.*, 66, 293
- Williams, I. P., Murray, C. D., & Hughes, D. W. 1979, *MNRAS*, 189, 483

Online Material

Table A.1. Observations for the orbital link with double-weighted C2 and single-weighted C3 residuals. Data were taken from *MPEC 1999-R19* and *MPEC 2004-J59*.

Coronagraph	Date and time [UT]	RA	Dec	Δ RA [arcsec]	Δ Dec [arcsec]
C2	1999 09 04.89595	10 48 46.9	+07 17 23	4.4+	2.0–
	1999 09 04.91256	10 48 37.0	+07 19 44	2.2–	2.0–
	1999 09 04.92090	10 48 32.3	+07 20 56	1.9–	.54–
	1999 09 04.93756	10 48 22.8	+07 23 13	3.8–	4.2–
	1999 09 04.97923	10 47 60.0	+07 29 05	.01+	1.3–
	1999 09 04.99590	10 47 50.8	+07 31 22	2.1–	2.8–
	1999 09 05.00423	10 47 46.1	+07 32 40	5.3–	6.2+
	1999 09 05.02090	10 47 37.9	+07 34 48	5.0+	3.2–
	1999 09 05.03756	10 47 28.7	+07 36 58	1.3–	9.6–
	1999 09 05.06344	10 47 15.2	+07 40 36	2.9–	1.8–
	1999 09 05.07924	10 47 08.1	+07 42 39	10.8+	5.9–
C3	1999 09 05.08756	10 47 03.3	+07 43 42	1.6+	9.5–
	1999 09 05.22100	10 45 59.6	+08 01 05	29.6–	10.0+
	1999 09 05.26261	10 45 44.2	+08 06 04	8.9–	10.6+
	1999 09 05.27923	10 45 38.6	+08 07 32	2.0+	17.3–
	1999 09 05.32090	10 45 24.5	+08 12 25	13.8+	5.9–
	1999 09 05.34590	10 45 13.9	+08 15 14	21.6–	.65+
	1999 09 05.36256	10 45 10.5	+08 16 40	4.6+	18.7–
	1999 09 05.38756	10 45 02.4	+08 19 39	7.7–	6.8+
	1999 09 05.40423	10 44 58.6	+08 21 14	2.9+	2.5+
	1999 09 05.42927	10 44 53.1	+08 23 32	14.4+	3.7–
	1999 09 05.47098	10 44 42.5	+08 27 15	8.5–	7.4–
	1999 09 05.48789	10 44 39.8	+08 28 45	1.6–	4.3–
	1999 09 05.51257	10 44 36.4	+08 30 55	8.4+	4.2+
	1999 09 05.52923	10 44 33.0	+08 32 17	6.6–	7.7+
	1999 09 05.57090	10 44 29.6	+08 34 57	12.4+	15.4–
	1999 09 05.59591	10 44 27.3	+08 36 59	6.8+	5.7+
	1999 09 05.61256	10 44 25.5	+08 37 51	6.5–	5.6–
	1999 09 05.63756	10 44 25.1	+08 39 12	.38–	14.0–
	1999 09 05.65423	10 44 24.2	+08 40 17	11.1–	4.8–
	1999 09 05.67924	10 44 25.1	+08 41 16	2.0–	23.7–
	1999 09 05.69590	10 44 26.4	+08 42 10	9.2+	17.9–
	1999 09 05.72090	10 44 26.7	+08 43 29	6.4–	5.5–
	1999 09 05.76261	10 44 30.3	+08 45 13	6.7–	2.2+
	1999 09 05.77923	10 44 32.8	+08 45 31	2.1+	13.1–
	1999 09 05.82090	10 44 37.8	+08 47 07	11.4–	11.7+
	1999 09 05.84590	10 44 42.8	+08 47 31	.77–	1.3+
	1999 09 05.86256	10 44 46.0	+08 47 29	.04–	20.3–
	1999 09 05.88757	10 44 51.6	+08 47 52	6.5+	21.9–
	1999 09 05.90424	10 44 54.6	+08 48 14	4.1–	13.1–
	1999 09 05.92923	10 45 00.8	+08 48 42	.65–	.35–
	1999 09 05.97099	10 45 10.5	+08 48 42	19.7–	14.0–
	1999 09 05.98756	10 45 15.3	+08 48 49	18.1–	8.5–
	1999 09 06.01257	10 45 23.8	+08 48 47	2.1–	8.7–
	1999 09 06.07091	10 45 42.2	+08 48 53	7.7–	19.3+
	1999 09 06.09591	10 45 50.0	+08 48 06	20.0–	11.1–
	1999 09 06.11256	10 45 57.2	+08 47 57	1.1–	6.8–
	1999 09 06.13758	10 46 05.5	+08 47 24	13.9–	16.5–
	1999 09 06.15423	10 46 11.5	+08 46 48	17.7–	35.0–
	1999 09 06.17923	10 46 20.3	+08 46 58	30.0–	4.2+
	1999 09 06.19590	10 46 28.0	+08 46 26	13.2–	6.3–
	1999 09 06.26262	10 46 54.3	+08 44 53	28.9–	.64+

Table A.1. continued.

Coronagraph	Date and time [UT]	RA	Dec	Δ RA [arcsec]	Δ Dec [arcsec]
C2	2003 09 08.20423	11 01 14.7	+06 06 52	11.5+	10.8–
	2003 09 08.21256	11 01 08.3	+06 08 03	9.0–	12.5–
	2003 09 08.22923	11 00 59.6	+06 10 29	10.5+	11.7–
	2003 09 08.24590	11 00 49.0	+06 13 11	.54+	5.6+
	2003 09 08.25423	11 00 44.6	+06 14 04	8.5+	13.5–
	2003 09 08.27117	11 00 34.5	+06 16 31	6.7+	12.7–
	2003 09 08.28756	11 00 25.7	+06 18 60	18.1+	4.5–
	2003 09 08.31362	11 00 10.0	+06 22 35	8.1+	12.1–
	2003 09 08.32923	11 00 00.5	+06 24 51	1.0–	8.5–
	2003 09 08.33756	10 59 56.2	+06 25 58	5.0+	11.9–
	2003 09 08.35423	10 59 46.7	+06 28 47	2.5+	16.9+
	2003 09 08.37090	10 59 38.3	+06 31 21	14.7+	31.7+
	2003 09 08.37923	10 59 27.5	+06 31 41	78.3–	17.5–
	2003 09 08.39590	10 59 23.1	+06 35 38	9.2–	81.8+
	2003 09 08.41256	10 59 17.0	+06 36 45	32.4+	12.3+
C3	2003 09 08.52923	10 58 13.8	+06 51 21	45.0–	30.5–
	2003 09 08.57090	10 57 53.8	+06 58 07	65.9–	66.5+
	2003 09 08.59591	10 57 48.0	+07 00 42	4.8+	41.8+
	2003 09 08.61256	10 57 39.2	+07 02 10	25.7–	12.7+
	2003 09 08.63756	10 57 29.8	+07 04 38	21.1–	11.0–
	2003 09 08.65423	10 57 24.2	+07 06 43	12.5–	2.2+
	2003 09 08.67925	10 57 18.3	+07 09 36	31.0+	12.0+
	2003 09 08.72090	10 57 05.3	+07 12 56	37.3+	47.4–
	2003 09 08.76258	10 56 49.0	+07 17 28	29.8–	18.7–
	2003 09 08.77923	10 56 46.9	+07 19 09	1.9+	10.0–
	2003 09 08.80452	10 56 40.6	+07 21 30	4.2–	4.0–
	2003 09 08.82090	10 56 35.3	+07 22 51	31.4–	6.8–
	2003 09 08.86257	10 56 26.7	+07 26 32	46.5–	13.7+
	2003 09 08.88774	10 56 24.8	+07 28 32	19.6–	21.7+
	2003 09 08.90424	10 56 24.1	+07 29 39	.81+	19.0+
	2003 09 08.92925	10 56 21.1	+07 31 00	5.1–	.01–
	2003 09 08.97099	10 56 17.5	+07 33 33	15.4–	1.4+
	2003 09 08.98756	10 56 17.3	+07 34 29	8.6–	2.5+
	2003 09 09.01256	10 56 16.1	+07 35 27	19.3–	16.7–
	2003 09 09.02923	10 56 12.6	+07 36 37	71.8–	5.6+
	2003 09 09.07090	10 56 19.1	+07 38 08	6.3+	9.8–
	2003 09 09.09590	10 56 19.6	+07 39 30	9.1–	17.1+
	2003 09 09.11259	10 56 20.3	+07 39 42	18.6–	4.2–
	2003 09 09.13756	10 56 24.0	+07 40 48	.31–	17.2+
	2003 09 09.15423	10 56 24.4	+07 40 59	23.3–	1.8+
	2003 09 09.17923	10 56 28.5	+07 41 27	12.3–	4.9–
	2003 09 09.19590	10 56 32.7	+07 41 55	12.6+	3.3+
	2003 09 09.22090	10 56 34.5	+07 42 08	23.0–	8.8–
	2003 09 09.23758	10 56 37.1	+07 42 11	29.9–	19.5–
	2003 09 09.26256	10 56 45.1	+07 42 49	15.2+	2.4+
	2003 09 09.27923	10 56 47.2	+07 42 49	6.5–	5.4–
	2003 09 09.32090	10 56 56.0	+07 42 47	20.1–	17.4–
	2003 09 09.34618	10 57 01.3	+07 42 49	36.9–	15.0–
	2003 09 09.36256	10 57 03.6	+07 42 41	67.7–	20.3–
	2003 09 09.38757	10 57 13.8	+07 42 59	19.7–	5.5+
	2003 09 09.40424	10 57 22.7	+07 42 42	40.7+	4.0–
	2003 09 09.43055	10 57 27.9	+07 42 30	.13+	.55–
	2003 09 09.44590	10 57 31.4	+07 42 17	19.0–	2.5–
	2003 09 09.47090	10 57 41.2	+07 41 52	7.2+	6.6–
	2003 09 09.48758	10 57 45.3	+07 41 30	13.9–	12.6–
	2003 09 09.51256	10 57 51.6	+07 40 57	46.5–	18.8–
	2003 09 09.52923	10 58 01.2	+07 41 34	9.8+	37.9+
	2003 09 09.57090	10 58 10.6	+07 40 22	73.4–	21.4+