

## GCIRS34W: an irregular variable in the Galactic Centre<sup>★</sup>

S. Trippe<sup>1</sup>, F. Martins<sup>1</sup>, T. Ott<sup>1</sup>, T. Paumard<sup>1</sup>, R. Abuter<sup>1</sup>, F. Eisenhauer<sup>1</sup>,  
S. Gillessen<sup>1</sup>, R. Genzel<sup>1,2</sup>, A. Eckart<sup>3</sup>, and R. Schödel<sup>3</sup>

<sup>1</sup> Max-Planck-Institut für extraterrestrische Physik, Giessenbachstrasse, Postfach 1312, 85741 Garching, Germany  
e-mail: trippe@mpe.mpg.de

<sup>2</sup> Department of Physics, University of California, Berkeley, USA

<sup>3</sup> 1. Physikalisches Institut, Universität zu Köln, Zùlpicher Strasse 77, 50937 Köln, Germany

Received 1 April 2005 / Accepted 17 October 2005

### ABSTRACT

We report the results of time-resolved photometric and spectroscopic near-infrared observations of the Ofpe/WN9 star and LBV candidate GCIRS34W in the Galactic Centre star cluster. Diffraction limited resolution photometric observations obtained in *H*- and *K*-bands show a strong, non-periodic variability on time scales from months to years in both bands, accompanied by variations of the stellar colour. Three *K*-band spectra obtained in 1996, 2003, and 2004 with integral field spectrometers are identical within their accuracies and exclude significant spectroscopic variability. The most probable explanation of the stellar photometric variability is obscuration by circumstellar material ejected by the star. The approximated position of GCIRS34W in an HR diagram is located between O supergiants and LBVs, suggesting that this star is a transitional object between these two phases of stellar evolution.

**Key words.** Galaxy: center – stars: variables: general – stars: winds, outflows – stars: mass loss – stars: individual: GCIRS34W

### 1. Introduction

The class of Ofpe/WN9 stars has received attention in the last decade. Initially defined as stars with spectroscopic characteristics of both the most extreme O supergiants and late WN stars (Walborn 1982), these objects may be in an intermediate evolutionary state between these two types of stars (Bohannon & Walborn 1989; Nota et al. 1996; Pasquali et al. 1997). Since the observation by Stahl (1983) of the Ofpe/WN9 star R127 turning into a Luminous Blue Variable (LBV), a link between these two classes of objects is suspected. Stahl (1986) also showed that the LBV AG Car displays a typical Ofpe/WN9 star in its minimum phase, and Crowther et al. (1995) associated a few Ofpe/WN9 stars with dormant LBVs from comparison of their chemical composition and physical parameters.

LBVs are well known for their strong, episodic variability (Conti 1984). These stars are usually very luminous ( $L \approx 10^6 L_{\odot}$ ). They can experience “great eruptions” with *V* increasing by more than 2 mag (e.g.  $\eta$  Car, P Cyg), or they can stay in a relatively quiescent phase with low *V* band variation ( $\approx 0.5$  mag) and suddenly erupt with brightening of 1–2 mag (see Humphreys & Davidson 1994, for a review).

All mechanisms developed to explain the behaviour of LBVs (Glatzel & Kiriakidis 1993; Dorfi & Gautschy 2000; Humphreys & Davidson 1984; Lamers & Fitzpatrick 1988; Nugis & Lamers 2002) are associated with strong mass outflows (with  $\dot{M}$  as high as  $0.1 M_{\odot}/\text{yr}$  in case of  $\eta$  Car, Massey 2003), which is confirmed by the detection of nebulae around LBVs (Voors et al. 2000). Hence spectroscopic variability due to changes in both  $T_{\text{eff}}$  and  $\dot{M}$  is also common in LBVs.

In this article we present the results of photometric and spectroscopic long-term observations of the star GCIRS34W (hereafter referred to as IRS34W), which is part of the star cluster surrounding the supermassive black hole in the centre of our Milky Way. This star cluster, located at a distance of 7.6 kpc (Eisenhauer et al. 2005), is a system of dynamically relaxed old stars with an admixture of young, hot, non-relaxed stars; some of the young, hot stars show strong He I emission (Krabbe et al. 1991). Most of the young and massive stars appear to reside in two thin rotating disks centred on the Galactic Centre supermassive black hole Sgr A\* (Genzel et al. 2003; Paumard et al. 2005). Among these objects, around 30 have been identified as Wolf-Rayet stars and LBV candidates (Paumard et al. 2004). Another group of early- and late-type B main sequence stars can be found in the immediate vicinity (few tens of light days) of Sgr A\* (Genzel et al. 2003; Eisenhauer et al. 2005), leading to the so-called “paradox of youth”, i.e. the question, of how these massive and young stars managed to reside in an area that is so close to the black hole (Ghez et al. 2003).

<sup>★</sup> Based on observations at the New Technology Telescope (NTT) and the Very Large Telescope (VLT) of the European Southern Observatory (ESO), Chile, the 2.2-m-telescope of the Max-Planck-Gesellschaft (MPG) in La Silla, Chile, and the Gemini North telescope of the Gemini Observatory on Mauna Kea, Hawaii.

The star IRS34W is one of the young He I stars, and it was already identified as an Ofpe/WN9 star in earlier examinations (Krabbe et al. 1995; Najarro et al. 1997); additionally, IRS34W has been classified as an LBV candidate by Paumard et al. (2001). Therefore this star is of great interest for seeking an understanding of star formation and evolution in the Galactic Centre.

In this article we describe the observations and data reduction in Sect. 2, present and discuss the results in Sect. 3 and summarise our conclusions in Sect. 4.

## 2. Observations

### 2.1. Photometry

Photometric observations of IRS34W were obtained regularly since 1992. From 1992 to 2002 we used the camera SHARP I (Hofmann et al. 1992) at the 3.5-m-NTT of ESO in La Silla, Chile, in speckle-imaging mode to obtain diffraction limited resolution  $K$  band images. All images were sky-subtracted, bad-pixel-, and flat-field-corrected and then deconvolved with a Lucy-Richardson-algorithm (Lucy 1974; Richardson 1972); the spatial resolution was 130 milli-arcseconds (mas). On the processed images we applied aperture photometry, using a set of four stars located in the observed field as calibrators. Additionally in 1992, 1997, and 1998 we obtained  $H$  band images with a spatial resolution of 100 mas. These images were processed (except deconvolution) and analysed like the  $K$  images.

Since 2002, we have used the detector system NAOS/CONICA (NACO for short), consisting of the adaptive optics system NAOS (Roussel et al. 2003) and the NIR camera CONICA (Hartung et al. 2003) at the 8.2-m-UT4 (Yepun) of the ESO-VLT on Cerro Paranal, Chile. We obtained AO-corrected diffraction limited resolution images in  $H$  (spatial resolution 40 mas) and  $K$  bands (spatial resolution 55 mas). After sky subtraction, bad-pixel-, and flat-field-correction, all images were deconvolved with a Wiener filtering algorithm (Ott et al. 1999). On the deconvolved images, we applied aperture photometry, here using an ensemble of 12 stars in the field of view as reference sources.

Along with our observations, we included images from the Galactic Centre Demonstration Science Data Set obtained in 2000 with the 8-m-telescope Gemini North on Mauna Kea, Hawaii, using the AO system Hokupa'a in combination with the NIR camera Quirc. These images were processed by the Gemini team and released to be used freely. We analysed one  $H$  and one  $K$  band image obtained in July 2000 by applying aperture photometry calibrated with a set of three stars.

### 2.2. Spectroscopy

For a time-resolved spectroscopic analysis of IRS34W, we compared three spectra obtained in 1996, 2003, and 2004. The first spectrum was obtained in March 1996 using the integral field spectrometer 3D (Weitzel et al. 1996) at the 2.2-m-MPG-telescope in La Silla, Chile. The data output is structured as data cubes with two spatial axes, each 16 pixels

long, and one spectral axis of 256 pixels length, so 3D provides seeing-limited  $16 \times 16$  pixel images with a spectrum for each image pixel. The March 1996 observations covered the wavelength interval from 1.97 to 2.21  $\mu\text{m}$  (the short wavelength half of  $K$  band) with a spectral resolution of  $R = 1500$ . The seeing was around 0.6'', and the spatial pixelscale was 300 mas/pixel. After sky subtraction, bad-pixel-, and flat-field-correction, the spectrum was calibrated spectrally with a neon lamp. Atmospheric features were removed by dividing by the normalised spectrum of a calibration star, only in the range of low atmospheric transmission and high noise below 2.02  $\mu\text{m}$  the correction was incomplete.

The second spectrum was obtained in April 2003 with the integral field spectrometer SPIFFI (Eisenhauer et al. 2003a,b) at the VLT-UT2 (Kueyen). SPIFFI produced seeing-limited data cubes with 32 pixels in each spatial axis and 1024 pixels in the spectral axis. The data were corrected and calibrated (sky, bad pixels, flat field, atmosphere, wavelength calibration) like the 3D spectrum. Seeing was around 0.3'', and the spatial pixelscale was 100 mas/pixel. The final spectrum covered the complete  $K$  band from 1.95 to 2.45  $\mu\text{m}$  with a spectral resolution of  $R = 3600$ .


The third spectrum was obtained in August 2004 using SINFONI, a combination of SPIFFI and the adaptive optics (AO) system MACAO (Bonnet et al. 2003, 2004) at VLT-UT4. As SPIFFI was equipped with a new  $2K \times 2K$  detector in early 2004, the data cubes now had a dimension of  $64 \times 32$  pixels in the two spatial dimensions and 2048 pixels along the spectral axis. The data were processed and reduced like the earlier SPIFFI data. The pixel scale of the images was  $100 \times 50$  mas/pixel, meaning that each image pixel is rectangular. Observations were obtained at a seeing of around 0.8'', while the spatial resolution was improved to 0.3'' by the AO system. The spectrum covered the  $K$  band from 1.95 to 2.45  $\mu\text{m}$  with a spectral resolution of  $R = 4500$ .

We corrected each spectrum for background flux by subtracting a spectrum obtained in an empty field 0.9'' north of IRS34W. To assure the comparability of the spectra and to estimate the influence of contamination by the neighbouring stars (especially by IRS34E located 0.4'' north east of IRS34W), we obtained the 2003 SPIFFI spectrum twice: once on the original data cube, and once on a data cube in which (1) the spatial resolution (0.3'') was reduced to 3D resolution (0.6'') by convolving the spatial images with a Gaussian; (2) the image pixels (pixel scale 100 mas/pixel) were combined to pixels the size of 3D pixels (300 mas/pixel); and (3) the spectral resolution ( $R = 3600$ ) was reduced to that of 3D ( $R = 1500$ ) by convolving the spectral axis with a Gaussian.

## 3. Results and discussion

The  $H$  and  $K$  band light curves and colour of IRS34W obtained from the stellar photometry are presented in Fig. 1. The magnitudes used in this discussion were not corrected for the (not well-known) extinction by interstellar matter, and the given times are days after JD 2 448 000.

Both lightcurves showed variations in magnitude on timescales between few 100 and around 1500 days, while a



**Fig. 1.** *Top panel:*  $H$  and  $K$  band lightcurves of IRS34W. The time is given as reduced Julian Date; the first data point was taken in March 1992, the last in September 2004. The vertical dashed lines indicate the times when the three spectra were obtained in 1996, 2003, and 2004. Strong non-periodic variability on time scales from months to years is clearly visible in both light curves. *Bottom panel:*  $H-K$  derived from the lightcurves. The comparison to the lightcurves shows that the colour gets redder when the magnitudes decrease, and bluer when the magnitudes increase.

selection of representative data points (one value per calendar year) is presented in Table 1. The  $K$  lightcurve in particular, showing an overall variation of  $\Delta K \approx 1.5$ , did not show any periodicity or regularity, including periods of nearly constant magnitude, as well as steplike changes in brightness.

The stellar colour  $H - K$  shows variations that are clearly correlated to the lightcurves. From day 845 to day 3731  $H - K$  got redder by about 0.8, whereas in the same time  $K$  decreased by 1.5. Within days 3731 and 4516,  $K$  increased again by around 0.4 and the colour got bluer by roughly 0.8. We did not observe, within the errors, a significant variation in the time from day 4516 to day 5271, while in the same time span  $H$  and  $K$  both varied by around 0.5 mag.

Variations on small timescales (few days) are not observed. Although the SHARP  $K$  lightcurve shows groups of data points that are not resolved in time (obtained in observing runs lasting 4–10 days,) which appeared to be somewhat scattered, these values are identical within their errors. One should note

**Table 1.** Selected  $H$  and  $K$  magnitudes (one data point per calendar year) and colours of IRS34W. The values are not corrected for extinction, the time is given in days after JD 2 448 000. A months- to years time-scale variability can be seen in  $H$  and  $K$ , including variations of the stellar colour.

Time	$H$	$K$	$H - K$
705		$10.56 \pm 0.10$	
			$2.64 \pm 0.20^a$
845	$13.20 \pm 0.17$		
1203		$10.33 \pm 0.12$	
1465		$10.77 \pm 0.10$	
1917		$10.75 \pm 0.12$	
2241		$10.81 \pm 0.11$	
2649	$14.67 \pm 0.10$	$11.55 \pm 0.11$	$3.12 \pm 0.15$
2949	$14.87 \pm 0.08$	$11.60 \pm 0.16$	$3.27 \pm 0.18$
3350		$11.95 \pm 0.13$	
3728		$12.46 \pm 0.09$	
			$3.44 \pm 0.15^b$
3731	$15.90 \pm 0.12$		
4094		$12.09 \pm 0.10$	
4516	$14.83 \pm 0.11$	$12.19 \pm 0.13$	$2.64 \pm 0.17$
4806	$14.29 \pm 0.11$	$11.66 \pm 0.10$	$2.63 \pm 0.15$
5271	$14.51 \pm 0.07$	$11.85 \pm 0.02$	$2.67 \pm 0.07$


<sup>a</sup> This colour is computed from the magnitudes at days 705 and 845.

<sup>b</sup> This value is computed from the magnitudes at days 3728 and 3731. Both  $H$  and  $K$  were extracted from the Gemini North data set.

that the Gemini  $K$  point at day 3728 appears to be slightly offset compared to the SHARP points; but taking into account both the error bars and the fact that these points were extracted from two different, separately processed data sets, this is no significant indication of short time variability either.

The normalized spectra of IRS34W obtained in 1996, 2003, and 2004 are presented in Fig. 2, with all wavelengths used below given in microns. All spectra show as prominent features the lines He I ( $\lambda$  2.058), He I ( $\lambda$  2.112), N III ( $\lambda$  2.116), He I ( $\lambda$  2.162), He I ( $\lambda$  2.164), and H I ( $\lambda$  2.166) in emission. Due to line-blending the lines He I ( $\lambda$  2.112), and N III ( $\lambda$  2.116) on the one hand and He I ( $\lambda$  2.162), He I ( $\lambda$  2.164), and H I ( $\lambda$  2.166) on the other, are not separated and are therefore treated here integrally as line complexes He I ( $\lambda$  2.112)+N III ( $\lambda$  2.116), resp. He I ( $\lambda$  2.162)+He I ( $\lambda$  2.164)+H I ( $\lambda$  2.166). All lines resp. line complexes show P Cygni profiles identifying IRS34W as a wind source. We especially note a remarkable similarity between the spectra of IRS34W and the spectrum of the Ofpe/WN9 star HDE 269445 presented in Morris et al. (1996).

Comparison of the spectra shows only slight variations in the equivalent widths (EWs) of all emission lines (see Table 2), which are too small to be significant. A direct graphical comparison by overplotting the spectra (see Fig. 2, bottom panel) shows that they appear to be identical within their typical accuracies.



**Fig. 2.** *Top panel:* normalised  $K$  band spectra of IRS34W obtained in March 1996 with 3D, in April 2003 with SPIFFI, and in August 2004 with SINFONI (spectra shifted along the flux axis). *Bottom panel:* the same spectra, smoothed and overplotted. Obviously, the spectra are identical within their typical accuracies.

Obviously, the photometric and spectroscopic observations of IRS34W lead to ambivalent results. On the one hand, the photometric magnitude of the star is highly variable on timescales from months to years. The shape of the light curve and the time scales of the variability rule out the possibility that IRS34W is an eclipsing binary. The amplitude of the variation, up to  $\approx 75\%$  in flux, shows that an eclipsing star would have to cover at least that fraction of IRS34W’s surface on the sky. The lack of significant variation in the line to continuum ratios shows that the continuum of this eclipsing star would have to be small compared to the fraction of the flux of IRS34W that remains to be seen. In that case, the temperature of the eclipsing star would be below  $\approx 5000$  K, and it would have CO absorption lines, which we would be able to detect even if diluted in the continuum of the blue supergiant. Therefore even a coincidental eclipse by an unrelated star can be excluded.

Another possible effect to be examined is the obscuration of IRS34W by interstellar matter. The central parsec is known to contain several gas patches of various dimensions that can be responsible for local enhancements of the extinction by  $\Delta K \approx 1$  (Paumard et al. 2004), and could therefore, in principle, account for the variability of IRS34W. Such clouds are, however,

**Table 2.** Equivalent widths of the dominant emission lines in the March 1996, April 2003, and August 2004 spectra. Column “A” is He I ( $\lambda$  2.058), “B” He I ( $\lambda$  2.112) + N III ( $\lambda$  2.116), and “C” He I ( $\lambda$  2.162) + He I ( $\lambda$  2.164) + H I ( $\lambda$  2.166). The wavelengths of the lines are given in microns, and the equivalent widths are given in  $\text{\AA}$ .

	A	B	C
1996	$22.1 \pm 1.9$	$3.2 \pm 1.6$	$18.7 \pm 1.9$
2003 <sup>a</sup>	$26.0 \pm 0.6$	$5.4 \pm 0.7$	$18.5 \pm 0.6$
2003 <sup>b</sup>	$25.4 \pm 0.9$	$3.5 \pm 0.8$	$18.5 \pm 0.5$
2004	$25.8 \pm 0.8$	$4.4 \pm 0.9$	$15.6 \pm 0.9$

<sup>a</sup> Spectrum taken from SPIFFI cube reduced to pixel scale, spatial and spectral resolution of 3D data.

<sup>b</sup> Spectrum obtained from original SPIFFI cube.

several arcseconds wide, and should therefore effect not only IRS34W but also neighbouring stars. Indeed, the lightcurves of some faint stars ( $K \approx 16$ ) taken from the original automatically generated NACO photometric data set (Trippe 2004) in a distance up to  $0.9''$  from IRS34W showed similar, but weaker variations. Therefore we re-analysed the magnitudes of IRS34W and its neighbouring stars in a set of selected good NACO images, applying additional background flux subtraction and Lucy-Richardson deconvolution instead of Wiener filtering. This analysis confirmed the magnitudes and variations found for IRS34W and made it possible to trace irregularities in some faint stars back to flux spill-over effects from IRS34W to stars located in its seeing halo. This rules out as well a cloud in the line of sight but much closer to the Earth, as it would be even larger in projection, and cause similar variability in a large number of stars, which would be obvious in our data. We finally could conclude that the photometric variability is indeed bound to the star, resp. its immediate vicinity (in terms of the given spatial resolution), and not to an external source.

On the other hand, the absence of significant spectroscopic variability complicates the understanding of the stellar behaviour. As the emission lines we observe in our spectra are sensitive especially to the stellar parameters  $T_{\text{eff}}$  and  $\dot{M}$ , significant variations of these parameters should be visible in the spectra. An example for this is the LBV AFGL 2298, which shows photometric variations in the  $K$  band of around 0.3 mag and variations in the EWs of He I ( $\lambda$  2.058) and H I ( $\lambda$  2.166) by factors of 3 to 4 within one year (Clark et al. 2003a). Indeed, our observations indicate that at least  $T_{\text{eff}}$  and  $\dot{M}$  remain rather constant, while the star gets dimmer or brighter.

Given that we do not see significant changes in the spectral lines, variations of the wind properties are highly improbable, since this would mean a change of the density (by modifications either of maximum velocity at the top of the wind,  $v_{\infty}$ , or  $\dot{M}$ ), which would then lead to significant observable spectroscopic variations.

Taking this into account, we find two possible interpretations for the photometric variability of IRS34W:

- (1) The variability is connected to the physics of the stellar photosphere. Since  $T_{\text{eff}}$  did not change, this assumption would lead to the conclusion that the stellar radius


(as  $L \propto R_*^2 \cdot T_{\text{eff}}^4$ ) changed by a factor of around 2. Indeed Dorfi & Gatschy (2000) observe comparable variations in stellar radii (by a factor up to  $\approx 1.6$ ) in their simulations of the pulsations of massive stars, but on timescales of days, not years, as in the case of IRS34W. More interestingly, they also note that the *rise* of a pulsating mode can be accompanied by a change in stellar radius (a factor of  $\approx 1.3$  in their model) and only a little increase in  $T_{\text{eff}}$  (2700 K on top of 18 000 K, thus probably not observable for us), on the timescale of two years. Although in this particular case the overall change in luminosity is smaller than the variations we observe ( $\Delta K \approx 0.44$  as calculated from a black-body distribution for the given changes in temperature and radius), it makes it plausible that we may have witnessed a phase transition between a hydrostatic equilibrium and a pulsating mode.

This explanation is complicated by the fact that a change in stellar radius would change the observed EWs, which scale as  $\dot{M}/R_*^{1.5}$ . Thus the constancy of the EWs would require a corresponding increase in  $\dot{M}$  parallel to the rise of the radius.

Apart from this rather improbable scenario, it is hard to understand how a significant change in stellar radius, accompanied by a new hydrostatic quasi-equilibrium that would last for years, would not be accompanied by a significant change in  $T_{\text{eff}}$ .

- (2) The photometric variations of IRS34W are not caused by the photosphere but from above the wind. The colour of the star, significantly redder than other stars of similar brightness (e.g. our photometric reference star ID185 located 3'' NE of IRS34W with  $K = 12.09 \pm 0.08$  and  $H - K = 2.26 \pm 0.13$ ) in the field, also suggests that it is affected by a higher extinction. This is supported by Clenet et al. (2001), who derive a colour of  $K - L = 2.4$  that is about 1 mag redder than the colours of other He I, resp. Ofpe/WN9 stars observed in the field. Additionally, Viehmann et al. (2005) find the  $H - K$  colours of other Galactic Centre He I stars are clearly bluer than what we find for IRS34W. As we were able to exclude external effects like stars or interstellar clouds, the variability would then be due to variations in the column density of circumstellar material expelled by the star.

Both cases – from which scenario (2) is obviously the more probable one – suggest the identification of IRS34W as a LBV – or close to this stage. It must be noted that a slightly elongated nebular emission feature is visible around IRS34W on an  $H/K/L$  NACO colour map that traces the dust emission (Genzel et al. 2003, Fig. 1 right) and that this feature could just as likely be an interstellar structure as circumstellar material from IRS34W that might have been ejected in earlier LBV type eruptions. It is also interesting to note that the shape of the  $K$  band light curve of IRS34W is very similar to the  $V$  light curve of the LBV AG Car (see Humphreys & Davidson 1994): a sudden decrease is followed by a plateau and then a slight increase in a time span of around six years. But contrary to IRS34W, in case of AG Car the photometric variability is accompanied by strong spectroscopic variability.



**Fig. 3.** The approximated position of IRS34W (shaded area) in an HRD compared to similar stars already known. Triangles represent Ofpe/WN9 stars (Pasquali et al. 1997; Crowther et al. 1995; Bianchi et al. 2004; Bresolin et al. 2002), circles LBVs (Humphreys & Davidson 1994; Clark et al. 2003a), and squares O supergiants (Bohannan & Crowther 1999; Repolust et al. 2004; Crowther et al. 2002); filled symbols mark Galactic stars, empty symbols are stars in the Large Magellanic Cloud and NGC 300; the solid line is the Humphreys-Davidson limit. This diagram shows that IRS34W is located between O supergiants and LBVs.

Another interesting point supporting the “dust scenario” is given by Clark et al. (2003b). Their mid-IR study of two ring nebulae shows that they contain stars with properties that are very similar to candidate LBVs. The  $K$  band spectra clearly identify them as luminous evolved massive stars. They are also similar to the spectrum of IRS34W, being dominated by H and He I emission lines. The surrounding nebulae are explained by direct ejection of material by the star and/or by interaction of ejected material with the interstellar medium. Of the two stars, G24.73+0.69 is especially interesting since it is photometrically variable ( $\Delta K \approx 0.3$ ) over  $\approx 1.5$  years, but shows only little spectroscopic variability (see Fig. 10 of Clark et al. 2003b). This similarity favours the explanation of the observed behaviour of IRS34W by the presence of circumstellar dust.

In addition, Clark et al. (2005) include IRS34W in their list of confirmed LBVs, although this is partly based on the assumed variation of stellar and wind parameters that we show is not likely. However, they mention the possibility of a dusty circumstellar environment to explain the magnitude variations of IRS34W, in agreement with the present study. They note that only LBVs, B[e] supergiants, and late WC stars are known to be dust producers. If dust is indeed the explanation for the variability of IRS34W, this is then an indication that it is closely related to Luminous Blue Variables, despite the lack of variation in stellar properties.

Figure 3 shows the approximate position of IRS34W in a Hertzsprung-Russell diagram. The luminosity was estimated as

follows:  $A_K$  was derived from the  $V$  band extinction map of Scoville et al. (2003) and the extinction law of Moneti et al. (2001), and was used together with the observed  $K$  magnitude to estimate  $M_K$ . Then, bolometric corrections computed from our preliminary modelling of Ofpe/WN9 stars covering a wide range of physical parameters ( $L$ ,  $T_{\text{eff}}$ ,  $\dot{M}$ ) were used to finally derive  $L$ . The extension of the area takes into account uncertainties in the extinction and the bolometric corrections. As for the effective temperature, we only show a range of values for which, within the  $L$  range, He I ( $\lambda$  2.058) shows a P-Cygni profile in our preliminary models. So far this is a rather rough estimate, but it gives some interesting indications on the evolutionary status of this star. Compared to already known Ofpe/WN9 stars, LBVs, and O supergiants, IRS34W is located between the O supergiant and LBV stages (like other known Ofpe/WN9 stars, see Fig. 3) and therefore appears to be an object in transition between these two phases.

The estimate of the luminosity of IRS34W presented above shows that it may be intrinsically less luminous than the other He I stars (see Najarro et al. 1997). As discussed above, it is also redder, regardless of the variations reported here. Given this and the fact that dust formation is the most likely explanation for the observed variability, we can imagine the following scenario: due to its lower luminosity, IRS34W is able to experience dust formation in its atmosphere. Such episodes of dust formation makes the star fainter and redder; at the end of each episode, dust is destroyed, but only partially, leading to a long-term accumulation of dust around the star so that on average, IRS34W is redder (and also even fainter) than the other He I stars. The key point of this scenario is the lower intrinsic luminosity of the star before any dust-formation episode.

This is very speculative, however, since the formation of dust in atmospheres of hot stars is still poorly understood, although Cherchneff & Tielens (1995) argue that special geometries, such as equatorial disks, can lead to the high densities necessary for dust formation. Other properties such as a thick wind or inhomogeneities in the atmosphere may also be necessary. In the other He I stars of the Galactic Centre, the luminosity may be too high, so that dust formation is inhibited by the radiation that is too strong. In IRS34W, the photon flux may be low enough to allow dust formation.

#### 4. Summary

In this article we have reported the results of time-resolved photometry and spectroscopy of the Galactic Centre star IRS34W and possible interpretations, which can be summarized as follows:

1. The photometric magnitude of IRS34W is highly variable on timescales from months to years with an overall variation of  $\Delta K \approx 1.5$ .
2. Variations in the stellar colour  $H - K$  track the photometric variability. When the brightness decreases, the colour gets redder; when the brightness increases, the colour gets bluer.
3. The spectra of the star do not show any significant variations over a time span of 8 years.
4. The most probable reason for the stellar behaviour is an obscuration by circumstellar material ejected by the star.
5. IRS34W is a star in transition between the O supergiant and LBV phases, while only the lack of spectroscopic variability prevents us from safely identifying it as an LBV.

IRS34W and five additional stars, which are very similar to IRS34W, have been classified as LBV candidates by Paumard et al. (2001). Among these additional stars, only one, IRS16SW, shows strong, periodic variability (Ott et al. 1999). Spectrophotometric monitoring of these stars is currently under way; detailed quantitative analysis and modelling of this group of stars, including IRS34W, should shed more light on the physics of this class of objects.

*Acknowledgements.* We thank the referee, F. Najarro, for helpful suggestions and comments that helped to improve the quality of the paper. F.M. acknowledges support from the Alexander von Humboldt Foundation.

#### References

- Bianchi, L., Bohlin, R., & Massey, P. 2004, *ApJ*, 601, 228  
 Bresolin, F., et al. 2002, *ApJ*, 577, L110  
 Bohannan, B., & Walborn, N. A. 1989, *PASP*, 101, 639  
 Bohannan, B., & Crowther, P. A. 1999, *ApJ*, 511, 374  
 Bonnet, H., Ströbele, S., Biancat-Marchet, F., et al. 2003, *SPIE*, 4839, 329  
 Bonnet, H., et al. 2004, *The ESO Messenger*, 117, 17  
 Cherchneff, I., & Tielens, A. G. G. M. 1995, *IAUS*, 163, 346  
 Conti, P. S. 1984, *IAUS*, 105, 233  
 Clark, J. S., Larionov, V. M., Crowther, P. A., Egan, M. P., & Arkharov, A. 2003a, *A&A*, 403, 653  
 Clark, J. S., Egan, M. P., Crowther, P. A., et al. 2003b, *A&A*, 412, 185  
 Clark, J. S., Larionov, V. M., & Arkharov, A. 2005, *A&A*, 435, 239  
 Clenet, Y., Rouan, D., Gendron, E., et al. 2001, *A&A*, 376, 124  
 Crowther, P. A., Hillier, D. J., & Smith, L. J. 1995, *A&A*, 293, 172  
 Crowther, P. A., Hillier, D. J., Evans, C. J., et al., 2002, *ApJ*, 579, 774  
 Dorfi, E. A., & Gautschy, A. 2000, *ApJ*, 545, 982  
 Eisenhauer, F., Abuter, R., Bickert, K., et al. 2003a, *SPIE*, 4814, 1548  
 Eisenhauer, F., Tecza, M., Thatte, N., et al. 2003b, *The ESO Messenger*, 113, 17  
 Eisenhauer, F., et al. 2005, *ApJ*, in press  
 Genzel, R., Schödel, R., Ott, T., et al. 2003, *ApJ*, 594, 812  
 Ghez, A. M., Duchêne, G., Matthews, K., et al. 2003, *ApJ*, 586, L127  
 Glatzel, W., & Kiriakidis, M. 1993, *MNRAS*, 263, 375  
 Hartung, M., Lenzen, R., Hofmann, R., et al. 2003, *SPIE*, 4841, 425  
 Hofmann, R., Blietz, M., Duhoux, Ph., et al. 1992, *Proceedings of the ESO Conference on Progress in Telescope and Instrumentation Technologies*, Garching, Germany, 617  
 Humphreys, R. M., & Davidson, K. 1984, *Science*, 223, 243  
 Humphreys, R. M., & Davidson, K. 1994, *PASP*, 106, 1025  
 Krabbe, A., Genzel, R., Drapatz, S., & Rotaciuc, V. 1991, *ApJ*, 382, L19  
 Krabbe, A., Genzel, R., Eckart, A., et al. 1995, *ApJ*, 447, L95  
 Lamers, H. J. G. L. M., & Fitzpatrick, E. L. 1988, *ApJ*, 324, 279  
 Lucy, L. B. 1974, *AJ*, 79, No. 6, 745  
 Massey, P. 2003, *ARA&A*, 41, 15  
 Moneti, A., Stolovy, S., Blommaert, J. A. D. L., Figer, D. F., & Najarro, F. 2001, *ApJS*, 102, 383

- Morris, P. W., Eenens, P. R. J., Hanson, M. M., Conti, P. S., & Blum, R. D. 1996, *ApJ*, 470, 597
- Najarro, F., Krabbe, A., Genzel, R., et al. 1997, *A&A*, 325, 700
- Nota, A., Pasquali, A., Drissen, L., et al. 1996, *ApJS*, 102, 383
- Nugis, T., & Lamers, H. J. G. L. M. 2002, *A&A*, 389, 162
- Ott, T., Eckart, A., & Genzel, R. 1999, *ApJ*, 523, 248
- Pasquali, A., Langer, N., Schmutz, W., et al. 1997, *ApJ*, 478, 340
- Paumard, T., Maillard, J. P., Morris, M., & Rigaut, F. 2001, *A&A*, 366, 466
- Paumard, T., Maillard, J.-P., & Morris, M. 2004, *A&A*, 426, 81
- Paumard, T., et al. 2005, in preparation
- Repolust, T., Puls, J., & Herrero, A. 2004, *A&A*, 415, 349
- Richardson, W. H. 1972, *J. Opt. Soc. Am.*, 62, 55
- Rousset, G., Lacombe, F., Puget, P., et al. 2003, *SPIE*, 4839, 140
- Scoville, N. Z., Stolovy, S. R., Rieke, M., Christopher, M., & Yusef-Zadeh, F. 2003, *ApJ*, 594, 294
- Stahl, O. 1986, *A&A*, 164, 321
- Stahl, O., Wolf, B., Klare, G., et al. 1983, *A&A*, 127, 49
- Trippe, S. 2004, master thesis (Diplomarbeit), Ludwig-Maximilians-Universität München
- Viehmann, T., Eckart, A., Schödel, R., et al. 2005, *A&A*, 433, 117
- Voors, R. H. M., Waters, L. B. F. M., de Koter, A., et al. 2000, *A&A*, 356, 501
- Walborn, N. R. 1982, *ApJ*, 256, 452
- Weitzel, L., Krabbe, A., Kroker, H., et al. 1996, *A&AS*, 119, 531