

Ultra low frequencies phenomena in Jovian decametric radio emission

O. V. Arkhipov¹ and H. O. Rucker²

¹ Institute of Radio Astronomy, National Academy of Sciences of Ukraine, Chervonopraporna 4, 61002 Kharkiv, Ukraine
e-mail: rai@ira.kharkov.ua

² Space Research Institute, Austrian Academy of Sciences, Schmidlstrasse 6, 8042 Graz, Austria
e-mail: rucker@oeaw.ac.at

Received 11 October 2005 / Accepted 24 January 2006

ABSTRACT

System of periodic peaks is found in the power spectra of ULF flux oscillations of Jovian decametric emission, which is observed at fixed frequencies. This result could be interpreted in terms of an ionospheric Alfvén resonator near Jupiter. Standing Alfvén wave could explain the system of emission bands in the dynamic spectrum of decametric emission. The space period of such bands and the frequency of standing waves have been used to estimate electron number density in the low magnetosphere of Jupiter. The result is consistent with radio occultation data. Hence, the ULF modulation hypothesis appears validated.

Key words. planets and satellites: individual: Jupiter – magnetic fields – radiation mechanisms: non-thermal – plasmas – waves – magnetohydrodynamics (MHD)

1. Introduction

It is widely accepted that waves of ultra low frequencies (ULF) in the Jovian magnetosphere can be studied only in situ, with space probes. Only electromagnetic, mainly decameter (DAM) radio emission is observable from the Earth. However, some theories connect the planetary radio emission with ULF Alfvén waves, which accelerate electrons and stimulate plasma instabilities with electromagnetic wave generation near Jupiter and other planets (Belcher 1987; Su et al. 2005; Zarka 1998). It has been supposed that Alfvén waves modulate the Jovian decametric emission (DAM) with the Io orbital period (42.46 h, Belcher 1987) and the wave bounce period (~ 10 min, Bagenal & Leblanc 1988).

Much higher frequencies (~ 20 Hz) are predicted for DAM modulation in an ionospheric Alfvén resonator near Jupiter (Su et al. 2005). As near the Earth, this resonator could operate in vertical direction with shear Alfvén waves, which are trapped between the ionospheric boundary and the peak of the Alfvén phase speed. There are verifiable consequences for the resonator hypothesis: (a) the space modulation of DAM by a standing Alfvén wave; (b) multiple resonant eigenfrequencies in DAM flux variations. We search for these effects to verify the hypothesis of DAM Alfvén modulation and a Jovian resonator.

2. S/NB-emission as detectors of ULF waves

There are various theoretical mechanisms for DAM modulation by ULF Alfvén waves. For example, Ergun et al. (2005) suggested that electron acceleration or modulation may provide the physical mechanism that transfers energy from the Alfvén wave to the S-burst. It has been supposed that the time interval between S-bursts reflects the period of the stimulating Alfvén wave (Su et al. 2005).

Another effect is the periodic inclination of the narrow radio beam by Alfvénic disturbance of a magnetic field (Arkhipov 2002). Zarka et al. (1997) argue that the average intensity profile of a S-burst is consistent with emission beaming in a widely open (70° – 80°) hollow cone of $\sim 2^\circ$ thickness, decreasing to zero at both (inner and outer) edges within $\sim 0.2^\circ$. The symmetry axis of such a radiation pattern is the magnetic vector at the position of the radio source. Hence, even $\sim 0.2^\circ$ variations of the magnetic direction could modulate the radio flux towards the Earth with this geometry.

The time-delay variations for Alfvénic displacements of the radio source in an inhomogeneous magnetic field are important (Arkhipov 2002). A typical S-burst (Fig. 1a) has a fixed-frequency duration of a few milliseconds and an instantaneous frequency band of a few tens of kHz (Ellis 1982). Although their origin is the subject of much debate, there are arguments for a small source dimension (~ 10 km, Ellis 1982; Zarka 1998). Moreover, the standard element of S-theories is the low group velocity of emission in the vicinity of its source for effective amplification of a radio wave by plasma instabilities (e.g. Melrose 1986; Zaitsev et al. 1986; Willes 2002). These properties could provide detectable effects by periodic displacements of radio sources by magnetospheric hydro-magnetic waves of very low frequencies. Thus time delay and frequency variations are possible.

S-bursts usually appear as lines in the dynamic spectrum, i.e. on the time-frequency plane. Such a form of the S-burst reflects the quasi-straight motion of a radio source along a magnetic line with a decrease of the emission frequency in accordance with the local gyrofrequency of electrons. However, there are S-bursts with a sinusoidal shape (Fig. 1a). In this case it is not possible to explain the drift rate using only the classical model based on the hypothesis of adiabatic motion of trapped electrons (Boudjada et al. 1997). The problem could be solved taking into consideration a parallel electric field (Galopeau et al. 1999)




Fig. 1. **a)** The train of S-bursts with sinusoidal shapes seen July 21, 1994 (Ryabov et al. 1997). **b)** The typical NB-event with oscillations seen November 12, 1988 (Riihimaa 1991). **c)** NB hooks (Riihimaa 1992) demonstrate the time-delay effect.

or an Alfvénic distortion of an active magnetic line (Arkhipov 2002). As Alfvén waves have parallel electric fields, these interpretations do not contradict each other. Parallel electric fields of Alfvén waves are considered as able to stimulate S-emission (Su et al. 2005).

Sometimes S-bursts are associated with narrow band (NB) emission at quasi constant frequency (Boudjada et al. 2000). Apparently, in this case the radio source is fixed at some height above the planet. Therefore, it generates narrow band emission (~ 100 kHz) near the local gyrofrequency of electrons. In many cases such emission oscillates around the mean frequency in dynamic spectra (Fig. 1b). There is an obvious time-delay effect in NB-events: NB-hooks have been seen, where the emission from one source is observed simultaneously at 2 to 3 different frequencies (Fig. 1c).

Such ambiguity of registration is known for S-bursts, too, as f, g, h, l, n, r and u forms of J.J.Riihimaa's classification of S-spectra (Riihimaa 1991). The dispersion delay effects, the phase-bunching model have been proposed to explain this ambiguity (Willes 2002). However, the sinuous shapes of S/NB-events (Ryabov et al. 1997) can not be explained with this approach. Nevertheless, the wave-like forms of S/NB events are natural for harmonic displacements of a radio source (Arkhipov 2002). Alfvén waves are examples of such harmonic displacements in magnetized plasma.

Thus there are various reasons for the search for Alfvén modulation of DAM. Although the nature of the S/NB source is still unknown, its emission could be used as a probe to study of ULF waves in the Jovian magnetosphere.

3. Space modulation

Particle acceleration, DAM generation or modulation are likely more effective in the antinodes (maxima) of the standing wave.




Fig. 2. The modulation of the mean probability of S-bursts in UTR-2 DAM spectra (Ryabov et al. 1985).

The DAM frequency decreases above Jupiter in accordance with the decrease in the local electron gyrofrequency f_{ce} . Hence, the ULF waves could be found in DAM dynamical spectra as quasi-periodic (in frequency) bands of radio emission. Indeed, it is well known that S/NB-emission in dynamic spectra is organized in bands near certain frequencies. The typical interval between such bands is 2–4 MHz (Riihimaa 1991; Ryabov et al. 1997).

Ryabov et al. (1985) found a quasi-periodic modulation of the mean probability of S-bursts in UTR-2 spectra (Fig. 2). This periodicity could be explained in terms of a standing wave. As its wavelength λ is equal to the one of propagating waves with the same frequency f_w , we can estimate the phase speed of the ULF disturbance:

$$V_{ph} = \lambda f_w, \quad (1)$$

where λ is the wavelength. This velocity is related to the electron number density for the Alfvén wave (e.g.: Nicholson 1983) as:

$$V_{ph} = \frac{1}{\sqrt{1/V_A^2 + 1/c^2}}, \quad (2)$$

where c is the speed of light; $V_A = B/(4\pi m_i n_i)^{1/2}$ is the Alfvén velocity. As a result, the electron density N_e in the radio source can be estimated for hydrogen plasma ($N_e = n_i$).

There are the following frequencies of maximum/minimum probability of S-emission in Fig. 2: 11.75, 13, 15, 16.5, 18.5, 20, 21.4, 22.5, 22.8 MHz. We use the frequencies f_1 and f_2 of some neighboring maxima (or minima) to calculate the corresponding planetocentric radius-vectors \vec{r}_1 and \vec{r}_2 of radio sources, which are on the same Io activated magnetic line, with the VIP4 magnetic model (Connerney et al. 1998) and the standard $f = f_{ce}$ approximation. Then the half-wavelength of the standing wave is estimated as $\frac{\lambda}{2} = |\vec{r}_1 - \vec{r}_2|$. The corresponding wavelengths λ are shown in Table 1. The estimations N_e are calculated with Eqs. (1), (2) and the most probable frequency of a S-burst recurrence of $f_w = 20$ Hz (Carr & Reyes 1999). The obtained N_e values are shown in Fig. 3. To estimate the scale height (H) correctly, the geometric altitude must be modified considering the spherical geometry and the influence of centrifugal force on co-rotating plasma. This modification is the special Z altitude (Melrose 1967):

$$Z = R_j(1 - R_j/r) - (\Omega R_j)^2(r^3/R_j^3 - 1)/(2Lg), \quad (3)$$

Table 1. Parameters of ULF waves.

f_1 [MHz]	f_2 [MHz]	Hemi- sphere	λ [km]	V_{ph} [km s ⁻¹]	N_e [cm ⁻³]	$Z - Z_0$ [km]
11.75	15	N	13 934	278 684	148	11 945
		S	13 241	264 811	264	12 541
13	16.5	N	13 138	262 755	347	10 295
		S	12 466	249 310	513	10 945
15	18.5	N	11 018	220 363	1302	8053
		S	10 449	208 973	1619	8773
16.5	20	N	9772	195 441	2505	6502
		S	9275	185 508	2984	7269
18.5	21.4	N	7134	142 686	7948	4830
		S	6779	135 578	9054	5644
20	22.5	N	5623	112 453	16 615	3638
		S	5349	106 972	18 647	4480
21.4	22.8	N	2976	59 525	75 893	2873
		S	2835	56 698	83 968	3731




Fig. 3. Electron number density N_e (cm⁻³) at different altitudes Z is estimated in terms of an Alfvén standing wave. The calculations are made for the northern (black squares) and southern (open squares) sources of Io-DAM with the VIP4 magnetic model, Io’s Jovigraphical longitude of 260° and a lead angle of 20°. The straight line approximation is found by a least squares fit using Eq. (4).

where: $R_J = 71\,372$ km is the Jupiter radius; $r = (|\vec{r}_1| + |\vec{r}_2|)/2$ is the effective planetocentric distance; $\Omega = 1.759 \times 10^{-4}$ rad/s is the angular velocity of Jupiter’s rotation; $L = 6$ is the McIlwain parameter for the Io torus; $g = 25.9$ m/s² is the acceleration of gravity in the Jovian exobase.

According to Melrose (1967), the electron density near Jupiter in the approximation of diffusion equilibrium can be described in the form:

$$N_e = N_0 \exp[-(Z - Z_0)/(2H)], \quad (4)$$

where N_0 is the electron number density at height Z_0 ; $H = k_b T / (m_i g)$ is the scale height (k_b is the Boltzmann constant; T is the temperature). Indeed, the obtained $\lg N_e - Z$ relation (Fig. 3) satisfactorily corresponds to this equation. The optimal scale height $H = 828 \pm 66$ km and $\lg N_0 = 5.41 \pm 0.16$ (cm⁻³) near the ionospheric maximum ($Z_0 = 2000$ km) are close to the estimates from radio occultations (400 km $< H < 960$ km; $4.2 < \lg N_0 < 5.6$; Strobel & Atreya 1983).




Fig. 4. **a)** The resonant peaks (arrowed) in the power spectrum of DAM flux variations at the fixed frequency 22.35 MHz during the S/NB-event of March 30, 2000. **b)** Another S/NB-band at 25.8 MHz simultaneously shows different resonances. The horizontal lines are the borders of the 0.98-significance interval.

4. Time modulations of Jovian radio emission

ULF properties of S/NB-emission have been rarely studied. Riihimaa (1991) found that the bursts in S-trains are repeated in rapid succession at quasi-periodic rates of 20–40 s⁻¹. Carr & Reyes (1999) note that typical S-bursts are quasi-repetitive, recurring at rates between about 2 and 400 Hz over intervals of a few seconds, the most probable rate being 20 Hz. At the frequencies below 20 Hz, Calvert et al. (1988) investigated the random (Poisson) statistics for the cumulative percentage occurrence of S-burst spacing. However, their experimental points (Calvert et al. 1988, Fig. 5) show the systematic divergences from the Poisson approximation. Unfortunately, the significance of these spectral details is unknown.

To study S/NB periods in more detail, we used the dynamical spectrum of the S-storm of March 30, 2000, which was recorded by an observation campaign within the frame of the INTAS project 03-51-5727 with the UTR-2 radio telescope at 2 ms time resolution. The Fourier power spectrum $S(\omega)$ of the radio flux-time function at a fixed receiver frequency f is calculated. It has been found that the relation between smoothed $\lg S(\omega)$ and the ULF frequency ω is quasi-linear with slightly positive curvature. Hence, this Fourier spectrum, smoothed with Bartlett spectral window, at 10 Hz $< \omega < 200$ Hz can be approximated quadratically as $\lg S(\omega) \approx a + b\omega + c\omega^2$, where the constants a , b and c are found by the least squares method. To visualize the fine spectral details, the residuals $\lg(S(\omega)) - (a + b\omega + c\omega^2)$ in dB are shown in Fig. 4.

The significance level is calculated for a standard χ^2 -distribution:

$$F_{\chi^2}(x) = x^{v/2-1} [2^{v/2} \Gamma(v/2)]^{-1} \exp(x/2), \quad (5)$$

Table 2. Comparison of found spectral peaks with predictions.

Figure	ω_{obs} [Hz]	ω_{cal}^a [Hz]	N_i^a [cm ⁻³]	H^a [km]
2a	45	51	2×10^4	780
2a	90	89	2×10^4	780
2b	27	27.2	2×10^5	800
2b	56	50.2	2×10^5	800

^a Su et al. (2005), Fig. 7.

where $\nu = 3N\eta$; N is the number of processed realizations; $\eta = 7$ is the ratio of the duration of one spectrum to the width of the Bartlett window.


The regular patterns of ULF spectral maxima are seen in Fig. 4a. Another S/NB-band at 25.8 MHz observed simultaneously shows different resonances (Fig. 4b). There is a concurrence between the observed significant frequencies ω_{obs} and the calculated eigenmodes ω_{cal} of the ionospheric resonator in some models of Su et al. 2005 (Table 2). Although there is a difference in the model ionospheric density, N_i , the scale height H is practically the same.

According to the direct measurements by space probes in the polar magnetosphere of the Earth, there are field-aligned regions (cavities) of very depleted plasma density about 10 times lower than in the surrounding medium. Moreover, remote sensing experiments argue for the existence of analogous cavities near Jupiter, Saturn, Uranus and Neptune (Zarka 1998). If two radio sources are in two separate magnetic tubes with different N_i and consequently V_{ph} , they are modulated by these ionospheric Alfvén resonators with different frequencies, even that of the same eigenmode. Therefore, the magnetospheric cavities could explain the differences in modulation frequencies and N_i estimations in Table 2.

To estimate the frequency range of the ionospheric resonator, the power spectra are calculated for Riihimaa's NB events (Riihimaa 1991, spectra Nos. 4, 6, 8–10, 12, 14, 15, 17–23, 26, 27, 29, 30, 39, 40, 43, 44, 48, 49, 59, 61–64, 70, 73–76). Figure 5 shows the distribution of frequencies of the main spectral peaks, calculated for each dynamical NB spectrum. Apparently, the ionospheric resonator works mainly on the frequencies 5–45 Hz.

5. Conclusions

The analysis of observational data shows that the best candidates for ULF modulation are the oscillations of Jovian narrow band emissions (S-bursts and NB-events). Such oscillations are multi-frequency with significant spectral peaks near the resonant frequencies of the ionospheric Alfvén resonator (eigenmodes of 5–45 Hz). S-bursts are repeated in trains with a typical frequency of 20 Hz. It is suggested that this dominating frequency reflects the period of the Alfvénic wave, which accelerate electrons and stimulates or modulates S-emission. The hypothesis of a ionospheric resonator predicts a standing wave that stimulates S-bursts in many spectral bands. These bands must be localized quasi-periodically at different frequencies according to the local electron gyrofrequencies in the maxima of the standing wave. Predicted bands are found as the periodic frequency dependence of the S-burst occurrence probability. The Alfvén velocity and electron number density (N_e) are estimated from the

**Fig. 5.** Histogram of frequencies of the main spectral peak (black) and three maximum peaks (grey) in power spectra of NB-events.

wavelength and known period (20 Hz). The obtained N_e profile ($\lg N_e [\text{cm}^{-3}] = 5.41 \pm 0.16$; $H = 828 \pm 66$ km) is consistent with radio occultation data. Hence, the obtained results are realistic, and the ULF modulation hypothesis appears valid.

Acknowledgements. We used the dynamical spectrum of the S-storm of March 30, 2000, which was recorded by an observation campaign within the frame of the INTAS project 03-51-5727 (A. Lecacheux, H. R. Rucker, A. A. Konvalenko and Yu. Tokarev) with the UTR-2 radio telescope. We thank Dr. V. E. Shaposhnikov for fruitful discussions.

References

- Arkhipov, A. V. 2002, *Kinematika i Fizika Nebesnykh Tel*, 18(5), 410
- Bagenal, F., & Leblanc, Y. 1988, *A&A*, 197, 311
- Belcher, J. W. 1987, *Science*, 238, 170
- Boudjada, M. Y., Rucker, H. O., Galopeau, P. H. M., Kleewein, P., & Mostetschnig, V. 1997, The contribution of the Riihimaa classification to the study of Jovian millisecond radio bursts, in *Planetary Radio Emissions IV*, ed. H. O. Rucker, S. J. Bauer, & A. Lecacheux (Vienna: Austrian Academy of Sciences Press), 91
- Boudjada, M. Y., Galopeau, P. H. M., Rucker, H. O., & Lecacheux, A. 2000, *A&A*, 363, 316
- Calvert, W., Leblanc, Y., & Ellis, G. R. A. 1988, *ApJ*, 335, 976
- Carr, T. D., & Reyes, F. 1999, *J. Geophys. Res.*, 104, A11, 25127
- Connerney, J. E. P., Acuna, M. H., Ness, N. F., & Satoh, T. 1998, *J. Geophys. Res.*, 103, A6, 11929
- Ellis, G. R. A. 1982, *Austr. J. Phys.*, 35, 2, 165
- Ergun R. E., Su, Y.-J., Andersson, L., et al. 2005, *Geophys. Res. Lett.*, in press
- Galopeau, P. H. M., Boudjada, M. Y., & Rucker, H. O. 1999, *A&A*, 341, 918
- Melrose, D. B. 1967, *Planet. Space Sci.*, 15, 2, 381
- Melrose, D. B. 1986, *J. Geophys. Res.*, 91, A7, 7970
- Nicholson, D. R. 1983, *Introduction to plasma Theory* (N.Y.: J. Wiley & Sons), 163
- Riihimaa, J. J. 1991, *Earth, Moon and Planets*, 53, 2, 157
- Riihimaa, J. J. 1992, *Wide-range high-resolution S-burst spectra of Jupiter* (Oulu: Univ. of Oulu), 24
- Ryabov, B. P., Arkhipov, A. V., & Shevchenko, V. A. 1985, *Astron. Vestn.*, 19, 4, 296
- Ryabov, B. P., Zarka, P., Rucker, H. O., Ryabov, V. B., & Boudjada, M. Y. 1997, Recurrent fine structures in Jovian S-burst emission, in *Planetary Radio Emissions IV*, ed. H. O. Rucker, S. J. Bauer, & A. Lecacheux (Vienna: Austrian Academy of Sciences Press), 65
- Strobel, D. F., & Atreya, S. K. 1983, *Ionosphere*, in *Physics of the Jovian magnetosphere*, ed. A. J. Dessler (Cambridge: Cambridge University Press), 51, Fig. 2.4
- Su, Y.-J., Jones, S. T., Ergun, R. E., et al. 2005, *J. Geophys. Res.*, in press
- Willes, A. J. 2002, *J. Geophys. Res.*, 107, A5
- Zaitsev, V. V., Zlotnik, E. Y., & Shaposhnikov, V. E. 1986, *A&A*, 169, 345
- Zarka, P. 1998, *J. Geophys. Res.*, 103, E9, 20159
- Zarka, P., Ryabov, B. P., Ryabov, V. B., et al. 1997, On the origin of Jovian decameter radio bursts, in *Planetary Radio Emissions IV*, ed. H. O. Rucker, S. J. Bauer, & A. Lecacheux (Vienna: Austrian Academy of Sciences Press), 51