

Self-consistent theory of turbulent transport in the solar tachocline

II. Tachocline confinement*

N. Leprovost and Eun-Jin Kim

Department of Applied Mathematics, University of Sheffield, Sheffield S3 7RH, UK
e-mail: n.leprovost@sheffield.ac.uk

Received 24 March 2006 / Accepted 5 May 2006

ABSTRACT

Aims. We provide a consistent theory of the tachocline confinement (or anisotropic momentum transport) within a hydrodynamical turbulence model. The goal is to explain helioseismological data, which show that the solar tachocline thickness is at most 5% of the solar radius, despite the fact that, due to radiative spreading, this transition layer should have thickened to a much more significant value during the sun's evolution.

Methods. Starting from the first principle with the physically plausible assumption that turbulence is driven externally (e.g. by plumes penetrating from the convection zone), we derive turbulent (eddy) viscosity in the radial (vertical) and azimuthal (horizontal) directions by incorporating the crucial effects of shearing due to radial and latitudinal differential rotations in the tachocline.

Results. We show that the simultaneous presence of both shears effectively induces a much more efficient momentum transport in the horizontal plane than in the radial direction. In particular, in the case of strong radial turbulence (driven by overshooting plumes from the convection zone), the ratio of the radial to horizontal eddy viscosity is proportional to $\mathcal{A}^{-1/3}$, where \mathcal{A} is the strength of the shear due to radial differential rotation. In comparison, in the case of horizontally driven turbulence, this ratio becomes of order $-\epsilon^2$, with negative radial eddy viscosity. Here, ϵ ($\ll 1$) is the ratio of the radial to latitudinal shear. The resulting anisotropy in momentum transport could thus be strong enough to operate as a mechanism for the tachocline confinement against spreading.

Key words. turbulence – Sun: interior – Sun: rotation

1. Introduction

One of the outstanding problems in solar physics is to understand the dynamics of the tachocline, a thin layer where the transition from latitudinal rotation in the convective zone to uniform rotation in the radiative interior takes place. This is particularly true since the tachocline links two regions of very different transport properties, thereby playing a crucial role in the overall angular momentum transport and chemical mixing on the course of the solar evolution. The tachocline is also crucial for the solar dynamo since the strong shear in the tachocline is believed to take part in the process by which the magnetic field of the sun is created. Helioseismic data (Charbonneau et al. 1999) constrain the thickness of this layer to be only a few percent of the solar radius. This poses a challenging problem since a radially localised shear flow (associated with the radial differential rotation) naturally tends to spread during the course of solar evolution, as shown by Spiegel & Zahn (1992), and it should have reached a much broader thickness than what is observed today.

Some physical processes have been proposed as responsible for the tachocline confinement. First, Spiegel & Zahn (1992) have shown that, if the turbulence is highly anisotropic (being much more vigorous in the horizontal direction than in the radial one), the spreading of the solar tachocline could be limited. More precisely, it is the turbulent transport of angular momentum that has to be anisotropic, meaning that the horizontal turbulent viscosity has to be much larger than the radial one. Physically,

this would enable the differential rotation to be smoothed out by transport in the horizontal direction rather than in the vertical one. An alternative mechanism for the stabilisation of the tachocline is due to magnetic field (Rüdiger & Kichatinov 1997; Gough & McIntyre 1998; MacGregor & Charbonneau 1999): even a rather weak poloidal magnetic field (of the order of 10^{-4} G) in the solar interior could prevent the downward flow from the convection zone (driven, for example, by rotation-induced meridional circulation) from penetrating deeply into the radiative interior, thereby confining the differential rotation to a thin region of space and leading to a uniformly rotating interior.

The relevance of these models (and any other that can be constructed with a physical process) to the sun is, however, not asserted very well. In particular, the success of these models strongly depends on the values of transport coefficients such as turbulent viscosities and magnetic diffusivities, which are often crudely parameterised unless their molecular values are used. Furthermore, the Gough & McIntyre model is far too complex to be analysed thoroughly, and only a simplified version of it is actually tractable (Garaud 2003). In the Spiegel & Zahn scenario, it is very important to identify physical mechanisms that can lead to anisotropic momentum transport and then to derive eddy viscosities from first principles (or from Navier-Stokes equation). Note that, while Spiegel & Zahn invoked a strong stable stratification as a source of anisotropic turbulence, they did not derive the values of eddy viscosities.

The purpose of this paper is to provide a consistent theory of the anisotropic momentum transport. The physical mechanism that we invoke is the turbulence regulation by shearing effect,

* Appendices A and B are only available in electronic form at <http://www.edpsciences.org>

the so-called shear stabilisation (Burrell 1997; Hahm 2002; Kim 2004). It is basically because a shear flow acting on a turbulent eddy creates small scales in the direction orthogonal to the flow and thus enhances dissipation, thereby reducing the transport and turbulence intensity (Kim & Dubrulle 2001; Kim & Diamond 2003; Kim et al. 2004; Kim 2006). Indeed, Kim (2005) has shown that a shear due to a stable radial differential rotation in the tachocline not only suppresses turbulent transport and turbulence intensity, but also leads to an “effectively” anisotropic transport of particles via stronger reduction in radial transport compared to the horizontal one. However, the key question of anisotropic momentum transport, necessary for the tachocline confinement, was not addressed in Kim (2005), since the effect of latitudinal differential rotation in the tachocline was neglected for simplicity. In the present paper, we provide a theory of anisotropic momentum transport in the tachocline within a hydrodynamical turbulence model, by taking into account the crucial effects of both radial and latitudinal differential rotations in the tachocline. Specifically, we compute the transport properties of turbulence self-consistently under the physically plausible assumption that turbulence arises by an external forcing (e.g. from plumes penetrating from the convection zone). Most results are derived in the limit of strong shear, but with latitudinal shear weaker than radial shear, as relevant to the tachocline. As in Kim (2005), we neglect the effects of stratification, magnetic field, and rotation to elucidate the crucial role of shearing due to radial and latitudinal differential rotations in the momentum transport. In particular, we show that shear alone can lead to anisotropic momentum transport, with a more efficient horizontal transport. While stratification, magnetic field, and rotation can also contribute to anisotropic turbulence, these are outside the scope of this study and will be addressed in future papers.

The remainder of the paper is organised as follows: we provide our model of the tachocline and solve the governing equations in Sect. 2, then study the turbulence amplitude and eddy viscosities in Sects. 3 and 4, respectively. Section 5 is devoted to our conclusions and discussions of the implications for the solar tachocline dynamics. Appendices contain some details of the algebra involved in solving our model and calculating the turbulent amplitude and viscosity.

2. Model

To elucidate the effect of radial and latitudinal differential rotation on turbulent transport, we adopt a simplified model for the tachocline as in Kim (2005) and use a local Cartesian reference frame where x , y , and z denote local radial, azimuthal, and latitudinal directions, respectively. We capture the radial and latitudinal differential rotation by a large-scale velocity field given by $\mathbf{U}_0(x, z) = -(x\mathcal{A}_x + z\mathcal{A}_z)\hat{y}$. Here, $\mathcal{A}_x = \partial_x U_0 > 0$ and $\mathcal{A}_z = \partial_z U_0 > 0$ are the strength of the radial and latitudinal shear due to radial and latitudinal differential rotation. In the quasi-linear approximation (Moffatt 1978), the Navier-Stokes equation for the small-scale fluctuating velocity can be written:

$$\partial_t \mathbf{v} - (x\mathcal{A}_x + z\mathcal{A}_z)\partial_y \mathbf{v} - (\mathcal{A}_x v_x + \mathcal{A}_z v_z)\hat{y} = -\nabla p + \nu \nabla^2 \mathbf{v} + \mathbf{f}, \quad \nabla \cdot \mathbf{v} = 0. \quad (1)$$

Here \mathbf{f} is the forcing on a small scale and ν is the viscosity of the fluid. We use a Fourier-transform with a time-dependent wave number to account non-perturbatively for the effect of shearing on eddies (e.g. Kim 2005):

$$\mathbf{v}(\mathbf{x}, t) = \int d^3 \mathbf{k}(t) e^{i\mathbf{k}(t) \cdot \mathbf{x}} \tilde{\mathbf{v}}(\mathbf{k}(t), t). \quad (2)$$

Here, both radial and latitudinal wave-numbers evolve in time as follows:

$$k_x(t) = k_x(0) + \mathcal{A}_x t \quad \text{and} \quad k_z(t) = k_z(0) + \mathcal{A}_z t. \quad (3)$$

To absorb the viscosity term, we set $\hat{\mathbf{V}} = \tilde{\mathbf{V}} \exp[\nu Q(t)]$ where $Q(t) = [k_x^3/3k_y\mathcal{A}_x + k_y^2 t + k_z^3/3k_y\mathcal{A}_z]$. The quasi-linear equations for the fluctuating part of the velocity field then become:

$$\begin{aligned} \partial_t \hat{v}_x &= -ik_x \hat{p} + \hat{f}_x, \\ \partial_t \hat{v}_y - (\mathcal{A}_x \hat{v}_x + \mathcal{A}_z \hat{v}_z) &= -ik_y \hat{p} + \hat{f}_y, \\ \partial_t \hat{v}_z &= -ik_z \hat{p} + \hat{f}_z, \\ 0 &= k_x \hat{v}_x + k_y \hat{v}_y + k_z \hat{v}_z. \end{aligned} \quad (4)$$

To solve Eq. (4), we change the time variable from t to $\tau = k_x(t)/k_y$ and introduce new variables: $\mathcal{A} = \mathcal{A}_x$, $\epsilon = \mathcal{A}_z/\mathcal{A}_x$, and $\phi = [k_z(0) - \epsilon k_x(0)]/k_y$. For parameter values typical of the solar interior, we have $\mathcal{A} = \Delta U_0/\Delta X \sim \Delta\Omega/(h/R_\odot) = 3 \times 10^{-6} \text{ s}^{-1}$, for the tachocline of thickness 4% of the solar radius, and $\mathcal{A}_z = \Delta U/\Delta z \sim 4\Delta\Omega \sim 8 \times 10^{-8} \text{ s}^{-1}$. With these values, the ratio of the azimuthal and radial shear is a small parameter $\epsilon \sim 2.7 \times 10^{-2}$. We thus calculate the turbulent amplitude (Sect. 3) and eddy viscosities (Sect. 4) up to order ϵ^2 . In terms of the new variables, Eq. (4) becomes

$$\begin{aligned} \mathcal{A} \partial_\tau \hat{v}_x &= -ik_y \tau \hat{p} + \hat{f}_x, \\ \mathcal{A} \partial_\tau \hat{v}_y - \mathcal{A}(\hat{v}_x + \epsilon \hat{v}_z) &= -ik_y \hat{p} + \hat{f}_y, \\ \mathcal{A} \partial_\tau \hat{v}_z &= -ik_y(\epsilon \tau + \phi) \hat{p} + \hat{f}_z, \\ 0 &= \tau \hat{v}_x + \hat{v}_y + (\epsilon \tau + \phi) \hat{v}_z. \end{aligned} \quad (5)$$

The solution of this system of equations can be found after a long but straightforward algebra with the result (see Appendix A for details):

$$\begin{aligned} \hat{v}_x &= \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}} \left\{ \frac{\gamma + \epsilon \phi \tau}{(\gamma + \epsilon^2)R(\tau)} + \frac{\epsilon \phi}{(\gamma + \epsilon^2)^{3/2}} [T(\tau) - T(\tau_1)] \right\} \\ &\quad - \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_2(\tau_1)}{\mathcal{A}} \frac{\epsilon}{\gamma + \epsilon^2}, \\ \hat{v}_y &= \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}} \left\{ \frac{\phi^2}{(\gamma + \epsilon^2)^{3/2}} (T(\tau) - T(\tau_1)) \right. \\ &\quad \left. - \frac{(\epsilon^2 + 1)\tau + \epsilon \phi}{(\gamma + \epsilon^2)R(\tau)} \right\} - \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_2(\tau_1)}{\mathcal{A}} \frac{\phi}{\gamma + \epsilon^2}, \\ \hat{v}_z &= \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}} \left\{ \frac{\epsilon - \phi \tau}{(\gamma + \epsilon^2)R(\tau)} - \frac{\phi}{(\gamma + \epsilon^2)^{3/2}} [T(\tau) - T(\tau_1)] \right\} \\ &\quad + \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_2(\tau_1)}{\mathcal{A}} \frac{1}{\gamma + \epsilon^2}. \end{aligned} \quad (6)$$

Here,

$$R(\tau) = (\epsilon^2 + 1)\tau^2 + 2\epsilon\phi\tau + \gamma, \quad \gamma = 1 + \phi^2, \quad (7)$$

$$T(\tau) = \arctan\left(\frac{(\epsilon^2 + 1)\tau + \epsilon\phi}{\sqrt{\gamma + \epsilon^2}}\right),$$

$$\hat{h}_1(\tau) = [\gamma + \phi\epsilon\tau]\hat{f}_x - [(1 + \epsilon^2)\tau + \phi\epsilon]\hat{f}_y + (\epsilon - \phi\tau)\hat{f}_z,$$

$$\hat{h}_2(\tau) = -\epsilon\hat{f}_x(\tau) - \phi\hat{f}_y(\tau) + \hat{f}_z(\tau).$$

3. Turbulence amplitude

We first examine how turbulence amplitude is affected by latitudinal shear, comparing the results with Kim (2005). In order to compute $\langle v_i^2 \rangle$, we assume the forcing to be incompressible (thus all the quantities can be expressed in terms of the x

and z components alone) with the statistics that are spatially homogeneous and temporally short correlated with the correlation time τ_f . Specifically, we take

$$\langle \tilde{f}_i(\mathbf{k}_1, t_1) \tilde{f}_j(\mathbf{k}_2, t_2) \rangle = \tau_f (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) \delta(t_1 - t_2) \psi_{ij}(\mathbf{k}_2), \quad (8)$$

for i and $j = 1, 2$ or 3 . The ψ_{ij} functions are the power spectrum of the forcing.

A typical solar value of the molecular viscosity is $\nu \sim 10^2 \text{ cm}^2 \text{ s}^{-1}$ and thus the parameter $\xi = \nu k_y^2 / \mathcal{A}$ is a small quantity provided that $k_y < 10^{-4} \text{ cm} \sim 10^{-6} / H_0$ where H_0 is the pressure scale height at the bottom of the convection zone. That the relevant length scale in the azimuthal direction is probably larger than 10^4 cm permits us to consider the strong shear limit $\xi \ll 1$. In addition, since ϵ is a small parameter, as noted previously, we expand the result up to ϵ^2 , keeping only the dominant term in ξ and ϵ for each component of the forcing ψ_{ij} . A long but straightforward algebra, very similar to that of the calculation of the turbulent viscosities (see next section and Appendix B), then gives us the turbulence amplitude in the strong shear limit $\xi \ll 1$ to leading order in ξ and ϵ :

$$\langle v_x^2 \rangle = \frac{\tau_f}{(2\pi)^3 \mathcal{A}} \int d^3 \mathbf{k} \left\{ \frac{g+a^2}{2g} \left[-a + \frac{g+a^2}{\sqrt{g}} \kappa_0 \right] \psi_{11}(\mathbf{k}) + \frac{(g+a^2)\epsilon}{g} \left[-a + \frac{a^2-g}{\sqrt{g}} \kappa_0 \right] \psi_{13}(\mathbf{k}) + \epsilon^2 \mathcal{G}_0 \psi_{33}(\mathbf{k}) \right\}, \quad (9)$$

$$\langle v_y^2 \rangle = \frac{\tau_f}{(2\pi)^3 \mathcal{A}} \int d^3 \mathbf{k} \mathcal{G}_0 \left\{ \frac{b^4}{g^3} \left((g+a^2)^2 \kappa_0^2 + g a^2 \right) \psi_{11}(\mathbf{k}) + \frac{2b^3 a}{g} \psi_{13}(\mathbf{k}) + b^2 \psi_{33}(\mathbf{k}) \right\},$$

$$\langle v_z^2 \rangle = \frac{\tau_f}{(2\pi)^3 \mathcal{A}} \int d^3 \mathbf{k} \mathcal{G}_0 \left\{ \frac{b^2}{g^3} \left((g+a^2)^2 \kappa_0^2 + g a^2 \right) \psi_{11}(\mathbf{k}) + \frac{2ba}{g} \psi_{13}(\mathbf{k}) + \psi_{33}(\mathbf{k}) \right\}.$$

Here,

$$\begin{aligned} a &= k_x / k_y, & b &= k_z / k_y, & g &= 1 + b^2, \\ \kappa_0 &= \pi/2 - \arctan(a/\sqrt{g}) = \arctan(\sqrt{g}/a), \\ \mathcal{G}_0 &= \Gamma(1/3)(3/2\xi)^{1/3}/3, & \xi &= \nu k_y^2 / \mathcal{A}. \end{aligned} \quad (10)$$

In Eq. (9), we kept only the leading order contributions in ξ and ϵ for the terms proportional to ψ_{11} , ψ_{13} , and ψ_{33} . For instance, for the component proportional to ψ_{33} in the radial turbulence amplitude, we expanded the quantities up to order $O(\epsilon^2)$. It is worth noting that $\langle v_x^2 \rangle$ is always positive, as it should be. While the second term proportional to ψ_{13} is not always positive, in order for the sum of the two terms proportional to ψ_{11} and ψ_{13} to be negative, ψ_{13} has to be at least ϵ^{-1} larger than ψ_{11} ; in that case, ψ_{33} would be at least of order ϵ^{-2} , so the last term would dominate the other two terms, making $\langle v_x^2 \rangle$ positive (recall that $\xi \ll 1$ and thus $\mathcal{G}_0 \gg 1$). In comparison, the horizontal turbulent velocities in Eq. (9) are obviously positive and are the same as that obtained by Kim (2005) in the case without a latitudinal shear.

Equation (9) indicates that the amplitude of the three components of the velocity depends not only on the radial and latitudinal shear (shearing rate \mathcal{A} and $\epsilon \mathcal{A}$) but also on the typical wavenumber \mathbf{k} and the power spectrum of the forcing $\psi_{ij}(\mathbf{k})$. We first examine how various components are affected by the radial and latitudinal shear. As can easily be seen from Eq. (9), the amplitude of all three components of the turbulent velocity is reduced due to radial shear, becoming very small as \mathcal{A}




Fig. 1. Sketch of the coordinate transformation from (x, z) to (x', z') . In the new referential, the shear is purely radial (see main text for details).

increases. Specifically, the horizontal turbulence is reduced by a factor $\mathcal{A} \mathcal{G}_0^{-1} \propto \mathcal{A}^{2/3}$, the inclusion of the latitudinal shear having no effect on the horizontal turbulence amplitude. In comparison, the radial turbulence velocity has a small contribution of the order of $O(\epsilon^2)$ from ψ_{33} because of a latitudinal shear. In order to understand this, it is instructive to consider a new reference frame such that the shear flow $\mathbf{U}_0 \hat{y}$, depending on both x and z , becomes a function of only one coordinate x' . This coordinate transformation can easily be obtained by rotating x and z axis by the angle $\theta = \arctan(\mathcal{A}_z / \mathcal{A}_x) \sim \epsilon$ around the y axis (see Fig. 1) as follows:

$$\begin{pmatrix} x' \\ z' \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} x \\ z \end{pmatrix} \sim \begin{pmatrix} 1 & \epsilon \\ -\epsilon & 1 \end{pmatrix} \begin{pmatrix} x \\ z \end{pmatrix}, \quad (11)$$

where we used $\epsilon \ll 1$. In the new (x', z') coordinates, the large-scale velocity field is given by $\mathbf{U}'_0 = -(\mathcal{A}'_x + \mathcal{A}'_z)^{1/2} x' \hat{y}'$, depending only on the radial direction, thus becoming a purely radial shear. In the (x', z') coordinate, we can therefore use the results of Kim (2005) that $\langle v_z'^2 \rangle \propto \mathcal{A}'^{-1} \mathcal{G}_0 \sim \mathcal{A}'^{-2/3}$ and $\langle v_x'^2 \rangle \propto \mathcal{A}'^{-1}$. By relating these to $\langle v_x^2 \rangle$ and $\langle v_z^2 \rangle$ via Eq. (11), we easily obtain the term proportional to ψ_{33} in $\langle v_x^2 \rangle$ in Eq. (9). That is, the latter is the result of the dependence of $\langle v_z^2 \rangle$ on the shear in the (x', z') reference frame, with an additional factor of $\epsilon^2 \sim \sin^2 \theta$ coming from the projection of this term onto the x axis. We also note that $\langle v_x'^2 \rangle = \langle v_z'^2 \rangle$ for $\epsilon \ll 1$, consistent with Eq. (9). We will further use this geometrical argument when discussing the turbulent viscosities.

Keeping these in mind, we now discuss the influence of the different types of forcing on the turbulent intensity. If the turbulent motions are primarily driven in the radial direction ($\psi_{33} = \psi_{13} = 0$), the latitudinal shear has no effect, and the results are the same as in the case of only a radial shear: the radial turbulence is reduced by a factor \mathcal{A} and the ratio of the radial to horizontal turbulence amplitude (taking the y direction as well as the z direction) is thus of the order of $\mathcal{A}^{-1/3}$. Alternatively, if the small-scale flow is mainly driven in the horizontal direction ($\psi_{11} = \psi_{13} = 0$), this component $\langle v_x^2 \rangle$ is reduced by a factor of $\epsilon^2 \mathcal{A}^{-2/3}$, with the ratio $\langle v_x^2 \rangle / \langle v_z^2 \rangle \propto \epsilon^2$, for the reason that was just discussed. Again, note that this is different from the case without latitudinal shear where the radial turbulence intensity was null (in the order of the approximation). The ratio $\langle v_x^2 \rangle / \langle v_z^2 \rangle \propto \epsilon^2$ is still small in the case of the sun, but it interestingly only depends on the ratio of the strength of the azimuthal and radial shear but not on their absolute magnitudes.

4. Turbulent viscosity

We now proceed to the calculation of the radial v_T^{xx} and horizontal viscosity v_T^{zz} defined as: $\langle v_x v_y \rangle = -v_T^{xx} \partial_x U_0 = v_T^{xx} \mathcal{A}$ and $\langle v_z v_y \rangle = -v_T^{zz} \partial_z U_0 = v_T^{zz} \epsilon \mathcal{A}$. From the previous section, one might naively estimate the ratio of these two to be:

$$\frac{\langle v_x v_y \rangle}{\langle v_y v_z \rangle} \sim \sqrt{\frac{\langle v_x^2 \rangle}{\langle v_z^2 \rangle}} \sim \xi^{1/6} \quad \text{or} \quad \epsilon, \quad (12)$$

for a x -dominant and z -dominant forcing, respectively. However, the momentum fluxes $\langle v_x v_y \rangle$ and $\langle v_y v_z \rangle$ depend not only on the velocity amplitudes, but also on how closely different components of the velocity are correlated (i.e. the phase-relation). That is, the ratio of the two should be given as follows:

$$\frac{\langle v_x v_y \rangle}{\langle v_y v_z \rangle} = \sqrt{\frac{\langle v_x^2 \rangle \cos \delta_{xy}}{\langle v_z^2 \rangle \cos \delta_{zy}}}, \quad (13)$$

where $\cos \delta_{xy}$ and $\cos \delta_{zy}$ (the so-called cross phase) represent the phase-relation between the x and y and the z and y components of the velocity, respectively. Thus, the estimate given in Eq. (12) may not be true if the cross-phase $\cos \delta_{xy}$ and $\cos \delta_{zy}$ are affected in an anisotropic way due to the different strengths in the radial and latitudinal shear. We shall show that this is indeed the case. Specifically, $\cos \delta_{xy} / \cos \delta_{zy} \propto \xi^{1/6}$ when the turbulence is driven mainly in the x -direction, while $\cos \delta_{xy} / \cos \delta_{zy} = -1$ when driven in the horizontal direction.

After a long, but straightforward algebra, we can obtain eddy viscosities in the following form (see Appendix B for details):

$$\begin{aligned} v_T^{xx} &= \frac{\tau_f}{(2\pi)^3 \mathcal{A}^2} \int d^3 \mathbf{k} \left\{ \frac{g+a^2}{2g^2} [b^2(g^2+a^2)\kappa_0^2 - g \right. \\ &\quad \left. - \frac{4\epsilon b^3 a}{\sqrt{g}} \kappa_0 \mathcal{G}_0] \psi_{11}(\mathbf{k}) \right. \\ &\quad \left. - \frac{(g+a^2)2\epsilon b^2}{g^{3/2}} \kappa_0 \mathcal{G}_0 \psi_{13}(\mathbf{k}) - \frac{2\epsilon^2 b^2 (g+a^2)}{g^{3/2}} \kappa_0 \mathcal{G}_0 \psi_{33}(\mathbf{k}) \right\}, \\ v_T^{zz} &= \frac{\tau_f}{(2\pi)^3 \mathcal{A}^2} \int d^3 \mathbf{k} \frac{1}{\epsilon} \left\{ \frac{g+a^2}{2g^2} \left[ba + \frac{4\epsilon b^4}{\sqrt{g}} \kappa_0 \mathcal{G}_0 \right] \psi_{11}(\mathbf{k}) \right. \\ &\quad \left. + \frac{2b^2(g+a^2)}{g^{3/2}} \kappa_0 \mathcal{G}_0 \left(\psi_{13}(\mathbf{k}) + \epsilon \psi_{33}(\mathbf{k}) \right) \right\}. \end{aligned} \quad (14)$$

Here, again, $a = k_x/k_y$, $b = k_z/k_y$, $g = 1 + b^2$, $\kappa_0 = \pi/2 - \arctan(a/\sqrt{g}) = \arctan(\sqrt{g}/a)$, $\mathcal{G}_0 = \Gamma(1/3)(3/2\xi)^{1/3}/3$, and $\xi = vk_y^2/\mathcal{A}$.

Equation (14) clearly shows that both v_T^{xx} and v_T^{zz} are reduced by shearing, becoming very small as \mathcal{A} increases. The exact scalings of the eddy viscosities, however, depend on the properties of the forcing. Since $\mathcal{G}_0 \propto \xi^{-1/3} \propto \mathcal{A}^{1/3}$ appears in some of the coefficients proportional to ψ_{11} in Eq. (14) with both positive and negative signs, the estimate is rather complex in the case where the forcing is mainly driven radially by ψ_{11} . To obtain a transparent scaling relation in this case, it is illuminating to consider a physically plausible case where the turbulent plumes coming from the convection zone are highly elongated in the x -direction, with small-scale structures in the radial direction being mainly created by radial shear (see Fig. 2). In this case, it is very likely that the forcing will select small wave numbers in the radial direction, permitting us to crudely set $a = k_x/k_y \sim 0$ in Eq. (14). Mathematically, this is justified if the support of the ψ_{ij} functions is localised near the line $k_x/k_y = 0$. In that case, the radial




Fig. 2. Effect of the large-scale shear (*left*) on plumes coming from the convection zone (*right*). In addition to the tilting induced by the large-scale shear, the turbulent velocity field induces stochastic distortion of the eddies, which is not drawn here for the sake of simplicity.

viscosity does not contain terms involving ϵ or ξ , while the horizontal component is proportional to $\mathcal{G}_0 = \Gamma(1/3)(3/2\xi)^{1/3}/3$. Therefore, the ratio of these two viscosities is of the order of $v_T^{xx}/v_T^{zz} \propto \xi^{1/3} \ll 1$. In other words, the momentum transport becomes anisotropic with a much larger horizontal transport compared to the radial one.

Interestingly, the contribution from ψ_{33} to v_T^{xx} comes with the opposite sign to that of ψ_{11} , while the contribution from ψ_{33} to v_T^{zz} comes with the same sign as that of ψ_{11} . Thus, as ψ_{33} takes a non-vanishing value due to horizontal forcing, it will further decrease v_T^{xx} , while increasing v_T^{zz} until v_T^{xx} vanishes and then becomes negative. In other words, the anisotropy in momentum transport becomes even stronger with $v_T^{xx}/v_T^{zz} \ll \xi^{1/3}$. In the extreme limit where the turbulence is solely driven horizontally with $\psi_{11} = \psi_{13} = 0$, $v_T^{xx}/v_T^{zz} \propto -\epsilon^2$. Note that this is true in general, regardless of the form of power spectrum of ψ_{33} . Since $\epsilon^2 \ll 1$, the ratio of the magnitude of the two is small. It is important to note that the two viscosities have different signs, with the radial (vertical) viscosity being negative, while the horizontal one is positive. To understand these results, it is useful to consider the coordinate (x', z') given in Eq. (11) where the shear flow depends only on x' . By an elementary calculation, we can easily show that the turbulent viscosity $v_T^{x'x'}$ in the (x', z') coordinate can be expressed in terms of v_T^{xx} and v_T^{zz} in the (x, z) coordinate as follows:

$$v_T^{x'x'} = \frac{1}{\mathcal{A}_x^2 + \mathcal{A}_z^2} \left[\mathcal{A}_x^2 v_T^{xx} + \mathcal{A}_z^2 v_T^{zz} \right]. \quad (15)$$

We now recall that in the (x', z') coordinate with a purely radial shear, $v_T^{x'x'} = 0$ in the case of a purely horizontal forcing ($\psi_{11} = \psi_{13} = 0$), as shown by Kim (2005). Thus, requiring that the right-hand side of Eq. (15) be zero immediately gives us the result $v_T^{xx}/v_T^{zz} \propto -\epsilon^2$, as found in this paper. This is an interesting result, since the negative viscosity here is not related to the 2-dimensionality of the flow in any sense, being merely due to the fact that a special rotation of the Cartesian frame maps our problem to the case of a purely radial shear.

To summarise, for a reasonable choice of the forcing, $v_T^{xx}/v_T^{zz} \propto \xi^{1/3} \ll 1$ and $-\epsilon^2$ for radial and horizontal forcing, respectively. In these two limiting cases, the ratio of the cross-phase given in Eq. (13) satisfies $\cos \delta_{xy} / \cos \delta_{zy} \propto \xi^{1/6}$ and -1 . It is worth noting that the small value of $\cos \delta_{xy} / \cos \delta_{zy} \propto \xi^{1/6}$ in the former case physically makes sense since a stronger decorrelation between velocity components in the radial direction and horizontal plane is expected because of a stronger radial shear.

These results thus suggest that horizontal momentum transport can be much more efficient than radial transport due to radial and latitudinal differential rotations.

5. Conclusions and discussion

We presented a novel mechanism for the anisotropic angular momentum transport to explain tachocline confinement. Starting from the first principle, we have consistently derived eddy viscosities and turbulence amplitude by incorporating the crucial effects of both radial and latitudinal differential rotations. We have shown that even in the absence of stratification and magnetic field, the shear induced by differential rotation alone can lead to anisotropic momentum transport. Specifically, when the turbulence is mainly driven radially by overshooting plumes coming from the convection zone, the ratio of the radial to horizontal eddy viscosities can be proportional to $\xi^{1/3}$, becoming very small as the radial shear \mathcal{A} increases. Here, $\xi = \nu k_y^2 / \mathcal{A}$ and \mathcal{A} is the shearing rate in the radial direction. In comparison, in the case where the turbulence is mainly driven horizontally, this ratio is proportional to $-\epsilon^2$, where ϵ ($\ll 1$) is the ratio of the shearing rate in the latitudinal direction to that in the radial one. In this case, the radial eddy viscosity becomes negative. These results thus suggest that horizontal momentum transport can be much more efficient than radial transport, with a strong anisotropic momentum transport, as a result of the shearing effects due to radial and latitudinal differential rotations.

As mentioned in the introduction, the anisotropic momentum transport has important implications for the dynamics of the tachocline. Spiegel & Zahn (1992) have shown that the radiative spreading of the tachocline could be limited if the ratio of the radial to horizontal diffusivities satisfies the following relation:

$$\frac{\nu_T^{xx}}{\nu_T^{zz}} \ll \left(\frac{h}{r_0}\right)^2 \sim 3 \times 10^{-3}, \quad (16)$$

where r_0 is the radius at which the tachocline is located and h its thickness. The estimate on the right-hand side has been obtained using helioseismologic values of Charbonneau et al. (1999). Indeed, this criterion can easily be satisfied for a reasonable choice of the forcing as follows. First, in the case of radial forcing that is dominated by the components with $k_x/k_y \ll 1$, the Spiegel & Zahn criterion is easily met for a characteristic length-scale in the azimuthal direction $L_y = 2\pi k_y^{-1} \gg 2\pi(r_0/h)^3(\nu/\mathcal{A})^{1/2} \sim 3 \times 10^5$ m for typical values of the solar tachocline $\mathcal{A} \sim 3 \times 10^{-6}$ s $^{-1}$. In the case of horizontal forcing, the ratio of radial to horizontal eddy viscosities becomes $\epsilon^2 \sim 7 \times 10^{-4}$. Furthermore, in this case, the radial eddy viscosity becomes negative, reinforcing the presence of radial shear with a sharp gradient in a localised region. Therefore, these results suggest that the anisotropy in momentum transport could be strong enough to operate as a mechanism for the tachocline confinement against spreading. Our results, however, cannot offer a mechanism for a uniform rotation in the solar interior. As noted in the introduction, this could be due to the presence of magnetic fields in this region (Rüdiger & Kichatinov 1997; Gough & McIntyre 1998; MacGregor & Charbonneau 1999).

We note that in this paper we have neglected some effects that could be important in the lower part of the tachocline. For example, the presence of stratification or rotation can generate a wave-like turbulence and consequently severely affects its

transport properties. The effect of Coriolis forces was studied in the context of the transport of angular momentum in the convection zone to elucidate the permanence of a differential rotation profile in that region. It has been shown that, in the strong rotation limit, the turbulent viscosity is more reduced by a factor of 4 (Kichatinov et al. 1994) in the direction perpendicular to the rotation axis compared to the one parallel to the rotation axis. Furthermore, rotation induces a non-diffusive part in the turbulent viscosity (the so-called Λ effect), which prevents the solid rotation to be a solution of the turbulent momentum equation. We can expect a density stratification to have a similar effect to the Coriolis force, as the linear equations are the same (Greenspan 1968) once the vertical and horizontal directions are interchanged. In that case, we expect the component in the direction of the stratification to be more reduced than in the plane of constant density. In fact, such an anisotropic eddy viscosity has been observed in a recent numerical simulation of a stably stratified turbulence driven by penetrative convection with an imposed shear (Miesch 2003). In particular, the angular momentum transport was found to be diffusive in the latitudinal direction but anti-diffusive in the radial direction, with turbulence mixing potential vorticity (see also McIntyre 2003). Another important effect is due to magnetic fields in the tachocline (as suggested by solar-dynamo models) by imposing Alfvén-like structure in the properties of turbulence. Furthermore, the simultaneous presence of toroidal magnetic field and latitudinal differential rotation can lead to a large-scale joint instability (Gilman & Fox 1997; Dikpati et al. 2004), which could further enforce an anisotropic turbulence. The interplay between shear and rotation is now being investigated (Leprovost & Kim 2006) and the related problems of stratification and magnetic field will be published in future papers.

Acknowledgements. This work was supported by U.K. PPARC Grant No. PP/B501512/1.

References

- Burrell, K. H. 1997, *Phys. Plasmas*, 4, 1499
- Charbonneau, P., Christensen-Dalsgaard, J., Henning, R., et al. 1999, *ApJ*, 527, 445
- Dikpati, M., Cally, P. S., & Gilman, P. A. 2004, *ApJ*, 610, 597
- Garaud, P. 2003, *Dynamics of the solar tachocline*, ed. M. J. Thompson, & J. Christensen-Dalsgaard (CUP)
- Gilman, P. A., & Fox, P. 1997, *ApJ*, 484, 439
- Gough, D. O., & McIntyre, M. E. 1998, *Nature*, 394, 755
- Gradshteyn, I. S., & Ryzhik, I. M. 1965, *Table of integrals series and products* (Academics press)
- Greenspan, H. P. 1968, *The theory of rotating fluids* (CUP)
- Hahm, T. S. 2002, *Plasma Phys. Control. Fusion A*, 44, 87
- Kichatinov, L. L., Pipin, V. V., & Rüdiger, G. 1994, *Astron. Nachr.*, 315, 157
- Kim, E. 2004, *Mod. Phys. Lett. B*, 18, 551
- Kim, E. 2005, *A&A*, 441, 763
- Kim, E. 2006, *Phys. Rev. Lett.*, 96, 084504
- Kim, E., & Diamond, P. H. 2003, *Phys. Rev. Lett.*, 90, 185006
- Kim, E., Diamond, P. H., & Hahm, T. S. 2004, *Phys. Plasmas*, 11, 4554
- Kim, E., & Durrulle, B. 2001, *Phys. Plasmas*, 8, 813
- Leprovost, N., & Kim, E. 2006, in preparation
- MacGregor, K. B., & Charbonneau, P. 1999, *ApJ*, 519, 911
- McIntyre, M. E. 2003, *Solar tachocline dynamics: eddy viscosity, anti-friction or something in between*, ed. M. J. Thompson, & J. Christensen-Dalsgaard (CUP)
- Miesch, M. S. 2003, *ApJ*, 586, 663
- Moffatt, H. K. 1978, *Magnetic field generation in fluids* (CUP)
- Rüdiger, G., & Kichatinov, L. L. 1997, *Astron. Nachr.*, 318, 273
- Spiegel, E. A., & Zahn, J.-P. 1992, *A&A*, 265, 106

Online Material

Appendix A: Solution of system (4)

Using the new field $\hat{u} = \hat{v}_x + \epsilon \hat{v}_z$, we obtain the following equations:

$$\begin{aligned} \mathcal{A} \partial_\tau [R(\tau) \hat{u}] &= \hat{h}_1, \\ \mathcal{A} [\gamma + \phi \epsilon \tau] \partial_\tau \hat{v}_x &= -\mathcal{A} \partial_\tau [\tau^2 \hat{u}] + [\gamma + \phi \epsilon \tau] \hat{f}_x - \tau \hat{f}_y - \phi \tau \hat{f}_z, \end{aligned} \quad (\text{A.1})$$

where $\gamma = 1 + \phi^2$, $R(\tau) = (\epsilon^2 + 1)\tau^2 + 2\epsilon\phi\tau + \gamma$ and $\hat{h}_1 = [\gamma + \phi \epsilon \tau] \hat{f}_x - [(1 + \epsilon^2)\tau + \phi \epsilon] \hat{f}_y + (\epsilon - \phi \tau) \hat{f}_z$. The first equation can be readily integrated to obtain $\hat{u} = \int d\tau_1 \hat{h}_1(\tau_1) / \mathcal{A} R(\tau_1)$. Then, the second equation can be used to obtain \hat{v}_x . Here, we provide some of the main steps in the calculation. First, we calculate the first term on the RHS of Eq. (A.1):

$$\partial_\tau [\tau^2 \hat{u}] = \frac{\tau^2 \hat{h}_1(\tau)}{\mathcal{A} R(\tau)} + \frac{2\tau[\gamma + \epsilon\phi\tau]}{R(\tau)^2} \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}}. \quad (\text{A.2})$$

Then plugging Eq. (A.2) into the second equation of (A.1), we obtain:

$$\begin{aligned} \hat{v}_x &= - \int_{\tau_0}^{\tau} d\tau_2 \frac{2\tau_2}{R(\tau_2)^2} \int_{\tau_0}^{\tau_2} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}} \\ &\quad + \int_{\tau_0}^{\tau} d\tau_1 \frac{[R(\tau_1) - \tau_1^2] \hat{f}_x - \tau_1 \hat{f}_y - \tau_1(\phi + \epsilon\tau_1) \hat{f}_z}{\mathcal{A} R(\tau_1)}, \\ &= - \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}} \int_{\tau_1}^{\tau} d\tau_2 \frac{2\tau_2}{R(\tau_2)^2} \\ &\quad + \int_{\tau_0}^{\tau} d\tau_1 \frac{[R(\tau_1) - \tau_1^2] \hat{f}_x - \tau_1 \hat{f}_y - \tau_1(\phi + \epsilon\tau_1) \hat{f}_z}{\mathcal{A} R(\tau_1)}, \\ &= \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_1(\tau_1)}{\mathcal{A}} \left\{ \frac{\gamma + \epsilon\phi\tau}{(\gamma + \epsilon^2)R(\tau)} + \frac{\epsilon\phi}{(\gamma + \epsilon^2)^{3/2}} [T(\tau) - T(\tau_1)] \right\} \\ &\quad - \int_{\tau_0}^{\tau} d\tau_1 \frac{\hat{h}_2(\tau_1)}{\mathcal{A}} \frac{\epsilon}{\gamma + \epsilon^2}. \end{aligned} \quad (\text{A.3})$$

Here we used the formula (2.175) and (2.172) of Gradshteyn & Ryzhik (1965) to calculate the integrals; $T(\tau) = \arctan[(\epsilon^2 + 1)\tau + \epsilon\phi] / \sqrt{\gamma + \epsilon^2}$ and $\hat{h}_2(\tau) = -\epsilon \hat{f}_x(\tau) - \phi \hat{f}_y(\tau) + \hat{f}_z(\tau)$. The two other components of the velocity can be easily obtained by using $\hat{v}_z = (\hat{u} - \hat{v}_x) / \epsilon$ and $\hat{v}_y = -\tau \hat{v}_x - (\epsilon\tau + \phi) \hat{v}_z$.

Appendix B: Calculation of the turbulent viscosities

We symbolically write $\langle v_x v_y \rangle$ and $\langle v_z v_y \rangle$ as follows:

$$\langle v_x v_y \rangle = \frac{\tau_f}{(2\pi)^3 \mathcal{A}} \int d^3 \mathbf{k} \{ V_{11} \phi_{11} + V_{12} \phi_{12} + V_{22} \phi_{22} \}, \quad (\text{B.1})$$

$$\langle v_z v_y \rangle = \frac{\tau_f}{(2\pi)^3 \mathcal{A}} \int d^3 \mathbf{k} \{ H_{11} \phi_{11} + H_{12} \phi_{12} + H_{22} \phi_{22} \},$$

where the ϕ_{ij} functions are the power spectra associated with the forcing functions h_1 and h_2 :

$$\langle \tilde{h}_i(\mathbf{k}_1, t_1) \tilde{h}_j(\mathbf{k}_2, t_2) \rangle = \tau_f (2\pi)^3 \delta(\mathbf{k}_1 + \mathbf{k}_2) \delta(t_1 - t_2) \phi_{ij}(\mathbf{k}_2), \quad (\text{B.2})$$

for i and $j = 1, 2$. The functions V_{ij} and H_{ij} (for i and $j = 1, 2$) are expressed as integrals over the variable τ ; for example, V_{12} can be written as

$$\begin{aligned} V_{12} &\equiv \int_a^{+\infty} d\tau e^{-2\nu(Q(\tau) - Q(a))} \left\{ \frac{(\epsilon^2 - \gamma)\phi + \epsilon[\epsilon^2 + 1 - \phi^2]\tau}{(\gamma + \epsilon^2)^2} \right. \\ &\quad \left. - \frac{2\epsilon\phi^2}{(\gamma + \epsilon^2)^{5/2}} [T(\tau) - T(a)] \right\}, \end{aligned} \quad (\text{B.3})$$

where, $a = k_x/k_y$, $\phi = (k_z - \epsilon k_x)/k_y$ and $\gamma = 1 + \phi^2$. Following Kim (2005), this integral is estimated in the strong shear limit, i.e. for $\xi \equiv \nu k_y^2 / \mathcal{A} \ll 1$. Thus, $\nu[Q(\tau) - Q(a)] = \xi[(\epsilon^2 + 1)\tau^3/3 + \epsilon\phi\tau^2 + \gamma\tau - \{\gamma a + (\epsilon^2 + 1)a^3/3 + \epsilon a^2\}]$, where $b = k_z/k_y$. Taking the limit $\xi \rightarrow 0$, we evaluate each term in (B.3) by keeping only terms to leading order in ξ . The results are

$$\begin{aligned} V_{11} &= -\frac{\gamma + \epsilon\phi a}{2(\gamma + \epsilon^2)^2 R(a)} - \frac{\epsilon\phi}{2(\gamma + \epsilon^2)^{5/2} \kappa} + \frac{(\gamma - \epsilon^2)\phi^2}{2(\gamma + \epsilon^2)^3} \kappa^2 \\ &\quad + \frac{\epsilon\phi^3}{(\gamma + \epsilon^2)^3} \kappa^2 \mathcal{G} + \frac{\epsilon\phi(\phi^2 - 1 - \epsilon^2)}{(\gamma + \epsilon^2)^{5/2}(\epsilon^2 + 1)} \kappa \mathcal{L}, \\ V_{12} &= \frac{(\epsilon^2 - \gamma)\phi}{(\gamma + \epsilon^2)^{5/2}} \kappa - \frac{2\epsilon\phi^2}{(\gamma + \epsilon^2)^{5/2}} \kappa \mathcal{G} + \frac{\epsilon(1 + \epsilon^2 - \phi^2)}{(\gamma + \epsilon^2)^2(\epsilon^2 + 1)} \mathcal{L}, \\ V_{22} &= \frac{\phi\epsilon}{(\gamma + \epsilon^2)^2} \mathcal{G}, \\ H_{11} &= \frac{\phi a - \epsilon}{2(\gamma + \epsilon^2)^2 R(a)} + \frac{\phi}{2(\gamma + \epsilon^2)^{5/2}} \kappa + \frac{\epsilon\phi^2}{(\gamma + \epsilon^2)^3} \kappa^2 \\ &\quad - \frac{\phi^3}{(\gamma + \epsilon^2)^3} \kappa^2 \mathcal{G} + \frac{(\epsilon^2 + 1 - \phi^2)\phi}{(\gamma + \epsilon^2)^{5/2}(\epsilon^2 + 1)} \kappa \mathcal{L}, \\ H_{12} &= -\frac{2\epsilon\phi}{(\gamma + \epsilon^2)^{5/2}} \kappa + \frac{2\phi^2}{(\gamma + \epsilon^2)^{5/2}} \kappa \mathcal{G} + \frac{\phi^2 - 1 - \epsilon^2}{(\gamma + \epsilon^2)^2(\epsilon^2 + 1)} \mathcal{L}, \\ H_{22} &= -\frac{\phi}{(\gamma + \epsilon^2)^2} \mathcal{G}. \end{aligned} \quad (\text{B.4})$$

Here, $\kappa = \pi/2 - T(a)$. To derive Eq. (B.4), we used formula (2.173), (2.175), and (2.18) in Gradshteyn & Ryzhik (1965). The results are written in terms of new integrals by making the substitution $y = 2\xi(\epsilon^2 + 1)\tau^3/3$ and taking the limit $\xi \rightarrow 0$:

$$\begin{aligned} \mathcal{G} &= \frac{1}{3} \left(\frac{3}{2\xi(\epsilon^2 + 1)} \right)^{1/3} \int_0^\infty \frac{dy}{y^{2/3}} \\ &\quad \times \exp \left[-y - \frac{3^{5/3}\epsilon\phi}{(\epsilon^2 + 1)} \left(\frac{2\xi(\epsilon^2 + 1)}{3} \right)^{1/3} y^{2/3} + O(\xi^{4/3}) \right] \\ &= \frac{\Gamma(1/3)}{3} \left(\frac{3}{2\xi(\epsilon^2 + 1)} \right)^{1/3} - \frac{3^{2/3}\epsilon\phi}{(\epsilon^2 + 1)} + O(\xi^{2/3}) \\ &\equiv \mathcal{G}_0 - \frac{3^{2/3}\epsilon\phi}{(\epsilon^2 + 1)} + O(\xi^{2/3}), \\ \mathcal{L} &= \frac{1}{3} \int_{\xi \rightarrow 0}^\infty \frac{dy}{y} \exp \left[-y - \frac{3^{5/3}\epsilon\phi}{(\epsilon^2 + 1)} \left(\frac{2\xi(\epsilon^2 + 1)}{3} \right)^{1/3} y^{2/3} + O(\xi^{4/3}) \right] \\ &= \frac{-\ln(\xi)}{3} + O(\xi^{1/3} \ln(\xi)) \equiv \mathcal{L}_0 + O(\xi^{1/3} \ln(\xi)). \end{aligned} \quad (\text{B.5})$$

We simplify (B.4) by keeping terms up to at most second order in ϵ . To this order, we have

$$\begin{aligned} \phi &= b - \epsilon a, \quad \gamma = 1 + b^2 - 2\epsilon b a + \epsilon^2 a^2, \\ R(a) &= (\gamma + a^2), \end{aligned} \quad (\text{B.6})$$

$$\kappa = \left[\pi/2 - \arctan \left(\frac{a}{\sqrt{\gamma}} \right) \right] - \frac{\epsilon\phi\sqrt{\gamma}}{\gamma + a^2} + O(\epsilon^2) \equiv \kappa_0 - \frac{\epsilon b}{\sqrt{g}} + O(\epsilon^2).$$

For simplicity, we assume that $\phi_{ij}(k_y) = \phi_{ij}(-k_y)$ and consequently neglect all the terms proportional to an odd power of b only (but keep those proportional to ba). Recalling that ξ is a small parameter, we will keep only the terms proportional to $\xi^{-1/3}$ and ξ^0 and the leading order term in the ϵ expansion. The dominant contributions then give us

$$V_{11} = -\frac{1}{2g(g + a^2)} + \frac{b^2}{2g^2} \kappa_0^2 - \frac{2\epsilon^2 b^4}{g^{7/2}} \kappa_0 \mathcal{G}_0 + \dots, \quad (\text{B.7})$$

$$V_{12} = -\epsilon \frac{2b^2}{g^{5/2}} \kappa_0 \mathcal{G}_0 + \dots,$$

$$V_{22} = -\frac{3^{2/3} \epsilon^2 b^2}{g^2} + \dots,$$

$$H_{11} = \frac{ba}{2g^2(g+a^2)} + \frac{2\epsilon b^4}{g^{5/2}(g+a^2)} \kappa_0 \mathcal{G}_0 + \dots,$$

$$H_{12} = \frac{2b^2}{g^{5/2}} \kappa_0 \mathcal{G}_0 + \dots,$$

$$H_{22} = \frac{3^{2/3} \epsilon b^2}{g^2} + \dots$$

Here, $g = 1 + b^2 = k_H^2/k_y^2$, and the dots stand for higher order terms in ξ and ϵ compared to those that were kept.

We now express ϕ_{ij} in terms of ψ_{ij} by assuming the forcing to be incompressible, for simplicity. In that case, the following relations hold: $\hat{h}_1(\tau) = R(\tau)\{\hat{f}_x + \epsilon \hat{f}_z\}$, and $\hat{h}_2(\tau) = [\phi\tau - \epsilon]\hat{f}_x + [\gamma + \epsilon\phi\tau]\hat{f}_z$. Keeping only the terms at most proportional to ϵ , we obtain

$$\begin{aligned} \phi_{11} &= (g+a^2)^2\psi_{11} + 2\epsilon(g+a^2)^2\psi_{13} + O(\epsilon^2), \\ \phi_{12} &= (g+a^2)[ba\psi_{11} + g\psi_{13}] \\ &\quad + \epsilon\{-(a^2+1)(g+a^2)\psi_{11} + g(g+a^2)\psi_{33}\} + O(\epsilon^2), \\ \phi_{22} &= b^2a^2\psi_{11} + 2gba\psi_{13} + g^2\psi_{33} + \epsilon\{-2ba(a^2+1)\psi_{11} \\ &\quad - 2[a^2b^2 + g(1+a^2)]\psi_{13} - 2gba\psi_{33}\} + O(\epsilon^2). \end{aligned} \tag{B.8}$$

By using (B.8) and Eq. (B.7) for the turbulent viscosities and by keeping only the dominant term for each component of the forcing, we obtain the Eqs. (14) given in the main text.