

ELODIE metallicity-biased search for transiting Hot Jupiters[★]

III. A hot Jupiter orbiting the star HD 185269

C. Moutou¹, B. Loeillet¹, F. Bouchy^{2,3}, R. Da Silva⁴, M. Mayor⁴, F. Pont⁴, D. Queloz⁴, N. C. Santos^{4,5,6},
D. Ségransan⁴, S. Udry⁴, and S. Zucker⁷

¹ Laboratoire d'Astrophysique de Marseille, Traverse du Siphon, 13013 Marseille, France

e-mail: Claire.Moutou@oamp.fr

² IAP, 98bis Bd Arago, 75014 Paris, France

³ Observatoire de Haute Provence, 04870 St Michel l'Observatoire, France

⁴ Observatoire de Genève, 51 ch. des Maillettes, 1290 Sauverny, Switzerland

⁵ Lisbon Observatory, Tapada da Ajuda, 1349-018 Lisboa, Portugal

⁶ Centro de Geofísica de Evora, Rua Romão Ramalho 59, 7002-554 Evora, Portugal

⁷ Weizmann Institute of Science, PO Box 26, Rehovot 76100, Israel

Received 13 July 2006 / Accepted 9 August 2006

ABSTRACT

We present new results of a search for extrasolar planets. The survey uses radial-velocity techniques and focuses on metal-rich stars. We used radial velocity measurements obtained with the echelle spectrograph ELODIE at the Observatoire de Haute Provence. New data have revealed a planetary companion to the slightly evolved star HD 185269. It belongs to the hot Jupiter class of planets, with a projected mass $M \sin i = 1.0 M_{\text{Jup}}$ and period $P = 6.84$ days. We describe the measurements that led to this discovery and discuss the orbital solution.

Key words. stars: individual: HD 185269 – planetary systems – techniques: radial velocities

1. Introduction

There are approximately 40 known extrasolar planets with a semi-major axis smaller than 0.1 AU, the so-called hot Jupiters. Among them, seven have a mass of less than about $0.1 M_{\text{Jup}}$ and should better be called hot Neptunes. These hot Neptunes could be very abundant as only recently have they become available above the detection limit of instruments. 85% of these short-period planets have a metal-rich parent star, in comparison to 70% for the whole sample of extrasolar planets. There is thus an indication that the frequency of planets versus the metallicity of the parent star relationship is stronger for short-period planets (see also Sozzetti 2004). To understand whether this trend is significant, we need data from more short-period planets orbiting metal-rich stars. This is the goal of two observational programs using radial-velocity techniques; both surveys started in 2003 and several planets have been discovered: HD 88133 (Fischer et al. 2005), HD 118203 (Da Silva et al. 2006), HD 149026 (Sato et al. 2005), HD 149143 and HD 109749 (Fischer et al. 2006; Da Silva et al. 2006) and HD 189733 (Bouchy et al. 2005). The observational strategy of these programs, introducing a bias in the selection of target stars based on their metallicity, allows a more efficient detection of planets, and will clarify the link between the metallicity of stars and the planet formation and migration. Also, the discovery of short-period planets opens the possibility to discover new transiting systems (as e.g. HD 149026 and HD 189733, see references above), with precise

determinations of their mass and radius. However, non-biased target sample surveys are still needed in order to explore other parameters of the origins of planetary systems.

In this paper, we report the detection of a new planetary system around the main-sequence star HD 185269 at the Observatoire de Haute Provence. In Sect. 2 the observations are described; in Sect. 3 we derive the stellar parameters and the interpretation is discussed in Sect. 4.

2. Observations

Radial-velocity measurements of the main-sequence star HD 185269 were obtained with ELODIE from June 2005 to July 2006 in the framework of the survey of metal-rich stars as described in Da Silva et al. (2006). ELODIE is a fiber-fed echelle spectrograph (Baranne et al. 1996) mounted on the 193-cm telescope at the Observatoire de Haute Provence (France), designed to obtain simultaneous spectra of a star and a calibration ThAr lamp, for precise radial-velocity measurements.

A first observation of HD 185269 shows its enrichment, with a metallicity deduced from the cross-correlation function of 0.15. Further measurements of this star revealed fluctuations in the radial velocity over a few days. The star was then regularly monitored. Individual measurements of HD 185269 have errors of the order of $10\text{--}12 \text{ m s}^{-1}$ (Table 1) while the standard deviation of the series is about 90 m s^{-1} . A 6.8-d periodicity is clearly visible.

[★] Based on observations collected with the ELODIE spectrograph on the 1.93-cm telescope (Observatoire de Haute Provence, France).

Table 1. Radial-velocity measurements of HD 185269 obtained with ELODIE from June 2005 to June 2006.

JD-2 400 000. day	VR km s ⁻¹	error ms ⁻¹
53551.55350	0.535	0.011
53552.57440	0.639	0.013
53553.54090	0.697	0.012
53554.55910	0.690	0.008
53555.54490	0.663	0.009
53574.55610	0.687	0.012
53576.51360	0.631	0.011
53577.53300	0.532	0.010
53587.46980	0.672	0.011
53588.45620	0.677	0.009
53589.46500	0.606	0.012
53590.45090	0.631	0.014
53591.48810	0.530	0.011
53592.41560	0.543	0.011
53594.47180	0.672	0.010
53596.43110	0.677	0.010
53610.43430	0.652	0.010
53613.41040	0.593	0.008
53624.43320	0.630	0.008
53625.37140	0.551	0.010
53626.37360	0.508	0.008
53627.36100	0.597	0.008
53628.32940	0.672	0.008
53667.33630	0.502	0.008
53720.25140	0.617	0.010
53723.22240	0.603	0.011
53808.68270	0.640	0.012
53869.60890	0.672	0.008
53871.55960	0.538	0.008
53872.61180	0.538	0.012
53873.56570	0.589	0.015
53875.60370	0.716	0.010
53894.58770	0.648	0.010
53896.56560	0.683	0.009
53897.56800	0.681	0.009
53901.58760	0.661	0.014
53932.46560	0.609	0.015
53934.42560	0.518	0.009

3. Stellar characteristics of HD 185269

HD 185269 has been observed by HIPPARCOS (ESA 1997). It is a G0IV star with a visual magnitude $V = 6.67$ and color index $B - V = 0.58$ and is located at 47.37 pc from the Sun. This results in an absolute V magnitude of 3.29. Strömgren narrow-band photometry on this star derives a metallicity of 0.02, i.e. almost solar (Nordström et al. 2004). Comparatively, the metallicity derived from the cross-correlation function (0.15) is slightly higher. Direct spectroscopic analysis as described in Santos et al. (2004) allows us to derive: $T_{\text{eff}} = 5983$ K, $\log g = 4.05$ and $[\text{Fe}/\text{H}] = 0.10$ (Table 2).



By comparison with the evolutionary tracks calculated by Girardi et al. (2002), we obtain a stellar mass of $1.33 \pm 0.07 M_{\odot}$ and an age of 4.0 ± 1.0 Gyr. The star is slightly evolved, as also suggested by the lithium content in the spectrum.

The projected rotational velocity derived from a calibration of the cross-correlation function is $v \sin i = 5.5$ km s⁻¹.

The stellar chromospheric activity of HD 185269 is low, as deduced from the absence of emission in the core of the CaII lines (Fig. 1). This excludes large amplitude radial-velocity jitter due to intrinsic activity of the star.

Table 2. Stellar parameters of HD 185269.

Spectral Type	G0 IV
V	6.67
$B - V$	0.58
Distance [pc]	47.37
M_V	3.29
BC	-0.06
M_{bol}	3.228
T_{eff} [K]	5983 ± 62
$\log g$	4.05 ± 0.22
$[\text{Fe}/\text{H}]$	0.10 ± 0.08
$v \sin i$ [km s ⁻¹]	5.5
Star mass [M_{\odot}]	1.33 ± 0.07
Age [Gyr]	4.0 ± 1.0

**Fig. 1.** The CaII K absorption line of the ELODIE spectrum of HD 185269, which shows no emission feature in its core.**Fig. 2.** The inverse slope of the bisector is shown against the radial-velocity measurements of HD 185269. No correlation is observed.

The bisector analysis of the cross-correlation functions shows no correlation with the measured velocimetric variations. The shape of stellar lines is therefore not related to the observed fluctuations in radial velocity (Fig. 2).




Fig. 3. Radial-velocity measurements of HD 185269 superimposed on the best Keplerian solution. Error bars represent the photon-noise uncertainties.

Table 3. Parameters for the Keplerian solution and the planetary companion.

Period [days]	6.8399 ± 0.0013
Periastron epoch [JD-2453000]	797.152 ± 0.12
Orbital eccentricity	0.225 ± 0.025
Radial velocity semi-amplitude [m s^{-1}]	93.56 ± 2.5
Systemic velocity [km s^{-1}]	0.617 ± 0.002
O-C residuals [m s^{-1}]	16
Orbital semi-major axis [AU]	0.077
Planet mass [M_J]	1.03 ± 0.03

4. Planetary system

A careful check of stellar properties shows that the radial-velocity fluctuations of HD 185269 were not likely of stellar origin. A Keplerian orbit was then adjusted on the 38 data points with photon noise uncertainty below 15 m s^{-1} (Fig. 3); the best parameters of the fit are a period of 6.84 days and an eccentricity of 0.23. The final parameters of the orbital solution are presented in Table 3. The residuals are 16 m s^{-1} .

With a stellar mass of $1.33 M_{\odot}$, the deduced minimum mass is $1.03 \pm 0.03 M_{\text{Jup}}$ for the planetary companion in an orbit with a 6.8399 ± 0.0015 day period; the folded orbit is shown in Fig. 4.

Like HD 118203 (Da Silva et al. 2006), HD 185269 has a short-period planet in eccentric orbit. The period of HD 185269 (6.84 days) is above the limit of circularization as expressed in Halbwegs et al. (2005) and evidently, circular orbits are extremely rare when the period is larger than about 5 days (Schneider 2006).




Fig. 4. Phase-folded radial velocity measurements of HD 185269 superimposed on the best Keplerian solution. Error bars represent the photon-noise uncertainties. Bottom plot: residuals as a function of time.

A short-period planet is a good target for transit detectability; for HD 185269, the transit probability is about 8%. However, the eccentricity makes the error on the time reference large and therefore the transit ephemeris is not precise. A photometric observation sequence will nevertheless be scheduled in the following months, in parallel with further radial-velocity measurements aimed at refining the orbital solution.

Note added in proofs: After this paper was submitted, an independent discovery of the same planet was announced by Johnson et al. (2006). Both sets of measurements and analysis are compatible and no transit is visible.

Acknowledgements. We thank the Programme National de Planetologie (PNP), the Swiss National Science Foundation (FNSRS) and the Geneva University for their continuous support of our planet-search programs. NCS thanks the Fundação para a Ciência e a Tecnologia (Portugal) for the scholarship SFRH/BPD/8116/2002 and grant POCI/CTE-AST/56453/2004.

References

- Baranne, A., Queloz, D., Mayor, M., et al. 1996, *A&AS*, 119, 373
- Bouchy, F., Udry, S., Mayor, M., et al. 2005, *A&A*, 444, L15
- Da Silva, R., Udry, S., Bouchy, F., et al. 2006, *A&A*, 446, 717
- ESA 1997, *VizieR Online Data Catalog*, 1239, 0
- Fischer, D. A., Laughlin, G., Butler, P., et al. 2005, *ApJ*, 620, 481
- Fischer, D. A., Laughlin, G., Marcy, G. W., et al. 2006, *ApJ*, 637, 1094
- Girardi, L., Bertelli, G., Bressan, A., et al. 2002, *A&A*, 391, 195
- Halbwachs, J. L., Mayor, M., & Udry, S. 2005, *A&A*, 431, 1129
- Johnson, J., Marcy, G., Fischer, D., et al. 2006, *ApJ*, in press
- Nordström, B., Mayor, M., Andersen, J., et al. 2004, *A&A*, 418, 989
- Santos, N. C., Israelian, G., & Mayor, M. 2004, *A&A*, 415, 1153
- Sato, B., Fischer, D. A., Henry, G. W., et al. 2005, *ApJ*, 633, 465
- Schneider, J. 2006, www.obspm.fr/planets