

The ultra-compact binary 4U 1850–087 observed with *INTEGRAL*: hard X-ray emission from an X-ray burster[★]

L. Sidoli¹, A. Paizis¹, A. Bazzano², and S. Mereghetti¹

¹ INAF-IASF, Istituto di Astrofisica Spaziale e Fisica Cosmica Milano, via E. Bassini 15, 20133 Milano, Italy
e-mail: sidoli@iasf-milano.inaf.it

² INAF-IASF, Istituto di Astrofisica Spaziale e Fisica Cosmica Roma, via del Fosso del Cavaliere 100, 00133 Roma, Italy

Received 5 June 2006 / Accepted 2 September 2006

ABSTRACT

Context. The X-ray burster 4U 1850–087, located in the Galactic globular cluster NGC 6712, is an ultracompact binary (orbital period ~ 21 min), likely harbouring a degenerate companion.

Aims. The source has been observed at soft γ -rays several times with the *INTEGRAL* satellite, during the monitoring of the Galactic plane, with an unprecedented exposure time. We analysed all available *INTEGRAL* observations, with the main aim of studying the long-term behaviour of this Galactic bulge X-ray burster.

Methods. The spectral results are based on the systematic analysis of all *INTEGRAL* observations covering the source position performed between March 2003 and November 2005.

Results. The source X-ray emission is hard and is observed, for the first time, up to 100 keV. A broad-band spectrum obtained combining the *INTEGRAL* spectrum together with a quasi-simultaneous *XMM-Newton* observation performed in September 2003 is well modeled with a disk-blackbody emission (with an inner disk temperature of ~ 0.8 keV) together with a power-law (with a photon index, Γ , of 2.1). The 2–100 keV luminosity is 1.5×10^{36} erg s⁻¹ (assuming a distance of 6.8 kpc).

Conclusions. *INTEGRAL* observations reveal for the first time that this X-ray burster displays a very hard X-ray spectrum, with a cut-off at energies higher than 100 keV, and that the source spends most of the time in this low luminosity and hard state. Indeed, a previous *BeppoSAX* observation in April 1997 observed high energy emission from 4U 1850–087 only up to 50 keV.

Key words. X-rays: stars: individual: 4U 1850–087 – X-rays: binaries

1. Introduction

4U 1850–087 (Swank et al. 1976) is an X-ray burster located in the galactic globular cluster NGC 6712. The likely optical counterpart displays an UV modulation (Anderson et al. 1993) with a period of 20.6 min (interpreted as the orbital period), implying a degenerate companion of $0.04 M_{\odot}$ (Homer et al. 1996).

The first broad-band spectrum from this source was obtained with *BeppoSAX* in the energy range 0.3–50 keV during a pointing performed in 1997 (Sidoli et al. 2001). The spectrum was well described with a disk-blackbody and Comptonized continuum with $N_{\text{H}} = 3.9 \times 10^{21}$ cm⁻², an inner disk temperature, kT_{in} , of ~ 0.6 keV, an inner projected radius of ~ 5 km (for an assumed NGC 6712 distance of 6.8 kpc, Harris 1996), a temperature, kT_0 , of the input “seed” photons, of $0.8^{+0.1}_{-0.2}$ keV, an electron temperature, kT_e , of 70 keV, and an optical depth, τ , of 1.7. The estimated 0.1–100 keV luminosity was 1.9×10^{36} erg s⁻¹ (at 6.8 kpc).

Other X-ray observations are reported for 4U 1850–087 limited to the energy range below 10 keV (*ASCA*, Juett et al. 2001; *XMM-Newton*, Sidoli et al. 2005; *Chandra*, Juett & Chakrabarty 2005).

We report here the first detection of 4U 1850–087 above 50 keV with the *INTEGRAL* satellite, obtained with an unprecedented exposure time collected during the monitoring of the Galactic plane.

2. Observations and results

The ESA *INTEGRAL* gamma-ray observatory, launched in October 2002, carries three co-aligned coded mask telescopes: the imager IBIS (Ubertini et al. 2003), which allows high angular resolution imaging over a large field of view ($29^{\circ} \times 29^{\circ}$) in the energy range 15 keV–10 MeV, the spectrometer SPI (Vedrenne et al. 2003; 20 keV–8 MeV) and the X-ray monitor JEM-X (Lund et al. 2003; 3–35 keV). IBIS is composed of a low-energy CdTe detector (ISGRI; Lebrun et al. 2003), sensitive in the energy range from 15 keV to 1 MeV, and a CsI detector (PICsIT; Labanti et al. 2003), designed for optimal performance at 511 keV, and sensitive in the 175 keV–10 MeV energy range.

We analyzed all public and Core Program IBIS observations pointed within 10° of the source. This resulted in 909 individual pointings (Science Windows, SWs) performed between March 2003 and November 2005. All the data have been processed using version 5.1 of the OSA *INTEGRAL* analysis software, and analysed with the corresponding rebinned response matrices.

A mosaic image of all 909 pointings revealed the sources that were active in this field during the period of interest. The list of all the detected sources (detection significance $>5\sigma$) was then used in the spectral step, where spectra for all the detected

[★] Based on observations with *INTEGRAL*, an ESA project with instruments and the science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), Czech Republic and Poland, and with the participation of Russia and the USA.

Table 1. Summary of all *INTEGRAL* observations of 4U 1850–087 analysed here. The observations have been grouped together in 4 data-sets for brevity. The 3rd and 4th columns list the Start and Stop Time of the four groups of observations, the fifth column reports the number of SWs in each data-set. Data-set n. 4 includes Core Program data still not publicly available.

Data set	Temporal window	Start Time (MJD)	End Time (MJD)	Num of SWs
1	Mar. 2003–May 2003	52 708.7	52 772.0	258
2	Sep. 2003–Nov. 2003	52 910.9	52 963.4	210
3	Mar. 2004–May 2004	53 075.1	53 126.5	84
4	Aug. 2004–Nov. 2005	53 238.1	53 684.0	357




Fig. 1. RXTE/ASM lightcurve (day average) of 4U 1850–087. The small solid circles in the upper region show the times of the *INTEGRAL* observations analysed here.

sources in the field of view were simultaneously extracted with the standard spectral extraction method.

4U 1850–087 is a relatively faint source, therefore a meaningful spectral analysis needs to be performed adding together several observations. For this reason, we grouped the pointings in four data-sets (listed in Table 1) from which we extracted four different IBIS/ISGRI spectra.

The 2–10 keV flux of 4U 1850–087 shows a brightening event peaking on 25 February 2004, and lasting ~ 10 days, as visible from the ASM/RXTE lightcurve reported in Fig. 1. Only part of its decay is covered by the *INTEGRAL* observations of data-set 3.

When we are averaging the spectra of 4U 1850–087 using data that span over different mission times, we use a systematic of 5% in the spectral fit process, to take into account for modifications in the instrument evolution (Lubinski et al. 2005).

The best-fit spectral results, using a single power-law are reported in Table 2 for the four data-sets. The fits display a high χ^2 (with a null hypothesis probability in the range 0.1–2.4%) which can be explained by the spectral distortions present when extracting IBIS/ISGRI spectra spanning very different mission times (Lubinski et al. 2005), especially for faint sources, and by possible intrinsic source spectral changes on smaller timescales, which cannot be a priori excluded. Nevertheless, the global spectral shape is rather constant on timescales of months-years.

We tried also to extract JEM-X spectra, but the smaller field of view, the faintness of the source, together with some instrumental issues (e.g. JEM-X switched-off in several occasions), severely reduced the number of useful observations. Thus data could be used only for the integrated spectrum covering the data-set 3 and part of the data-set 4. The JEM-X spectra correspond to

Table 2. Results of the spectral analysis of the four temporal-selected IBIS/ISGRI spectra. Data-set numbers are the same as in Table 1. The 2nd column reports the net exposure time of each spectrum. The 3rd column lists the IBIS/ISGRI average rates in the energy range 20–100 keV. Fluxes are in units of 10^{-10} erg cm^{-2} s^{-1} (20–100 keV).

Data-set	Exp. (ks)	IBIS/ISGRI rate (s^{-1})	Photon Index	Flux	$\chi^2/\text{d.o.f.}$
1	372	1.25 ± 0.05	2.5 ± 0.2	0.8	21.6/8
2	383	2.36 ± 0.05	2.1 ± 0.1	1.8	18.8/7
3	114	1.42 ± 0.09	2.2 ± 0.3	1.1	11.6/7
4	490	1.59 ± 0.05	2.2 ± 0.1	1.2	16.1/7




Fig. 2. 4U 1850–087 combined JEM-X and IBIS/ISGRI simultaneous spectrum, fit with a single power-law (photon index of 2.2 ± 0.2). In the lower panel, the residuals in units of standard deviations are shown.

76.7 ks net exposure time for JEM-X1 and 55.5 ks for JEM-X2, with a source count rate of 0.23 ± 0.02 and 0.22 ± 0.03 , respectively in the two JEM-X units (5–20 keV).

The combined JEM-X plus IBIS/ISGRI simultaneous spectrum (taken from the data-set 3) is shown in Fig. 2, fitted with an absorbed power-law, with absorbing column density fixed at 2.5×10^{21} cm^{-2} (the expected interstellar absorption towards the globular cluster NGC 6712). The fit is a good deconvolution of the spectrum ($\chi^2/\text{d.o.f.} = 27.9/23$) and results in a photon index of 2.2 ± 0.2 and in a 5–100 keV flux corrected for the absorption of $(2.4 \pm 0.3) \times 10^{-10}$ erg cm^{-2} s^{-1} (based on the IBIS/ISGRI response matrix). Other simple models, like bremsstrahlung or a cut-off power-law, never result in a better fit.

4U 1850–087 has been observed with *XMM-Newton* on 27 September 2003 (see Sidoli et al. 2005), i.e. within the period covered by *INTEGRAL* data-set 2, with the main aim to study the low energy absorption intrinsic to the source. Indeed the *XMM-Newton* spectrum below ~ 2 keV is complex and we refer to Sidoli et al. 2005 for its detailed description (and for the *XMM-Newton* data reduction). EPIC PN operated in Small Window mode, and with a net exposure time of 8.1 ks. A combined spectral analysis EPIC/IBIS interestingly allows us to extend the spectral study of this source in the soft X-rays, below 5 keV, and to significantly refine the spectral parameters. During the *XMM-Newton* observation, only EPIC PN data did not suffer from pile-up problems, thus we use this spectrum (1.7–12 keV, see Sidoli et al. 2005 for the details) to perform a broad-band analysis of the quasi-simultaneous EPIC-IBIS observations.

Among the different models considered by Sidoli et al. (2005) when fitting the *XMM-Newton* spectrum, we tried their best-fit, a disk-blackbody emission together with a power-law.

Table 3. Results of the spectral analysis of the broad-band quasi-simultaneous spectrum (EPIC/PN–IBIS/ISGRI). Fluxes (2–100 keV) are corrected for the absorption, are in units of 10^{-10} erg cm $^{-2}$ s $^{-1}$, and are based on the EPIC pn response matrix. The assumed distance is 6.8 kpc (Harris et al. 1996). The COMPTT model assumes a spherical geometry. i is the inclination angle of the disk.

Model	N_{H} (10^{22} cm $^{-2}$)	Parameters	Flux	$\chi^2/\text{d.o.f.}$
DISKBB+POW	0.4 ± 0.2	$\Gamma = 2.07^{+0.07}_{-0.15}$ $kT_{\text{in}} = 0.8 \pm 0.1$ keV $r_{\text{in}} \times (\cos(i))^{0.5} = 1.7^{+1.1}_{-0.4}$ km	2.8 ± 0.1	147.4/203
DISKBB+COMPTT	0.25 fixed	$kT_{\text{e}} > 40$ keV $kT_0 = 0.67 \pm 0.05$ keV $\tau = 2.2 \pm 0.5$ $kT_{\text{in}} = kT_0$ $r_{\text{in}} \times (\cos(i))^{0.5} = 4 \pm 1$ km	2.9 ± 0.1	156.3/203




Fig. 3. Broad-band 4U 1850–087 EPIC/PN spectrum (obtained in September 2003) combined with the quasi-simultaneous IBIS/ISGRI (dataset 2). On the left, the count spectra are displayed, together with residuals in units of standard deviations, when fitted with a disk-blackbody plus a power-law. On the right, the corresponding 1.7–100 keV energy spectrum is shown.

This gives already a very nice result ($\chi^2/\text{d.o.f.} = 147.4/203$), with the spectral parameters reported in Table 3 and the spectrum shown in Fig. 3. The normalization constant for the IBIS/ISGRI spectrum relative to PN (constant factor fixed at 1) was $1.8^{+0.2}_{-0.4}$. We note that this normalization factor between the two instruments is in agreement with that reported for the Crab spectrum (Kirsch et al. 2005). The broad band flux (2–100 keV, corrected for the absorption) is 2.8×10^{-10} erg cm $^{-2}$ s $^{-1}$ (based on the EPIC PN response matrix). Fixing the column density to the best-fit (5.7×10^{21} cm $^{-2}$) found by Sidoli et al. (2005), does not change the spectral results.

In order to obtain physical information, we then adopted the best-fit model used for the *BeppoSAX* 4U 1850–087 spectrum (Sidoli et al. 2001), a disk-blackbody plus a Comptonized emission (COMPTT model in XSPEC; Titarchuk 1994). Since the data extend only down to 1.7 keV, we linked the temperature of the seed photons, kT_0 , to the inner disk temperature, kT_{in} . This is also justified by the empirical finding that in ultra-compact binaries the two temperatures are similar, within the uncertainties (Sidoli et al. 2001). We obtain a good fit ($\chi^2/\text{d.o.f.} = 156.3/203$), with the parameters reported in Table 3. To obtain better constraints to the spectral parameters, in this case the absorbing column density has been fixed to the NGC 6712 interstellar absorption value. With this model, the normalization constant of the IBIS/ISGRI spectrum with respect to EPIC PN lies in the range 1.5–1.9, again in agreement with the standard value (Kirsch et al. 2005). Adopting a cut-off power-law instead of a power-law in the two-component model ($\chi^2/\text{d.o.f.} = 172.2/202$), the cut-off energy resulted in $E_{\text{c}} > 110$ keV (90% confidence level), with a best-fit photon index of 1.9 ± 0.1 .

3. Discussion and conclusions

Previously, the source broad-band spectrum was observed only with *BeppoSAX* in April 1997. At that time, the PDS non-imaging instrument, covering the range from 20 to 200 keV, detected the source only up to 50 keV (Sidoli et al. 2001).

We report here for the first time the discovery of hard (50–100 keV) X-ray emission from the X-ray burster 4U 1850–087 and a long-term study of its X-ray spectral behaviour. It is now possible to directly measure the high energy luminosities of this source: they are $\sim 1.3 \times 10^{36}$ erg s $^{-1}$ and 7×10^{35} erg s $^{-1}$ respectively in the 1–20 keV and 20–200 keV energy ranges (at 6.8 kpc). This implies that 4U 1850–087 falls into the so-called “burster box” (Barret et al. 2000).

In order to allow a proper comparison with the *INTEGRAL* spectrum we re-analysed the *BeppoSAX* observation (April 1997; refer to Sidoli et al. 2001 for the details of the data reduction) adopting the same model used here, a disk-blackbody emission together with a power-law. This resulted in the following parameters ($\chi^2/\text{d.o.f.} = 170.4/158$): an absorbing column density of $(0.46 \pm 0.03) \times 10^{22}$ cm $^{-2}$, an inner disk blackbody temperature, kT_{in} , of 0.66 ± 0.03 keV, and a powerlaw photon index of 1.96 ± 0.06 . Thus, there is a good agreement with the *INTEGRAL* spectroscopy.

The new observations with *INTEGRAL* allow us to put much more stringent limits on the presence of a high energy cut-off. In Fig. 4 we compare the confidence contour levels for the high energy cut-off and the power-law photon index, obtained with the EPIC–IBIS joint spectrum (solid contours) and with *BeppoSAX* (dashed contours). During the *BeppoSAX* observation a lower




Fig. 4. Comparison of the confidence contour levels for the high energy cutoff (in units of keV): solid contours have been derived analysing EPIC–IBIS joint spectrum, while the dashed contours mark the *BeppoSAX* results.




Fig. 5. Comparison of the *INTEGRAL* spectrum of the source 4U 1850–087 (data-set 2, the same displayed in Fig. 3), with a spectrum of a typical Z-source, GX 349+2 (data from Paizis et al. 2006).

limit to the high energy cut-off could be placed at $E_c > 60$ keV (90% level), while now with *INTEGRAL* we can interestingly shift it towards much higher energies, with $E_c > 100$ keV. This allows us to include 4U 1850–087 among the few tens of low-luminosity low-mass X-ray binaries (LMXRBS) for which the spectrum has been observed up to 100 keV (Barret et al. 2000; Di Salvo & Stella 2002; Bazzano et al. 2006). More interestingly, 4U 1850–087 is now among the hardest type I X-ray bursters in our Galaxy.

The *INTEGRAL* spectrum displays properties which fits well into the classification of low luminosity ($\sim 0.01 L_{\text{Edd}}$) weakly magnetized neutron stars with “hard spectra” (Barret et al. 2001): a broad-band spectrum extending up to 100 keV,

well described by a soft disk-blackbody emission together with a hard Comptonized component, with a optical depth of the Comptonizing corona of ~ 2 , seed photons temperature below 1 keV, a high electron temperature, above 40 keV. A few LMXRBs appear to spend most of the time in this “hard state” (Di Salvo & Stella 2002), like 4U 0614+091 (Piraino et al. 1999), which also displays spectral parameters similar to 4U 1850–087, with a photon index in the range 2.3–2.4. Also for 4U 1850–087 the four different *INTEGRAL* spectra suggest that the source spend most of its lifetime in this spectral state (on timescales of months or years, if compared with *BeppoSAX*).

The very hard emission from 4U 1850–087 is clearly evident also when compared (see Fig. 5) to a typical spectrum of a Z-source, GX 349+2, reported in an *INTEGRAL* study of the spectral behaviour of the persistent LMXRBs hosting a neutron star (Paizis et al. 2006).

Acknowledgements. We acknowledge the Italian Space Agency financial and programmatic support via contract I/R/046/04.

References

- Anderson, S. F., Margon, B., Deutsch, E. W., & Downes, R. A. 1993, *AJ*, 106, 1049
- Barret, D. 2001, in *Adv. Space Res.*, 28, 307
- Barret, D., Olive, J. F., Boirin, L., et al. 2000, *ApJ*, 533, 329
- Bazzano, A., et al. 2006, Proceedings of the conference, A life with stars, Amsterdam, 22–26 August 2005, *New Astronomy Reviews*, ed. Kaper, Van der Klis, & Wijers, in press
- Di Salvo, T., & Stella, L. 2002, proceedings of the XXII Moriond Astrophysics Meeting, The Gamma-Ray Universe (Les Arcs, March 9–16, 2002), ed. A. Goldwurm, D. Neumann, & J. Tran Thanh Van (Vietnam: The Gioi Publishers) [arXiv:astro-ph/0207219]
- Harris, W. E. 1996, *AJ*, 112, 1487
- Homer, L., Charles, P. A., Naylor, T., et al. 1996, *MNRAS*, 282, L37
- Juett, A. M., & Chakrabarty, D. 2005, *ApJ*, 627, 926
- Juett, A. M., Psaltis, D., & Chakrabarty, D. 2001, *ApJ*, 560, L59
- Kirsch, M. G. F., Ubruel, U. G., Burrows, D., et al. 2005, *SPIE*, 5898, 22 [arXiv:astro-ph/0508235]
- Lubinski, P., Dubath, P., & Paltani, S. 2005, available at http://isdc.unige.ch/Soft/download/osa/osa_doc/prod/osa_cross_cal-1.0.pdf
- Labanti, C., Di Cocco, G., Ferro, G., et al. 2003, *A&A*, 411, L149
- Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141
- Lund, N., Budtz-Jørgensen, C., Westergaard, et al. 2003, *A&A*, 411, L231
- Merloni, A., Fabian, A. C., & Ross, R. R. 2000, *MNRAS*, 313, 193
- Paltrinieri, B., Ferraro, F. R., Paresce, F., et al. 2001, *AJ*, 121, 3114
- Paizis, A., Farinelli, R., Titarchuk, L., et al. 2006, *A&A*, 559, 187
- Parmar, A. N., Stella, L., & Giommi, P. 1989, *A&A*, 222, 96
- Piraino, S., Santangelo, A., Ford, E. C., et al., 1999, *A&A*, 349, L77
- Sidoli, L., Parmar, A. N., Oosterbroek, T., et al. 2001, *A&A*, 368, 451
- Sidoli, L., La Palombara, N., Oosterbroek, T., & Parmar, A. N. 2005, *A&A*, 443, 223
- Swank, J. H., Becker, R. H., Pravdo, S. H., et al. 1976, *IAU Circ.*, 3010
- Titarchuk, L. 1994, *ApJ*, 434, 570
- Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131
- Vedrenne, G., Roques, J.-P., Schönfelder, V., et al. 2003, *A&A*, 411, L63
- Verbunt, F., Bunk, W., Hasinger, G., & Johnston, H. M. 1995, *A&A*, 298, 21