

## *Erratum*

# Effective collision strengths for electron impact excitation of C II<sup>★</sup>

N. J. Wilson, K. L. Bell, and C. E. Hudson

Department of Applied Mathematics and Theoretical Physics, The Queen's University of Belfast, Belfast BT7 1NN, Northern Ireland  
e-mail: c.hudson@am.qub.ac.uk

A&A 432, 731-736 (2005), DOI: 10.1051/0004-6361:20041855

### ABSTRACT

We correct data published in the above paper as Table 2 and replot Figs. 2 and 4.

**Key words.** atomic processes – line: formation – methods: analytical – errata, addenda

## 1. Introduction

In the paper “Effective collision strengths for electron impact excitation of C II” published in A&A 432, 731–736 (2005), an error was made in processing the collision strengths to obtain the Maxwellian-averaged effective collision strengths. Table 2 (available at the CDS) containing the non-zero effective collision strengths has now been updated with the corrected data.

The figures presented here are the updated versions of Figs. 2 and 4. The effective collision strengths are plotted as a function of  $\log(T)$  where  $T$  is the electron temperature in kelvin, and are compared with data from Blum & Pradhan (1992). The upper figure corresponds to the  $2s2p^2\ ^4P_{1/2}-2s2p^2\ ^2D_{5/2}$  transition. The lower figure is the  $2s2p^2\ ^2D_{3/2}-2s^23p\ ^2P_{3/2}^o$  transition. In both these plots the dashed line represents the Blum & Pradhan (1992) data and the solid lines are the current updated values.

## 2. Conclusions

Our corrected data shows improved agreement with the work of Blum & Pradhan for the two transitions here noted. However, we still find discrepancies for other transitions.

## References

Blum, R. D., & Pradhan, A. K. 1992, ApJS, 80, 425




Fig. 2.




Fig. 4.

\* Table 2 is only available in electronic form at the CDS via anonymous ftp to [cdsarc.u-strasbg.fr](http://cdsarc.u-strasbg.fr) (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/461/765>