

“Expansion” around the vacuum: how far can we go from Λ ?

J. S. Alcaniz¹ and H. Štefancić^{2,*}

¹ Observatório Nacional, Rua Gal. José Cristino 77, 20921-400 Rio de Janeiro - RJ, Brasil
e-mail: alcaniz@on.br

² Departament d'Estructura i Constituents de la Matèria, Universitat de Barcelona, Av. Diagonal 647, 08028 Barcelona, Catalonia, Spain
e-mail: stefancic@ecm.ub.es

Received 22 May 2006 / Accepted 20 September 2006

ABSTRACT

The cosmological constant (Λ), i.e., the energy density stored in the true vacuum state of all existing fields in the Universe, is the simplest and the most natural possibility for describing the current cosmic acceleration. However, despite its observational successes, such a possibility exacerbates the well known Λ problem, requiring a natural explanation for its small, but nonzero, value. In this paper we discuss how different our Universe may be from the Λ CDM model by studying observational aspects of a kind of “expansion” around the vacuum given by the equation of state (EOS) $p_d = -\rho_d - A\rho_d^q$. In different parameter regimes, such a parametrization is capable of describing both quintessence-like and phantom-like dark energy, transient acceleration, and various (non)singular possibilities for the final destiny of the Universe, including singularities at finite values of the scale factor, the so-called “Big Rip”, as well as sudden future singularities. By using some of the most recent cosmological observations we show that, if the functional form of the dark energy EOS has additional parameters, very little can be said about their values from the current observational results, which postpones a definitive answer to the question posed above until the arrival of more precise observational data

Key words. cosmology: theory – cosmology: miscellaneous – distance scale

1. Introduction

The understanding of the global evolution of the observationally amenable universe, mathematically encoded in the dynamics of its scale factor a , is of utmost importance in explaining practically all cosmological phenomena. One of the most intriguing aspects of this evolution is the recently established late-time transition from a decelerated to an accelerating regime of the expansion of the Universe. Evidence confirming the low-redshift onset of the accelerating expansion phase comes directly from current type Ia supernovae (SNIa) measurements (Riess et al. 1998, 2004; Perlmutter et al. 1999), and is indirectly supported by various other cosmological observations such as the measurements of the cosmic microwave background (CMB) anisotropies (de Bernardis et al. 2000; Spergel et al. 2003, 2006), and the current large scale structure (LSS) data (Tegmark et al. 2004).

Although the actual nature of the mechanism behind the current cosmic acceleration, as well as its link with fundamental interactions, remains as a completely open question, there are many, often very ingenious, attempts to model the current stage of the cosmic evolution. Some of the most important questions that these models try to answer are:

1. *What is the mechanism causing the accelerating expansion of the universe?*

The idea of a late time accelerating universe is usually associated with unknown physical processes involving either new fields in high energy physics or modifications of gravity on very large scales. In following the former route, the predominant concept nowadays is the existence of a new

component with sufficiently negative pressure, named *dark energy* (Sahni & Starobinsky 2000; Peebles & Ratra 2003; Padmanabhan 2003), which is fully characterized by its equation of state (EOS), $w(t) = p_d(t)/\rho_d(t)$, where $p_d(t)$ and $\rho_d(t)$ are, respectively, the dark component pressure and energy density. The value $w = -1$ characterizes the *vacuum energy* (Λ), which is conceptually the simplest model (Λ CDM). Although these scenarios constitute a kind of benchmark model in the analysis of observational data, the large discrepancy between the theoretically predicted and “observed” values of Λ has incited the study of dynamical dark energy (DDE) models.

In this case, Λ is treated as a dynamical quantity, whereas its constant EOS, $w = -1$, is preserved. This includes, among others, models based on renormalization group running Λ (Shapiro & Sola 2000, 2002; Babic et al. 2002; Shapiro et al. 2005; Sola & Štefancić 2005) and vacuum decay (Özer & Taha 1986; Bertolami 1986; Cunha et al. 2002; Alcaniz & Lima 2005). Other DDE models, in which the potential energy density associated with a dynamical scalar field (ϕ) dominates the dynamics of the low-redshift Universe, have also been extensively discussed in the current literature (see, e.g., Ratra & Peebles 1988; Wetterich 1988; Caldwell et al. 1998). The EOS parameter in this class of models is necessarily a function of time and may take values > -1 [*quintessence*] or < -1 [*phantom*] (Caldwell 2002; Faraoni 2002; Alcaniz 2004). In this regard, it is still worth mentioning that some recent observational analyses (Melchiorri et al. 2003), implying that a transition from $w > -1$ to $w < -1$ might have happened at low redshifts, have also instigated the development of models with a Λ boundary

* On leave of absence from The Theoretical Physics Division, Rudjer Bošković Institute, Zagreb, Croatia.

crossing (Nojiri & Odintsov 2005; Feng et al. 2005; Stefancic 2005).

The second route to the acceleration problem involves modifications of gravity at cosmological scales (without dark energy) (Randall & Sundrum 1999; Dvali et al. 2000; Maia et al. 2005). An interesting example of this approach is the so-called braneworld models, which offer an intriguing mechanism for the acceleration of the universe and have been successfully tested by a number of observational analyses (Alcaniz 2002; Jain et al. 2002; Lue 2003).

2. What is the duration of the accelerating expansion phase?

Although the transition from an initially decelerated to a late-time accelerating expansion is becoming observationally well established, the acceleration redshift z_{acc} , as well as the duration of the accelerating phase, depends crucially on the cosmological scenario (see, e.g., Bassett et al. 2004). Several models imply an eternal acceleration (e.g. Λ CDM model) or even an accelerating expansion until the onset of a cosmic singularity (e.g., phantom models with constant EOS). However, an eternally accelerating universe seems not agree with string/M-theory since it is endowed with a cosmological event horizon, which prevents the construction of a conventional S-matrix describing particle interactions within the framework of these theories (Fischler et al. 2001; see also Ellis et al. 2001). In that sense, cosmological models that predict the possibility of a transient acceleration phenomenon are of special interest.

3. What is the fate of the presently accelerating Universe?

As is well-known, information on the fate of the Universe is encoded in the form of its past expansion and fundamentally depends on the nature of its dominant energy component. As an example, the so-called phantom energy models ($w < -1$) predict a very dramatic fate for the Universe, i.e., a universe that ends with a future singularity, the so-called *Big Rip* (Caldwell et al. 2003; Nesseris & Perivolaropoulos 2004a). As shown by Lima & Alcaniz (2004), the thermodynamic fate of a dark-energy dominated universe may also be considerably modified, with the possibility of an increasingly hot expanding universe.

In this paper we aim at addressing some of the above questions in the context of a dark energy model characterized by the following EOS (Stefancic 2005; Nojiri & Odintsov 2004)

$$p_d = -\rho_d - A\rho_d^\alpha. \quad (1)$$

In this simple framework, by using the current observational data to place limits on the parameters of (1), one can simultaneously test (i) whether the nature of the dark energy is quintessence-like ($A < 0$) or phantom-like ($A > 0$); (ii) if the acceleration is eternal or transient; and (iii) which form of future asymptotic (non)singular behavior is implied observationally. The organization of the paper is the following. Section 2 presents a theoretical analysis and discusses the most interesting phenomenological implications of the model. Section 3 is devoted to a comparison of the model against the observational data. We end this paper by summarizing our main results in the concluding section.

2. The model

The EOS (1) can be understood as a sort of “expansion” around the vacuum EOS. The *a priori* motivation for the study of such a dark energy EOS is twofold. First, the Λ CDM cosmology, characterized by the “true” cosmological constant, is the standard in analysis of the observational data. This means

that the measurement of a possible deviation of the dark energy EOS from the vacuum EOS is presently one of the highest priorities in Cosmology. The model (1) provides a suitable framework for such a study. Second, from the theoretical side, the model (1) represents a dynamical dark energy model with a time-dependent EOS parameter, which can describe both quintessence and phantom types of dark energy behavior. A detailed analysis of the dynamics of the universe with a dark energy component (1) was made by Stefancic (2005). In what follows, we summarize the main results of this analysis.

The conservation of the energy-momentum tensor for the dark energy model (1) yields the scaling of dark energy density with the scale factor of the universe, i.e.,

$$\rho_d = \rho_{d,0} \left(1 + 3\tilde{A}(1 - \alpha) \ln a\right)^{1/(1-\alpha)}, \quad (2)$$

where we set $a_0 = 1$ and the abbreviation $\tilde{A} = A\rho_{d,0}^{\alpha-1}$ was introduced. The parameter $w = p_d/\rho_d$ of the EOS (1) is given by

$$w = -1 - \frac{\tilde{A}}{1 + 3\tilde{A}(1 - \alpha) \ln a}. \quad (3)$$

This equation reveals that, for different signs of \tilde{A} , we may expect different behavior of the dark energy component, which presently generates the acceleration of the Universe. For example, for $\tilde{A} > 0$, w remains below -1 during the entire evolution of the universe, whereas for $\tilde{A} < 0$ the EOS parameter always remains above the Λ barrier ($w = -1$) and may become even larger than $-1/3$, which means that the dark energy component can change its action from accelerating to decelerating. The expression for $p_d = w\rho_d$ can be obtained directly from Eqs. (2) and (3) in a straightforward manner.

Here we assume that the universe contains two components, namely, the dark energy component and the nonrelativistic matter component. Thus, the Friedmann equation is given by

$$\left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2} = \frac{8\pi G}{3}(\rho_d + \rho_m). \quad (4)$$

In the case where the dark energy density dominates over the energy density of nonrelativistic matter and the spatial curvature term k/a^2 , Eq. (4) can be solved analytically to obtain the asymptotic time-dependence of the scale factor a . For $\alpha \neq 1/2$ the time-dependence of the scale factor is given by

$$\begin{aligned} & \left(1 + 3\tilde{A}(1 - \alpha) \ln a_1\right)^{(1-2\alpha)/(2(1-\alpha))} \\ & - \left(1 + 3\tilde{A}(1 - \alpha) \ln a_2\right)^{(1-2\alpha)/(2(1-\alpha))} = \\ & \frac{3}{2}\tilde{A}(1 - 2\alpha)\Omega_{d,0}^{1/2}H_0(t_1 - t_2), \end{aligned} \quad (5)$$

whereas for $\alpha = 1/2$ one obtains

$$\ln \frac{1 + \frac{3}{2}\tilde{A} \ln a_1}{1 + \frac{3}{2}\tilde{A} \ln a_2} = \frac{3}{2}\tilde{A}\Omega_{d,0}^{1/2}H_0(t_1 - t_2). \quad (6)$$

In the expressions above, we have introduced the notation $\Omega_{d,0} = \rho_{d,0}/\rho_{c,0}$, where $\rho_{c,0} = 3H_0^2/(8\pi G)$ is the present critical energy density. Throughout this paper, the subscript 0 refers to the present epoch of the evolution of the Universe.

Using Eqs. (2), (5), and (6) it is possible to determine the characteristic values of the model parameters which separate qualitatively different behaviors of the model. The characteristic value of the parameter \tilde{A} is $\tilde{A} = 0$. Thus, for $\tilde{A} > 0$ the

model has a phantom-like behavior, for $\tilde{A} < 0$ the model behaves as quintessence, while for $\tilde{A} = 0$ the model is equivalent to the Λ CDM scenario. The characteristic values for the second parameter α are $\alpha = 1$ and $\alpha = 1/2$. The behavior of the model in the intervals delimited by these two values is described below. The description of the rich variety of phenomena realized for different values of the parameters \tilde{A} and α is divided into cases $\tilde{A} > 0$ and $\tilde{A} < 0$.

2.1. $\tilde{A} > 0$

1. For $\alpha > 1$ the dark energy density (ρ_d) and pressure (p_d) diverge at a *finite value of the scale factor*, which is also reached in a finite time interval from today. This type of singular ending of the universe, first introduced by Štefančić (2005) and Nojiri & Odintsov (2004), represents a kind of singularity that is stronger than the so-called “Big Rip” singularity¹.
2. The case $\alpha = 1$ corresponds to the well studied phantom energy with a constant EOS parameter, which leads to the well known “Big Rip” event (Caldwell et al. 2003; Nesseris & Perivolaropoulos 2004a).
3. In the interval $1/2 < \alpha < 1$ the scale factor a diverges at a finite time, as do the dark energy density ρ_d and pressure p_d . The value of the EOS parameter w remains bounded. The singular fate of the universe in this parameter range is qualitatively similar to the “Big Rip” singularity (Caldwell et al. 2003; Nesseris & Perivolaropoulos 2004a).
4. For $\alpha \leq 1/2$ the future expansion is nonsingular and the scale factor a reaches infinity in a infinite future.

2.2. $\tilde{A} < 0$

1. The case $\alpha > 1$ is characterized by the nonsingular future expansion. The dark energy density in this case is a decreasing function of the scale factor (the cosmic time) and vanishes for $t \rightarrow \infty$.
2. The case $\alpha = 1$ corresponds to the case of quintessence-like dark energy with a constant EOS parameter $w = -1 - \tilde{A}$.
3. For $\alpha < 1$ the dark energy density reaches 0 at a finite value of the scale factor $a_{\text{NULL}} > a_0$ (and at a finite time). At this scale-factor value the Hubble parameter, H and the total energy density are constant, whereas the parameter of dark energy EOS diverges, $w \rightarrow +\infty$. The behavior of the dark energy pressure and the acceleration of the universe differs for cases $0 < \alpha < 1$, $\alpha = 0$, and $\alpha < 0$.
 - 3.1. In the case where $0 < \alpha < 1$, the dark energy pressure vanishes at a_{NULL} . For $a > a_{\text{NULL}}$ the dark energy pressure can no longer be properly defined by (1). Since H is finite at a_{NULL} , the question is how the expansion is continued. One possibility is that for $a > a_{\text{NULL}}$ the universe expands with $\rho_d = 0$ and $p_d = 0$, which also satisfies (1).
 - 3.2. For $\alpha = 0$ the dark energy density changes its sign at a_{NULL} and the dark energy pressure is positive for $a > a_{\text{NULL}}$. The expansion continues until the Hubble parameter vanishes at some finite $a_{\text{STOP}} > a_{\text{NULL}}$.
 - 3.3. For $\alpha < 0$ the dark energy pressure diverges at a_{NULL} , $p_d \rightarrow +\infty$, and $\ddot{a} \rightarrow -\infty$. This is a concrete realization of a sudden future singularity recently introduced by Barrow (Barrow 2004).

¹ For a detailed classification of singularities in phantom cosmologies, see, e.g., Nojiri et al. (2005).




Fig. 1. The deceleration parameter as a function of redshift for some selected combinations of the pair (\tilde{A}, α) and $\Omega_{m,0} = 0.3$. The horizontal dashed line divides decelerating and accelerating universes. Note that a number of different possibilities for the recent past and future of the Universe can be achieved from parametrization (1). The standard Λ CDM model (thick line) is also shown for the sake of comparison.

The behavior of the model for $\tilde{A} < 0$ and $\alpha < 1$ is characterized by the finite duration of the phase of the accelerated expansion of the universe (see Fig. 1). The transition from the decelerated to the accelerated expansion of the universe is the result of the interplay of the evolution of the dark energy density and the energy density of nonrelativistic matter. The exit from the accelerated phase is, however, the consequence of the intrinsic dynamics of the dark energy model (1) in this parameter regime. Finally, a scenario in which the phase of accelerated expansion starts and ends in the recent past is also allowed in this parameter regime. This nonstandard possibility, closely related to condition (8), is also discussed in our observational analysis.

3. Observational aspects

The description of the model (1) discussed in the previous section clearly shows that it comprises a multitude of cosmological solutions. In a model with such a wealth of different possibilities constraints on the parameter space arising from current observational data are likely to rule out many of the possible scenarios for the evolution/fate of the Universe. This section investigates such observational constraints by studying (i) the influence of parameters \tilde{A} and α on the epoch of the deceleration/acceleration (D/A) transition, as well as on the duration of the accelerating phase and (ii) by placing observational bounds on the parametric space $\tilde{A} - \alpha$ from a statistical analysis involving a large set of cosmological observations.

3.1. D/A transition

An important characteristic of the current SNe Ia data is the evidence of a possible D/A transition at $z_* \sim 0.5$ (Riess et al. 2004). If confirmed by the upcoming data, the precise value of z_* will become an important observational constraint for decisively deciding between competing cosmological models. In order to study the bounds on parameters \tilde{A} and α from the current




Fig. 2. The transition redshift z_* as a function of the parameter \tilde{A} for values of α lying in the interval $[\frac{1}{2}, 2]$. Horizontal lines stand for the 1σ interval $0.33 \leq z_* \leq 0.59$, as provided by current SNe Ia observations (Riess et al. 2004). Note that our current standard model, i.e., the Λ CDM scenario, predicts a transition epoch $\approx 2\sigma$ off from the central value obtained in Riess et al. (2004).

estimates of z_* we first derive the deceleration parameter as a function of the scale factor a , i.e.,

$$q(a) = -1 + \frac{3\Omega_{m,0}a^{-3} - \Omega_{d,0}\tilde{A}F(a, \alpha, \tilde{A})^{\alpha/(1-\alpha)}}{2\Omega_{m,0}a^{-3} + \Omega_{d,0}F(a, \alpha, \tilde{A})^{1/(1-\alpha)}}, \quad (7)$$

where $\Omega_{m,0} = \rho_{m,0}/\rho_{c,0}$ and $F(a, \alpha, \tilde{A}) = 1 + 3\tilde{A}(1-\alpha)\ln(a)$. As can be checked, an immediate constraint on the model parameters arises by combining Eq. (7) with the fact that the universe is currently accelerated [$q(a = a_0 = 1) < 0$], i.e.,

$$\tilde{A} > \frac{\Omega_{m,0} - 2/3}{1 - \Omega_{m,0}}. \quad (8)$$

Note that the above expression does not depend on parameter α , a feature of the model that will be encountered in the statistical analysis performed in the next section. This means that for values of $\Omega_{m,0} \approx 0.3$, as indicated by dynamical estimates at scales up to about $2 h^{-1}$ Mpc (Calberg et al. 1996) [and also confirmed by current CMB measurements (Spergel et al. 2004)], we find that parameter \tilde{A} is restricted to the interval $\tilde{A} > -0.52$. Examples of the behavior of the deceleration parameter as a function of redshift are shown in Fig. 1 for selected values of \tilde{A} and α and $\Omega_{m,0} = 0.3$. As expected from Eq. (9), irrespective of the value of α , models with $\tilde{A} > (<) -0.52$ are currently accelerated (decelerated), while for the critical case $\tilde{A} = -0.52$, the Universe expands presently with a constant Hubble flow ($q_0 = 0$) and constitutes new examples of coasting cosmologies (Kolb 1989). As mentioned earlier, note that the possibility of a presently decelerated universe that experienced a recent accelerated phase is also contained in parametrization (1). In the particular example shown in Fig. 1 ($\tilde{A} = -0.58$ and $\alpha = 0.3$) the Universe would have entered an accelerated regime at $z \approx 0.55$, then remained accelerated for $2.04 h^{-1}$ Gyr, and have switched again to a new decelerated period at $z \approx 0.17$.

Another interesting behavior that is also achieved from some combinations of the pair (\tilde{A}, α) has to do with the intriguing

possibility that the current cosmic acceleration may not be a lasting feature. As can also be seen from Fig. 1, some scenarios of type (1) predict a universe that was decelerated in the past, began to accelerate at $z_* \approx 0.5$, is currently accelerated, but will eventually decelerate in the future. Such behavior, according to some authors (Fischler et al. 2001), seems to fully agree with the requirements of string or M-theory, in that eternally accelerating universes are endowed with a cosmological event horizon that prevents the construction of a conventional S-matrix describing particle interactions within these frameworks². A typical example of an eternally accelerating universe, i.e., the Λ CDM model, is also shown in Fig. 1 for the sake of comparison.

Figure 2 shows the transition redshift z_* as a function of parameter \tilde{A} . Note that the behavior of z_* can be easily obtained from Eq. (7) or, more specifically, from the expression

$$\frac{1}{2}\Omega_{m,0}(1+z_*)^3 - \Omega_{d,0}\mathcal{F}^{\alpha/(1-\alpha)}\left[\frac{3}{2}\tilde{A} + \mathcal{F}\right] = 0, \quad (9)$$

where $\mathcal{F} = 1 - 3\tilde{A}(1-\alpha)\ln(1+z_*)$. It also shows the interval $0.33 \leq z_* \leq 0.59$, which corresponds to $\pm 1\sigma$ of the estimate for the transition redshift given by Riess et al. (2004). For values of α lying in the interval $[\frac{1}{2}, 2]$, we find two branches that are compatible with the current estimate for z_* , i.e., $-0.52 < \tilde{A} \leq -0.17$ and $0.46 \leq \tilde{A} \leq 4.35$. In agreement with recent results, we also found that the Λ CDM scenario ($\tilde{A} = 0$) is $\approx 2\sigma$ off from the central value of the current estimate for the redshift of the D/A transition.

3.2. Statistical analysis

As mentioned earlier, the best way to check the viability of parametrization (1) and, based on it, to discuss how different our Universe may be from the standard Λ CDM scenario, is to perform a statistical analysis involving the currently available sets of cosmological observations. To this end, we use here some of the most recent observational data such as the Supernova Legacy Survey (SNLS) set of 115 SNe Ia, recently published by Astier et al. (2006). These distance measurements are combined with the growth-of-structure index and current estimates of the age of the Universe. For the LSS, we use the linear growth rate $f(z_{2df}) = 0.58 \pm 0.11$, measured by the 2 degree field (2dF) galaxy redshift survey (2dFGRS) (Verde et al. 2002; Lahav et al. 2002; Hawkins et al. 2003), where $z_{2df} = 0.15$ is the effective redshift of this survey and $f = d \ln D / d \ln a$ is determined by solving the equation for the linear growth rate (Lahav et al. 1991)

$$D'' + 2E(z)D' - \frac{3}{2}\Omega_{m,0}(1+z)^3D = 0, \quad (10)$$

and primes denote $d/d(H_0t)$. The age of the Universe is assumed to be $t_0^{\text{obs}} = 13.6 \pm 0.2$ Gyr, as obtained by MacTavish et al. (2005) from a joint analysis involving recent LSS data (the matter power spectra from the 2dFGRS and SDSS redshift surveys) and results of the most recent CMB experiments (WMAP, DASI, VSA, ACBAR, MAXIMA, CBI, and BOOMERANG) (for more details on statistical analyses we refer the reader to Lima et al. 2003; Choudhury & Padmanabhan 2005; Nesseris & Perivolaropoulos 2004b; Capozziello et al. 2005;

² Another interesting example of transient acceleration is provided by the brane-world scenarios discussed by Sahni & Shtanov (2002).




Fig. 3. The linear growth rate $f(z_{2df})$ as a function of index α for some selected values of parameter \tilde{A} . Note that, while the quantity $f(z_{2df})$ is an insensitive function of α , it strongly depends on the parameter \tilde{A} .

Alam & Sahni 2005.)³. In Fig. 3 we display the linear growth rate $f(z_{2df})$ as a function of the index α for some selected values of the parameter \tilde{A} and $\Omega_{m,0} = 0.27$. Note that while the quantity $f(z_{2df})$ is almost insensitive to the index α , it depends considerably on the parameter \tilde{A} .

In Fig. 4 we show the resulting likelihood from our statistical analysis. Contours of 68.3%, 95.4% and 99.73% confidence levels (c.l.) are displayed in the $\tilde{A} - \alpha$ plane by considering a joint analysis involving the observational data described above. Note that at least two important conclusions may be drawn from this figure. First, that the allowed interval for parameter \tilde{A} is considerably restricted around the value $\tilde{A} = 0$, which is formally equivalent to the cosmological constant Λ (see Eq. (3)). In other words, such a result amounts to saying that we cannot go very far from the vacuum EOS ($w_{\text{eff}} = -1$), in order to be consistent with the presently available cosmological observations. This is very probably true, although the second aspect of Fig. 3 also shows that almost no constraint can be placed on the additional parameter α from the available sets of observational data, which hampers any definitive answer to our initial question. Such a conclusion may also be generalized for other EOS parametrizations as, for instance, the so-called Chaplygin gas EOS (Kamenshchik et al. 2001; Bilic et al. 2002; Bento et al. 2002; Dev et al. 2003) or still for EOS parametrizations that explicitly depend on time/redshift (Goliath et al. 2001; Efstathiou 1999; Chevallier & Polarski 2001; Linder 2003; Padmanabhan & Choudhury 2003; Jain et al. 2006; see also Maor et al. 2002, for a discussion of this topic). For the analysis performed here, the best-fit model is found for values of $\tilde{A} = 0.03$ and $\alpha = 0.98$, which corresponds to a 13.6-Gyr-old, currently accelerated

³ Note that in the analysis of the observational data one needs to take into account the peculiarities of the behaviour of the dark energy density (1) at positive redshifts. Namely, when $\tilde{A}(1 - \alpha) > 0$, the dark energy density is no longer a well defined function of redshift (for a sufficiently high redshift). A possible interpretation in this case is that the dark energy EOS (1) is just an effective description valid at lower redshifts and that the DE density is negligible at higher redshifts. These considerations are taken into account in our statistical analysis of the observational data.




Fig. 4. The plane $\tilde{A} - \alpha$ for the EOS parametrization discussed in this paper. The curves correspond to confidence regions of 68.3% and 95.4% for a joint analysis involving SNe Ia, growth of structure index, and the age of the Universe, as discussed in the text. Note that, although the parameter \tilde{A} is restricted to the interval $-0.06 \leq \tilde{A} \leq 0.12$ at 95.4% c.l., parameter α is completely unconstrained by the current observational data.

universe with $q_0 = -0.63$. Note that, according to our discussion presented in Sect. 2, these values for \tilde{A} and α also characterize a singular fate of the universe qualitatively similar to the “Big Rip” singularity (Caldwell et al. 2003; Nesseris & Perivolaropoulos 2004a), in which the scale factor a diverges at a finite time, as do the dark energy density ρ_d and pressure p_d . Naturally, such a scenario cannot be taken as a prediction from the current observational data since positive and negative values of \tilde{A} are possible at 1σ level and the value of α is completely unconstrained by current observations. In particular, at 95.4% c.l. the values of the parameter \tilde{A} are restricted to the interval $-0.06 \leq \tilde{A} \leq 0.12$.

4. Concluding remarks

Although some questions about the current value of the dark energy EOS still remain to be answered, it is nowadays becoming a consensus that its effective value must be some very close to the cosmological constant prediction, i.e., $w \equiv p_d/\rho_d = -1$. However, if the functional form of the EOS of the dark component responsible for the present cosmic acceleration has additional parameters, very little can be said about their values from the available sets of observational data. This important aspect in our search for a better understanding of the physical properties of dark energy is exactly the focus of our discussion in this paper. By considering a kind of “expansion” around the vacuum of the type $p_d = -\rho_d - A\rho_d^\alpha$, we have first discussed the rich variety of phenomena predicted for the past and future of the Universe when different values (and/or combinations) of the parameters \tilde{A} and α are considered. We also studied the D/A transition and shown that for a large interval of parameter \tilde{A} , the parametrization discussed here fully agrees with the current estimate for the acceleration redshift (z_*), as given by Riess et al. (2004). Another interesting aspect of parametrization (1) is its prediction of models with a transient acceleration phase (D/A/D transition), which seems to be in accordance with the requirements of the string

or M-theory (Fischler et al. 2001) (see also Ellis et al. 2001). Finally, we also performed a statistical analysis involving some of the most recent cosmological observations to show that, while the parameter \tilde{A} is tightly restricted to values around the cosmological constant ($\tilde{A} = 0.03 \pm 0.09$ at 95.4% c.l.), the value of α is completely unconstrained by the current observational data, which hampers a definitive answer to our initial question. We expect that in the near future new sets of observations, along with more theoretical effort, will be able to shed some light on our (so far) *dark* search to measure how far we can really go from Λ .

Acknowledgements. H.S. acknowledges the support of the Secretaría de Estado de Universidades e Investigación of the Ministerio de Educación y Ciencia of Spain within the program “Ayudas para movilidad de Profesores de Universidad e Investigadores españoles y extranjeros”. He is also partially supported by MEC and FEDER under project 2004-04582-C02-01 and by the Dep. de Recerca de la Generalitat de Catalunya under contract CIRIT GC 2001SGR-00065 and would like to thank the Departament E.C.M. of the Universitat de Barcelona for the hospitality. J.S.A. acknowledges support from Conselho Nacional de Desenvolvimento Científico e Tecnológico (Brazilian Research Agency) grants CNPq/307860/2004-3 and CNPq/475835/2004-2 and from FAPERJ No. E-26/171.251/2004.

References

- Alcaniz, J. S. 2002, *Phys. Rev. D*, 65, 123514
 Alcaniz, J. S. 2004, *Phys. Rev. D*, 69, 0, 83521
 Alcaniz, J. S., Jain, D., & Dev, A. 2003, *Phys. Rev. D*, 67, 043514
 Alcaniz, J. S., & Maia, J. M. F. 2003, *Phys. Rev. D*, 67, 043502
 Alcaniz, J. S., & Lima, J. A. S. 2005, *Phys. Rev. D*, 72, 063516
 Allen, S. W., Schmidt, R. W., Ebeling, H., Fabian, A. C., van Speybroeck, L. 2004, *MNRAS*, 353, 457
 Astier, P., Guy, J., Regnault, N., et al. 2006, *A&A*, 447, 31
 Babic, A., Guberina, B., Horvat, R., & Stefancic, H. 2002, *Phys. Rev. D*, 65, 085002
 Barrow, J. D. 2004, *Class. Quant. Grav.*, 21, L79
 Bassett, B. A., Corasaniti, P. S., & Kunz, M., *ApJ*, 617, L1
 Bento, M. C., Bertolami, O., Sen, A. A. 2002, *Phys. Rev. D*, 66, 043507
 Bertolami, O. 1986, *Nuovo Cimento Soc. Ital. Fis. B*, 93, 36
 Calberg, R. G., Yee, H. K. C., Ellingson, E., et al. 1996, *ApJ*, 462, 32
 Caldwell, R. R. 2002, *Phys. Lett. B*, 545, 23
 Caldwell, R. R., Dave, R., & Steinhardt, P. J. 1998, *Phys. Rev. Lett.*, 80, 1582
 Caldwell, R. R., Kamionkowski, M., & Weinberg, N. N. 2003, *Phys. Rev. Lett.*, 91, 071301
 Capozziello, S., Cardone, V. F., Elizalde, E., Nojiri, S., & Odintsov, S. D. 2006, *Phys. Rev. D*, 73, 043512
 Chevallier, M., & Polarski, D. 2001, *Int. J. Mod. Phys. D*, 10, 213
 Choudhury, T. R., & Padmanabhan, T. 2005, *A&A*, 429, 807
 Corasaniti, P. S., Kunz, M., Parkinson, D., Copeland, E. J., & Bassett, B. A. 2004, *Phys. Rev. D*, 70, 083006
 Cunha, J. V., Lima, J. A. S., & Alcaniz, J. S. 2002, *Phys. Rev. D*, 66, 023520
 de Bernardis, P., Ade, P. A. R., Bock, J. J., et al. 2000, *Nature*, 404, 955
 Dev, A., Jain, D., & Alcaniz, J. S. 2003, *Phys. Rev. D*, 67, 023515
 Dev, A., Jain, D., & Alcaniz, J. S. 2004, *A&A*, 417, 847
 Dvali, G., Gabadadze, G., & Porrati, M. 2000, *Phys. Lett. B*, 485, 208
 Efstathiou, G. 1999, *MNRAS*, 310, 842
 Eisenstein, D. J., Zehavi, I., Hogg, D. W., et al. 2005, *ApJ*, 633, 560
 Ellis, J. R., Mavromatos, N. E., & Nanopoulos, D. V. 2001 [[hep-th/0105206](#)]
 Faraoni, V. 2002, *Int. J. Mod. Phys. D*, 11, 471
 Feng, B., Wang, X.-L., & Zhang, X.-M. 2005, *Phys. Lett. B*, 607
 Fischler, W., Kashani-Poor, A., McNees, R., & Paban, S. 2001, *JHEP*, 0107, 3
 Freedman, W. L., Madore, B. F., Gibson, B. K., et al. 2001, *Astrop. J.*, 553, 47
 Goliath, M., Amanullah, R., Astier, P., Goobar, A., & Pain, R. 2001, *A&A* 380, 6
 Hawkins, E., Maddox, S., Cole, S., et al. 2003, *MNRAS*, 346, 78
 Jain, D., Dev, A., & Alcaniz, J. S. 2002, *Phys. Rev. D*, 66, 083511
 Jain, D., Alcaniz, J. S., & Dev, A. 2006, *Nucl. Phys. B*, 732, 379 123003
 Kamensshchik, A., Moschella, U., & Pasquier, V. 2001, *Phys. Lett. B*, 511, 265
 Kolb, E. W. 1989, *ApJ*, 344, 543
 Lahav, O., Lilje, P. B., Primack, J. R., & Rees, M. J. 1991, *MNRAS*, 251, 128
 Lahav, O., Bridle, S. L., Percival, W. J., et al. 2002, *MNRAS*, 333, 961
 Lima, J. A. S., & Alcaniz, J. S. 2004, *Phys. Lett. B*, 600, 191
 Lima, J. A. S., Cunha, J. V., & Alcaniz, J. S. 2003, *Phys. Rev. D*, 68, 023510
 Linder, E. V. 2003, *Phys. Rev. Lett.*, 90, 091301
 Lue, A., & Starkman, G. D. 2003, *Phys. Rev. D*, 67, 064002
 MacTavish, C. J., Ade, P. A. R., Bock, J. J., et al. 2005 [[arXiv:astro-ph/0507503](#)]
 Maor, I., Brustein, R., McMahon, J., & Steinhardt, P. J. 2002, *Phys. Rev. D*, 65, 12300
 Maia, M. D., Monte, E. M., Maia, J. M. F., & Alcaniz, J. S. 2005, *Class. Quant. Grav.*, 22, 1623
 Melchiorri, A., Mersini, L., Odman, C. J., & Trodden, M. 2003, *Phys. Rev. D*, 68, 043509
 Nesseris, S., & Perivolaropoulos, L. 2004a, *Phys. Rev. D*, 70, 043531
 Nesseris, S., & Perivolaropoulos, L. 2004b, *Phys. Rev. D*, 70, 123529
 Nojiri, S., & Odintsov, S. D. 2005, *Phys. Rev. D*, 72, 023003
 Nojiri, S., & Odintsov, S. D. 2004, *Phys. Rev. D*, 70, 103522
 Nojiri, S., Odintsov, S. D., & Tsujikawa, S. 2005, *Phys. Rev. D*, 71, 063004
 Özer, M., & Taha, M. O. 1986, *Phys. Lett. B*, 171, 363
 Padmanabhan, T. 2003, *Phys. Rep.*, 380, 235
 Padmanabhan, T., & Choudhury, T. R. 2003, *MNRAS*, 344, 823
 Peebles, P. J. E., & Ratra, B. 2003, *Rev. Mod. Phys.*, 75, 559
 Perlmutter, S., Aldering, G., Goldhaber, G., et al. 1999, *ApJ*, 517, 565
 Randall, L., & Sundrum, R. 1999, *Phys. Rev. Lett.*, 83, 3370
 Ratra, B., & Peebles, P. J. E. 1988, *Phys. Rev. D*, 37, 3406
 Riess, A. G., Strolger, L.-G., Tonry, J., et al. 2004, *ApJ*, 607, 665
 Sahni, V., & Starobinsky, A. A. 2000, *Int. J. Mod. Phys. D*, 9, 373
 Sahni, V., & Shtanov, Y. 2002, *IJMP D*, 11, 1515
 Sahni, S., & Shtanov, Y. 2003, *JCAP*, 0311, 014
 Shapiro, I. L., & Sola, J. 2000, *Phys. Lett. B*, 475, 236.
 Shapiro, I. L., & Sola, J. 2002, *JHEP*, 0202, 006
 Shapiro, I. L., & Sola, J., & Stefancic, H. 2005, *JCAP*, 0501, 012
 Spergel, D. N., Verde, L., Peiris, H. V., et al. 2003, *ApJS*, 148, 175
 Sola, J., & Stefancic, H. 2005, *Phys. Lett. B*, 624, 147
 Sola, J., & Stefancic, H. 2006, *Mod. Phys. Lett. A*, 21, 479
 Stefancic, H. 2005, *Phys. Rev. D*, 71, 124036
 Stefancic, H. 2005, *Phys. Rev. D*, 71, 084024
 Tegmark, M., Strauss, M. A., Blanton, M. R., et al. 2004, *Phys. Rev. D*, 69, 103501
 Verde, L., Heavens, A. F., Percival, W. J., et al. 2002, *MNRAS*, 335, 432
 Wang, P., & Meng, X.-H. 2005, *Class. Quant. Grav.*, 22, 283
 Wetterich, C. 1988, *Nucl. Phys. B*, 302, 668