

Energy levels and radiative rates for inner shell transitions of Fe XVI[★]

K. M. Aggarwal and F. P. Keenan

Astrophysics Research Centre, School of Mathematics and Physics, Queen's University, Belfast BT7 1NN, Northern Ireland, UK
e-mail: K. Aggarwal@qub.ac.uk

Received 11 May 2006 / Accepted 16 November 2006

ABSTRACT

Aims. In this paper we report calculations for energy levels and radiative rates for transitions in Fe XVI.

Methods. The General purpose Relativistic Atomic Structure Package (GRASP) has been adopted for the calculations of energy levels and radiative rates.

Results. Energies for the lowest 134 levels among the $(1s^2 2s^2) 2p^6 3\ell$, $2p^5 3s^2$, $2p^5 3s 3p$, $2p^5 3s 3d$, $2p^5 3p 3d$, and $2p^5 3p^2$ configurations of Fe XVI are reported. Additionally, radiative rates, oscillator strengths, and line strengths are reported for all electric dipole (E1), magnetic dipole (M1), electric quadrupole (E2), magnetic quadrupole (M2), and electric octupole (E3) transitions among these levels. Comparisons are made with the available results in the literature, and the accuracy of the present results is assessed. Finally, lifetimes for all excited levels are also listed, although no measurements are presently available with which to compare these.

Key words. atomic data – atomic processes

1. Introduction

In a recent paper (Aggarwal & Keenan 2006) we reported energy levels, radiative rates, collision strengths, and excitation rate coefficients for transitions among 39 levels of the $(1s^2 2s^2 2p^6) nl$; $n \leq 7$, $\ell \leq 4$ configurations of Fe XVI. However, many lines of Fe XVI in the 15–17 Å X-ray range are frequently observed in astrophysical and laboratory plasmas, and some of these arise from the inner (L) shell excitation of Fe XVI, such as: $2p^6 3s + e^- \rightarrow 2p^5 3\ell 3\ell'$ (see, for example, Dere et al. 2001). These emission lines particularly arise at comparatively higher temperatures, and hence atomic data for such transitions are also useful in fusion plasmas. Therefore, in this paper we focus on the inner shell electron impact excitation of Fe XVI.

Earlier calculations for this ion have been performed by Cornille et al. (1994) and Bautista (2000). Cornille et al. adopted the SuperStructure (SS) code of Eissner et al. (1974) for the generation of wavefunctions, and the distorted-wave (DW) code of Eissner & Seaton (1972) for the computations of collision strengths (Ω). They included 44 levels of the $2p^6 3s$, $2p^5 3s^2$, $2p^5 3s 3p$ and $2p^5 3s 3d$ configurations in their calculations, and reported values of Ω at a single energy of 60 Ryd. The main deficiency of their work is that they neither considered the variation of Ω with energy, nor included the contribution of resonances in the determination of excitation rates. Both of these deficiencies were removed in the subsequent work of Bautista, who adopted a semi-relativistic R -matrix approach (Berrington et al. 1995). In his calculations, he included 59 LS terms of the $2p^6 3\ell$, $2p^5 3s^2$, $2p^5 3s 3p$, $2p^5 3s 3d$, $2p^5 3p 3d$, and $2p^5 3p^2$ configurations, and obtained results for *fine-structure* transitions through an algebraic transformation. In this process, he also included the relativistic effects through *term coupling coefficients* (TCC). Since Fe XVI is a moderately heavy ion ($Z = 26$), both configuration interaction

(CI) and relativistic effects are supposed to be important in the determination of wavefunctions, and subsequently of Ω . He observed that the contribution of resonances has enhanced the *effective* collision strengths (Υ) by up to three orders of magnitude, depending on the temperature, and the relativistic effects have enhanced these by several orders of magnitude, for some of the transitions. Since both of these observed enhancements are unusually high, one of our *aims* is to assess the accuracy of his reported results, apart from making other improvements.

In the present work, we include 134 levels among the $(1s^2 2s^2) 2p^6 3\ell$, $2p^5 3s^2$, $2p^5 3s 3p$, $2p^5 3s 3d$, $2p^5 3p 3d$, and $2p^5 3p^2$ configurations of Fe XVI. Additionally, our approach is fully relativistic, as we employ the GRASP (General purpose Relativistic Atomic Structure Package) code of Dyaal et al. (1989) for the generation of wavefunctions, and the Dirac Atomic R -matrix Code (DARC) of Norrington & Grant (2007) for computations of Ω , and subsequently of Υ . However, it may be noted that the $2p^5 3d^2$ configuration has been deliberately excluded from the calculations, because it generates 45 levels and all of which lie *above* the lowest 134 already included. The same argument applies to other higher configurations, such as $2p^5 3\ell 4\ell$ and $2p^5 4\ell^2$. Since all of these configurations give rise to energy levels *above* the 134 levels included presently, their inclusion in a calculation will not significantly affect the results for energy levels and/or the radiative rates among the lowest 134 levels. For example, a calculation performed with the additional inclusion of the $2p^5 3d^2$, $2p^5 3\ell 4\ell$ and $2p^5 4\ell^2$ configurations, resulting in a total of 1201 levels, yields energy levels within 1 eV (or equivalently within 0.1%) with those listed in Table 1. Therefore, their specific inclusion in a calculation may affect the results of effective collision strengths (at higher electron temperatures), due to the inclusion of resonances arising from the higher levels, but will not affect the reported results of energy levels and/or the radiative rates. Furthermore, the calculations for Ω (and subsequently for Υ) are computationally demanding, as the size of the Hamiltonian matrix is large (13 330), and hence will take a considerably long time to conclude. Therefore, in this

[★] Tables 2, 3 and 5 are only available in electronic form at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/463/399>

Table 1. Energy levels (in Ryd) and mixing coefficients of Fe XVI.

| Index | Configuration | Level | NIST | GRASP ^a | GRASP ^b | FAC ^c | Mixing coefficients ^d |
|-------|---------------------------------------------------|--------------------------------------------|----------|--------------------|--------------------|------------------|--------------------------------------|
| 1 | 2p ⁶ 3s | ² S _{1/2} | 0.00000 | 0.00000 | 0.00000 | 0.00000 | 0.999(1) |
| 2 | 2p ⁶ 3p | ² P ^o _{1/2} | 2.52598 | 2.56393 | 2.56618 | 2.54757 | 0.999(2) |
| 3 | 2p ⁶ 3p | ² P ^o _{3/2} | 2.71688 | 2.75844 | 2.75486 | 2.73706 | 0.999(3) |
| 4 | 2p ⁶ 3d | ² D _{3/2} | 6.15562 | 6.20499 | 6.19591 | 6.16949 | 0.999(4) |
| 5 | 2p ⁶ 3d | ² D _{5/2} | 6.18209 | 6.23332 | 6.22081 | 6.19510 | 0.999(5) |
| 6 | 2p ⁵ 3s ² | ² P ^o _{3/2} | 52.60745 | 52.36353 | 52.31794 | 52.41746 | 0.985(6) |
| 7 | 2p ⁵ 3s ² | ² P ^o _{1/2} | 53.51871 | 53.30895 | 53.24028 | 53.33762 | 0.985(7) |
| 8 | 2p ⁵ 3s3p | ⁴ S _{3/2} | | 54.13058 | 54.08374 | 54.17709 | -0.886(8), 0.431(14) |
| 9 | 2p ⁵ 3s3p | ⁴ D _{5/2} | 54.51199 | 54.37708 | 54.32940 | 54.42538 | 0.828(9), -0.416(13) |
| 10 | 2p ⁵ 3s3p | ⁴ D _{7/2} | | 54.43349 | 54.38145 | 54.47626 | 0.998(10) |
| 11 | 2p ⁵ 3s(¹ P)3p | ² D _{3/2} | 54.79442 | 54.43554 | 54.38938 | 54.48851 | 0.676(17), 0.536(11) |
| 12 | 2p ⁵ 3s(¹ P)3p | ² P _{1/2} | 54.68514 | 54.56418 | 54.52045 | 54.61812 | 0.582(12), -0.541(16), 0.478(18) |
| 13 | 2p ⁵ 3s3p | ⁴ P _{5/2} | | 54.67382 | 54.62489 | 54.72388 | 0.810(13), 0.481(21) |
| 14 | 2p ⁵ 3s3p | ⁴ P _{3/2} | 55.55084 | 54.67609 | 54.62869 | 54.72922 | 0.543(11), 0.507(14), 0.431(22) |
| 15 | 2p ⁵ 3s(¹ P)3p | ² S _{1/2} | 55.05876 | 54.89407 | 54.84442 | 54.93897 | 0.646(15), -0.537(18) |
| 16 | 2p ⁵ 3s3p | ⁴ D _{1/2} | 55.35947 | 55.28057 | 55.21306 | 55.30909 | 0.797(16), 0.453(12) |
| 17 | 2p ⁵ 3s3p | ⁴ D _{3/2} | 55.48705 | 55.37720 | 55.30739 | 55.40431 | 0.696(17), -0.430(11) |
| 18 | 2p ⁵ 3s3p | ⁴ P _{1/2} | 55.48705 | 55.39547 | 55.33684 | 55.42765 | 0.648(18), 0.545(15) |
| 19 | 2p ⁵ 3s(³ P)3p | ² D _{5/2} | 54.79442 | 55.49867 | 55.44467 | 55.52949 | -0.908(19) |
| 20 | 2p ⁵ 3s(³ P)3p | ² P _{3/2} | 55.85156 | 55.50353 | 55.43583 | 55.52862 | 0.663(14), -0.521(20) |
| 21 | 2p ⁵ 3s(¹ P)3p | ² D _{5/2} | 55.67842 | 55.57474 | 55.50469 | 55.60006 | -0.788(21), 0.464(9), |
| 22 | 2p ⁵ 3s(¹ P)3p | ² P _{3/2} | | 55.63602 | 55.58697 | 55.67262 | -0.642(22), 0.561(20), -0.418(24) |
| 23 | 2p ⁵ 3s(³ P)3p | ² P _{1/2} | | 56.31892 | 56.25643 | 56.34105 | 0.737(23), -0.420(12), -0.419(15) |
| 24 | 2p ⁵ 3s(³ P)3p | ² D _{3/2} | 56.65347 | 56.48207 | 56.40948 | 56.48980 | 0.761(24), -0.412(11) |
| 25 | 2p ⁵ 3s(³ P)3p | ² S _{1/2} | 57.10911 | 57.04715 | 56.99370 | 57.07898 | 0.894(25) |
| 26 | 2p ⁵ 3p ² | ⁴ P ^o _{3/2} | | 57.11374 | 57.06488 | 57.15728 | -0.579(26), 0.471(30) |
| 27 | 2p ⁵ 3p ² (¹ D) | ² P ^o _{1/2} | | 57.11721 | 57.06704 | 57.16091 | 0.708(27), -0.447(63) |
| 28 | 2p ⁵ 3p ² | ⁴ P ^o _{5/2} | | 57.18875 | 57.13944 | 57.22878 | 0.861(28), -0.457(35) |
| 29 | 2p ⁵ 3p ² | ² F ^o _{7/2} | | 57.20144 | 57.14806 | 57.24572 | -0.828(29) |
| 30 | 2p ⁵ 3p ² (¹ D) | ² P ^o _{3/2} | | 57.27558 | 57.22768 | 57.32069 | -0.517(26), -0.415(30) |
| 31 | 2p ⁵ 3p ² (¹ D) | ² D ^o _{5/2} | | 57.40167 | 57.35118 | 57.44791 | -0.624(31), 0.442(42) |
| 32 | 2p ⁵ 3p ² (³ P) | ² D ^o _{3/2} | | 57.43847 | 57.39127 | 57.48571 | 0.783(32), 0.420(43) |
| 33 | 2p ⁵ 3p ² | ⁴ P ^o _{1/2} | | 57.45866 | 57.40547 | 57.49659 | -0.765(33), 0.490(45) |
| 34 | 2p ⁵ 3p ² | ⁴ D ^o _{7/2} | | 57.47691 | 57.42105 | 57.51299 | 0.944(34) |
| 35 | 2p ⁵ 3p ² | ⁴ D ^o _{5/2} | | 57.49089 | 57.43729 | 57.53061 | 0.591(51), 0.555(35), 0.404(28) |
| 36 | 2p ⁵ 3p ² | ⁴ D ^o _{1/2} | | 57.93073 | 57.87458 | 57.96637 | 0.788(36), -0.440(45) |
| 37 | 2p ⁵ 3p ² (¹ D) | ² D ^o _{3/2} | | 57.97229 | 57.91773 | 58.01218 | 0.617(48), -0.465(37) |
| 38 | 2p ⁵ 3s3d | ⁴ P ^o _{1/2} | | 57.98889 | 57.93888 | 58.03972 | -0.979(38) |
| 39 | 2p ⁵ 3s3d | ⁴ P ^o _{3/2} | | 58.05947 | 58.00538 | 58.10584 | 0.938(39) |
| 40 | 2p ⁵ 3s3d | ⁴ F ^o _{9/2} | | 58.15573 | 58.09555 | 58.19396 | 1.000(40) |
| 41 | 2p ⁵ 3s3d | ⁴ P ^o _{5/2} | 58.25730 | 58.17918 | 58.11794 | 58.22565 | 0.799(41), -0.449(59) |
| 42 | 2p ⁵ 3p ² | ² F ^o _{5/2} | | 58.19900 | 58.12751 | 58.21310 | -0.641(42), -0.442(31) |
| 43 | 2p ⁵ 3p ² | ⁴ D ^o _{3/2} | | 58.21215 | 58.14156 | 58.23244 | 0.655(43), -0.410(37) |
| 44 | 2p ⁵ 3s3d | ⁴ F ^o _{7/2} | | 58.21941 | 58.16102 | 58.25588 | 0.885(44) |
| 45 | 2p ⁵ 3p ² | ² S ^o _{1/2} | | 58.28129 | 58.21189 | 58.30146 | 0.712(45), 0.538(33), 0.447(36) |
| 46 | 2p ⁵ 3s3d | ⁴ F ^o _{5/2} | 58.37577 | 58.30062 | 58.24386 | 58.33759 | 0.747(46), 0.431(50) |
| 47 | 2p ⁵ 3s(¹ P)3d | ² D ^o _{3/2} | | 58.40180 | 58.34352 | 58.43276 | -0.576(58), 0.444(47) |
| 48 | 2p ⁵ 3p ² | ⁴ S ^o _{3/2} | | 58.43197 | 58.36742 | 58.45592 | 0.479(48), 0.443(26), 0.437(52) |
| 49 | 2p ⁵ 3s3d | ⁴ D ^o _{7/2} | 58.52157 | 58.45282 | 58.39564 | 58.48497 | 0.815(49), 0.497(61) |
| 50 | 2p ⁵ 3s(¹ P)3d | ² F ^o _{5/2} | 58.54891 | 58.48128 | 58.42242 | 58.51233 | 0.570(50), 0.482(59) |
| 51 | 2p ⁵ 3p ² (³ P) | ² D ^o _{5/2} | | 58.51272 | 58.43827 | 58.53265 | -0.673(51), 0.506(35) |
| 52 | 2p ⁵ 3p ² (¹ S) | ² P ^o _{3/2} | | 58.51295 | 58.45493 | 58.54002 | 0.694(52) |
| 53 | 2p ⁵ 3s(¹ P)3d | ² P ^o _{1/2} | 58.53068 | 58.53541 | 58.48029 | 58.57178 | 0.622(53), -0.589(55) |
| 54 | 2p ⁵ 3s(¹ P)3d | ² P ^o _{3/2} | 58.64915 | 58.71984 | 58.66234 | 58.74794 | 0.550(54), -0.548(56) |
| 55 | 2p ⁵ 3s3d | ⁴ D ^o _{1/2} | | 58.82057 | 58.76090 | 58.83807 | 0.646(55), 0.541(63), -0.420(67) |
| 56 | 2p ⁵ 3s3d | ⁴ D ^o _{3/2} | 58.98632 | 59.10868 | 59.04324 | 59.12486 | -0.521(56), -0.405(65) |
| 57 | 2p ⁵ 3s(³ P)3d | ² F ^o _{7/2} | 58.73116 | 59.19215 | 59.13497 | 59.21341 | 0.772(57), 0.489(29), -0.402(61) |
| 58 | 2p ⁵ 3s3d | ⁴ F ^o _{3/2} | 59.25058 | 59.21520 | 59.14077 | 59.23393 | 0.700(58), 0.568(47) |
| 59 | 2p ⁵ 3s3d | ⁴ D ^o _{5/2} | 59.25058 | 59.23119 | 59.15303 | 59.24131 | 0.654(59), 0.535(46) |
| 60 | 2p ⁵ 3s(³ P)3d | ² D ^o _{5/2} | 58.90430 | 59.27727 | 59.20474 | 59.28913 | -0.653(60) |
| 61 | 2p ⁵ 3s(¹ P)3d | ² F ^o _{7/2} | 59.38727 | 59.33240 | 59.25137 | 59.33982 | -0.666(61), 0.492(49), 0.461(44) |
| 62 | 2p ⁵ 3s(¹ P)3d | ² D ^o _{5/2} | 59.37816 | 59.40018 | 59.34082 | 59.42136 | 0.677(62) |
| 63 | 2p ⁵ 3p ² (³ P) | ² P ^o _{1/2} | | 59.40747 | 59.34593 | 59.43010 | 0.527(53), -0.482(27) |
| 64 | 2p ⁵ 3p ² (¹ S) | ² P ^o _{1/2} | | 59.54384 | 59.47971 | 59.55638 | 0.881(64), |

Table 1. continued.

| Index | Configuration | Level | NIST | GRASP ^a | GRASP ^b | FAC ^c | Mixing coefficients ^d |
|-------|---------------------------------------------------|--------------------------------------------|----------|--------------------|--------------------|------------------|---------------------------------------|
| 65 | 2p ⁵ 3p ² (³ P) | ² P _{3/2} ^o | | 59.61159 | 59.55298 | 59.63031 | 0.601(65), 0.462(30) |
| 66 | 2p ⁵ 3s(³ P)3d | ² D _{3/2} ^o | | 59.75825 | 59.68822 | 59.75204 | -0.506(76), 0.506(66), -0.493(54) |
| 67 | 2p ⁵ 3s(³ P)3d | ² P _{1/2} ^o | 59.90670 | 60.02441 | 59.95160 | 60.00957 | -0.802(67), -0.429(53) |
| 68 | 2p ⁵ 3p3d(³ S) | ⁴ D _{1/2} | | 60.09502 | 60.04611 | 60.14409 | -0.730(68), -0.569(120) |
| 69 | 2p ⁵ 3p3d(³ S) | ⁴ D _{3/2} | | 60.17116 | 60.11829 | 60.21617 | 0.740(69), 0.517(119) |
| 70 | 2p ⁵ 3s(³ P)3d | ² F _{5/2} ^o | 59.74267 | 60.20698 | 60.12842 | 60.20253 | 0.619(70), 0.407(60) |
| 71 | 2p ⁵ 3p3d(³ S) | ⁴ D _{5/2} | | 60.29006 | 60.23238 | 60.32960 | 0.759(71), 0.410(99) |
| 72 | 2p ⁵ 3p3d(³ D) | ⁴ G _{7/2} | | 60.41131 | 60.35233 | 60.44618 | 0.616(72), 0.429(74), 0.403(82) |
| 73 | 2p ⁵ 3p3d | ⁴ G _{9/2} | | 60.43500 | 60.37302 | 60.47084 | 0.884(73) |
| 74 | 2p ⁵ 3p3d(³ S) | ⁴ D _{7/2} | | 60.44591 | 60.38502 | 60.50865 | -0.681(74) |
| 75 | 2p ⁵ 3p3d | ⁴ G _{11/2} | | 60.47640 | 60.41011 | 60.47567 | 1.000(75) |
| 76 | 2p ⁵ 3s(³ P)3d | ² P _{3/2} ^o | 60.09806 | 60.47817 | 60.40230 | 60.47991 | 0.667(76), 0.408(66), 0.402(30) |
| 77 | 2p ⁵ 3p3d(³ D) | ² D _{5/2} | | 60.50269 | 60.44777 | 60.53828 | -0.484(113) |
| 78 | 2p ⁵ 3p3d(³ D) | ² D _{3/2} | | 60.54189 | 60.48675 | 60.58196 | -0.451(112), 0.428(78) |
| 79 | 2p ⁵ 3p3d(³ D) | ² F _{7/2} | | 60.60902 | 60.54820 | 60.64577 | 0.680(79), 0.473(108) |
| 80 | 2p ⁵ 3p3d(¹ P) | ² F _{5/2} | | 60.61475 | 60.55605 | 60.64870 | -0.559(101), 0.426(80) |
| 81 | 2p ⁵ 3p3d(¹ P) | ² P _{1/2} | | 60.65721 | 60.60183 | 60.70206 | -0.540(97), 0.509(81), 0.480(126) |
| 82 | 2p ⁵ 3p3d(¹ D) | ² G _{7/2} | | 60.72887 | 60.67077 | 60.76248 | 0.531(82) |
| 83 | 2p ⁵ 3p3d(³ D) | ⁴ P _{1/2} | | 60.78928 | 60.73533 | 60.83237 | 0.762(83), 0.531(115) |
| 84 | 2p ⁵ 3p3d(³ D) | ⁴ P _{3/2} | | 60.82790 | 60.77171 | 60.86720 | 0.744(84), -0.550(86) |
| 85 | 2p ⁵ 3p3d(³ P) | ⁴ F _{9/2} | | 60.82979 | 60.76830 | 60.85897 | 0.615(85), 0.525(109), 0.500(91) |
| 86 | 2p ⁵ 3p3d(³ D) | ⁴ S _{3/2} | | 60.87238 | 60.81615 | 60.90864 | 0.609(86) |
| 87 | 2p ⁵ 3p3d(³ P) | ⁴ D _{7/2} | | 60.88786 | 60.82874 | 60.92297 | 0.693(87), -0.464(94) |
| 88 | 2p ⁵ 3p3d(³ D) | ⁴ F _{5/2} | | 60.91229 | 60.85344 | 60.94462 | -0.451(89), 0.406(114), 0.406(99) |
| 89 | 2p ⁵ 3p3d(³ D) | ⁴ P _{5/2} | | 60.91779 | 60.85914 | 60.94992 | 0.742(89) |
| 90 | 2p ⁵ 3p3d(³ D) | ⁴ F _{3/2} | | 60.97310 | 60.91494 | 60.99975 | -0.561(90) |
| 91 | 2p ⁵ 3p3d(³ D) | ⁴ F _{9/2} | | 61.01944 | 60.95573 | 61.04967 | 0.795(91), -0.591(85) |
| 92 | 2p ⁵ 3p3d(¹ P) | ² P _{3/2} | | 61.02141 | 60.96448 | 61.05502 | 0.443(92) |
| 93 | 2p ⁵ 3p3d(¹ P) | ² D _{5/2} | | 61.07479 | 61.01801 | 61.10393 | -0.506(114), 0.433(93), |
| 94 | 2p ⁵ 3p3d(³ D) | ⁴ F _{7/2} | | 61.09930 | 61.03764 | 61.12961 | 0.513(94), -0.511(106), 0.429(121) |
| 95 | 2p ⁵ 3p3d(¹ D) | ² F _{5/2} | | 61.12272 | 61.06369 | 61.15376 | 0.610(95), -0.458(102) |
| 96 | 2p ⁵ 3p3d(³ D) | ⁴ D _{7/2} | | 61.14633 | 61.08498 | 61.16957 | 0.826(96) |
| 97 | 2p ⁵ 3p3d(³ P) | ² P _{1/2} | | 61.16838 | 61.11025 | 61.19611 | 0.525(68), -0.486(120) |
| 98 | 2p ⁵ 3p3d(¹ D) | ² D _{3/2} | | 61.23950 | 61.18234 | 61.26807 | 0.595(98), -0.507(119) |
| 99 | 2p ⁵ 3p3d(³ P) | ⁴ D _{5/2} | | 61.26175 | 61.20205 | 61.28725 | -0.509(99), 0.455(122) |
| 100 | 2p ⁵ 3p3d(¹ P) | ² D _{3/2} | | 61.31016 | 61.24893 | 61.33183 | -0.544(112), 0.484(100) |
| 101 | 2p ⁵ 3p3d(³ D) | ⁴ G _{5/2} | | 61.32200 | 61.24585 | 61.33821 | 0.606(101), 0.502(107), 0.410(80) |
| 102 | 2p ⁵ 3p3d(³ P) | ⁴ F _{5/2} | | 61.41068 | 61.34445 | 61.43000 | 0.451(80) |
| 103 | 2p ⁵ 3p3d(³ D) | ⁴ D _{1/2} | | 61.41192 | 61.35321 | 61.42990 | -0.573(103), 0.501(126) |
| 104 | 2p ⁵ 3p3d(³ P) | ⁴ F _{3/2} | | 61.47289 | 61.40208 | 61.49645 | 0.461(104), 0.411(100) |
| 105 | 2p ⁵ 3p3d(¹ D) | ² S _{1/2} | | 61.48154 | 61.42443 | 61.50666 | -0.646(105), -0.628(115) |
| 106 | 2p ⁵ 3p3d(³ P) | ⁴ F _{7/2} | | 61.50415 | 61.42221 | 61.51326 | 0.605(72) |
| 107 | 2p ⁵ 3p3d(³ D) | ² F _{5/2} | | 61.55458 | 61.47347 | 61.56445 | -0.490(107), 0.466(125) |
| 108 | 2p ⁵ 3p3d(¹ P) | ² F _{7/2} | | 61.64063 | 61.56056 | 61.64670 | 0.499(87), -0.418(108), 0.414(106) |
| 109 | 2p ⁵ 3p3d(¹ D) | ² G _{9/2} | | 61.66765 | 61.58175 | 61.66985 | -0.662(109), 0.457(85), 0.445(73) |
| 110 | 2p ⁵ 3p3d(³ D) | ⁴ D _{3/2} | | 61.68685 | 61.61493 | 61.69331 | 0.604(110) |
| 111 | 2p ⁵ 3p3d(³ D) | ² G _{9/2} | | 61.75682 | 61.69334 | 61.77370 | 0.919(111) |
| 112 | 2p ⁵ 3p3d(³ S) | ² D _{3/2} | | 61.79240 | 61.72111 | 61.81103 | 0.588(90), -0.423(104) |
| 113 | 2p ⁵ 3p3d(³ S) | ² D _{5/2} | | 61.86910 | 61.80037 | 61.88206 | 0.494(113), 0.439(118) |
| 114 | 2p ⁵ 3p3d(³ D) | ⁴ D _{5/2} | | 61.94324 | 61.86516 | 61.94608 | 0.469(114), 0.439(93) |
| 115 | 2p ⁵ 3p3d(³ P) | ⁴ P _{1/2} | | 61.95561 | 61.88979 | 61.96959 | 0.469(132), -0.466(83), 0.462(115) |
| 116 | 2p ⁵ 3p3d(³ P) | ² F _{7/2} | | 61.96033 | 61.89980 | 61.97643 | -0.520(129), 0.496(116), 0.444(79) |
| 117 | 2p ⁵ 3p3d(³ P) | ⁴ P _{3/2} | | 61.99275 | 61.91457 | 61.99405 | 0.470(84), 0.456(123) |
| 118 | 2p ⁵ 3p3d(³ P) | ⁴ P _{5/2} | | 62.01950 | 61.94888 | 62.02740 | 0.494(118), -0.423(93) |
| 119 | 2p ⁵ 3p3d(³ P) | ⁴ D _{3/2} | | 62.05674 | 61.97802 | 62.06391 | 0.467(119) |
| 120 | 2p ⁵ 3p3d(³ P) | ⁴ D _{1/2} | | 62.05790 | 61.97998 | 62.06754 | -0.610(120), -0.444(81) |
| 121 | 2p ⁵ 3p3d(¹ D) | ² F _{7/2} | | 62.07990 | 61.99687 | 62.08230 | -0.559(121) |
| 122 | 2p ⁵ 3p3d(¹ D) | ² D _{5/2} | | 62.09892 | 62.02163 | 62.10258 | 0.499(130) |
| 123 | 2p ⁵ 3p3d(³ P) | ² P _{3/2} | | 62.11304 | 62.05018 | 62.11879 | -0.492(123), -0.425(78), -0.418(127) |
| 124 | 2p ⁵ 3p3d(³ P) | ² D _{3/2} | | 62.34616 | 62.27680 | 62.32272 | 0.589(124), 0.457(78) |
| 125 | 2p ⁵ 3p3d(³ P) | ² F _{5/2} | | 62.35587 | 62.28342 | 62.34591 | -0.485(125), 0.446(122) |
| 126 | 2p ⁵ 3p3d(¹ D) | ² P _{1/2} | | 62.39653 | 62.31859 | 62.38120 | -0.549(97), -0.463(126) |
| 127 | 2p ⁵ 3p3d(¹ D) | ² P _{3/2} | | 62.58476 | 62.50435 | 62.55824 | 0.492(127), 0.467(131) |
| 128 | 2p ⁵ 3p3d(³ D) | ² P _{1/2} | | 62.78038 | 62.70084 | 62.74799 | -0.577(128), 0.520(132) |

Table 1. continued.

| Index | Configuration | Level | NIST | GRASP ^a | GRASP ^b | FAC ^c | Mixing coefficients ^d |
|-------|---------------------------------------|-------------------------------|----------|--------------------|--------------------|-------------------------|----------------------------------|
| 129 | 2p ⁵ 3p3d(³ D) | ² G _{7/2} | 62.80742 | 62.72376 | 62.79937 | 0.656(129), 0.482(116) | |
| 130 | 2p ⁵ 3p3d(³ P) | ² D _{5/2} | 62.90893 | 62.83399 | 62.88355 | 0.533(130), 0.479(134) | |
| 131 | 2p ⁵ 3p3d(³ D) | ² P _{3/2} | 62.93802 | 62.86270 | 62.92609 | 0.526(124), -0.494(131) | |
| 132 | 2p ⁵ 3p3d(³ D) | ² S _{1/2} | 63.11751 | 63.03638 | 63.09278 | 0.630(132), 0.574(128) | |
| 133 | 2p ⁵ 3p3d(¹ S) | ² D _{3/2} | 63.27350 | 63.21048 | 63.29593 | 0.875(133) | |
| 134 | 2p ⁵ 3p3d(¹ S) | ² D _{5/2} | 63.34064 | 63.27358 | 63.37006 | 0.850(134) | |

NIST: <http://physics.nist.gov/PhysRefData>.

^a energies *without* Breit and QED corrections;

^b energies *with* Breit and QED corrections;

^c energies from the Flexible Atomic Code of Gu (2003);

^d mixing coefficient of the level (in bracket).

paper we report our results for energy levels and radiative rates (A -values) for five types of transitions, namely electric dipole (E1), electric quadrupole (E2), magnetic dipole (M1), magnetic quadrupole (M2) and electric octupole (E3), as all of these are required in the modelling of plasmas. Additionally, we report lifetimes for all excited levels, although no measurements are presently available with which to compare these results.

2. Energy levels

The 2p⁶3ℓ, 2p⁵3s², 2p⁵3s3p, 2p⁵3s3d, 2p⁵3p3d, and 2p⁵3p² configurations of Fe XVI give rise to 134 fine-structure levels, listed in Table 1. Our calculated energies obtained from the GRASP code, *with* and *without* including Breit and QED effects, are given in this table along with those from the experimental compilations of NIST (<http://physics.nist.gov/PhysRefData>). For our calculations, we have used the option of *extended average level* (EAL), in which a weighted (proportional to $2j + 1$) trace of the Hamiltonian matrix is minimized. This produces a compromise set of orbitals describing closely lying states with moderate accuracy. The inclusion of Breit and QED effects *lowers* the energies up to a maximum of 0.08 Ryd ($\leq 0.2\%$) – see for example, levels 61, 101 and 131. In general, the theoretical energies agree well in magnitude with the experimental ones, but differences are up to 0.4 Ryd ($\leq 0.7\%$) for some levels, such as: 11, 57 and 70. Additionally, our energies without the Breit and QED corrections are comparatively in better agreement with the corresponding experimental results, although the latter are available for only a few levels.

However, the level orderings in theory and experiment are slightly different in a few instances, such as for levels: 11/12 and 61/62. Most of these levels have energies very close to each other, and differing amount of configuration interaction (CI) may result in slightly different level orderings. However, a more serious problem is of level identification. Some of the levels of a configuration with the *same* J value are highly mixed, such as levels 11/17 (2p⁵3s3p ²D_{3/2}, ⁴D_{3/2}), 26/30 (2p⁵3p² ⁴P_{3/2}^o, ²P_{3/2}^o) and 66/76 (2p⁵3s3d ²D_{3/2}^o, ²P_{3/2}^o). For these (and other similar) levels the identification of a particular level is not unique, and is not based on the strength of the dominant eigenvector alone. This is a common problem in all atomic structure calculations, especially when the levels of a configuration are highly mixed, and the addition of more CI complicates the identifications further rather than making the task easier. Therefore, the configuration and the J values given in Table 1 are definite, but the corresponding LSJ designation provided for a level is only for guidance, and is liable to interchange. This is the main reason that some of the experimental levels, such as 2p⁵3s(¹P)3p ²D_{3/2}

and 2p⁵3s(³P)3p ²D_{3/2}, i.e. levels 11 and 24, appear to be listed at the wrong places.

We have taken all precautions to label the levels in Table 1, although some *ambiguity* in their designations cannot be avoided as discussed above. Furthermore, in the absence of any other theoretical or experimental energies to compare with, we have performed another calculation from the *Flexible Atomic Code* (FAC) of Gu (2003), which is easily available from the website <http://kipac-tree.stanford.edu/fac>. Energy levels obtained from this code are also listed in Table 1, and the excellent agreement (within 0.1 Ryd) between the GRASP and FAC energy levels gives confidence in our results. The level orderings are also (nearly) the same in both calculations, although some closely spaced levels, such as 41/42, 69/70 and 102/103, have switched positions. Finally, in the last column of Table 1 we have included the mixing coefficients (largest three where appropriate), which may be helpful for further comparisons and redesignations of the levels.

3. Radiative rates

The absorption oscillator strength (f_{ij}) and radiative rate A_{ji} (in s⁻¹) for a transition $i \rightarrow j$ are related by the following expression:

$$f_{ij} = \frac{mc}{8\pi^2 e^2} \lambda_{ji}^2 \frac{\omega_j}{\omega_i} A_{ji} = 1.49 \times 10^{-16} \lambda_{ji}^2 (\omega_j/\omega_i) A_{ji} \quad (1)$$

where m and e are the electron mass and charge, respectively, c is the velocity of light, λ_{ji} is the transition energy/wavelength in Å, and ω_i and ω_j are the statistical weights of the lower (i) and upper (j) levels, respectively. Similarly, the oscillator strength f_{ij} (dimensionless) and the line strength S (in atomic unit, 1 a.u. = 6.460 × 10⁻³⁶ cm² esu²) are related by the following standard equations:

for the electric dipole (E1) transitions:

$$A_{ji} = \frac{2.0261 \times 10^{18}}{\omega_j \lambda_{ji}^3} S^{E1} \quad \text{and} \quad f_{ij} = \frac{303.75}{\lambda_{ji} \omega_i} S^{E1}, \quad (2)$$

for the magnetic dipole (M1) transitions:

$$A_{ji} = \frac{2.6974 \times 10^{13}}{\omega_j \lambda_{ji}^3} S^{M1} \quad \text{and} \quad f_{ij} = \frac{4.044 \times 10^{-3}}{\lambda_{ji} \omega_i} S^{M1}, \quad (3)$$

for the electric quadrupole (E2) transitions:

$$A_{ji} = \frac{1.1199 \times 10^{18}}{\omega_j \lambda_{ji}^5} S^{E2} \quad \text{and} \quad f_{ij} = \frac{167.89}{\lambda_{ji}^3 \omega_i} S^{E2}, \quad (4)$$

Table 4. Transition wavelengths (λ_{ij} in Å), radiative rates (A_{ji} in s^{-1}), and oscillator strengths (f_{ij} , dimensionless) for some ($f \geq 10^{-8}$) magnetic dipole (M2) and electric octupole (E3) transitions of Fe XVI. ($a \pm b \equiv a \times 10^{\pm b}$).

| (a) Magnetic quadrupole (M2) transitions | | | | | | | | | |
|------------------------------------------|-----|----------------|---------------|---------------|-----|-----|----------------|---------------|---------------|
| i | j | λ_{ij} | A_{ji}^{M2} | f_{ij}^{M2} | i | j | λ_{ij} | A_{ji}^{M2} | f_{ij}^{M2} |
| 1 | 6 | 17.418 | 2.449+05 | 2.228-08 | 2 | 113 | 15.384 | 1.941+05 | 2.066-08 |
| 1 | 26 | 15.969 | 2.681+05 | 2.050-08 | 2 | 122 | 15.327 | 2.224+05 | 2.350-08 |
| 1 | 30 | 15.924 | 3.256+05 | 2.475-08 | 2 | 125 | 15.260 | 1.266+05 | 1.325-08 |
| 1 | 37 | 15.734 | 2.622+05 | 1.946-08 | 2 | 134 | 15.011 | 3.341+05 | 3.386-08 |
| 1 | 39 | 15.710 | 7.363+05 | 5.449-08 | 3 | 10 | 17.651 | 2.057+05 | 1.922-08 |
| 1 | 41 | 15.680 | 3.053+06 | 3.376-07 | 3 | 72 | 15.821 | 9.035+05 | 6.781-08 |
| 1 | 42 | 15.677 | 1.791+06 | 1.980-07 | 3 | 74 | 15.812 | 1.943+06 | 1.457-07 |
| 1 | 46 | 15.646 | 6.013+05 | 6.620-08 | 3 | 77 | 15.795 | 6.577+05 | 3.690-08 |
| 1 | 47 | 15.619 | 2.542+05 | 1.860-08 | 3 | 78 | 15.784 | 6.177+05 | 2.307-08 |
| 1 | 48 | 15.613 | 1.624+05 | 1.187-08 | 3 | 79 | 15.768 | 7.334+05 | 5.467-08 |
| 1 | 50 | 15.598 | 6.502+05 | 7.115-08 | 3 | 80 | 15.766 | 7.812+05 | 4.366-08 |
| 1 | 51 | 15.594 | 2.082+05 | 2.277-08 | 3 | 81 | 15.753 | 1.576+06 | 2.931-08 |
| 1 | 52 | 15.589 | 1.101+06 | 8.021-08 | 3 | 82 | 15.734 | 4.683+05 | 3.476-08 |
| 1 | 54 | 15.534 | 1.515+06 | 1.096-07 | 3 | 86 | 15.695 | 6.068+05 | 2.241-08 |
| 1 | 56 | 15.434 | 1.547+06 | 1.105-07 | 3 | 87 | 15.692 | 1.408+06 | 1.039-07 |
| 1 | 58 | 15.408 | 2.042+05 | 1.453-08 | 3 | 88 | 15.685 | 1.678+06 | 9.285-08 |
| 1 | 59 | 15.405 | 3.876+05 | 4.137-08 | 3 | 89 | 15.683 | 6.410+05 | 3.546-08 |
| 1 | 62 | 15.356 | 1.283+05 | 1.361-08 | 3 | 90 | 15.668 | 1.012+06 | 3.725-08 |
| 1 | 65 | 15.302 | 5.515+05 | 3.872-08 | 3 | 92 | 15.655 | 5.517+05 | 2.027-08 |
| 1 | 70 | 15.155 | 1.136+05 | 1.173-08 | 3 | 93 | 15.641 | 6.914+05 | 3.804-08 |
| 1 | 76 | 15.087 | 2.396+05 | 1.635-08 | 3 | 94 | 15.635 | 3.434+05 | 2.517-08 |
| 2 | 8 | 17.688 | 1.598+05 | 1.500-08 | 3 | 96 | 15.623 | 2.638+05 | 1.931-08 |
| 2 | 9 | 17.605 | 1.461+05 | 2.037-08 | 3 | 97 | 15.616 | 7.073+05 | 1.293-08 |
| 2 | 69 | 15.834 | 3.251+05 | 2.444-08 | 3 | 98 | 15.597 | 6.094+05 | 2.222-08 |
| 2 | 71 | 15.802 | 1.680+06 | 1.887-07 | 3 | 103 | 15.551 | 1.061+06 | 1.924-08 |
| 2 | 78 | 15.733 | 1.734+06 | 1.287-07 | 3 | 104 | 15.538 | 1.555+06 | 5.628-08 |
| 2 | 80 | 15.714 | 8.790+05 | 9.763-08 | 3 | 105 | 15.532 | 2.034+06 | 3.678-08 |
| 2 | 84 | 15.656 | 2.694+05 | 1.980-08 | 3 | 106 | 15.533 | 2.467+05 | 1.785-08 |
| 2 | 86 | 15.644 | 2.070+06 | 1.519-07 | 3 | 107 | 15.519 | 2.687+05 | 1.455-08 |
| 2 | 88 | 15.634 | 1.172+05 | 1.289-08 | 3 | 113 | 15.433 | 6.397+05 | 3.426-08 |
| 2 | 89 | 15.633 | 1.301+06 | 1.430-07 | 3 | 114 | 15.416 | 4.763+05 | 2.546-08 |
| 2 | 92 | 15.604 | 1.758+06 | 1.284-07 | 3 | 115 | 15.410 | 1.322+06 | 2.352-08 |
| 2 | 95 | 15.578 | 3.409+05 | 3.720-08 | 3 | 116 | 15.407 | 3.057+05 | 2.176-08 |
| 2 | 99 | 15.541 | 5.121+05 | 5.563-08 | 3 | 118 | 15.395 | 3.764+05 | 2.006-08 |
| 2 | 101 | 15.530 | 7.033+05 | 7.628-08 | 3 | 119 | 15.387 | 8.663+05 | 3.075-08 |
| 2 | 102 | 15.503 | 3.432+05 | 3.710-08 | 3 | 123 | 15.368 | 5.654+05 | 2.002-08 |
| 2 | 107 | 15.470 | 1.912+05 | 2.057-08 | 3 | 129 | 15.196 | 2.146+05 | 1.486-08 |
| 2 | 110 | 15.432 | 1.563+05 | 1.116-08 | 3 | 134 | 15.058 | 6.979+05 | 3.558-08 |
| 2 | 112 | 15.405 | 3.331+05 | 2.370-08 | 5 | 40 | 17.567 | 1.784+05 | 1.376-08 |

| (b) Electric octupole (E3) transitions | | | | | | | | | |
|----------------------------------------|-----|----------------|---------------|---------------|-----|-----|----------------|---------------|---------------|
| i | j | λ_{ij} | A_{ji}^{E3} | f_{ij}^{E3} | i | j | λ_{ij} | A_{ji}^{E3} | f_{ij}^{E3} |
| 1 | 44 | 15.668 | 9.998+04 | 1.472-08 | 2 | 107 | 15.470 | 9.715+04 | 1.046-08 |
| 1 | 46 | 15.646 | 2.078+05 | 2.288-08 | 2 | 113 | 15.384 | 2.569+05 | 2.735-08 |
| 1 | 49 | 15.605 | 2.649+05 | 3.869-08 | 2 | 121 | 15.333 | 1.483+05 | 2.091-08 |
| 1 | 50 | 15.598 | 3.122+05 | 3.416-08 | 2 | 134 | 15.011 | 1.183+05 | 1.199-08 |
| 1 | 57 | 15.410 | 2.348+05 | 3.344-08 | 3 | 82 | 15.734 | 1.466+05 | 1.088-08 |
| 1 | 59 | 15.405 | 1.503+05 | 1.604-08 | 3 | 85 | 15.708 | 2.682+05 | 2.481-08 |
| 1 | 60 | 15.392 | 1.117+05 | 1.190-08 | 3 | 95 | 15.628 | 2.140+05 | 1.175-08 |
| 1 | 61 | 15.380 | 4.066+05 | 5.767-08 | 3 | 96 | 15.623 | 1.892+05 | 1.384-08 |
| 1 | 70 | 15.155 | 1.417+05 | 1.464-08 | 3 | 100 | 15.579 | 3.972+05 | 1.445-08 |
| 2 | 72 | 15.770 | 1.314+05 | 1.959-08 | 3 | 102 | 15.553 | 2.398+05 | 1.305-08 |
| 2 | 74 | 15.761 | 1.731+05 | 2.578-08 | 3 | 109 | 15.491 | 2.891+05 | 2.600-08 |
| 2 | 77 | 15.744 | 1.067+05 | 1.190-08 | 3 | 111 | 15.461 | 3.844+05 | 3.444-08 |
| 2 | 82 | 15.683 | 1.827+05 | 2.695-08 | 3 | 121 | 15.382 | 2.578+05 | 1.829-08 |
| 2 | 93 | 15.590 | 2.410+05 | 2.634-08 | 3 | 122 | 15.376 | 3.466+05 | 1.843-08 |
| 2 | 106 | 15.483 | 2.385+05 | 3.428-08 | 3 | 129 | 15.196 | 1.467+05 | 1.016-08 |

and for the magnetic quadrupole (M2) transitions:

$$A_{ji} = \frac{1.4910 \times 10^{13}}{\omega_j \lambda_{ji}^5} S^{M2} \quad \text{and} \quad f_{ij} = \frac{2.236 \times 10^{-3}}{\lambda_{ji}^3 \omega_i} S^{M2}. \quad (5)$$

In Table 2 we present transition energies/wavelengths (λ in Å), radiative rates (A_{ji} in s^{-1}), oscillator strengths (f_{ij} , dimensionless), and line strengths (S in a.u.), in length form only, for all 2676 electric dipole (E1) and 3418 magnetic quadrupole (M2) transitions among the 134 levels of Fe XVI. The indices used

to represent the lower and upper levels of a transition have already been defined in Table 1. Similar results for 3942 electric quadrupole (E2) and 3023 magnetic dipole (M1) transitions are listed in Table 3. Furthermore, sometimes the electric octupole (E3) transitions are also observed, and are required in plasma modelling. Therefore, the A - and f -values for all 3187 E3 transitions are also listed in Table 2. However, a majority of the E3 (and M2) transitions have very small oscillator strengths. Therefore, in Table 4 we have listed the A - and f -values of all those M2 and E3 transitions, whose f -values are $\geq 10^{-8}$. Since no other similar results are available in the literature, we hope that the presently reported A -values for all types of transitions will be useful in plasma modelling.

In the absence of other similar A -values being available in the literature with which to compare, it is difficult to assess the accuracy of our present results. A general criterion to assess the accuracy is to compare the length and velocity forms of the f - or A -values. Before we discuss these we would like to remind readers that such comparisons are only desirable, but are *not* a fully sufficient test to assess accuracy, as different calculations (or combinations of configurations) may give comparable f -values in the two forms, but entirely different results in magnitude – see Aggarwal et al. (2007) for further details and comparisons.

Since the electric dipole (E1) transitions are comparatively more important, we focus on the accuracy assessments of their A -values alone. Among the stronger transitions (i.e. $f \geq 0.01$), the length and velocity forms differ by over 20% for 267 (10%) transitions, whereas only 49 (2%) transitions have differing A -values of over 50%. Among these, four transitions, namely 25–66, 57–118, 65–115 and 66–128, have the ratio $A_L/A_V = f_L/f_V$ with values of up to 10. Therefore, based on a satisfactory agreement between the two forms, we may state that the accuracy of our listed A -values is better than 20% for a majority of the strong transitions. Differences between the two forms for weaker transitions ($f < 0.01$) can sometimes be several orders of magnitude. Particularly noteworthy are 5 transitions, namely 27–155 ($f \sim 10^{-7}$), 38–81 ($f \sim 10^{-8}$), 65–99 ($f \sim 10^{-6}$), 69–70 ($f \sim 10^{-7}$), and 76–120 ($f \sim 10^{-8}$), for which the ratio A_L/A_V is very large. However, all such transitions are very weak, and hence sensitive to mixing coefficients, but do not affect the overall accuracy of the calculations. Before concluding we make one more comparison with our calculations from FAC.

A comparison made between the A -values from GRASP and FAC (not listed in Table 2) indicates excellent agreement for a majority of the strong transitions. However, 143 (5%) of the transitions have f -values differing by over 20%, whereas only 42 (1.6%) transitions differ by over 50%. Among these, only three transitions, namely 34–88, 35–88 and 41–88, have the f -values differing by an order of magnitude. Considering that the calculations have been performed for a large number of transitions, such differences for a few transitions do not affect the overall accuracy of our results, and we may state with confidence that our listed A -values are probably accurate to better than 20% for a majority of the strong transitions.

4. Lifetimes

The lifetime τ for a level j is defined as follows:

$$\tau_j = \frac{1}{\sum_i A_{ji}}. \quad (6)$$

Since this is a measurable parameter, it provides a check on the accuracy of the calculations. However, to our knowledge no measurement of lifetimes for the levels of Fe XVI are so far available in the literature. Nevertheless, in Table 5 we list our calculated lifetimes, which *include* the contributions from four types of transitions, i.e. E1, E2, M1 and M2. These results may be helpful for future comparisons with theory or experiments. Most of the τ values are very small (\sim ps), but the lifetime of the $2p^5 3s 3p \ ^4D_{7/2}$ level is $\sim \mu$ s, and therefore it may be comparatively easier to measure.

5. Conclusions

In the present work, results for energy levels, radiative rates, oscillator strengths, and line strengths for all transitions among the lowest 134 levels of Fe XVI have been presented. Additionally, results for radiative rates have been presented for five types of transitions, namely E1, E2, E3, M1 and M2. A complete set of results are likely to be useful for the modelling of a variety of plasmas.

Based on the comparison made between the length and velocity forms of the oscillator strengths, as well as the comparison between the GRASP and FAC calculations, we assess that our radiative rates are accurate to better than 20% for a majority of strong transitions. Similarly, based on the comparison between the GRASP and FAC calculations, as well as with the available compiled experimental results, our energy levels are assessed to be accurate to better than 1%. However, the level designations are likely to fluctuate for a few of them, for which further work is recommended.

Lifetimes for all excited levels of Fe XVI are listed, but no comparisons could be possible due to the paucity of other theoretical or experimental results.

Our calculations for collision strengths, and subsequently for excitation rates, are in progress. These calculations, being highly computationally demanding, will take a considerably long time to conclude. We hope to present our results for these highly desired and important parameters in future.

Acknowledgements. This work has been financed by the Engineering and Physical Sciences and Particle Physics and Astronomy Research Councils of the United Kingdom, and FPK is grateful to AWE Aldermaston for the award of a William Penney Fellowship. Finally, we thank an anonymous referee for a critical reading of the earlier version of the paper, and for providing some constructive comments.

References

- Aggarwal, K. M., & Keenan, F. P. 2006, *A&A*, 450, 1249
- Aggarwal, K. M., Tayal, V., Gupta, G. P., & Keenan, F. P. 2007, *ADNDT*, in press
- Bautista, M. A. 2000, *J. Phys. B*, 33, 71
- Berrington, K. A., Eissner, W. B., & Norrington, P. H. 1995, *Comput. Phys. Commun.*, 92, 290
- Cornille, M., Dubau, J., Faucher, P., Bely-Dubau, F., & Blancard, C. 1994, *A&AS*, 105, 77
- Dere, K. P., Landi, E., Young, P. R., & Del Zanna, G. 2001, *ApJS*, 134, 331
- Dyall, K. G., Grant, I. P., Johnson, C. T., Parpia, F. A., & Plummer, E. P. 1989, *Comput. Phys. Commun.*, 55, 424
- Eissner, W., & Seaton, M. J. 1972, *J. Phys. B*, 5, 2187
- Eissner, W., Jones, M., & Nussbaumer, H. 1974, *Comput. Phys. Commun.*, 8, 270
- Gu, M. F. 2003, *ApJ*, 582, 1241
- Norrington, P. H., & Grant, I. P. 2007, *Comput. Phys. Commun.*, in preparation