

Erratum

EROs found behind lensing clusters

**I. Stellar populations and dust properties of optical dropout EROs
and comparison with related objects**

D. Schaerer^{1,2}, A. Hempel¹, E. Egami³, R. Pelló², J. Richard^{2,4}, J.-F. Le Borgne², J.-P. Kneib⁵, M. Wise⁶, and F. Boone⁷

¹ Geneva Observatory, University of Geneva, 51, chemin des Maillettes, 1290 Sauverny, Switzerland
e-mail: [daniel.schaerer;angela.hempel]@obs.unige.ch

² Observatoire Midi-Pyrénées, Laboratoire d'Astrophysique, UMR 5572, 14 Avenue E.Belin, 31400 Toulouse, France

³ Steward Observatory, University of Arizona, 933 North Cherry Street, Tucson, AZ 85721, USA

⁴ Caltech Astronomy, MC105-24, Pasadena, CA 91125, USA

⁵ OAMP, Laboratoire d'Astrophysique de Marseille, UMR 6110 traverse du Siphon, 13012 Marseille, France

⁶ Astronomical Institute Anton Pannekoek, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

⁷ Observatoire de Paris, LERMA, 61 Av. de l'Observatoire, 75014 Paris, France

A&A 469, 47–60 (2007), DOI: 10.1051/0004-6361:20077127

Key words. galaxies: high-redshift – cosmology: early Universe – galaxies: evolution – galaxies: starburst – infrared: galaxies – errata, addenda

The stellar masses derived from the SED fits using *Hyperz* and given in Tables 5–8 are overestimated by a factor $(1 + z)$. Stellar masses derived using Eq. (1) and other quantities are not affected by this error.

Article published by [EDP Sciences](http://www.aanda.org) and available at <http://www.aanda.org> or <http://dx.doi.org/10.1051/0004-6361:20077127e>