A young double stellar cluster in an HII region, emerging from its parent molecular cloud^{★,★★}

J. L. Yun¹, A. A. Djupvik², A. J. Delgado³, and E. J. Alfaro³

¹ Universidade de Lisboa – Faculdade de Ciências Centro de Astronomia e Astrofísica da Universidade de Lisboa, Observatório Astronómico de Lisboa, Tapada da Ajuda, 1349-018 Lisboa, Portugal

e-mail: yun@oal.ul.pt

² Nordic Optical Telescope, Apdo 474, 38700 Santa Cruz de La Palma, Spain

e-mail: akaas@not.iac.es

³ Instituto de Astrofísica de Andalucía, CSIC, Apdo 3004, 18080-Granada, Spain

e-mail: [\[delgado;emilio\]@iaa.es](mailto:[delgado;emilio]@iaa.es)

Received 16 November 2007 / Accepted 22 February 2008

ABSTRACT

Aims. We investigate the star formation occurring in the region towards IRAS 07141-0920 contained in the HII region Sh2-294 (S294). We report the discovery^{***} of a new young double stellar cluster, and we describe its properties.

Methods. High-resolution optical *UBVRI* and *H α* images obtained with ALFOSC mounted on the Nordic Optical Telescope (NOT), near-infrared *JHK_s* images obtained with NOTCam at the Nordic Optical Telescope, and VLT/ISAAC images obtained through the *H₂* (2.12 μ m) filter were all used to make photometric and morphological studies of the point sources and the nebula seen towards Sh2-294. Identification of PMS members was done using three different methods: comparison with isochrones in optical colour–magnitude diagrams and detection of either near-IR excess or *H α* emission.

Results. The optical images reveal an emission nebula rich in morphological details. The nebula is composed mainly of ultraviolet scattered light and of *H α* emission with local morphological features conveying an impression of strong dynamical complexity. Contrasting with the bright parts of the nebula, opaque patches are seen. They appear elongated, filamentary, or clumpy. Our optical photometry confirms that the identification of the illuminator of the nebula is likely to be a B0.5V star located at a distance of about 3.2 kpc. Our high-resolution near-IR images reveal an embedded cluster, extending about 2 pc and exhibiting sub-clustering: a denser, more condensed, subcluster surrounding the optical high-mass B0.5V illuminator star; and a more embedded, optically invisible, subcluster located towards the eastern, dark part of the nebula and including the luminous MSX source G224.1880+01.2407, a massive protostellar candidate that could be the origin of jets and extended features seen at 2.12 μ m. The double cluster appears to be clearing the remaining molecular material of the parent cloud, creating patches of lower extinction and allowing some of the least reddened members to be detected in the optical images. We find 12 main-sequence (MS) and 143 pre-main sequence (PMS) members. The most massive star fits a 4 Myr, post-MS isochrone. The age of the optically selected PMS population is estimated to be 7–8 Myr. The IR-excess population shows sub-clustering on scales as small as 0.23 pc and is probably much younger.

Key words. stars: formation – stars: pre-main sequence – ISM: HII regions – ISM: dust, extinction – ISM: individual objects: IRAS 07141-0920 – ISM: individual objects: Sh2-294 (S294)

1. Introduction

Galactic nebulae are often clear signposts for regions of star formation due to the proximity of the young stars to the parent material of the molecular cloud that originated the stars. The presence of nebulosity has been used to find and identify star formation regions (e.g. Yun & Clemens 1994; Tapia et al. 1997; Rubio et al. 1998). In addition, young stars are the source of

energetic winds, jets, or outflows, whose interaction with their surrounding cloud material produces shocks, which are denounced by nebulae with characteristic shapes (e.g. bow shocks, HH objects; Reipurth & Bally 2001) and which ultimately lead to the clearing of the cloud material and emergence of the young stars out of the cloud core.

The morphology of nebulae seen towards star-forming regions can reveal the processes that have been occurring in the interior of the parent cloud. Diffuse reflection nebulae, both in the optical and in the near-infrared, are the result of illumination from bright sources more or less embedded in the cloud. Optical emission nebulae corresponding to excited gas can be detected using narrow-line filters, the most common one being *H α* (e.g. Fich et al. 1990).

Small groups or isolated young low-mass stars have a limited dynamical effect on their surroundings. On the other hand, rich clusters of young stars or high-mass young stars will often produce spectacular nebulae. Massive star formation is currently being addressed via a variety of the new instruments made

* Based on observations made with the Nordic Optical Telescope, operated jointly on the island of La Palma by Denmark, Finland, Iceland, Norway, and Sweden, in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. Based on observations collected at the ESO 8.2-m VLT-UT1 Antu telescope (programme 66.C-0015A).

** Table 1 is only available in electronic form at the CDS via anonymous ftp to [cdsarc.u-strasbg.fr](ftp://cdsarc.u-strasbg.fr) (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/483/209>

*** A new publication (Samal et al. 2007) with a similar analysis of a field in this region appeared when our paper was in the refereeing process.

available in the last years. Unlike the low-mass family of young stars, the relatively short timescale for their accretion and pre-main sequence evolution implies that massive stars reach the main sequence still embedded in their parent molecular cloud cores. On the other hand, their strong hard radiation ionizes their surroundings, creating an ultra-compact HII (UCHII) region, which may represent the earliest recognisable stage in the life of massive stars (Churchwell 2002). The UCHII regions have been identified using $H\alpha$ surveys (e.g. Rodgers et al. 1960), radio surveys, and looking for characteristic IRAS colours (e.g. Wood & Churchwell 1989). More recently, they have been studied using a combination of near and thermal infrared surveys and K -band spectroscopy (Bik et al. 2003, 2006).

The IRAS source IRAS 07141-0920 has colours close to those of ultra-compact HII regions (Wood & Churchwell 1989). It is seen towards a large, extended optical nebula. Rodgers et al. (1960) detected $H\alpha$ emission, and Fich et al. (1990) detected the $H\alpha$ line at $v(\text{LSR})$ of 33.5 km s^{-1} . The source was first cataloged as an HII region by Sharpless (1959) in his catalogue of HII regions (source Sh2-294 or S294). Using UBV photoelectric photometry, Moffat et al. (1979) derived a photometric spectral type of B0.5V for the hottest star in this HII region, which they assume to be the ionizing source. Their estimated distance was 4.6 kpc. Extended radio continuum emission from S294 at 1.46 GHz has been detected and mapped, resulting in a diameter for the HII region of at least $7 \times 7 \text{ arcmin}^2$ (Fich 1993). A strong CO $J = 1-0$ molecular line transition towards IRAS 07141-0920 has also been detected, at a radial velocity $v(\text{LSR})$ of 32 km s^{-1} (Blitz et al. 1982; Wouterloot & Brand 1989). They derived a kinematic distance of 3 kpc, much less than the value of $4.6 \pm 1.5 \text{ kpc}$ derived by optical photoelectric photometry (Moffat et al. 1979). Wouterloot & Brand (1989) also note that the CO line is flat-topped (non-Gaussian), a hint that the molecular gas is not in a quiescent state, possibly with turbulence or other line broadening mechanism present. Because they have a single observation and have not mapped the cloud, there is no estimate of the extent of this molecular cloud seen towards IRAS 07141-0920.

Searches for masers have resulted in non-detections: water masers – Henkel et al. (1986); methanol masers – MacLeod et al. (1998), Codella et al. (1995). Bica et al. (2003) reported the presence of a 2MASS infrared, loose star cluster towards Sh2-294. Samal et al. (2007) provide a multivalength analysis of the Sh2-294 region describing the distribution and nature of the young stellar objects and the morphology of the thermal ionized gas and dust emissions¹. Here we present an optical and near-infrared study of the extended optical nebula and the embedded stellar population seen towards Sh2-294 (S294)/IRAS 07141-0920. We confirm the presence of a young embedded cluster that appears to be actively in the process of destroying its cradle (dissipating the surrounding parent molecular cloud). Most cluster members are still embedded and are either not seen or are very faint in our deep optical CCD images.

Section 2 describes the observations and data reduction. In Sect. 3, we describe the properties of the extended nebula. Section 4 describes the stellar density distribution and the presence of a young embedded cluster. In Sect. 5, we present our photometric results, estimate the reddening, distance, and age of the cluster, and make cluster membership assignments. In Sect. 6, we discuss the spatial distribution of the cluster members

and present a possible star formation scenario for this region. A summary is given in Sect. 7.

2. Observations and data reduction

2.1. Optical observations

Optical ($UBVRI$ and $H\alpha$) observations were carried out during the night of 2006 November 29 with the NOT (Nordic Optical Telescope), operating at the ORM (La Palma) under very good seeing conditions (0.6 arcsec). The ALFOSC (Andalucia Faint Object Spectrograph and Camera) CCD camera was used in imaging mode (with a plate scale of 0.19 arcsec/pixel). Short and long exposure images were obtained through each band: short exposures with integration times of 10, 10, 5, 5, 5, and 60 s for $UBVRI$ and $H\alpha$ ($FWHM = 3.3 \text{ nm}$) bands, respectively; and long exposures with integration times of 600, 300, 300, 300, 300, and 1200 s for the same bands, respectively.

The observations were reduced with the adequate routines in the IRAF² package. In each of the $UBVRI$ -bands, three long exposures were co-added into a single image. Aperture and (point spread function (PSF) photometry was performed in both the short and the long exposure images. The resulting magnitudes were then joined to produce a final list of instrumental magnitudes.

Frames of a standard field were also secured at different airmasses. The field includes 12 stars in the SA98, around star 652 (Landolt 1992). These observations were used to transform the instrumental magnitudes to the standard system, with the procedure described by Delgado et al. (1998), based upon the assumption of multilinear dependence of standard magnitudes on airmass, instrumental magnitude, and an appropriate colour index.

2.2. Near-infrared observations

2.2.1. NOTCam JHK_s observations

Near-infrared (J , H , and K_s) images were obtained on 2007 January 29 using the Nordic Optical Telescope near-IR Camera/spectrograph (NOTCam)³. The detector was the $1024 \times 1024 \times 18$ micron Hawaii engineering grade array. The wide field camera (0.234"/pix) was used, and the observations were performed with a ramp-sampling readout mode. Every sky position was integrated for 30 s, reading the array every 5 s and using the linear regression result of these 6 readouts. For each filter, 9 dithered sky positions were observed. Because of varying sky conditions, only the best frames were selected, resulting in total on-source integration times of 210, 180, and 150 s for J , H , and K_s , respectively.

The images were reduced using a set of own NOTCam scripts to correct for bad pixels, subtract the sky background, and flatfield the images. Differential twilight flats taken a few nights apart were used to correct for the pixel-to-pixel variations of the response. The selected images were then aligned, shifted, trimmed, and median-combined to produce a final JHK_s image of $3.7' \times 3.4'$ size.

² The Image Reduction and Analysis Facility (IRAF) is distributed by the National Optical Astronomical Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

³ See URL <http://www.not.iac.es/instruments/NOTCam/> for more information on NOTCam.

¹ This article was published while our manuscript was in the refereeing process.

Point sources were extracted using *daofind* with a detection threshold of 6σ , and a few additional sources were added in by hand. Aperture photometry was made with a small aperture (radius = 4 pix, which is about the measured *FWHM* of the PSF) and aperture corrections, found from 5 bright and isolated stars in each image, were used to correct for the flux lost in the wings of the PSF. The error in determining the aperture correction was <0.02 mag in all cases. These errors were added to the value *MERR*, which is created by the IRAF task *phot*. A total of 491, 496, and 494 sources were found to have fluxes in *J*, *H*, and *K_S*, respectively, and errors $\sigma_{K_S} < 0.35$ mag in the *K_S* band (the noisiest and least sensitive). All these are included in the table of photometric results for completeness (Table 1, see below), but we include only the 444 NOTCam sources with $\sigma_{K_S} < 0.2$ mag in our analysis in this paper.

The brightest star in the field S294-B0.5 (ID 717) had peak pixel values outside the linear range of the detector both in the *H* and *K_S* bands. We therefore used only the first of the 6 ramp-sampling readouts, with an effective integration time of 5 s for this source.

We used the 2MASS All-Sky Release Point Source Catalogue (Skrutskie et al. 2006; Cutri et al. 2003) to calibrate our observations both in terms of photometry and astrometry. The plate solution was found using 27 isolated and bright 2MASS stars over the FOV and the *iraf* task *ccmap* with the “*tnx*” projection allowing for geometric distortion and a 3rd order polynomial with full cross terms. NOTCam XY positions were then transformed to RA and Dec, and all 2MASS sources were matched to all NOTCam sources using a search radius of $0.36''$, which corresponds to three times the positional accuracy from the plate solution. We excluded all 2MASS sources that do not have photometric quality flag from A to D in all bands. Also a few multiples and one clearly variable star were excluded, leaving a total of 85 sources that could be compared in terms of fluxes and positions. The *JHK_S* zeropoints were determined using 2MASS stars brighter than *K_S* = 13.5 mag after excluding 2 cases of possible variability (i.e. 17 stars). The standard deviation of the offset between NOTCam and 2MASS photometry is 0.07, 0.05, and 0.05 mag in *J*, *H*, and *K_S*, respectively, for sources with *K_S* < 13.5 mag. We estimate the completeness limit of the observations to be roughly 18.5, 18.0, and 17.5 mag in *J*, *H*, and *K_S*, respectively.

2.2.2. ISAAC H₂ observations

Near-infrared observations, using the H₂ filter centred at the $\nu = 1-0$ line at $2.12 \mu\text{m}$, were conducted on 2000 November 10 using the ESO Antu (VLT Unit 1) telescope equipped with the short-wavelength arm (Hawaii Rockwell) of the ISAAC instrument. The ISAAC camera (Moorwood et al. 1998) contains a 1024×1024 pixel near-infrared array and was used with a plate scale of $0.147 \text{ arcsec/pixel}$ resulting in a field of view of $2.5 \times 2.5 \text{ arcmin}^2$ on the sky. At each of 5 different jitter positions (with the length of the jitter box equals to about half the field of view), one image with 60 s exposure time were taken.

Standard procedures for near-infrared image reduction were applied (e.g. Yun & Clemens 1994) resulting in a final mosaic image for each band. A sky image was computed for each dithered frame. Each frame was then sky-subtracted and flat-fielded using master sky flats. The final clean images were corrected for bad pixels while constructing the final mosaic which covers about $5 \times 5 \text{ arcmin}^2$ on the sky. Due to the presence of significant field distortion in these ISAAC images (two pixels at the edges and 2.5 pixels in the corners), the individual

Fig. 1. $H\alpha$ image towards IRAS 07141-0920 covering about $6 \times 6 \text{ arcmin}^2$ ($5.4 \times 5.4 \text{ pc}^2$, for our estimated distance of 3.2 kpc). Notice the bright nebula and the patchy and wispy dark cloud material.

frames were corrected before they were put together to form the mosaic. The correction for field distortion was performed using IRAF/GEOTRAN, together with the adequate correction files provided by ESO at their web page. The central, co-added region, with enhanced signal-to-noise ratio covers about $2.5 \times 2.5 \text{ arcmin}^2$ on the sky.

3. Extended nebula

3.1. Morphology and colours

Figure 1 shows the $H\alpha$ image centred at the IRAS 07141-0920 position and covering about $6 \times 6 \text{ arcmin}^2$. The high-resolution of ALFOSC allied to the very good seeing during the observation produced an image containing a wealth of detail. It reveals a large, extended nebula that appears to be illuminated by one or more objects possibly seen in this $H\alpha$ image.

Contrasting with the bright parts of the nebula, dusty, opaque patches are seen, especially on the northern and eastern sides. These dark clouds appear to be in the process of being shredded or blown apart by stellar winds, jets, or outflows phenomena. They appear elongated, filamentary, wispy, clumpy, or knotty. The overall image conveys an impression of streams, flows, turbulence, cavities, so a region of great dynamical complexity.

The main contributions to the nebula brightness are continuum ultraviolet scattered light (mostly contained in the *U*-band) and $H\alpha$ emission. This is shown in the colour composite image shown in Fig. 2. This figure shows a *U*-band (blue), *R*-band (green), and $H\alpha$ (red) colour composite image towards IRAS 07141-0920. With this colour code, the blue extended emission traces scattered light, while red extended emission traces $H\alpha$ nebular emission from excited gas. The green colour (*R*-band continuum) traces essentially point sources. Much of the nebular light seen is composed of $H\alpha$ emission. The contribution of continuum scattered light is stronger near the centre of the nebula where the illuminator is most likely to reside.

Fig. 2. *U*-band (blue), *R*-band (green), and $H\alpha$ (red) colour composite image towards IRAS 07141-0920, covering about 6×6 arcmin². North is up and east to the left. Blue extended emission traces scattered light. Red extended emission traces $H\alpha$ nebular emission from excited gas. The green colour (*R*-band continuum) traces essentially point sources.

3.2. The illuminator star

Given the presence of the HII region and the large, extended nebula, at least one O or B star is responsible for the ionization and excitation of the nebular gas. To identify the illuminator(s), we note that it should be located close to the central region of the nebula which also happens to be where the blue scattered light is brightest.

Because dark lanes of obscuring material can be seen in silhouette against the bright nebula (mostly in the northern part of the nebula), we infer that the illuminator (or illuminators) must be partially embedded in the cloud. The illuminator should not be too deeply embedded, with its light escaping from “holes” (regions of lower optical depth, lower extinction) after scattering off the walls of cavities on the near side of the cloud and exciting or ionizing the gas at the surface of the cloud.

In this central region, the brightest source is the star marked with the letter “B0.5V” in Fig. 3, and we identify this object as being the illuminator of this nebula. This object coincides with star No. 4 of Moffat et al. (1979), who first identified it as the hottest star in the region and thus the one responsible for the HII region. This identification has been confirmed by Samal et al. (2007). From here on, we refer to this object as S294-B0.5V.

Additional arguments support this conclusion. Figure 3 reveals nebular structures that display illuminated rims and shadows, implying that the incident light is along the directions indicated by the two solid line arrows. In all cases, the only bright object located within the bright scattered light region is object S294-B0.5V. In Sect. 5, we discuss optical and infrared photometry results for this object, and the parameters of the associated cluster.

Fig. 3. Zoomed-in $H\alpha$ image of the nebula covering 1.9×1.9 arcmin² of the southern and central regions of the nebula. B0.5V is the illuminator of the nebula. Notice the rim-bright features, some of them indicated by arrows. North is up and east to the left.

4. Stellar density distribution: a young stellar cluster

Together with this massive B0.5V star, this region hosts the birth of a young stellar cluster associated to the massive star. Figure 4 presents the NOTCam *JHK_s* colour composite image obtained towards IRAS 07141-0920. The optical nebula that dominated much of the *UBVR* and $H\alpha$ images is absent. Instead, a large number of stars are seen in the region of the optical nebula but also to the east, where a dark patch of extinction is present in the optical images.

A higher angular resolution image is provided by the VLT ISAAC H_2 image seen in Fig. 5. The stars appear to form two distinct groups or sub-clusters: a western group (the western sub-cluster), which includes S294-B0.5V, less extincted and located towards the optical nebula; an eastern group (the eastern sub-cluster), redder and located towards a dark optical region.

The eastern part of Fig. 5 also shows elongated, jet-like, or filamentary emission which suggest flows or material being blown away, most likely by powerful stellar winds, or affected by ionization fronts.

The presence of the two groups of stars is seen more clearly in the contour plot of the stellar surface density, shown in Fig. 6, obtained from the H_2 mosaic image. Two peaks of the stellar surface density can be identified. The stronger peak is located in the central region of the optical nebula, and we name it the western stellar density peak. It contains S294-B0.5V, the illuminator of the optical nebula, and a steep slope in the number of stars surrounding it. Thus, in a small region around the massive S294-B0.5V star, there is a large number of young lower-mass stars (see also the zoomed-in upper-right panel of Fig. 5). The stellar surface density at this western peak is about 14 times higher than the value of the sky background stellar surface density. The second peak is located in the dark region to the east of the nebula, so we name it the eastern stellar density peak. Together, the two peaks correspond to a young stellar cluster

Fig. 4. *J* (blue), *H* (green), and *K_s* (red) colour composite image towards IRAS 07141-0920. The image covers about 3.4×3.4 arcmin² (dashed-line box in Fig. 7). North is up and East to the left. The white cross indicate the position of the IRAS source ((0, 0) position of Fig. 1).

Fig. 5. Central region of the H₂ mosaic image towards IRAS 07141-0920. The axes give the coordinates relative to the IRAS source. The two zoomed-in views reveal elongated emission in the eastern part, and the enhanced concentration of stars around S294-B0.5V.

showing sub-structuring on a scale of about 2 pc, the size of the region. The position of the IRAS source, at offset (0, 0), is located between the two peaks with the IRAS uncertainty ellipse in the east-west direction.

Interestingly, the map of 1.46 GHz continuum emission (Fich 1993) also displays two peaks located approximately at the same positions of the peaks of the stellar surface density. However, the stronger radio continuum peak is associated with the eastern peak, as it coincides with the more embedded sub-cluster, invisible in the optical images. The 1280 MHz radio continuum map of Samal et al. (2007) contains a peak coincident with the stronger 1.46 map, and thus the eastern stellar density peak.

Fig. 6. Contour plot of the stellar surface density of the H₂ image. Two peaks of the stellar surface density can be identified: the strongest, in the west, is the “western peak” (where the illuminator is located, see text); the second one, in the east, is the “eastern peak”. The axes give the coordinates relative to the IRAS source. The lowest contour level is 9 sigma above the sky background stellar surface density. The highest contour level is about 14 times the value of the sky background stellar surface density.

5. Photometric results

5.1. Standard calibration

As mentioned above, observations of standard stars in the Landolt catalogue (Landolt 1992) were secured for standard optical calibration purposes. On the other hand, when additional reliable standard values are available for a region of the sky, an advantageous calibration procedure is at hand, by means of direct correlation between the instrumental values and those published standard values (Delgado & Alfaro 2000; Delgado et al. 2007). The only published photometric study including stars of this region (Moffat et al. 1979; MFJ in the following) contains *UBV* indices for 15 stars included in the field of our *UBVRI* observations. Of these 15 stars, we have good, non-saturated photometry for 14 stars. The correlation with the values from this study can be performed with some caution. Several stars included in their photoelectric study are located in the central region of the nebula, and their diaphragm photometry could be affected by the nebula brightness and some undetected crowding. In particular, this could be the case of the most luminous star, S294-B0.5V (MFJ No. 4), which is surrounded by several less luminous stars, more clearly visible in NIR frames. To avoid the influence of these effects, we used, for the correlation, 8 out of the 14 common stars, which are located outside the small central region marked in Fig. 7. This area encloses the concentration of stars seen in the *UBVRI* frames and, more clearly, in the NIR frames. The resulting correlation between *UBV* indices is very well-defined, and it provides a set of standard, MFJ-derived, $(U - B)$, $(B - V)$, and V values.

These values derived from photoelectric photometry are presumably closely tied to the original Johnson system. They are, however, systematically different from those obtained through calibration with Landolt standards. For the 8 stars used in our correlation, we obtained $V_{\text{Landolt}} - V_{\text{MFJ}} = -0.04 \pm 0.02$; $(U - B)_{\text{Landolt}} - (U - B)_{\text{MFJ}} = -0.07 \pm 0.03$; $(B - V)_{\text{Landolt}} - (B - V)_{\text{MFJ}} = 0.01 \pm 0.03$. These differences between *UBV* Johnson

Fig. 7. Schematic representation of the field covered by the *UBVR* observations. Only stars detected in the *V*-band are plotted. The smaller central square (solid-line) defines what we name *core-stars* (see text). The larger dashed-line box indicates the field of the *JHK_s* NIR images.

and Landolt systems were previously found and discussed by Delgado & Alfaro (2000). In our case here, the ZAMS line that we use for colour excess and distance determination (Schmidt-Kaler 1982) is likely to be more adequate for Johnson values, and we adopt the Johnson-derived set of standard indices. As for the calibration of $(V - R)$ and $(V - I)$ indices, it is obtained from the Landolt standard star measurements, following the multilinear model described by Delgado et al. (1998). The uncertainties of all the standard indices obtained, given by the root mean squared deviation of the residuals, amount to 0.02, 0.03, 0.03, 0.06, and 0.03, for V , $(U - B)$, $(B - V)$, $(V - R)$, and $(V - I)$, respectively.

5.2. Colour excess, distance, and age

The standard magnitudes of all the stars detected in our *UBVR* $H\alpha$ images are listed in Table 1. In this table, Cols. (1)–(3) identify the stars giving, respectively, the running ID number, and the equatorial coordinates (RA and Dec, epoch 2000); Cols. (4) through (14) give the photometric indices V , $(U - B)$, $(B - V)$, $(V - R)$, $(V - I)$, R , I , $(R - H\alpha)$, J , H , and K_s , and their corresponding uncertainties in parenthesis. Column (15) gives the cluster membership assignments from optical photometry (see Sect. 5.3). Column (16) indicates whether the source has near-IR excess. Finally, Col. (17) contains the identifications of the stars if it appears in the WEBDA data base⁴.

A schematic chart of the field covered by our optical observations is plotted in Fig. 7, where all stars with V measurements are shown. In the following discussion, we first concentrate on the stars located in the smaller square region at the centre of the

field, indicated in Fig. 7. For simplicity, we refer to these stars as the *core stars*.

In Fig. 8, we plot the colour–colour (CC) diagram $(U - B)$ vs. $(B - V)$, and the colour–magnitude (CM) diagrams V vs. $(B - V)$ and V vs. $(V - I)$ of all the observed stars. Among these stars, the core stars from Fig. 7 are represented by open circles and open squares. The latter appear to be part of a group of foreground stars at very low reddening. In the CC diagram (Fig. 8), the ZAMS line (Schmidt-Kaler 1982) shifted for a colour excess of $E(B - V) = 0.1$ (dashed line) fits well this group of stars. These foreground stars are homogeneously distributed in the observed field, and thus include some core-stars, namely, those we represent by squares.

On the other hand, the open circles appear to be a sequence of reddened core-stars. In fact, the solid line is the previous ZAMS line shifted to a location defined by $E(B - V) \approx 1.3$, with a reddening slope $\alpha = 0.72$. With these parameters the solid line fits well the shape of the most defining part of the sequence, namely that close to late B and early A type stars. Thus, we assume that the stars represented with open circles are very likely to be reddened stars associated with the nebula. They correspond to the visible members of the young stellar cluster, better identified in the spatial distribution seen in the NIR images. We refer to them as *nebula-associated stars*, or just simply *nebula stars*. The two stars represented by crossed circles, which are indeed among the core-stars, are not considered in principle MS nebula stars, judging from their locations in this diagram and in the CM diagrams. They belong to a conspicuous group of stars, which could be also associated with the nebula, and showing at the same time a possible excess in the $(U - B)$ index. This is a feature associated with accretion in forming stars, as discussed by Gullbring et al. (1998, 2000). A more detailed discussion of these stars is postponed to the discussion of cluster membership.

We are therefore left with 8 stars (listed in Table 2), that we consider as the most probable nebula stars. Table 2 also provides a quantitative estimate of the colour excess $E(B - V)$ for the nebula stars, using the value of the reddening slope $\alpha = 0.72$. The calculated value for the nebula illuminator, star MFJ No. 4 (No. 717 in Table 2), leads to a spectral type estimate of B0.5V for this star, in good agreement with MFJ and close to the spectroscopic type estimated by Russeil et al. (2007).

With the values of the colour excesses listed in Table 2, determination of the visual absorption is possible, once the visual absorption coefficient $R_V \equiv A_V/E(B - V)$ is obtained. Here we dispose of *JHK_s* observations, which allow the estimate of R_V for individual stars with well-defined $E(B - V)$ colour excess. For a particular star, an $E(V - K)$ value is calculated in the $V - K$ vs. $B - V$ diagram, by shifting the ZAMS line (Koornneef 1983) in the $B - V$ axis with the known $E(B - V)$, and calculating the distance to the line in the $V - K$ axis. For our nebula stars, we then obtain R_V values through the relation $R_V(K) = 1.12 * E(V - K)/E(B - V) + 0.02$ (Fitzpatrick 1999). These values provide an average $R_V(K) = 4.5 \pm 0.6$ from our NIR observations, and $R_V(K) = 4.3 \pm 0.6$ with K magnitudes from 2MASS. These high values of the absorption coefficient are, however, most probably affected by the NIR properties of the nebula stars. At least two of the nebula stars in Table 2 (520 and 660) show clear signs of NIR excess. In the presence of probable, or even suspected K_s -band excess, the use of the above relation is not adequate. On the other hand, a more reliable estimate of the absorption coefficient can be obtained from $(V - I)$ photometry (Guetter & Turner 1997; GT97 in the following). The simple relation used by these authors, $R_V(I) = 2.40 \times \frac{E(V - I)}{E(B - V)}$, is valid for $(V - I)$ indices in the Kron-Cousins system.

⁴ This research has made use of the WEBDA database, operated at the Institute for Astronomy of the University of Vienna.

Fig. 8. $(U - B)$ – $(B - V)$ CC diagram and V – $(B - V)$, V – $(V - I)$ CM diagrams. Core stars (see text) are shown as squares and circles. The ZAMS line is plotted twice in the CC panel, with slope $E(U - B)/E(B - V) = 0.72$: for $E(B - V) = 0.1$ (dashed line) and for $E(B - V) = 1.3$ (solid line). The second line fits well the sequence of reddened core-stars (circles), named nebula stars. The two crossed circles are core stars rejected as nebula stars (see text). The dotted line in the CC diagram encloses a region with stars possibly associated with the nebula and exhibiting $(U - B)$ excess. Note also the distinctly deeper magnitude limits as we move towards redder bands, from U to I .

Table 2. Parameters for nebula stars.

N^a	$E(B - V)$	$E(V - I)$	$R_V(I)$	DM_o
520	1.18	1.66	3.38	13.4
580	1.27	1.88	3.56	12.7
660	1.31	2.32	4.26	12.5
694	1.35	1.87	3.33	12.0
697	1.28	1.79	3.37	13.2
717	1.20	1.75	3.51	12.6
733	1.23	1.86	3.63	12.1
755	1.28	1.83	3.43	11.9

^a ID number given in Table 1 (only available on-line).

The resulting R_V values are adopted here. In Fig. 9, we have plotted the CC diagram $(V - I)$ vs. $(B - V)$. Symbols have the same meaning as in Fig. 8. The line plotted represents the ZAMS (Schmidt-Kaler 1982), shifted to account for the calculated average value of $E(B - V)$ and a reddening slope $\frac{E(V - I)}{E(B - V)} = 1.44$. We can see that this value of the slope is adequate for all nebula stars. The resulting values of $E(V - I)$ and $R_V(I)$ are listed in Table 2.

The values of the distance modulus obtained are also listed in Table 2. The average values $E(B - V) = 1.26 \pm 0.06$, and $DM = 12.5 \pm 0.5$ are obtained for the nebula stars listed in Table 2. Even with the large dispersion, this value is distinctly lower than the distance given by MFJ, but it agrees with the revised value of 3.2 kpc obtained by Russeil et al. (2007). The greater distance obtained by MFJ stems from their adopted value for the absorption coefficient, around $R_V = 3.1$, too low when compared to our calculated values. The presence of R_V values both lower and higher than those describing the average galactic extinction have been reported in the literature; see Turner (1994), Delgado et al. (1998) and references therein. GT97 find a similar value and discuss their significance as representative of the extinction law in their line-of-sight. They argue that the peculiar absorption coefficient indeed reflects the extinction law in their line of sight, rather than being related to the material physically associated to the stars. In our case, the dispersions of the

Fig. 9. $(V - I)$ vs. $(B - V)$ colour-colour plot. Symbols for core-stars and nebula stars are as in Fig. 8. The ZAMS line is shifted for a colour excess of $E(B - V) = 1.26$, and a reddening slope $E(V - I)/E(B - V) = 1.44$.

averages for reddening slope and absorption coefficient are higher than theirs, partly because of larger errors in our photometry, partly because some of the peculiarities of the extinction in the region could indeed be related to specific properties of the intracluster or circumstellar material in our region. This possibility is suggested by the higher $R_V(K)$ values found in our case than the ones by GT97, whereas the $E(V - I)$ excesses and $R_V(I)$ values are similar to theirs.

We notice that, in general, the presence in a region of R_V values higher than the standard value ($R_V = 3.1$) adopted for the average galactic extinction, might be caused by absorption due to material associated to the observed stars, and is usually

mentioned in connection with the observation of young star clusters (Turner 1994; de Winter et al. 1997). In line with this reasoning, the total reddening in the direction of the object would be composed of a foreground component that can be assumed to follow a normal reddening law, characterised by $R_V = 3.1$, and a component originating in the intracluster material, associated to the young stars, and responsible for the increase in the R_V value (Pandey et al. 2003; Sharma et al. 2007; Pandey et al. 2008). In the present case, the evidence from our CC diagrams (Figs. 8 and 9) indicates that only a small part (0.1 to 0.2 dex in $E(B - V)$) of the reddening affecting the nebula stars is to be considered as caused by foreground reddening. Assuming that the absorption due to this part is described by $R_V = 3.1$ would have little influence on the final distance estimate.

5.3. Cluster membership: optical photometry assignments

Using the procedures described in detail by Delgado et al. (2007), we analysed whether each star is likely to belong to the young cluster. This membership analysis is applied to the optical *UBVR* observations. It consists of calculating colour excesses and distance moduli for all stars with respect to the ZAMS and PMS isochrones. A star is then considered a cluster member candidate (MS or PMS), if its colour excess and distance modulus, with respect to any of the comparison lines, coincide (within the errors) with the corresponding average values calculated for a previously selected number of bona fide unevolved MS members. In the present calculation, the MS members are the 8 nebula stars mentioned above, which provide the values listed in Table 2, and the corresponding average value of colour excess and distance modulus of the cluster.

The reference lines used in the comparison are the ZAMS line (Schmidt-Kaler 1982), and the PMS isochrones from two different models: Palla & Stahler (1999); and Siess et al. (2000) (in the following, P99 and S00, respectively). When the distance modulus for a star is calculated from comparison with PMS isochrones, we obtain a PMS membership guess for it. The comparison can be performed in several CM diagrams. Some differences occur when using P99 or S00 models. The number of stars that fulfil the membership criteria are different: 58 stars turn out to be candidate PMS members with respect to P99 (assignment in 2 or 3 CM diagrams), whereas only 38 fulfil the requirements when compared to S00. The bulk of this difference arises in the number of stars assigned as members around the spectral types A and F. Assigned PMS members of spectral types around G and later coincide for both isochrone models. The difference stems from the S00 isochrones being more luminous (around 0.5 brighter in M_V around A and F spectral types) than the P99 isochrones. As a result, the average age of the PMS candidate members is also younger when measured with the P99 isochrones. On the other hand, practically all stars assigned with respect to S00, are also assigned according to P99 models. The assignments listed in Table 1 were considered with respect to either of the two models. The entries in Col. (16) are: 0 for no assignment, 1 for MS members, 2 for PMS members assigned in 2 CM diagrams, and 3 for PMS members assigned in 3 CM diagrams.

The CM diagrams in Fig. 10 show the membership assignments obtained in the analysis. In these plots, we show the post-MS isochrone of $\log \text{Age}(\text{yr}) = 6.6$ (Girardi et al. 2002), shifted to account for the parameters of colour excesses derived for S294B0.5V. PMS isochrones of 1 and 10 Myr from P99, shifted to account for the average parameters derived above, are also plotted. Stars indicated with larger dots in Fig. 10 are those

Fig. 10. CM diagrams V vs. $(B - V)$ and V vs. $(V - I)$, with indication of cluster candidates. Circles are MS and post-MS members. They include the nebula stars in Table 2. Large dots are PMS members assigned in the two CM diagrams V vs. $(V - R)$, and V vs. $(V - I)$. Squares are PMS members assigned in three CM diagrams, V vs. $(B - V)$, V vs. $(V - R)$ and V vs. $(V - I)$. The $\text{LogAge}(\text{yr}) = 6.6$ (≈ 4 Myr) post-MS isochrone of Girardi et al. (2002) is plotted (solid line), shifted to account for the colour excess and distance derived for star S294B0.5V (MFJ No. 4). The Palla & Stahler (1999) isochrones for 1 and 10 Myr are also plotted (dashed lines), shifted for the average colour excess and distance modulus computed from the values of the nebula stars (see Table 2). Isochrones are labelled with their ages in Myr.

with a PMS membership assignment in two CM diagrams (V vs. $(V - R)$ and $(V - I)$). This means (Delgado et al. 2007) that both colour excess and distance modulus calculated with respect to some PMS isochrone between 1 and 10 Myr, coincide (within errors) with the average values obtained from the nebula stars. Similarly, stars assigned as members in three CM diagrams are marked as squares. We detect PMS candidates down to masses around $0.7 M_{\odot}$. The average age value for these PMS candidates can be taken as an age estimate for the possible PMS sequence. The values obtained for stars with membership assignments in at least three CM diagrams, are 7.4 ± 1.3 Myr for P99 isochrones, and 8.3 ± 0.7 Myr for S00 isochrones. These values are higher than the age estimate obtained for the MS members. The significance of this estimate has yet to be assessed. If confirmed, it would indicate a sequential star formation in the region, with the most massive stars being formed last. This kind of trend is also suggested by the results on other regions of active formation; see Mamajek et al. (2002); Sartori et al. (2003); and Delgado et al. (2007).

5.4. Cluster members with near-IR excess

The $J - H/H - K_S$ diagram for 437 NOTCam sources with $\sigma_{K_S} < 0.2$ mag is shown in Fig. 11. The loci of giant, supergiant, and main-sequence stars (Koornneef 1983) are indicated with bold curves. We used the $A_{\lambda} \propto \lambda^{-1.7}$ parametrization of the NIR extinction law (Whittet 1988) to calculate a reddening slope of 1.6 in the $J - H/H - K_S$ diagram for the NOTCam JHK_S filter pass bands at $\lambda \lambda$ 1.247, 1.632, and $2.140 \mu\text{m}$, respectively. Most stars are located around the reddened loci of main-sequence and giant stars.

Fig. 11. $J - H/H - K_S$ diagram for the 437 sources with $\sigma_{K_S} < 0.2$ mag. The reddening slope of 1.6 results from applying the $A_\lambda \propto \lambda^{-1.7}$ parametrization of the NIR extinction law and the reddening vector of an A0 star is drawn (dashed line). The bona-fide IR-excess sources (big dots) are located more than 2σ to the right of and below the reddening line and the probable IR-excess sources (triangles) have a separation from the reddening line between 1σ and 2σ of the individual errors in the $J - H$ and $H - K_S$ indices. Large circles mark nebula stars listed in Table 2.

Sources located right- and downwards of the reddening vector of an A0 star have excess emission in the infrared, a clear sign of hot circumstellar dust. We define as bona-fide IR-excess sources those separated from the reddening vector down and right by more than 2σ of the individual errors in the $(J - H)$ and $(H - K_S)$ indices (48 sources) and as probable IR-excess sources those with a separation between 1σ and 2σ (17 sources). The 8 encircled sources in the figure are the so-called nebula stars discussed in Sect. 5.2. Two of them have clear or probable near-IR excesses (IDs 520, 660) and one (ID 733) is uncertain (see Table 2). In Table 1 the column IREX has the value 0 for no IR-excess, 1 for probable IR-excess, and 2 for bona-fide IR-excess sources. The sources with excess emission in the infrared owing to hot circumstellar dust are candidate young stellar objects (YSOs), either pre-main sequence (PMS) stars with circumstellar discs or protostars embedded in a more spherical dust distribution. We cannot separate clearly PMS stars from protostars only via near-IR photometry, but because of the different time scales of these populations, the majority of the IR-excess YSOs are probably PMS stars with discs.

There are six very red objects near the centre of the embedded cluster that are not detected in the J band. The lower limit $(J - H)$ values suggest that three of these have IR excesses and are probably very embedded YSOs in their earliest evolutionary phase. These have been given the IREX indicator value 1 in Table 1.

Only 144 (272) of the 444 near-IR sources have V (I) band counterparts, and these are typically the less extinguished sources with A_V less than 10 mag approximately. Of the 68 IR-excess selected YSOs only 10 (30) are detected in V (I), and of these, 2 have been designated PMS stars from the optical photometry fulfilling the criteria of two CM diagrams (ID 412, 473, 822). Also, 3 of the 12 optically selected MS stars have IR-excesses (ID 343, 520, 660). We note that our near-IR observations cover only the central $3.7' \times 3.4'$ part of the $6.5' \times 6.5'$ optical field (see Fig. 7).

The analysis of the optical photometry finds a population of 58 PMS sources, of which only 2 overlap with the 68 IR-excess YSOs. This shows that the two photometric methods are complementary and sample different parts of the YSO population

of this type of regions. The optically selected PMS stars are in general expected to be closer to the main-sequence phase and have less measureable circumstellar dust or none at all. It is also known that only about 60% of the classical T Tauri stars in Taurus-Auriga have measurable NIR excesses (see Fig. 7 in Strom et al. 1993). In addition, a large fraction of the IR-excess population is not optically visible, making a direct comparison difficult.

The fact that 25% of the optically selected *main-sequence* stars show IR-excesses is probably related to their all being relatively massive, and in theory they arrive on the main-sequence already while still accreting material, skipping the PMS phase. It is also possible that these 3 out of 12 main-sequence stars are unresolved binaries.

The IR-excess population may not be well-sampled by near-IR photometry, however. The mid-IR ISO⁵ surveys of young embedded clusters demonstrated that in all regions only about 50% of the IR-excess population is sampled by near-IR photometry (Kaa & Bontemps 2001). Mid-IR photometry can separate IR-excess from reddening better and detect hot dust that is obscured from view.

5.5. The luminous protostar candidate

Searching the IRSA archive we found one mid-IR source inside our near-IR NOTCam field in the MSX6C catalogue from the Midcourse Space Experiment. This is G224.1880+01.2407, which is located within $1.6''$ (with position angle 255 degrees) of an IR-excess source in our sample (ID = 1050, $K_S = 15.1$ mag). This is less than 3σ of the MSX6C positional uncertainty, and therefore we consider the source ID = 1050 a possible near-IR counterpart. According to its spectral energy distribution (SED) in the mid-IR (see Fig. 12), it is a protostar candidate. The SED index⁶ from 2 to $15 \mu\text{m}$ is as high as $\alpha_{2-15} = 2.7$, which according to the YSO classification scheme (Lada 1987; Greene et al. 1994), though strictly valid for low-mass stars only, means this is a protostar candidate of the type Class I or possibly in the transition phase from Class 0 to Class I. The brightness of the object in the mid-IR (0.8755 Jy at $8.28 \mu\text{m}$) indicates it is a luminous/massive protostar.

Combining the NOTCam JHK_S fluxes with the fluxes in the MSX bands ACDE at $\lambda\lambda$ 8.28, 12.13, 14.65, and $21.34 \mu\text{m}$, respectively, and the IRAS 12, 25, 60, and $100 \mu\text{m}$ fluxes, the SED is sampled over a sufficient wavelength interval to compare it to YSO models (see Fig. 12). The online YSO SED model-fitting tool provided by Robitaille et al. (2007) was used to find the best YSO model for the observed SED. The input ranges in interstellar extinction and distance was limited to $10 \text{ mag} < A_V < 40 \text{ mag}$ and $2.1 \text{ kpc} < D < 4.1 \text{ kpc}$, respectively. The models that fit the observed SED all give: a stellar mass from 8 to $12 M_\odot$, a total luminosity of 1 to $4 \times 10^3 L_\odot$, a stellar age of 40 000 yrs or less, and an envelope accretion rate of $\sim 5 \times 10^{-3} M_\odot/\text{yr}$. Adjustments for aperture mismatches of the different wavelengths do not significantly alter the derived values.

For all best-fitting models, the measured flux in the MSX D band ($13.5\text{--}15.9 \mu\text{m}$) is too low. We speculate that this is caused by very strong absorptions, such as the known feature at $15.2 \mu\text{m}$ that Alexander et al. (2003) found to be present for the youngest and most embedded YSOs they surveyed with the

⁵ Infrared Space Observatory.

⁶ The SED index is defined as $\alpha_{2-15} = -\frac{\log(v_{15}F_{v_{15}}) - \log(v_2F_{v_2})}{\log v_{15} - \log v_2}$ where F_ν are the flux densities (in Jy) at the frequencies ν (in Hz) corresponding to the wavelengths 2 and $15 \mu\text{m}$.

Fig. 12. The spectral energy distribution (SED) of the luminous proto-star candidate sampled by NOTCam JHK_s band fluxes, MSX ACDE band fluxes, and IRAS 12, 25, 60, and 100 μm fluxes. The best-fitting model found using the online SED fitter (Robitaille et al. 2007) is plotted as a continuous line that splits into two at longer wavelengths for small (*lower*) and large (*upper*) aperture sizes. The Y-axis is scaled to a distance of 1 kpc.

Fig. 13. MSX A band (8.28 μm) image with the error ellipse of IRAS 07141-0920 taken from the IRAS point source catalogue, as well as the location of the optical nebula illuminator, the massive main-sequence star S294-B0.5V (circle). The FOV is roughly the same as for the optical images, and the J2000 RA and Dec grid is shown. The intensity is in logarithmic scale.

ISOCAM CVF. This feature is believed to be attributed to the bending mode of CO_2 ice.

An MSX A band image (6.8–10.8 μm) covering the area of our optical and near-IR studies is shown in Fig. 13. In addition to the bright MSX point source, the 8 μm image shows large-scale diffuse emission extending as curved arms $\sim 2'$ or more from the point source. The diffuse emission is strong at 8 μm , but barely distinguished from the noise in the other MSX bands. Its main origin is likely the emission features at 7.7 and 8.6 μm , two of the emission bands of polycyclic aromatic hydrocarbons (PAHs). As shown in Fig. 15, the curved filaments in emission at 8 μm trace the borders of the $\text{H}\alpha$ emission well. This closely resembles the findings of Deharveng et al. (2003) for the geometrically simpler sources Sh 217 and Sh 219, two spherically symmetric HII regions surrounded by annular photodissociation regions traced by both 21 cm HI and 8 μm PAH emission. In our case the morphology is more complex.

Fig. 14. Plot of the instrumental index ($R - \text{H}\alpha$) versus V mag. Small dots represent all stars in our sample with valid R and $\text{H}\alpha$ measurements. Different symbols are used as follows: squares for stars with NIR excess; crosses for stars with possible emission in $\text{H}\alpha$ ($(R - \text{H}\alpha) > -3.4$); open circles for PMS candidate members, with respect to P99 or S00 models, selected in 2 CM diagrams; closed circles for PMS candidate members selected in 3 CM diagrams; triangles for stars with possible ($U - B$) excess.

5.6. PMS membership and $\text{H}\alpha$ emission

The combined evidence from all photometric diagrams indicates several stars associated to the cluster, which are not detectable as candidate members at optical wavelengths. We consider, in this Section, spectral features that could be present in several stars in our sample and have been identified in the literature as spectral signatures characteristic of PMS stars. The basic feature in this context is the presence of $\text{H}\alpha$ in emission, typical in the spectra of low-mass forming stars of the T Tauri type, both in classical (CTT) and weak-lined (WTT) stars, as well as in higher-mass forming stars of the HAeBe type. The value of the $R - \text{H}\alpha$ index has been used in previous studies of young clusters to describe this feature, although its value may be reflecting spectral properties other than those due to the PMS nature of the stars; see Sung et al. (1997), Sung et al. (2000), Sharma et al. (2007). For our stars, this index, calculated as the difference between instrumental magnitudes, is listed in Col. 11 of Table 1, and plotted versus V mag in Fig. 14. Different symbols are used to distinguish stars that show the different features or spectral signatures of interest: possible $\text{H}\alpha$ emission ($R - \text{H}\alpha > -3.4$ in the plot), PMS candidates from our $UBVRI$ photometry, NIR excess, as described in Sect. 5.4, and probable excess in $U - B$ (see Sect. 5.2 and Fig. 8).

In their spectroscopic study of the PMS population in the 4 Myr old cluster Tr 37, Sicilia-Aguilar et al. (2005) find 40% of stars with signs of accretion, a percentage that decreases to almost zero in the 10 Myr old cluster NGC 7160. Furthermore, the percentage of these accreting stars with signs of NIR excess, 50% in Tr 37, drastically decreases with increasing age. The corresponding numbers from our photometric analysis would be: 16 stars with $(R - \text{H}\alpha) > -3.4$ out of 58 stars selected as optical PMS candidates (27%), and 4 stars among these 16 (25%) with NIR excess. Considering $\text{H}\alpha$ emission as a possible tracer of accretion processes and given the age of our optically selected PMS sequence (around 8 Myr), the corresponding percentages in our sample are consistent with those results.

Fig. 15. The spatial distribution of the cluster members overlaid on the $H\alpha$ image (grayscale). The white contour lines represent the MSX $8\ \mu\text{m}$ isophote emission. The positions of main-sequence stars (blue diamonds), PMS stars selected from optical CM diagrams (yellow circles), IR-excess stars selected from the $J - H/H - K_S$ diagram (red squares), and $H\alpha$ emission sources selected from the $(R - H\alpha)$ index (green crosses) are shown. The massive MSX protostar candidate is located at the strongest peak of the $8\ \mu\text{m}$ emission. The spatial distribution of the IR-excess sources is strongly sub-clustered.

Figure 14 indicates that the presence of possible $(U - B)$ excess coincides frequently with the indication of $H\alpha$ emission, but it seems to exclude PMS candidates selected in 2 or 3 optical CM diagrams. In particular, those of earlier spectral type (around AF: stars marked with both an open and a closed circle), most do not show any sign of circumstellar activity or ongoing accretion, whereas several fainter PMS candidates, below spectral type G and later (marked with only open circles), do show signs of $H\alpha$ emission, but only two of them possible $(U - B)$ excess. This mutual exclusion of PMS nature (inferred from optical CM diagrams) and $(U - B)$ excess might have two possible causes. Stars with $(U - B)$ excesses might have other spectral anomalies, such as $(B - V)$ indices with some excess, and possible $H\alpha$ emission. These features would prevent them from being selected as optical PMS candidates by our CM diagrams method. On the other hand, and more interestingly, the disappearance of $(U - B)$ excesses as we move to earlier type stars could be due to magnetospheric accretion stopping operating in HAeBe stars due to weaker or absent magnetic fields (Wade et al. 2007) when compared to magnetic fields in lower-mass PMS stars.

In general, the progressive disappearance in our PMS sample of spectral features, such as $H\alpha$ emission and even $(U - B)$ excess, as we move to earlier spectral types, is a finding on line with arguments in favour of a faster dissipation of accreting discs and envelopes in stars of higher masses (Hernández et al. 2005). In fact, a decrease in number and intensity of emission features in stars of spectral type F and earlier has been found in comprehensive analyses of the spectral types of PMS stars

(Hernández et al. 2004). Furthermore, the results of Manoj et al. (2006), relative to HAeBe stars, show a rapid decrease in $H\alpha$ emission with age, but more interestingly, no F-type star in their sample exhibits an equivalent width of $H\alpha$ in emission above 10, independently of age. In this same context, Hernández et al. (2006) have found that the PMS population in Orion, between ages of 5 and 10 Myr, show signs of residual debris discs, without excess in the IRAC bands and without emission lines in their optical spectra, but with a varying degree of $24\ \mu\text{m}$ excess.

6. Discussion

6.1. Spatial distribution

A total of 133 cluster member candidates were found in the previous section using CM and CC diagrams based on $UBVRIJHK_S$ photometry. From the optical CM diagrams using the method of Delgado et al. (2007), we find 12 main-sequence stars and 58 PMS stars. Using IR-excess as a youth and member criterium we find 68 PMS members from the $J - H/H - K_S$ diagram. Only 5 of these coincide with the optically selected members. In addition, we find 43 sources with strong $H\alpha$ emission using the $R - H\alpha$ index. Of these, 21 coincide with members selected from optical CM diagrams or with NIR excess. The total number of cluster candidates found in this study is thus 155.

The spatial distribution of the cluster members can give valuable information about cluster evolution and star formation scenario. In Fig. 15 we show the location of the cluster candidates

with different symbols according to the selection method overlaid on the $H\alpha$ image. While it is clear from Fig. 6 that there are mainly two stellar density peaks at $2\ \mu\text{m}$, referred to as the eastern and western clusters, the spatial distribution of the cluster candidates, as selected by our previously described methods, gives a more nuanced impression. The density of cluster members is still highest at the location of the eastern and western clusters, but about half of the members are extended both towards SE and NW, tracing the dense middle lane of the region seen in absorption on the $H\alpha$ image. On a large scale, star formation seems to be going on along a filament or flattened density structure along a ridge oriented SE-NW.

As many as 8 of the 12 optically selected MS cluster members are located in the area of the strongest (western) stellar density peak seen at $2.12\ \mu\text{m}$ (see Fig. 6) around the source S294-B0.5V. Three are located in an area about $1.5'$ to the NW, and the last one about $1.5'$ to the NE. We note that there is only one MS member found towards the SE.

The optically selected PMS cluster members are mainly distributed in roughly three loosely clustered areas: one around the exciting source S294-B0.5V, one about $1.5'$ to the NW, and one about $1.5'$ to the SE.

The IR-excess selected YSOs have a spatial distribution mainly across the NW-SE diagonal of the $3.7' \times 3.4' JHK_s$ image. In addition to the more scattered component of IR-excess source distribution, about half of them are strongly clustered. Around the source S294-B0.5V and coinciding with the strongest stellar density peak of Fig. 6 as many as 8 IR-excess sources are inside $\sim 16''$ (see the zoom-up in Fig. 5 for high spatial resolution). Towards the eastern cluster, there is clustering of IR-excess sources in a dense part of the cloud practically void of optical sources. The projected separations between all the near-IR excess sources have a distribution with one peak at about 15 arcsec and another at 90 arcsec. The latter reflects the extent of the most active region of star formation, about 1.4 pc for an adopted distance of 3.2 kpc. The peak at 15 arcsec or 0.23 pc reflects the typical size of the smallest stellar groups. This is in good agreement with the smallest clustering scale of Class II sources (0.25 pc) found in the nearby Serpens star-forming region (Kaas et al. 2004). The youngest of the PMS cluster members, the luminous MSX source G224.1880+01.2407, a massive protostellar candidate, resides in the easternmost of these groups.

Whether the groups of the embedded eastern cluster are younger than the members in the western cluster around S294-B0.5V is not clear. The clustering of IR-excess sources is strong around S294-B0.5V as well. The age difference between the two most massive stars is clear, however. One is already on the main sequence, while the other one is still in its protostellar phase. It is also quite likely that the median age of the IR excess sources is less than the median age of the optically selected PMS sources, and for comparison the Serpens Class II sources with a similar clustering scale have an age estimate of about 2 Myr (Kaas et al. 2004). Kinematic data may be needed in order to better understand the nature of the sub-clustering processes observed in this star-forming region.

6.2. A possible star formation scenario for Sh2-294

The MSX A-band image shows diffuse emission along curved filaments that trace the borders of the $H\alpha$ emission (see Fig. 15). If the $8\ \mu\text{m}$ emission is mainly due to PAHs, as suggested in Sect. 5.5, this spatial distribution can be interpreted as a photodissociation region (PDR) forming a shell around the optical HII region, in a similar fashion as for Sh2-217 and

Sh2-219 (Deharveng et al. 2003), although the morphology is less symmetrical for Sh2-294. The bright MSX point source is observed in the direction of the eastern edge of the PDR region, only $\Delta\alpha = 24.3''$ and $\Delta\delta = 4''$ offset from the eastern and strongest of the two peaks in the 1.46 GHz VLA map (Fich 1993). The IRAS colours indicate an ultra-compact HII region, and the quoted $7'$ diameter of the VLA source (Fich 1993) is understood to include the free-free emission from the larger and optically visible HII region centred on the western radio peak close to S294-B0.5V.

The appearance in the near-IR images is that of a double cluster with the eastern cluster containing the MSX protostar, probably responsible for the ultra-compact HII region, and the western cluster centred on the massive star S294-B0.5V, responsible for the optical HII region. That the mid-IR emission traces the borders of the optical HII region so well argues strongly in favour of a physical connection and against a chance alignment of two clusters at different distances. Their projected separation is small, but the eastern, embedded cluster must be towards the front side of the optical HII region. Its shape formed by the sub-clustered IR-excess sources traces the east-west filament ($8\ \mu\text{m}$ emission and $H\alpha$ absorption). The spatial distribution of the bulk of the cluster members, the lower mass stars, is not distinctly bimodal, however, and the optically selected PMS stars (with typical ages of 7–8 Myr) are spatially mixed in many areas with the IR-excess sources (most of which are not seen in the optical, are sub-clustered, and probably much younger). This can be partly due to projection and spatially variable cloud extinction.

The ongoing massive star formation, traced by the MSX protostar G224.1880+01.2407, seems to take place in the PDR. It may have been triggered by the optical HII region excited by S294-B0.5V, estimated to have spent about 4 Myr on the main sequence. If this is the case, the age difference between the two generations is probably quite small. For the bulk of the cluster members, the low-mass stars, it is hard to argue for any distinct age difference between the eastern and western parts, although there is probably a difference between the median age of the IR excess sources and that of the optically selected PMS stars. To map the molecular cloud and gain more insight into the star formation scenario for Sh2-294, follow-up observations in the mm and radio wavelengths are needed.

7. Summary

We have presented an optical and near-infrared imaging and photometric study of the HII region Sh2-294 (S294) associated with the IRAS source PSC 07141-0920. High-resolution images in the *UBVRI* and $H\alpha$ bands reveal an emission nebula with very rich morphological details. The nebula is composed of ultraviolet scattered light and of $H\alpha$ emission with local morphological features conveying an impression of a complex web of streams, flows, shocks, turbulence, cavities, making it a region of great dynamical complexity. Opaque patches are seen in contrast with the bright parts of the nebula. These dark cloud regions appear to be in the process of being shredded or blown apart by stellar winds, jets, or outflows phenomena. They appear elongated, filamentary, wispy, clumpy, or knotty.

Our optical photometry confirms that the illuminator of the nebula is likely to be star No. 4 of Moffat et al. (1979), with a spectral type of B0.5V, but located at a distance of about 3.2 kpc, as recently revised by Russeil et al. (2007).

Our high-resolution near-IR images reveal an embedded cluster, extending for about 2 pc and exhibiting sub-clustering: a

denser, more condensed sub-cluster (western sub-cluster) surrounding the high-mass B0.5V illuminator star; and a more embedded sub-cluster (eastern sub-cluster) located towards the eastern dark part of the nebula and including the luminous MSX source G224.1880+01.2407, a massive protostellar candidate that could be the origin of jets and extended features seen at $2.12\ \mu\text{m}$. This double cluster appears to be clearing the remaining molecular material of the parent cloud, creating patches of lower extinction and allowing some of the least reddened members to be detected in the optical images.

Of the 155 stars considered as cluster members, 12 of them are MS members. The remaining are PMS candidates, inferred from three different types of indicators: membership assignment from comparison with PMS isochrones, $H\alpha$ emission deduced from the $(R - H\alpha)$ index, and the presence of NIR excess. About half of the IR-excess sources are spatially distributed in sub-clusters with typical sizes of about 15 arcsec or 0.23 pc. These are probably younger than the optically selected PMS stars. The age estimate for the massive MS member is about 4 Myr, and the average age of the optically selected PMS members around 7–8 Myr. Our results support time-scale estimates for the duration of accretion disks around 10 Myr for late G-type stars and later, and a remarkable decrease in this duration towards earlier spectral types.

We provide a catalogue containing the photometry of all 1344 stars detected at least in one of the *UBVR_IJHK_S* bands. Cluster membership is also included.

Acknowledgements. J.L.Y. thanks the Instituto de Astrofísica de Andalucía for hosting him during his sabbatical leave. This work has been partly supported by the Portuguese Fundação para a Ciência e Tecnologia (FCT). This work has been supported by the Spanish MEC through grant AYA2004-05395 and by the Consejería de Educación y Ciencia de la Junta de Andalucía, through TIC101. Part of the data presented here were taken with ALFOSC, which is owned by the Instituto de Astrofísica de Andalucía (IAA) and operated at the Nordic Optical Telescope under an agreement between IAA and the NBIfAFG of the Astronomical Observatory of Copenhagen. We made use of the NASA ADS Abstract Service and of the WEBDA data base, developed by Jean-Claude Mermilliod at the Laboratory of Astrophysics of the EPFL (Switzerland), and further developed and maintained by Ernst Paunzen at the Institute of Astronomy of the University of Vienna (Austria). This publication made use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research made use of data products from the Midcourse Space Experiment. Processing of the data was funded by the Ballistic Missile Defense Organization with additional support from NASA Office of Space Science. This research made use of the NASA/ IPAC Infrared Science Archive, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. This research also made use of the SIMBAD database, operated at the CDS, Strasbourg, France, as well as SAOImage DS9, developed by the Smithsonian Astrophysical Observatory.

References

Alexander, R. D., Casali, M. M., André, P., Persi, P., & Eiroa, C. 2003, *A&A*, 401, 613
 Bik, A., Lenorzer, A., Kaper, L., et al. 2003, *A&A*, 404, 249
 Bik, A., Kaper, L., & Waters, L. B. F. M. 2006, *A&A*, 455, 561
 Bica, E., Dutra, C. M., Soares, J., & Barbuy, B. 2003, *A&A*, 404, 223
 Blitz, L., Fich, M., & Stark, A. A. 1982, *ApJS*, 49, 183
 Codella, C., Palumbo, G. G. C., Pareschi, G., et al. 1995, *A&AS*, 80, 149

Churchwell, E. 2002, *Annu. Rev. Astron. Astrophys.*, 40, 27
 Cutri, R. M., Skrutskie, M. F., Van Dyk, S., et al. 2003, Explanatory Supplement to the 2MASS All Sky Data Release, <http://www.ipac.caltech.edu/2mass/releases/allsky/doc/explsup.html>
 Deharveng, L., Zavagno, A., Salas, L., et al. 2003, *A&A*, 399, 1135
 Delgado, A. J., & Alfaro, E. J. 2000, *AJ*, 119, 1848
 Delgado, A. J., Alfaro, E. J., Moitinho, A., & Franco, J. 1998, *AJ*, 116, 1801
 Delgado, A. J., Alfaro, E. J., & Yun, J. L. 2007, *A&A*, 467, 1397
 de Winter, D., Koulis, C., Thé, P. S., et al. 1997, *A&AS*, 121, 223
 Fich, M. 1993, *ApJS*, 86, 475
 Fich, M., Dahl, G. P., & Treffers, R. R. 1990, *AJ*, 99, 622
 Fitzpatrick, E. L. 1999, *PASP*, 111, 63
 Girardi, L., Bertelli, G., Bressan, A., et al. 2002, *A&A*, 391, 195
 Greene, T. P., Wilking, B. A., André, P., Young, E. T., & Lada, C. J. 1994, *ApJ*, 434, 614
 Guetter, H. H., & Turner, D. G. 1997, *AJ*, 113, 2116
 Gullbring, E., Hartmann, L., Briceño, C., & Calvet, N. 1998, *ApJ*, 492, 323
 Gullbring, E., Calvet, N., Muzzerolle, J., & Hartmann, L. 2000, *ApJ*, 544, 927
 Henkel, C., Guesten, R., & Haschick, A. D. 1986, *A&A*, 165, 197
 Hernández, J., Calvet, N., Briceño, C., Hartmann, L., & Berlind, P. 2004, *AJ*, 127, 1682
 Hernández, J., Calvet, N., Hartmann, L., et al. 2005, *AJ*, 129, 856
 Hernández, J., Briceño, C., Calvet, N., et al. 2006, *ApJ*, 652, 472
 Kaas, A. A., & Bontemps, S. 2001, in *From Darkness to Light: Origin and Evolution of Young Stellar Clusters*, ed. T. Montmerle, & P. André, *ASP Conf. Proc.*, 243, 367
 Kaas, A. A., Olofsson, G., Bontemps, S., et al. 2004, *A&A*, 421, 623
 Koornneef, J. 1983, *A&A*, 128, 84
 Lada, C. J. 1987, in *Star Forming Regions*, ed. M. Peimbert, & J. Jugaku (Dordrecht: D. Reidel Publishing Co.), *IAU Symp.*, 115, 1
 Landolt, A. U. 1992, *AJ*, 104, 340
 Mamajek, E. E., Meyer, M. R., & Liebert, J. 2002, *AJ*, 124, 1670
 Manoj, P., Bhatt, H. C., Maheswar, G., & Muneer, S. 2006, *ApJ*, 653, 657
 MacLeod, G. C., Van der Walt, D. J., North, A., et al. 1998, *AJ*, 116, 293
 Moffat, A. F. J., Jackson, P. D., & Fitzgerald, M. P. 1979, *A&AS*, 38, 197
 Moorwood, A., Cuby, J.-G., Biereichel, P., et al. 1998, *Messenger*, 94, 7
 Palla, F., & Staller, S. W. 1999, *ApJ*, 525, 772 (P99)
 Pandey, A. K., Upadhyay, K., Nakada, Y., & Ogura, K. 2003, *A&A*, 397, 191
 Pandey, A. K., Sharma, S., Ogura, K., et al. 2008, *MNRAS*, 383, 1241
 Reipurth, B., & Bally, J. 2002, *Annu. Rev. Astron. Astrophys.*, 39, 403
 Robitaille, T. P., Whitney, B. A., Indebetouw, R., & Wood, K. 2007, *ApJS*, 169, 328
 Rodgers, A. W., Campbell, C. T., & Whiteoak, J. B. 1960, *MNRAS*, 121, 103
 Rubio, M., Barba, R. H., Walborn, N. R., et al. 1998, *AJ*, 116, 1708
 Russeil, D., Adami, C., & Georgelin, Y. M. 2007, *A&A*, 470, 161
 Samal, M. R., Pandey, A. K., Ojha, D. K., et al. 2007, *ApJ*, 671, 555
 Sartori, M. J., Lépine, J. R. D., & Dias, W. S. 2003, *A&A*, 404, 913
 Schmidt-Kaler, T. 1982, in *Numerical Data and Functional Relationships in Science and Technology*, ed. K. Schaifers, & H. H. Voigt (Berlin: Springer), 2b, 14
 Sharma, S., Pandey, A. K., Ojha, D. K., et al. 2007, *MNRAS*, 380, 1141
 Sharpless, S. 1959, *ApJS*, 4, 257
 Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *ApJ*, 131, 1163
 Sicilia-Aguilar, A., Hartmann, L. W., Hernández, J., Briceño, C., & Calvet, N. 2005, *AJ*, 130, 188
 Siess, L., Dufour, E., & Forestini, M. 2000, *A&A*, 358, 59
 Strom, K. M., Strom, S. E., & Merrill, K. M. 1993, *ApJ*, 412, 223
 Sung, H., Bessell, M. S., & Lee, S.-W. 1997, *AJ*, 114, 2644
 Sung, H., Chun, M.-Y., & Bessell, M. S. 2000, *AJ*, 120, 333
 Tapia, M., Persi, P., Bohigas, J., & Ferrari-Toniolo, M. 1997, *AJ*, 113, 1769
 Turner, D. G. 1994, *RMA&A*, 29, 163
 Wade, G. A., Bagnulo, S., Drouin, D., Landstreet, J. D., & Monin, D. 2007, *MNRAS*, 376, 1145
 Whittet, D. C. B. 1988, *The observed properties of interstellar dust in the infrared*, in *Dust in the Universe*, ed. M. E. Bailey, & D. A. Williams (New York: Cambridge University Press), 25
 Wouterloot, J. G. A., & Brand, J. 1989, *A&AS*, 80, 149
 Wood, D. O. S., & Churchwell, E. 1989, 340, 265
 Yun, J. L., & Clemens, D. P. 1994, *AJ*, 108, 612