

Parallaxes and proper motions for 20 open clusters as based on the new Hipparcos catalogue

F. van Leeuwen

Institute of Astronomy, Madingley Road, Cambridge, UK
e-mail: fv1@ast.cam.ac.uk

Received 19 November 2008 / Accepted 2 February 2009

ABSTRACT

Context. A new reduction of the astrometric data as produced by the Hipparcos mission has been published, claiming that the accuracies for nearly all stars brighter than magnitude $H_p = 8$ are improved, by up to a factor 4, compared to the original catalogue. As correlations between the underlying abscissa residuals have also been reduced by more than an order of magnitude to an insignificant level, our ability to determine reliable parallaxes and proper motions for open clusters should be improved.

Aims. The new Hipparcos astrometric catalogue is used to derive mean parallax and proper motion estimates for 20 open clusters. The HR-diagrams of the nearest clusters are compared and combined to provide future input to sets of observational isochrones.

Methods. Three different methods are applied, according to the proximity of the cluster, to compensate, where needed, for projection effects, spread in position along the line of sight, and the internal velocity dispersion of the cluster stars.

Results. The new parallaxes have accuracies between 2 and 2.5 times higher than what had been derived from the original Hipparcos catalogue. At least two to three groups of clusters, mostly of similar ages, are observed to each occupy their own specific space in the HR diagram. A significant discrepancy in distance moduli from those obtained with isochrone-based main-sequence fitting remains, in particular for one of these groups, containing the Pleiades, NGC 2516, and Blanco 1. The difference in absolute magnitudes between this group and one containing the Hyades and Praesepe clusters appears to be correlated with systematic differences in the Strömgen Δc_0 index between those groups. The same dependency has been known to exist for a long time, and is again confirmed by the Hipparcos data, in variations in absolute magnitudes for field stars of the same effective temperature.

Conclusions. The positions of the cluster HR diagrams are consistent within different groups of clusters shown for example by the near-perfect alignment of the sequences for the Hyades and Praesepe, for Coma Ber and UMa, and for the Pleiades, NGC 2516, and Blanco 1. The groups are mutually consistent when systematic differences in Δc_0 are taken into account, where the effect of these differences on the absolute magnitudes has been calibrated using field-star observations.

Key words. methods: data analysis – astrometry – Hertzsprung-Russell (HR) and C-M diagrams – open clusters and associations: general – stars: distances

1. Introduction

The data accumulated by the Hipparcos mission allowed for the determination of large numbers of accurate absolute-parallax measurements for nearby stars (ESA 1997; Perryman et al. 1997; van Leeuwen 2007b). The new reduction (van Leeuwen 2007a) (VL7B from hereon), for example, provides some 27 000 stars with parallaxes known to better than 10 per cent, and over 10 000 known to better than 5 per cent. One of the many things learnt from this mission is the relatively large intrinsic spread in luminosities, even among stars which appear very similar in spectral type and photometric colour indices. An examination of G8V and K0V stars, for example, showed an intrinsic dispersion of 0.4 in absolute magnitudes for stars of the same colour index $B - V$ (VL7B). A smaller, but still very significant, level of spread in absolute magnitudes is also observed for stars which were selected by Nicolet (1981) to be very similar in colour indices within the Geneva photometric system, the so-called photometric boxes (VL7B). Such a spread contrasts markedly with the very narrow main sequences observed for the best-studied and nearest open clusters, the Hyades (Perryman et al. 1998; Madsen et al. 2002), Pleiades (Mitchell & Johnson 1957; van Leeuwen et al. 1986) and Praesepe (Crawford & Barnes 1969a). The ability to superimpose the HR diagrams of such

clusters on the HR diagram of the nearby stars, using parallax-based distance moduli rather than isochrone fitting, could provide quantitative information on some of the dependencies that cause the observed spread among the absolute magnitudes of apparently very similar field stars. Predictions on those dependencies are available from theoretical isochrones as based on a combination of models of stellar structure, stellar evolution and stellar atmospheres. However, observational evidence is required to test and eventually validate such models. The possibility of doing so has been hampered by the difficulty to determine with sufficient accuracy the parallaxes of star clusters. Even with the release of the Hipparcos data in 1997, some questions were left, most noticeably because of correlations between the errors on the underlying data from which the astrometric parameters for the clusters had been obtained. Doubts have in particular been raised on the effect that small areas of the sky with concentrations of bright stars may have locally on the measured parallaxes.

The new Hipparcos reduction has improved this situation in two ways: the error-correlation level on the underlying data has been reduced to an insignificant level, and the formal accuracies for most stars, and in particular the brighter stars, have been significantly improved. Both of these improvements have been made possible through a detailed study of the satellite dynamics and incorporating the findings of that study in the reconstruction

of the along-scan attitude, which provides the reference frame for all the Hipparcos astrometric measurements. A potential weakening of the overall rigidity of the reconstructed catalogue, which may be caused by small areas with many bright stars, has also been dealt with. This so-called “connectivity” issue has been explained further in VL7B and van Leeuwen (2007b) (VL7V from hereupon). Given these improvements in both formal errors and overall reliability of the data, a renewed analysis of cluster parallaxes seemed very needed, not least as an apparent discrepancy for the Pleiades parallax as determined from the Hipparcos data has been used on several occasions to allegedly demonstrate errors in that catalogue (Pinsonneault et al. 1998; Soderblom et al. 1998; Pan et al. 2004; Zwahlen et al. 2004; Southworth et al. 2005; Soderblom et al. 2005). The reasons used in those papers to explain the Hipparcos results should be unreliable for the Pleiades cluster, such as error correlations and poor connectivity, have been eliminated by the new reduction (VL7B, VL7V).

For the open clusters studied here we make a few basic assumptions. The first is that all cluster-member stars share the same space motion, have the same age and originate from the same chemical composition “cloud”. The stars are bound to the cluster, which means they will have a small internal dispersion in their velocities, decreasing towards the halo of the cluster. However, in particular the older clusters also tend to be surrounded by a wider halo of escaped former cluster members, which can complicate membership determinations. Furthermore, some of the youngest clusters can not easily be distinguished from fairly accidental concentrations in OB associations, and may not always be bound. However, in all cases the assumption of a shared overall velocity and distance has been applied.

The determination of astrometric parameters for clusters poses different problems as well as possibilities depending on the distance of the cluster. For the nearest cluster, the Hyades, the group is effectively three-dimensionally resolved, while its proximity causes it to cover a relatively large area (about 35 degrees diameter) on the sky. This creates the possibility of using the proper motions of individual cluster members to resolve the cluster along the line of sight. Intermediate-distance clusters can not be resolved this way, but astrometric studies still need to take into account projection effects on the proper motions, and a dispersion along the line of sight in the parallaxes of the individual cluster members. In defining the membership and determining the mean proper motion, the internal velocity dispersion also has to be taken into account, though this will often not be available as an independent measurement. Beyond about 250 pc, the dispersion of the parallaxes resulting from the finite size of the cluster becomes small with respect to formal errors on the measured parallaxes of the individual stars, and the same applies to the proper motion accuracies and the contribution from the internal velocity dispersion. The stars in these clusters can be treated as all having effectively the same parallax and proper motion, which can then be derived directly from a combined solution of the underlying data. This is the combined-*abscissae* method that was first presented by van Leeuwen & Evans (1998), and applied for all clusters by van Leeuwen (1999a) and Robichon et al. (1999). In those studies, however, the emphasis was on compensating for the error correlations in the underlying data, as well as in the proper combination of the two data streams that were used to create the old catalogue. These complications no longer apply when using of the new catalogue to derive astrometric parameters for open clusters.

This paper has been organized as follows. Section 2 presents a summary of the data, with particular emphasis on its use in

deriving star-cluster astrometric parameters. Section 3 reviews briefly attempts made by Makarov (2002, 2003) of applying corrections to the original Hipparcos astrometric data in the context of cluster-parallax determinations. Section 4 presents the different techniques applied to clusters at different distances. Section 5 presents an overview of other, isochrone independent distance modulus determinations for open clusters, and in particular introduces Hipparcos data to determine accuracies of these methods. This is followed by Sect. 6, where the results for individual clusters within 250 pc are presented, and Sect. 7 for clusters between 250 and 500 pc. In Sect. 8 the new set of distance moduli as derived from the parallaxes is compared with a range of earlier determinations, and the HR-diagram positions of cluster main sequences as obtained through ground-based photometry are compared. Section 9 briefly summarizes the space velocities of the clusters as derived from the new parallax and proper motion data. Finally, Sect. 10 presents conclusions from the current study.

2. The Hipparcos data

This section provides a summary of those aspects of the Hipparcos data that are of direct relevance to determining astrometric parameters for open clusters. For a more detailed description of the reductions that led to the new catalogue the reader is referred to VL7B, while VL7V provides a report on the validation tests applied to the data in the new catalogue.

The Hipparcos data originate from two fields of view, separated by a basic angle of 58 degrees, approximately scanning the sky along great circles. The transiting images seen through the two fields of view were projected on the same focal plane. The Hipparcos astrometric data have been derived from transit times over a modulating grid in this focal plane. Through the reconstruction of the along-scan attitude, these transit times are translated to scan phases. In the original reductions these scan phases are projected and measured on a reference great circle, with one such circle defined for each orbit (10.6 h period) of the satellite. The *abscissae* provided thus refer to a mean over all observations of a star over that period of time. These observations will therefore generally contain the combined information from between one and 8 transits of the field of view. In the new reduction the *abscissae* provided refer to individual field of view transits and are measured relative to the instantaneous scan direction. This allows for an improved recognition of faulty measurements, which usually are specific to just one field-of-view transit, and the ability to correct for small-scale grid distortions as a function of the across-scan coordinate.

With respect to the data presented in the first catalogue (ESA 1997), a number of very significant changes and overall improvements have been made. All improvements stem from the way the along-scan attitude is modelled, and how dynamic peculiarities of the satellite motion have been taken into account. There were two peculiarities in the satellite motion that were known but the extent and effect of which had been underestimated in the original reductions: (1) hits by very small particles, causing rotation-rate discontinuities, and (2) discrete satellite-non-rigidity events due to temperature changes affecting at least one of the solar panel mountings, causing so-called “scan-phase discontinuities”. In the original reductions by the two consortia, FAST (Kovalevsky et al. 1992) and NDAC (Lindegren et al. 1992), these events were left mostly unrecognized and effectively smoothed over. This caused locally strongly correlated errors in the reference frame as created by the along-scan attitude reconstruction, as well as a non-Gaussian noise added to




Fig. 1. *Top:* Abscissa-error correlation level (ρ) for all field-transit abscissa residuals (FTARs) with formal errors below 6 mas, as a function of separation along the scan direction. *Bottom:* coincidence fraction c of field-transit measurements as a function of separation on the sky. The coincidence data are based on a random selection covering 10 per cent of all single stars in the new catalogue (from VL7V).

the abscissa measurements. The consequences of this “smoothing over” were made worse still by the use of the great-circle reduction method (van der Marel 1988; van der Marel & Petersen 1992), which referred all measurements over a 10.6 h period to projections on a single reference great circle. This was initially needed in order to deal with the relatively large errors (compared to the Hipparcos target accuracies) on the a priori astrometry provided by the Hipparcos Input Catalogue (Turon et al. 1992; ESA 1992). The great-circle reduction process caused multiple replication of local attitude errors. The effects of a local scan-phase discontinuity could therefore be felt at many different positions on the reference great circle, all separated by an integer number times the basic angle between the two fields of view. A further and more detailed analysis of these effects can be found in van Leeuwen & Evans (1998), van Leeuwen (2005a) and VL7B.

In the new reductions the peculiarities in the dynamics of the satellite have been incorporated fully in the attitude model (van Leeuwen & Fantino 2003). The great-circle reduction process has been replaced by a global iterative solution, similar to what is planned for the Gaia mission (O’Mullane et al. 2007). The iterations are between the reconstruction of the along-scan attitude and the construction of the star catalogue, both using the abscissa residuals as input data. A calibrated set of instrument parameters provides the link between the two systems. The result is a reduction in the overall noise level for the reconstructed along-scan attitude by about a factor five, and a reduction of the abscissa-error correlations by at least an order of magnitude. In the new reduction, the error-correlation level for field-transit abscissae has been examined by VL7V (Fig. 1). Even for relatively bright stars, correlation levels stay in all but the most extreme cases well below 0.1. To see how these error correlations may accumulate to correlated errors in the astrometric parameters of neighbouring stars, the coincidence factor has been introduced (van Leeuwen 1999b). Considering the scan of a single field of




Fig. 2. Correlations (ρ) in the differences between the parallax determinations in the old and new solutions as a function of stellar separation on the sky (ϕ , in degrees). The figures show how initially the catalogue still tries to fit to the original 1997 catalogue, and correlations are small. After several iterations the memory of the original catalogue is lost and the correlations in the original catalogue become fully visible. By iteration 12, this process has stabilized, and memory of the “errors” in the original catalogue has been effectively lost. The upper curve in each figure is the situation for the final iteration. A comparison with the abscissa correlations (Fig. 1) shows clearly that the source of these correlations is entirely in the original reduction (ESA 1997).

view as contained within a single full rotation of the satellite, the coincidence factor determines how often the measurements of two stars are contained within the same scan, as a fraction of the total number of scans for each star. This fraction drops rapidly as a function of the separation of the stars on the sky, as can be seen in the lower graph of Fig. 1. Any correlations between astrometric parameters for neighbouring stars would have to be the result of the combined effects of the abscissa-error correlations and the coincidence fraction. Given the low level of the correlations and the small range of the coincidence factors, no significant level of correlation is expected for the astrometric parameters of neighbouring stars in the new reduction. Similarly, a comparison between the new and the old reduction exposes the correlations in the astrometric parameters for the old reduction. Following these correlations as iterations for the new solution proceeded showed how gradually the memory of the old solution was lost (Fig. 2).

The reduction of about a factor five in the noise level on the reconstructed along-scan attitude leads to a significant improvement in the formal errors on the parallax determinations for the brighter stars. In the published catalogue these accuracies were set by the reconstructed-attitude noise, while in the new reduction they are limited by primarily the photon noise on the abscissa data. For some of the brightest stars, this has resulted in accuracy improvements by up to a factor five. Also the reliability of the formal errors is much improved, as the errors on the field-of-view transit data can be directly related to the integrated photon count of the underlying measurements. Small corrections to the formal errors were derived and applied according to the across-scan position of a transit in the field of view, and the colour index of the star, in particular affecting the very red stars. Altogether, the new reduction provides a significant improvement in accuracy as well as overall reliability of the Hipparcos astrometric data, all of which should benefit the determination of astrometric parameters for open clusters.

3. Corrected Hipparcos data?

At this point it is worth noting the work done by Valeri Makarov on trying to remove the correlated errors caused by problems in

the attitude modelling from the abscissa residuals as presented in the 1997 reduction (Makarov 2002, 2003, 2006). This correction was based on an idea first presented by van Leeuwen (1999b), which mainly focussed on the issue of connectivity between the two fields of view, and how this can be affected by the presence of locally dense groups of bright stars. The idea is that if such a group, like the core of the Pleiades cluster, is dominating the information for the attitude reconstruction, then if the astrometric data for this group of stars contains systematic errors, these errors would also reflect in the abscissa measurements for stars measured simultaneously with the Pleiades stars, but in the other field of view. By then examining the abscissa residuals of such measurements, one may be able to detect these systematic errors in, say, the Pleiades field. A simple test on the abscissa residuals of stars that shared some measurements this way with the Pleiades showed, however, that the unit weight standard deviation of the residuals for coinciding measurements was in fact slightly smaller than for the non-coinciding measurements (van Leeuwen 1999b). In other words, there is no signal contained in those residuals. The reason that Makarov was still able to produce a different parallax for the Pleiades cluster based on such data is that the coinciding stars tend to be much fainter than the Pleiades stars, and add substantial amounts of photon noise to the “corrected” data. The actual correction factor and other criteria as used by Makarov were quite arbitrary (see also van Leeuwen 2005b), and, combined with the photon noise contributions, could be adjusted to provide an expected result. The new reduction has taken care of the attitude modelling effects Makarov was trying to adjust, and it has done so in the only correct way, i.e. by re-analysing in detail the satellite dynamics and implementing this in an iterative solution. The results from the new reduction therefore effectively supersede the earlier results obtained by Makarov, even when these appear to give a “more correct” (that is, as expected by some) result.

4. Techniques for determining the astrometric parameters

In this section we look at techniques used to extract astrometric parameters for open clusters under different conditions of observing. First a set of exact relations is derived, followed by a first-order approximation, and finally the combined-abscissae method, as used for distant clusters, is recalled.

4.1. Exact solution

We define a fiducial reference point which we associate with the centre of the cluster. Its position vector with respect to the Sun is defined as \mathbf{R} , which can be expressed in, for example, equatorial coordinates as:

$$\mathbf{R} = R \cdot \begin{bmatrix} \cos \alpha \cos \delta \\ \sin \alpha \cos \delta \\ \sin \delta \end{bmatrix}. \quad (1)$$

Similarly, the velocity vector $\dot{\mathbf{R}}$ can be expressed as:

$$\dot{\mathbf{R}} = \begin{bmatrix} \cos \alpha \cos \delta & -\sin \alpha & -\cos \alpha \sin \delta \\ \sin \alpha \cos \delta & \cos \alpha & -\sin \alpha \sin \delta \\ \sin \delta & 0 & \cos \delta \end{bmatrix} \cdot \begin{bmatrix} \dot{R} \\ R \dot{\alpha} \cos \delta \\ R \dot{\delta} \end{bmatrix}, \quad (2)$$

where the coefficients on the right relate to the observed radial velocity V_{rad} and proper motion ($\mu_{\alpha^*}, \mu_{\delta}$):

$$\dot{R} = V_{\text{rad}}$$

$$\begin{aligned} R \dot{\alpha} \cos \delta &= \kappa \frac{\mu_{\alpha^*}}{\varpi} \\ R \dot{\delta} &= \kappa \frac{\mu_{\delta}}{\varpi}. \end{aligned} \quad (3)$$

Here $\kappa = 4.74047$, and is the transformation factor from 1 mas yr^{-1} at 1 kpc to 1 km s^{-1} . As was introduced with the release of the Hipparcos catalogue, we use $\mu_{\alpha^*} \equiv \mu_{\alpha} \cos \delta$.

What we are interested in is how the proper motions for cluster members will vary as a result of position in the cluster. For this purpose we include Eq. (3) in Eq. (2) and invert it to give

$$\begin{bmatrix} V_{\text{rad},i} \\ \kappa \mu_{\alpha^*,i}/\varpi_i \\ \kappa \mu_{\delta,i}/\varpi_i \end{bmatrix} = \begin{bmatrix} \cos \alpha_i \cos \delta_i & \sin \alpha_i \cos \delta_i & \sin \delta_i \\ -\sin \alpha_i & \cos \alpha_i & 0 \\ -\cos \alpha_i \sin \delta_i & -\sin \alpha_i \sin \delta_i & \cos \delta_i \end{bmatrix} \cdot \dot{\mathbf{R}}, \quad (4)$$

where the index “ i ” has been added to indicate reference to different positions in the cluster, with $i = 0$ being the values for the assumed cluster centre. We now write Eqs. (2) and (3) as

$$\dot{\mathbf{R}} = \mathbf{A}_0^{-1} \mathbf{v}_0, \quad (5)$$

where $\mathbf{v} = (V_{\text{rad}}, \kappa \mu_{\alpha^*}/\varpi, \mu_{\delta}/\varpi)$, and similarly Eq (4) as

$$\mathbf{v}_i = \mathbf{A}_i \dot{\mathbf{R}} = \mathbf{A}_i \mathbf{A}_0^{-1} \mathbf{v}_0, \quad (6)$$

which fully describes the relation between the observed components of the space velocity as a function of position on the sky and distance of the star.

4.2. Convergent point

The velocity vector of the cluster points to a position referred to as the convergent point:

$$\dot{\mathbf{R}} = \dot{R} \begin{bmatrix} \cos \alpha_c \cos \delta_c \\ \sin \alpha_c \cos \delta_c \\ \sin \delta_c \end{bmatrix}. \quad (7)$$

The common cluster-velocity components of all cluster member proper motions point towards this convergent point. The position of the convergent point depends on the actual velocity vector, of which one component, the radial velocity, is measured directly, and the other component, the proper motion, is scaled with the parallax of the cluster. Therefore, its determination on the sky allows for a measurement of the cluster distance. This is the traditional method used for determining the distance of the Hyades cluster. With the accuracies available in the Hipparcos data, this is no longer the best method available for the Hyades, but still the same effects can be put to use in different ways, to learn about the spatial distribution in the cluster.

Equation (4) can be rotated such that the proper motion is aligned with the direction of the convergent point, by multiplying it left and right by the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \psi & \sin \psi \\ 0 & -\sin \psi & \cos \psi \end{bmatrix}, \quad (8)$$

where $\psi = \arctan(\mu_{\delta,p}/\mu_{\alpha^*,p})$. The additional index “p” shows that the proper motion used here is predicted, based on the assumed space velocity of the cluster and the position of the star on the sky. This rotation then creates the predicted proper motion direction towards the convergent point. Applying the same rotation to the observed proper motion gives the observed proper motions in the direction of the convergent point and perpendicular to it. The rotation also needs to be applied to the errors and error-correlation level for the proper motions. The length of the

observed proper motion vector in the direction of the convergent point is a measure of the distance of the star, given the assumed distance of the cluster centre. The predicted value for the perpendicular component is zero. The observed perpendicular proper motion components can therefore be further investigated for the internal velocity dispersion of the cluster. A rotation of the observed proper motions to the direction of the convergent point and a re-scaling as described above, provides so-called reduced proper motions (van Altena 1966), which are useful for membership recognition in the Hyades cluster.

4.3. First-order approximation

For clusters more distant than the Hyades, the differences between the stellar coordinates and those of the cluster centre are sufficiently small to be approximated in the relevant trigonometric functions. Thus, we develop the matrix \mathbf{A}_i in Eq. (6) as

$$\Delta \mathbf{A}_i \approx \frac{\partial \mathbf{A}_0}{\partial \alpha} \Delta \alpha_i + \frac{\partial \mathbf{A}_0}{\partial \delta} \Delta \delta_i, \quad (9)$$

where $\Delta \alpha_i = \alpha_i - \alpha_0$ and $\Delta \delta_i = \delta_i - \delta_0$. Substituting Eq. (9) in Eq. (6) gives

$$\Delta \mathbf{v}_i = \mathbf{v}_i - \mathbf{v}_0 \approx \left(\frac{\partial \mathbf{A}_0}{\partial \alpha} \Delta \alpha_i + \frac{\partial \mathbf{A}_0}{\partial \delta} \Delta \delta_i \right) \mathbf{A}_0^{-1} \mathbf{v}_0. \quad (10)$$

The two matrix products give

$$\frac{\partial \mathbf{A}_0}{\partial \alpha} \mathbf{A}_0^{-1} = \begin{bmatrix} 0 & \cos \delta_0 & 0 \\ -\cos \delta_0 & 0 & \sin \delta_0 \\ 0 & -\sin \delta_0 & 0 \end{bmatrix}, \quad (11)$$

and

$$\frac{\partial \mathbf{A}_0}{\partial \delta} \mathbf{A}_0^{-1} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{bmatrix}. \quad (12)$$

The first-order corrections for the proper motions and radial velocities of cluster members as a function of position on the sky are thus given by

$$\begin{aligned} \Delta V_{\text{rad},i} &\approx \Delta \alpha_i \frac{\kappa \mu_{\alpha*,0}}{\varpi_0} \cos \delta_0 + \Delta \delta_i \frac{\kappa \mu_{\delta,0}}{\varpi_0}, \\ \Delta \mu_{\alpha*,i} &\approx \Delta \alpha_i \left(\mu_{\delta,0} \sin \delta_0 - \frac{V_{\text{rad},0} \varpi_0}{\kappa} \cos \delta_0 \right), \\ \Delta \mu_{\delta,i} &\approx -\Delta \alpha_i \mu_{\alpha*,0} \sin \delta_0 - \Delta \delta_i \frac{V_{\text{rad},0} \varpi_0}{\kappa}, \end{aligned} \quad (13)$$

where

$$\begin{aligned} \Delta \mu_{\alpha*,i} &\equiv \mu_{\alpha*,i} \frac{\varpi_0}{\varpi_i} - \mu_{\alpha,0}, \\ \Delta \mu_{\delta,i} &\equiv \mu_{\delta,i} \frac{\varpi_0}{\varpi_i} - \mu_{\delta,0}. \end{aligned} \quad (14)$$

Using $\Delta \varpi = \varpi_i - \varpi_0$, this can be written as

$$\begin{aligned} \Delta \mu_{\alpha*,i} &\approx -\frac{\Delta \varpi_i}{\varpi_0} \mu_{\alpha*,0} + (\mu_{\alpha*,i} - \mu_{\alpha*,0}), \\ \Delta \mu_{\delta,i} &\approx -\frac{\Delta \varpi_i}{\varpi_0} \mu_{\delta,0} + (\mu_{\delta,i} - \mu_{\delta,0}). \end{aligned} \quad (15)$$

When the proper motion is large and the cluster is relatively nearby, the first term on the right will create an additional proper motion dispersion along the direction of the cluster proper motion. This effect is referred to as the relative secular parallax

(van Leeuwen 1983). In the extreme case of the Hyades cluster, the proper motions can provide precise measurements of $\Delta \varpi_i$. The precision of these determinations is ultimately limited by intrinsic noise on the proper motions that results from the internal motions in the cluster. The range within which the above approximations can be used is set by the amplitudes of the proper motion and radial velocity, relative to the formal errors on the measured proper motions.

When combining the astrometric data of different cluster members, the formal errors and parameter correlations need to be taken properly into account. There are additional noise contributions to the proper motions from the internal velocity dispersion and to the parallaxes from the finite size of the cluster. The main catalogue for the new reduction provides for each solution the upper triangular matrix \mathbf{U} with which the astrometric parameters are to be multiplied when combining data from different stars. This multiplication normalizes the errors on the parameters. In fact, the inverse of the matrix \mathbf{U} is a square root of the noise variance matrix \mathbf{N} :

$$\mathbf{N}_i = \mathbf{U}_i^{-1} \cdot \left(\mathbf{U}_i^{-1} \right)^T. \quad (16)$$

To this we have to add the ‘‘cosmic’’ noise contributions from the parallax and proper motion dispersions. The parallax dispersion is derived from the typical positional spread of stars in an open cluster, for which a value of $\sigma_r = 3$ pc is assumed. The transformation to a dispersion on the parallax is given by

$$\frac{\sigma_{\varpi,p}}{\varpi_0} = \frac{\sigma_r}{r_0}. \quad (17)$$

Thus, as a cluster becomes more distant, this effect rapidly loses significance. Even for the Pleiades the effect is small, at about $\sigma_{\varpi} \approx 0.2$ mas.

The additional noise on the proper motions originates from the internal velocity dispersion, which has a typical value of $\sigma_v \approx 0.8$ km s⁻¹ in the cluster centre, and the relative secular parallax, for which the contributions are described by Eq. (15). In detail, both contributions depend on the projected distance from the cluster centre, reflecting the space density distribution of the cluster. In a cluster like the Pleiades both effects are of order 1 mas s⁻¹, but for most other clusters the effects are smaller to much smaller. The contribution to the noise matrix for the parallax and proper motion measurements as due to the actual distance of a cluster star is given by

$$\mathbf{N}_p = \begin{bmatrix} \sigma_{\varpi,p}^2 & -\sigma_{\varpi,p}^2 \frac{\mu_{\alpha*,0}}{\varpi_0} & -\sigma_{\varpi,p}^2 \frac{\mu_{\delta,0}}{\varpi_0} \\ -\sigma_{\varpi,p}^2 \frac{\mu_{\alpha*,0}}{\varpi_0} & \sigma_{\varpi,p}^2 \left(\frac{\mu_{\alpha*,0}}{\varpi_0} \right)^2 & \sigma_{\varpi,p}^2 \left(\frac{\mu_{\alpha*,0} \mu_{\delta,0}}{\varpi_0^2} \right) \\ -\sigma_{\varpi,p}^2 \frac{\mu_{\delta,0}}{\varpi_0} & \sigma_{\varpi,p}^2 \left(\frac{\mu_{\alpha*,0} \mu_{\delta,0}}{\varpi_0^2} \right) & \sigma_{\varpi,p}^2 \left(\frac{\mu_{\delta,0}}{\varpi_0} \right)^2 \end{bmatrix}, \quad (18)$$

where $\sigma_{\varpi,p}$ represents the dispersion in parallaxes as caused by the spread in distances along the line of sight for cluster members (Eq. (17)). The noise contribution from the internal motions is simply given by

$$\mathbf{N}_q = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \sigma_{\mu,q}^2 & 0 \\ 0 & 0 & \sigma_{\mu,q}^2 \end{bmatrix}, \quad (19)$$

where $\sigma_{\mu,q}$ is the internal proper motion dispersion resulting from the internal velocity dispersion. The total noise as applicable to the parallax and proper motion of a cluster member when considering its contribution to the cluster parallax and proper motion is given by the sum of the three matrices:

$$\mathbf{N}_t = \mathbf{N}_i + \mathbf{N}_p + \mathbf{N}_q = \mathbf{W}_t^{-1} \cdot \left(\mathbf{W}_t^{-1} \right)^T. \quad (20)$$

Multiplication by the upper-triangular matrix W_t of the equations and observations for each cluster member, describing the corrections to a common cluster parallax and proper motion, provides a set of fully normalized observation equations that can be solved for in a least squares solution.

4.4. Distant clusters

For distant clusters, where the formal errors on the proper motions and parallaxes of the individual members are significantly larger than the corrections described above, the cluster parameters may be derived from a combined analysis of the abscissa residuals for the member stars. Thus, we repeat the astrometric parameter solutions for the cluster stars, but now solve for a single proper motion and parallax for all cluster members.

In order to make this common-parameter solution possible, the abscissa residuals for stars involved need to be re-calculated, to provide residuals relative to a chosen reference parallax and proper motion, which can be seen as the first estimate of the cluster parallax and proper motion. The abscissa residual for observation “ j ” of star “ i ” as provided in the epoch astrometry file in VL7B is represented by da_{ij} . A vector z_i represents the cluster parallax and proper motion, and similarly z_c the reference values for the cluster. The corrected abscissa residuals are then given by

$$da'_{ij} = da_{ij} + (z_i - z_c) \cdot \frac{\partial a_{ij}}{\partial z}. \quad (21)$$

The initial choice for z_c does not affect the linear solution for the astrometric parameters, which includes positional corrections for each star and a single parallax and proper motion for all stars together. The advantage of solving for the cluster parameters in this way is the decrease in degrees of freedom for the solution. This allows for a better recognition of outliers among the measurements and thus a potentially more robust solution. To compensate for abscissa-error correlations, as was described by van Leeuwen & Evans (1998), is no longer required using the new reduction data. Neither is there a need to combine the data from two partly correlated reductions. The process of deriving cluster parameters this way has become significantly simpler and because of that also more reliable, as any of the correlation corrections that had to be applied for processing the data from the 1997 release was also a source of uncertainty in the estimates of the astrometric parameters and their formal errors as derived for the clusters.

5. Some ground-based distance modulus or parallax determination methods, independent of isochrone fitting

In this section three methods that have been used for isochrone-independent cluster-distance determinations are reviewed by incorporating the new Hipparcos data. Some of the methods described below derived or used relations between observed parameters (generally photometric colour indices) and physical parameters of the observed stars, using stars with ground-based parallaxes, or open clusters and their assumed distances, for calibration. The uncertainty has been in distinguishing the error contributions from the calibration objects (such as the measured parallax) from the cosmic noise on such calibrations. The independent absolute parallaxes obtained from the Hipparcos data allow for an assessment of these noise levels.




Fig. 3. A comparison between the ground-based parallax values as used by Crawford (1975) and parallaxes for the same stars as obtained from the new Hipparcos reduction. The comparison shows a systematic difference of 3.4 mas and a noise on the ground-based parallaxes of 10 mas.

5.1. Crawford's calibrations of the Strömgen photometric system

Here an overview is presented on a set of calibrations that have been (and are still) used for luminosity calibrations based on Strömgen photometry. A more recent calibration (but still pre-Hipparcos) is also available (Jordi et al. 1997), but concerns more the calibration of relations between specific indices, and recognizes that there are significant uncertainties in empirical photometric luminosity calibrations. First calibrations incorporating the Hipparcos data have been presented by Domingo & Figueras (1999) for late A-type stars and Jordi et al. (2002) on preliminary results for F, late A and B type stars.

The empirical calibration of absolute magnitudes through observations in the Strömgen photometric system by Crawford was done in three stages, on the F type stars using the Pleiades, α Per and Coma Ber for establishing calibration slopes, and nearby stars with ground-based parallaxes for zero-point determination (Crawford 1975), on the B-type stars (Crawford 1978) and A type stars (Crawford 1979) using distances of the Pleiades and α Per as determined through the F-star calibration. Thus, ultimately, the zero point of the calibration of the system rests on the ground-based parallaxes of 17 nearby F-type stars, while the slopes has been determined by means of the combined data from open cluster stars.

As the ground-based parallaxes used by Crawford had relatively large errors, he applied a Lutz-Kelker correction (Lutz & Kelker 1973) to compensate for the bias in the derived distance moduli and thus the absolute magnitudes. Such corrections are provided for 48 of the 51 stars. Using the new Hipparcos parallaxes, the Lutz-Kelker correction is not needed, as parallax errors are for nearly all stars below 1 per cent. The direct comparison between the parallaxes (Fig. 3) shows a systematic difference of $\varpi_{\text{HIP}} - \varpi_{\text{GB}} = -3.4 \pm 1.4$ mas, and a noise level on this selection of ground-based parallaxes of 10.3 mas. Very similar results were obtained by Jordi et al. (2002) based on the 1997 reduction of the Hipparcos data. There it was also shown that the mean offset for the nearest stars, as used by Crawford, the correction was still larger, at -5 mas. A comparison between the absolute magnitudes as given by Crawford before ($M_V(\varpi)$) and after the Lutz-Kelker correction ($M_V(\text{cor})$) and those derived using the new Hipparcos parallaxes ($M_V(\text{HIP})$) gives the following results:

$$M_V(\text{HIP}) - M_V(\varpi) = 0.090 \pm 0.041,$$

$$M_V(\text{HIP}) - M_V(\text{cor}) = -0.093 \pm 0.041, \quad (22)$$

where in the first solution the noise level was determined as 0.29, and in the second solution as 0.31 mag. The offset in the first solution is caused entirely by the systematic differences of 3.4 mas between the ground-based and Hipparcos parallaxes. After applying the correction for the systematic offset in the ground-based parallaxes, the observed difference is a (rounding-off) residual of -0.013 ± 0.041 mag before application of the Lutz-Kelker correction. The latter ‘‘correction’’ seems to do no good to the data, possibly because corrections of this kind are derived for, and are strictly only applicable *statistically* to *groups* of stars with the same relative errors on the measured parallaxes. The relation between the absolute magnitude, $H\beta$ and Δc_1 is now given by

$$M_V(\text{HIP}) = (3.965 \pm 0.046) - (1.545 \pm 0.101) d\beta + (1.43 \pm 0.26) d\beta^2 - (11.3 \pm 0.7) \Delta c_0, \quad (23)$$

where $d\beta \equiv 10 \cdot (H\beta - 2.64)$, and for these nearby stars the reddening has been assumed zero ($\Delta c_0 = \Delta c_1$). The slope for Δc_1 is effectively the same as determined by Crawford, both from using field stars and from an application to NGC 752, where the turn-off for the main sequence is found in the F-star region. The latter confirmed the relation between Δc_0 and the surface gravity, $\log g$. An alternative solution was made in which the data for the Pleiades (Crawford & Perry 1976) and α Per (Crawford & Barnes 1974) was incorporated, with two additional terms to account for the distance moduli of these clusters, but with $\Delta c_0 = 0$, as was also done by Crawford in his determination of the distance moduli of those clusters. This gave the following solution:

$$M_V(\text{HIP}) = (3.958 \pm 0.041) - (1.668 \pm 0.076) d\beta + s(0.92 \pm 0.18) d\beta^2 - (10.5 \pm 0.6) \Delta c_0, \quad (24)$$

and a nominal distance modulus for the Pleiades of 5.621 ± 0.058 (against 5.54 found by Crawford) and for α Per of 6.045 ± 0.061 (6.1 found by Crawford). However, what is important here is that even after application of Eq. (24), the remaining uncertainty on M_V for the field stars is still 0.20 mag (compared to 0.29 mag in Crawford’s original calibration, and 0.25 mag in Jordi’s calibration), while for the cluster stars alone the remaining noise is significantly smaller, at around 0.14 mag. The distance moduli found here for those two clusters make the assumption that the stars in those clusters are most like the average field star, but as long as the source behind the variation of the field-star magnitudes has not been quantified, this assumption has little meaning. The actual uncertainty in the distance moduli for the two clusters is therefore likely to be significantly higher, and should provisionally be put at 0.20 mag.

In the solution to Eq. (23) also dependencies on Δm_0 and cross terms of Δc_0 and $\hat{\beta}$ were tested, but the result was not significant. The parameter Δm_0 is correlated with abundance variations (Nissen 1970b,a; Gustafsson & Nissen 1972; Crawford 1975), which suggests that such variations are not a significant contribution in the remaining spread in absolute magnitudes.

Table 1 shows the results of applying Eq. (24) to the mean β values derived by Crawford for the five intervals he used for his analysis. The M_V values derived from the new analysis differ by up to 0.14 mag from the values derived by Crawford for $M_c(\text{cor})$.

Conclusions on the Crawford calibrations are the following. The systematic errors on the ground-based parallaxes for the calibration stars were partly offset by the Lutz-Kelker correction, which by itself does not seem to do any good to the data. The Crawford calibration can, as a result be offset by up to 0.14 mag.

Table 1. Comparison between absolute magnitude calibrations for different mean values of β .

$\bar{\beta}$	$\bar{M}_c(\varpi)$	$\bar{M}_c(\text{cor})$	$\bar{M}_c(\text{HIP})$
2.604	4.77 ± 0.23	4.64 ± 0.22	4.677 ± 0.044
2.631	4.21 ± 0.25	4.03 ± 0.26	4.115 ± 0.038
2.648	3.92 ± 0.30	3.69 ± 0.30	3.830 ± 0.043
2.671	3.69 ± 0.52	3.43 ± 0.58	3.529 ± 0.050
2.711	3.53 ± 0.12	3.33 ± 0.17	3.237 ± 0.087

The first three columns are from Crawford (1975); the final column from the current paper, using the least squares solution that provided the coefficient in Eq. (24).

However, the remaining noise level for the calibration standards is much larger than what would be caused by measurement inaccuracies. It may be partly caused by duplicity, in which case the zero point of the calibration is also affected. When deriving distances of individual objects or star clusters using this calibration, this noise needs to be accounted for in the final accuracy figure until it can be quantified and related to independent observed parameters. For the calibrations of the A and B stars there are now available large numbers of calibration stars with well determined parallaxes, which should replace the current extrapolation based on the F stars and cluster distances derived therefrom. This concept is further evaluated in Sect. A. For the current paper the important aspect is the difference between the determination precision, usually quoted as accuracy, for the cluster-distance determinations, and the actual accuracy, which also takes into account the observed cosmic spread intrinsic to the calibration relation.

5.2. Photometric boxes in the Geneva photometric system

The method of the photometric boxes is based on the hypothesis that if two stars have nearly the same colours or derived parameters, then they will also have nearly the same physical and atmospheric properties (Nicolet 1981; Creze et al. 1980). This was applied by Nicolet (1981) to photometric data in the Geneva system for determining the distance moduli of 43 open clusters. The comparisons in the boxes were done directly between stars with ground-based parallax determinations and cluster stars, always sharing the same boxes. The accuracies for these parallaxes were still a major factor in these calculations, which can now be eliminated by substituting those data with the Hipparcos results. What can be measured is the difference in apparent magnitude between cluster members and parallax stars sharing the same photometric boxes. In the original solution two compatible noise sources had to be accounted for, the uncertainty of the measured parallaxes and the uncertainty in the hypothesis stated above. Using the Hipparcos parallaxes, the uncertainty in the hypothesis dominates fully. Nicolet estimated this to be at a level of 0.3 mag.

The detailed data for one cluster, NGC 752, are provided by Nicolet (1981). They are partly reproduced here in Table 2, together with the new parallax data. Based on these data, an estimate of the distance modulus for the cluster can be obtained:

$$dm = 10 - 5 \log \varpi + \Delta m + \varepsilon, \quad (25)$$

where ε is the error on Δm , which was estimated by Nicolet to have a σ value of 0.3 mag, and has now been determined to be around 0.27 to 0.28 mag. The (photometric) distance modulus is determined as 8.25 ± 0.06 . However, this is assuming that NGC 752 in its photometric properties behaves as the average of

Table 2. Data from Nicolet (1981) for the cluster NGC 752.

Id	HD	HIP	ϖ (mas)	Δm
88	160 915	86 736	56.65 ± 0.24	6.834
123	109 085	61 174	54.70 ± 0.17	6.815
139	139 664	76 829	57.35 ± 0.16	6.957
235	7439	5799	42.76 ± 0.30	6.165
259	22 001	16 245	46.12 ± 0.13	6.565
266	22 001	16 245	46.12 ± 0.13	6.430
273	43 318	29716	26.89 ± 0.31	5.727
273	201 772	104 738	30.34 ± 0.34	5.987
304	70 958	41 211	37.57 ± 0.33	6.188
401	4813	3909	63.48 ± 0.35	6.876
402	189 245	98 470	47.06 ± 0.41	6.295
401	191 862	99 572	36.10 ± 0.41	6.079
401	196 378	101 983	40.55 ± 0.27	7.032
402	40 136	28 103	67.21 ± 0.25	7.144
403	193 307	100 412	32.24 ± 0.47	6.144
403	221 356	116 106	38.29 ± 0.54	5.820
414	198 188	102 762	17.11 ± 0.88	4.759
416	88 742	50 075	43.77 ± 0.41	6.459
416	210 918	109 821	37.57 ± 0.33	6.186
428	104 731	58 803	39.49 ± 0.28	6.420
430	221 356	116 106	38.29 ± 0.54	5.905
432	23 754	17 651	56.73 ± 0.19	7.115

The identifier in Col. 1 comes from a proper motion study of this cluster by Ebbighausen (1939). The identifiers in Cols. 2 and 3 apply to the comparison star. Column 4 gives the parallaxes from the new Hipparcos reduction.

the comparison stars. In this respect, a star cluster, with its own very narrow distribution of luminosities, would be the equivalent of a single field star, and could itself in principle be anywhere within the Δm distribution.

The photometric boxes method could be explored further though. Using the now much more extensive volume of photometric data and the Hipparcos parallax information for many of these stars should allow for a much closer examination of the differential effects within these boxes, and the way these may be related to luminosity variations. This in turn could lead the way to an improved photometric calibration of distance moduli for distant star clusters.

5.3. Differential parallax determinations with the MAP

Before starting the analysis of the Hipparcos data for the open clusters, it is worthwhile to investigate the results obtained with differential (small-field) parallax measurements, in particular those obtained with the multichannel astrometric photometer (MAP) (see for example Gatewood et al. 1990) and Hubble Space Telescope, both of which use the same techniques to calibrate from measured relative to estimated absolute parallaxes. Of particular interest are those results obtained before the release of the Hipparcos catalogue, as these provide the most likely measurements not biased by an assumed outcome. A comparison of this kind was done by Gatewood et al. (1998) using the first release of the Hipparcos catalogue, where it was noticed that an additional noise component was present, but the origin and amplitude was not specified. Ground-based parallax measurements obtained mostly with the MAP were found for a total of 71 stars that have also data entries in the new Hipparcos catalogue. The errors as given for those measurements are explained by Gatewood as being precision rather than accuracy, and this becomes quite clear when examining the statistics of the differences with the Hipparcos measurements, shown in Fig. 4.




Fig. 4. Differences between MAP-based and Hipparcos parallaxes for 71 stars. The error bars contain the contributions of the formal errors on the Hipparcos parallaxes and the precisions as given for the MAP data.

Without adding absolute-calibration noise, the weighted mean of the differences and associated standard deviation are given by

$$\langle \Delta M \rangle = -0.501 \quad (26)$$

$$\sigma \Delta M = 2.425. \quad (27)$$

With correct weights, the expected value for the standard deviation equals 1.0. Adding a noise contribution to account for calibration uncertainties to the precision reduces the unit-weight standard deviation. A value close to 1.0 is reached when adding a calibration noise of 2.37 mas:

$$\langle \Delta M \rangle = -0.114 \quad (28)$$

$$\sigma \Delta M = 1.002 \quad (29)$$

$$\sigma(\langle \Delta M \rangle) = 0.333. \quad (30)$$

The added noise should be regarded as an average over the studies included here, and may well vary significantly depending on sources available for calibration. However, it seems unlikely that these parallaxes will have absolute accuracies any better than at best 1.5 to 2.0 mas. The source of this calibration uncertainty is likely the generally small number of calibration stars available and the large intrinsic spread in absolute magnitudes within either an interval in colour index or a spectral class. For the present paper, these results are important when comparing the Hipparcos results with MAP-based parallax determinations for the same clusters.

6. Nine clusters and a moving group within 250 pc

In this section the results for clusters within 250 pc are presented. In order of derived distance, these are the UMa moving group, the Hyades, Coma Ber, Pleiades, IC 2602, IC 2391, α Per, Praesepe, NGC 2451 and Blanco 1. NGC 2451, however, does not correspond to the group of stars that was originally given this designation (Röser & Bastian 1994). The cluster identified as Cr 359, assigned by Loktin & Matkin (1994) a distance within the same range, has been rejected, as it does not show from the Hipparcos data in either proper motion or distribution on the sky (see also Robichon et al. 1999). The indications within the Hipparcos data for the existence of a cluster at 170 pc in Ophiuchus, as suggested by Mamajek (2006), are too weak.




Fig. 5. Reduced proper motions in the selection field for the Hyades cluster. In this diagram, μ_x and μ_y represent the proper motion in the direction of the convergent point and the perpendicular direction respectively. The concentration of stars around $(-115, 0)$ (in red) are the cluster members. The distribution is stretched out in μ_x due to the significant effect of the cluster depth. This information is used to determine differential positions in the cluster along the line-of-sight.

In all cases where HR diagrams are presented, mean reddening corrections for the clusters have been applied according to what appears from the consulted literature to be a reliable estimate. In all cases these corrections are small. No individual corrections for the differential reddening in the Pleiades and α Per, or to data of individual field stars, have been applied. Corrections for metallicity variations have not been applied. For the small variations shown by the clusters examined here, there is no observational confirmation of significant luminosity variations with varying metallicity (see further Sect. A).

6.1. The Hyades

A major study of the Hyades cluster as based on the first Hipparcos catalogue was presented by Perryman et al. (1998), and further studies based on the same data by Madsen et al. (2000), Madsen et al. (2001) and de Bruijne et al. (2001). Differences between the results of these studies show one of the problems associated with most open-cluster studies, the statistical variations in results due to the relatively small number of member stars available, further complicated by uncertainties of membership. Combined with different selection criteria, results can look sometimes even significantly different. A major complication for the Hyades cluster, as well as other similar clusters, is the apparent presence of a halo of stars “evaporated” from the cluster. These can easily disturb estimates of the distance, space velocity and internal velocity dispersion of a cluster.

Based on the new reduction the Hyades cluster membership was reassessed, and similarly the internal dispersion in the proper motions as a function of the projected distance to the “centre” of the cluster, where it should be understood that in a loosely bound, and not very rich cluster like the Hyades, the




Fig. 6. Map of the selected 150 Hyades stars, showing the current positions (points to the *right* of each line) and those extrapolated by 100 000 years (left-side points).

centre is not very well defined. The position of the projected centre was assumed to be (in the ICRS)

$$\alpha_0 = 4^{\text{h}}26^{\text{m}}, \quad (31)$$

$$\delta_0 = 16^{\circ}54'. \quad (32)$$

The catalogue has been searched for potential Hyades members within a radius of 18.54° from this position. Furthermore, the distance from the Sun for the cluster centre was assumed to be 46.7 pc. Within this area on the sky, stars with parallax values were selected such that

$$16.2 \text{ mas} < (\varpi \pm 3\sigma_{\varpi}) < 31.5 \text{ mas}, \quad (33)$$

or between 31.6 and 61.7 pc distance from the Sun. The chosen limits imply that a complete sphere with radius 15 pc around the assumed cluster centre is covered. A total of 270 stars were thus selected. The reduced proper motions of the selected stars were then examined for further membership selection. The reduced proper motions (Fig. 5) describe the observed proper motions as the components towards, and perpendicular to, the convergent point of the cluster, and corrected for the angular separation from the convergent point (Sect. 4.2) (see also van Altena 1966). Based on the reduced proper motions 150 stars were selected as probable members (one additional star, HIP 20895, was rejected based on a comparison between the measured parallax and the kinematic distance). Figure 6 shows these stars and their extrapolated displacements over the next 100 000 years that would result from the observed proper motions. The mean space velocity relative to the Sun for the selected stars has been derived from the observed parallaxes and proper motions only, applying Eq. (4) in a least-squares solution, and is found to be

$$\dot{\mathbf{R}} = \begin{bmatrix} -5.99 \\ 44.73 \\ 4.93 \end{bmatrix} \pm \begin{bmatrix} 0.25 \\ 0.61 \\ 0.14 \end{bmatrix} \text{ km s}^{-1}, \quad (34)$$

with a unit-weight standard deviation of 1.86. The latter is a clear indication that the observed proper motions are significantly affected by internal motions. As a result, the formal errors on the




Fig. 7. Comparison between the formal errors on the distance moduli for the Hyades stars as measured directly through parallaxes (dots) and as derived from the space velocity and the observed proper motions (crosses).

proper motions are insufficient for representing their actual noise level. The formal errors on the input data to the solution do include the errors on the measured parallaxes.

The space velocity, together with the observed proper motions and their covariances can be used to derive, by means of Eq. (6), the kinematic (relative) parallaxes for the cluster members. The advantage of this for the Hyades cluster stems from the relatively large proper motion, combined with small formal errors, and can clearly be seen when comparing the error estimates on distance moduli derived from the measured and kinematic parallaxes (Fig. 7). The mean parallax of the selected stars was calculated from the kinematic parallax estimates, and after projecting their distances onto the line-of-sight for the cluster centre, to avoid any (though quite small) projection-induced bias. The value thus obtained is

$$\varpi_{\text{Hyad}} = 21.53 \pm 2.76 \ (\pm 0.23) \text{ mas}, \quad (35)$$

where the first error shows the standard deviation, representing the spread of stars in the cluster. The formal errors on the measurements hardly contribute to this value, being overall much smaller. The second error in Eq. (35) gives the error on the mean. Thus, based on the error on the mean, the weighted mean distance of the 150 plausible Hyades members selected here is 46.45 ± 0.50 pc, which is equivalent to a distance modulus of 3.334 ± 0.024 . The determination by Perryman et al. (1998) gave a distance of 46.34 ± 0.27 pc, based on a somewhat different selection of stars and reduction methods.

For the current study, the most important result is the position and shape of the Hyades HR diagram, representing an observed isochrone for stars of a specific age and chemical composition. The comparison between the diagrams as derived using the observed or the kinematic parallaxes has been shown on various previous occasions (Madsen 1999; Madsen et al. 2000; de Bruijne et al. 2001) and VL7B, so here is shown only the diagram using the distance moduli for individual stars as based on their kinematic parallaxes (Fig. 8). Clear from this figure is the narrow and well-defined main sequence, as well as the scattering of double stars up to 0.75 mag above it.




Fig. 8. The Hyades HR diagram for 150 probable members, using distance moduli for individual stars as based on kinematically derived parallaxes.

6.2. Coma Ber and UMA

Coma Ber is a sparse cluster at high galactic latitude, considered older than the Pleiades and younger than the Hyades and Praesepe. Crawford & Barnes (1969b) derived a distance modulus of 4.5 for this cluster, and showed in the same paper the results for a small number of members of the core of the UMA moving group, which have characteristics very similar to the Coma Ber cluster. For this group a distance modulus of 1.8 is stated. Both distance determinations used the calibrations of the parameters c_1 and β as presented by Strömgren (1966). That calibration is in fact based on the one presented by Fernie (1965), who used open cluster data for which distances had been derived from on UBV photometry by Johnson et al. (1961). Among those clusters was Coma Ber, for which a distance modulus of 4.5 was given. All distances given in that paper were tied-in to a distance of 40 pc for the Hyades, which we now know should be 16 per cent larger, at 46.45 pc. It is therefore reasonable to assume that the distance and distance-moduli estimates based on this material may have been underestimated. It would in fact be more surprising if they were correct.

There are few independent distance determinations for Coma Ber. Based on the so-called photometric boxes in the Geneva photometric system, and thereby effectively calibrating the cluster distance on ground-based parallaxes of nearby stars, Nicolet (1981) derived a distance modulus of 4.76 ± 0.15 . A differential parallax determination by Gatewood (1995) gave a value of 4.34 ± 0.09 for the distance modulus. This is an internal error, and considering the discussion in Sect. 5.3, an error of the order of 0.3 to 0.4 mag for the absolute value may be more realistic. Based on UBV photometry, Cameron (1985) derived a distance modulus of 4.65, while a more recent photometric distance determination by Pinsonneault et al. (1998) gave a value of 4.54 ± 0.04 . The parallax determinations using the first release




Fig. 9. A tangential projection of the distribution on the sky for members of the clusters Coma Ber (*top left*), Pleiades (*top right*), Praesepe (*bottom left*) and α Per (*bottom right*). The coordinates are in radians, relative to the assumed cluster centre.




Fig. 10. The distribution of the parallaxes against their formal errors for the clusters Coma Ber (*top left*), Pleiades (*top right*), Praesepe (*bottom left*) and α Per (*bottom right*). The central vertical line represents the mean parallax as determined in each cluster based on members identified in the Hipparcos catalogue. The black lines on either side show the 1σ error level, including the internal distance dispersion for the cluster. The outer grey lines similarly represent the 2σ level. This display of the data should best allow the reader to evaluate the consistency of the measured parallaxes for the member stars in each cluster.

Table 3. Summary of distance modulus determinations for the Pleiades over the past 30 years.

Reference	DM	σ	Type	Δ
An et al. (2007)	5.66	0.05	1	0.03
Soderblom et al. (2005)	5.63	0.02	3	0.00
Percival et al. (2005)	5.63	0.05	1	0.00
Southworth et al. (2005)	5.71	0.07	2	0.08
Zwahlen et al. (2004)	5.61	0.05	2	-0.02
Munari et al. (2004)	5.61	0.03	2	-0.02
Pan et al. (2004)	5.65	0.03	2	0.02
Stello & Nissen (2001)	5.61	0.03	1	-0.02
Gatewood et al. (2000)	5.59	0.12	3	-0.04
Narayanan & Gould (1999)	5.58	0.18	4	-0.05
Pinsonneault et al. (1998)	5.63	0.03	1	0.00
Giannuzzi (1995)	5.61	0.26	1	-0.02
van Leeuwen (1983)	5.57	0.08	1	-0.06
Nicolet (1981)	5.73	0.06	1	0.10

Column 4 gives a code for the type of determination: (1) photometric; (2) binary star; (3) differential parallax; (4) convergent point; Col. 5 gives the difference from the weighted mean value of 5.63.

of the Hipparcos data gave values of 4.70 ± 0.04 (Robichon et al. 1999) and 4.77 ± 0.05 (van Leeuwen 1999a).

Eight members of the UMa group were analyzed for the present paper, with individual parallax accuracies in the new Hipparcos reduction between 0.14 and 0.41 mas, or relative errors between 0.35 and 1.0 per cent. The selected stars are from the list of Crawford & Barnes (1969b) (only those with single five-parameter astrometric solutions), including HD 113139 (HIP 63504). The average distance modulus for these stars as derived from the new Hipparcos catalogue is 2.016 ± 0.051 , placing this group about 10 per cent further away than was derived by Crawford & Barnes (1969b). The main reason for including these stars in the present paper and in connection with the Coma Ber cluster is the similarity in age between the two groups. The UMa stars are not examined with the cluster-analysis tools shown in Sect. 4, but rather as individual stars, taking full advantage of the high relative accuracies of the individual parallaxes.

A selection of 27 cluster members has been used for determining the parallax and proper motion of the Coma Ber cluster. There may be two or three more possible members of Coma Ber in the Hipparcos catalogue, but these deviate either in proper motion or in parallax by more than what could be expected from the formal errors. Excluding these stars had only an insignificant effect on the final results. Deriving the astrometric parameters for this, as well as the remaining clusters in this section, has been done through an examination of the astrometric parameters as provided for the individual member stars in the new catalogue. The proper motions have been corrected for the differential projection effects described by Eq. (13), using a radial velocity as specified in Table 5, -1.2 km s^{-1} . The cluster members are spread sparsely over an area with a diameter of about 7.5 degrees on the sky (Fig. 9). Data correlations therefore did not play a major role when the parallax was derived using the 1997 catalogue, and it is thus not surprising that the results from the new reduction confirm and significantly strengthen the original findings by van Leeuwen (1999a) and Robichon et al. (1999). The distribution of the parallaxes and their formal errors is shown in Fig. 10. The summary data for Coma Ber are presented in Table 4. The parallax for Coma Ber is determined as $11.53 \pm 0.12 \text{ mas}$, which implies a distance modulus of 4.691 ± 0.023 . Thus, the same increase of 10 per cent in distance

as observed for the UMa group with respect to the determinations by Crawford & Barnes (1969b). Considering the calibration noise for ground-based differential parallaxes as determined in Sect. 5.3, the Hipparcos value is in good agreement with the determination by Gatewood (1995). There is a good agreement, too, with the values derived by Nicolet (1981) and Cameron (1985). The only marginally significant disagreement is with the determination by Pinsonneault et al. (1998), which differs by 3.4σ . The HR-diagram for Coma Ber is shown in Fig. 11 together with the results for the UMa stars and those obtained earlier for the Hyades cluster. At the parallaxes determined here, the HR diagrams of Coma Ber and the UMa group accurately overlap, but are slightly sub-luminous (0.15 mag.) with respect to the Hyades.

6.3. The Pleiades

More than any other result from the Hipparcos mission, the parallax estimate for the Pleiades cluster has been causing controversy (Pinsonneault et al. 1998; Narayanan & Gould 1999; van Leeuwen 1999a,b; Robichon et al. 1999; Stello & Nissen 2001; Percival et al. 2003; Pan et al. 2004; Percival et al. 2005; Southworth et al. 2005; van Leeuwen 2005b; An et al. 2007; Valls-Gabaud 2007). In many ways this has been highly beneficial, as it has stimulated research projects on the Pleiades. In total there have been at least 15 attempts to determine the parallax or distance modulus of the Pleiades since the publication of the Hipparcos data in 1997. It has also been one of many reasons to continue with a further detailed analysis of the Hipparcos data. With the results of a wide range of studies now available, including the new Hipparcos determination, the solution may not as yet have been identified, but the discrepancy between different approaches has become clearer, and will undoubtedly be resolved in the near future. As has been announced on a few earlier occasions (van Leeuwen 2008), the new reduction of the Hipparcos data did not change significantly earlier estimates for the cluster parallax.

The Pleiades cluster covers an area on the sky of about 9 to 10 degrees in diameter (van Leeuwen 1980, 1983), which is, contrary to statements made by An et al. (2007), much larger than the error-correlation scale length for the 1997 Hipparcos catalogue (van Leeuwen 1999b). In that area, the Hipparcos catalogue contains about 57 members of the cluster, but only 53 of these can be used for determining the cluster parallax. The parallax determinations for the remaining 4 stars may be disturbed by duplicity or orbital motion. Figure 9 shows the distribution on the sky of the selected stars. The distribution of parallax measurements and their formal errors is shown in Fig. 10, with respect to the measured parallax of $8.32 \pm 0.13 \text{ mas}$ (distance modulus 5.40 ± 0.03), as an internally consistent distribution of data points. In the new reduction, these data points are essentially uncorrelated (VL7B, VL7V). From the examination of the parallax zero point in the new catalogue (VL7V), there is no reason to suspect that all parallaxes in the catalogue are offset by anything more than the offset-detection limit of about 0.1 mas.

A comparison between the Pleiades parallax determination as presented above and some of the many determinations presented in the literature over the past 30 years looks, at first sight at least, alarming. Table 3 gives a summary for those not-Hipparcos-based observations. A couple of determinations have been left out on purpose: by Gatewood et al. (1990) (superseded by Gatewood et al. 2000); by Makarov (2002) (superseded by van Leeuwen 2007a, and by the present paper); by Fox Machado et al. (2006) and by Cameron (1985), where no formal error is

Table 4. Summary data for 8 clusters within 250 pc.

Cluster	α, δ degr.	Diam. Stars	$\log(\text{Age})$ $E(B - V)$	V_R (km s^{-1})	$\mu_{\alpha,*}$ (mas yr^{-1})	μ_δ	ϖ (mas)	DM	Dist. (pc)	Std Rej.
Coma Ber	186.0	7:5	8.78	-1.2	-11.75	-8.69	11.53	4.69	86.7	0.87
C1222+263	26.0	27	0.00		± 0.27	± 0.25	± 0.12	± 0.02	± 0.9	0
Pleiades	56.5	8:6	8.08	5.7	20.10	-45.39	8.32	5.40	120.2	1.45
C0344+239	24.1	53	0.04		± 0.28	± 0.27	± 0.13	± 0.03	± 1.9	0
Praesepe	130.0	4:5	8.90	33.6	-35.81	-12.85	5.49	6.30	181.5	1.29
C0937+201	19.7	24	0.01		± 0.29	± 0.24	± 0.18	± 0.07	± 6.0	0
α Per	52.5	8:0	7.55	-1.6	22.73	-26.51	5.80	6.18	172.4	1.28
C0318+484	49.0	50	0.09		± 0.17	± 0.17	± 0.09	± 0.03	± 2.7	0
IC 2391	130.0	2:0	7.88	19.0	-24.69	22.96	6.90	5.80	144.9	1.12
C0838-528	-53.1	11	0.01		± 0.32	± 0.31	± 0.12	± 0.04	± 2.5	0
IC 2602	160.2	2:9	7.83	16.0	-17.02	11.15	6.73	5.86	148.6	0.99
C1041-641	-64.4	15	0.03		± 0.24	± 0.23	± 0.09	± 0.03	± 2.0	0
Blanco 1	1.1	2:5	8.32	3.5	20.11	2.43	4.83	6.58	207.0	1.24
C0001-302	-30.1	13	0.01		± 0.35	± 0.25	± 0.27	± 0.12	± 12.0	0
NGC 2451	115.3	2:0	7.76	21.7	-21.41	15.61	5.45	6.32	183.5	1.42
C0743-378	-38.5	14	0.00		± 0.28	± 0.29	± 0.11	± 0.04	± 3.7	0

The first column identifies the cluster; Cols. 2 to 4 provide external input data, such as reddening, radial velocity, position on the sky; Cols. 5 to 7 provide the data obtained in the current study for the astrometric parameters of each cluster; Cols. 8 and 9 provide the derived parameters of the distance and distance modulus; Col. 10 gives the solution statistics, the unit-weight standard deviation of the residuals and the percentage of rejected observations. There are always three observations per star.




Fig. 11. The HR diagrams for Coma Ber (*left*), Pleiades (*centre*) and Praesepe (*right*), shown as full dots in each diagram. The Hyades stars are shown in the background as open circles. In addition, the Coma Ber diagram also shows the UMa stars (crosses), the Pleiades diagram shows the Coma Ber stars as crosses, and similarly for the Pleiades stars are shown in the Praesepe diagram.

given; and by [Chen \(1999\)](#), which contained serious flaws in the methods used (see also [Robichon et al. 1999](#)). The remaining papers do not provide independent observations either, which becomes clear when examining their spread with respect to the quoted errors. Taking only those determinations published after the publication of the Hipparcos data in 1997 (as shown in [Fig. 12](#)), the weighted mean and standard deviation for 11 observations are determined as 5.63 and 0.58 respectively. The latter value should be close to 1.0 instead. When applying a more realistic formal error to the differential parallax measurements, as described in [Sect. 5.3](#), the uncertainties on the distance-modulus

determinations by [Gatewood et al. \(2000\)](#) and [Soderblom et al. \(2005\)](#) are likely to be seriously underestimated (finding exactly the “expected value” does not mean that the standard deviation associated with the measurement is small), which would further lower the unit-weight standard deviation for these observations.

Of the external measurements, those based on photometry only do not constitute independent measurements of the distance modulus, but rather a fitting to a set of models. Completely independent determinations are needed to verify these models. There are only four sets of such observations available:




Fig. 12. Distribution of determinations (since 1997) for the Pleiades distance modulus and its standard error (other than those directly derived from the Hipparcos data). The distribution of the data points strongly suggests the presence of correlations between the different measurements. The central line represents the mean value of 5.63, the black diagonal lines on either side the 1σ , and the grey diagonal lines the 2σ error level.

1. the convergent point determination by [Narayanan & Gould \(1999\)](#), which has a high error associated with it (see also [Lindegren et al. 2000](#); [van Leeuwen 2005b](#));
2. the differential parallax measurements, for which the uncertainties in the calibration to absolute values have been seriously underestimated;
3. one binary system with radial velocity measurements and an astrometric orbit by [Zwahlen et al. \(2004\)](#);
4. the Hipparcos-data based parallax, derived from the measurements of 53 cluster members spread over some 50 square degrees on the sky.

The only meaningful discrepancy with the Hipparcos result (at 3.6σ) is between the results for this one binary star, when its parallax is interpreted as the parallax of the cluster. It drops to below 3σ when also the uncertainty of the position of the star within the cluster is accounted for. It should be noted that for the data on the individual star in question, there is no significant discrepancy with the Hipparcos data.

It has been stated by [Soderblom et al. \(1998\)](#), and re-iterated by [Stello & Nissen \(2001\)](#) and [An et al. \(2007\)](#), that at the parallax found from the Hipparcos data for the Pleiades cluster, the cluster members will occupy an area of the HR diagram in which no field stars, or more specifically no field stars that are assumed to be young, are found. This is not the case, as field stars contained in the Hipparcos catalogue are observed in the area occupied by the Pleiades. This is shown in Fig. 13, where the Pleiades are compared with the data for field stars with parallax accuracies better than 7 per cent. For those stars reddening corrections tend to be very small. At all areas covered by the Pleiades HR diagram field stars are also found. A detailed study of those stars occupying the same region would be of interest.

There are additional problems in the comparison with the active solar-type stars in [Soderblom et al. \(1998\)](#). The first is that this type of activity as observed in the Pleiades is restricted to stars of spectral type later than about K2 to K3 ([van Leeuwen & Alphenaar 1982](#); [van Leeuwen et al. 1987](#)). Observing similar activity among the more massive solar-type stars indicates an age for those stars considerably lower than the Pleiades age. They would therefore be expected to be found in or near star-forming regions, most likely as members of OB associations. However,




Fig. 13. The HR diagram for the Pleiades (full dots) superimposed on the HR diagram for field stars with parallax accuracies better than 7 per cent. The distance modulus applied is the value of 5.40 mag derived in the present study, and the applied reddening is $E(B - V) = 0.04$.

most of these stars are not linked to clusters or associations. As a consequence, they do not have an independently assessed age. The “young” ages assigned to these stars is almost entirely based on interpretation of their activity ([Soderblom et al. 1998](#)).

The difference between the distance moduli of the Hyades and the Pleiades does not require an explanation in terms of metallicity. As is shown in Appendix A, the observed differences in luminosity, when accepting the Hipparcos-based parallaxes, are reflected in systematic differences in the photometric indices $B - U$ and Strömgren c_1 for F-type stars in the same way as what is observed for a large sample of field stars.

The summary data for the Pleiades are shown in Table 4, and a comparison between the HR diagrams of the Pleiades, Coma Ber and the Hyades is presented in Fig. 11. Clearly visible in this figure is the difference in slope of the HR diagram for the Hyades and the Pleiades over the colour range 0.0 to 0.5 in $B - V$.

6.4. Praesepe

The Hipparcos data for Praesepe is relatively poor due to the closeness of the cluster to the ecliptic plane. Membership is not a major problem for this cluster. Figure 9 shows a map of the 24 Praesepe members included in the Hipparcos catalogue. The area covered by the cluster is about 4.5° in diameter. The parallax measurements for the selected stars are shown in Fig. 10. The accumulated information on the cluster can be found in Table 4. The distance modulus as derived from the new Hipparcos reduction is, at 6.30 ± 0.07 effectively identical to one determined by

An et al. (2007) (6.33 ± 0.04) and fully consistent with the value derived by Percival et al. (2003) (6.24 ± 0.04 , using $(V - I)_C$ data). It is larger, but not significantly, than the value of 6.15 derived by Cameron (1985) from UBV data, and the value of 6.10 ± 0.18 derived by Nicolet (1981) using photometric boxes. Also within the same range is the value derived from differential parallax measurements by (Gatewood & de Jonge 1994), 6.42 ± 0.33 . These results, in a way, mostly reflect the very strong similarity between the Hyades and Praesepe clusters. As shown in Fig. 11, the HR diagrams of the two clusters closely overlap when distance moduli are determined using the Hipparcos astrometric data. This is as expected, given the difference of 3.03 mag in distance modulus, as derived from the parallax measurements, between the Hyades and Praesepe. This is very close to the value of 3.0 which has been used in the past to determine photometrically the distance of the Praesepe cluster with respect to the Hyades (see for example Uggren et al. 1979).

6.5. α Per

In its distribution on the sky (Fig. 9), α Per resembles more a small remnant of an OB association than a bound open cluster. Still there is a very clear clustering in proper motion space (Fig. 14) and a well-defined HR diagram for the stars selected on proper motions. The Hipparcos catalogue contains 50 stars for which the membership is probable to very likely, providing a well-defined proper motion and parallax for the group (see Table 4). The radial velocity for this cluster was taken from Petrie & Heard (1969) as -1.6 km s^{-1} , which is based on the measurements of 77 stars in the field of the cluster. One star should be noted here in particular, HD 22192 (HIP 16826, ψ Per), which was considered as a non member by Petrie & Heard (1969), and also as a possible non-member by Abt (1978), but which star is found to have a well determined proper motion and parallax that are very close to the mean cluster values, and which therefore has been considered as a high-probability member in this study. The observed spread in the radial velocities of this star as reported by Petrie & Heard (1969) is larger than would be expected on the basis of the errors assigned to the 10 individual observations. In addition, the mean value quoted for the radial velocity of this star indicates that a straight (not weighted) mean was used, in which case the radial velocity found is $-5.0 \pm 2.5 \text{ km s}^{-1}$. For the weighted mean a value of $-3.1 \pm 2.6 \text{ km s}^{-1}$ is found, which is fully within the range of the mean cluster radial velocity. It is unclear how Petrie & Heard (1969) found an error on the mean of 1.7 km s^{-1} for what appears to be the un-weighted mean. But why pay all this attention to one star? As a member of the α Per cluster it is its brightest B star by more than 0.5 mag, and as such it can play an important role in the determination of the age of this system. The distribution of radial velocities indicates in addition that it could be an orbiting binary star.

Compared to the distance-modulus values derived from the 1997 Hipparcos data by Robichon et al. (1999) (6.40 ± 0.08) or van Leeuwen (1999a) (6.31 ± 0.08), the new determination (6.18 ± 0.03) brings it closer to other, all photometric, determinations. As an example, the value given by Dambis (1999) is 5.94 ± 0.05 , and by Pinsonneault et al. (1998) it is given as 6.23 ± 0.06 . The differences with the two earlier Hipparcos-based determination stem at least partly from the selections of member stars. In the new reduction, the higher accuracies of the proper motions allows for an improvement in selection, and shows that the selection of members by Robichon et al. (1999) contained at least two non-members (HIP 15160 and HIP 16880) and three




Fig. 14. Distribution of the proper motions for the selected member stars in the α Per cluster.

stars for which the membership is rather uncertain (HIP 14697, HIP 16340, and HIP 16625). Of these stars, HIP 16340 is by far the brightest, and its much smaller parallax will have affected the earlier results.

The HR-diagram for α Per is shown in Fig. 15 in comparison with the diagram of the Pleiades cluster. There is an indication that the Pleiades stars on the red-side of $(B - V) = 0.2$ are generally more blue than those of α Per, while for stars on the blue-side the different sequences overlap very well. This raises once again the question as to what part of the main sequence can be used to derive a reliable photometric distance estimate (see also Percival et al. 2005), as significantly different results will be obtained for different fitting criteria.

6.6. IC 2391 and IC 2602

The Southern Hemisphere clusters IC 2391 and IC 2602 are sometimes described as being like the Pleiades, though a little younger, and less populous. IC 2602 is embedded in the Scorpio-Centaurus OB association (Bertiau 1958; de Zeeuw et al. 1999). An extensive photometric and astrometric study by Braes (1961, 1962) defined probable membership and derived a distance estimate of 150 pc (distance modulus 5.88). A combined UBV photometric and spectroscopic study at the same time by Whiteoak (1961) gave a distance of 155 pc (distance modulus 5.95), using measurements for 40 probable and 29 possible cluster members. Based on $ubvy$ photometry for 33 stars, Hill & Perry (1969) determined a distance modulus of 5.85 ± 0.06 , and Nicolet (1981), using photometric boxes to connect the distance determination directly to stars with measured parallaxes, derived a distance modulus of 5.75 ± 0.47 . Finally, Pinsonneault et al. (1998) reported a distance modulus of 6.02 based on main-sequence fitting.

The proper motion and photometric study of IC 2391 by Hogg (1960) identified 21 stars as likely members and derived a distance of 153 ± 3 pc, from a distance modulus of 5.92 ± 0.05 . Measurements in $ubvy$ and $H\beta$ for 25 cluster members were obtained by Perry & Hill (1969), from which was derived a distance modulus of 6.00 ± 0.06 . They also derived a mean distance modulus of 5.90 for earlier determinations. Nicolet (1981) determined a distance modulus of 6.13 ± 0.32 based on the photometric boxes. Maitzen & Catalano (1986) conducted a study of CP2 stars in IC 2391. It is interesting to note that four stars identified as cluster members in that study were found not to




Fig. 15. The HR diagram for: (*left*) the α Per cluster (dots); (*middle*) IC 2602 (dots) and IC 2391 (crosses); (*right*) NGC 2451 (dots) and Blanco 1 (crosses). In each diagram, the Pleiades data are shown as open circles in the background as comparison.

be so, based on their proper motions, though the parallaxes of these stars are close to the parallax for IC 2391. These stars are HD 74195 (HIP 42536), HD 74762 (HIP 42829), HD 74665 (HIP 42767) and HD 75202 (HIP 43071). Finally, [Pinsonneault et al. \(1998\)](#) derived a distance modulus for IC 2391 of 5.99, based on main-sequence fitting to theoretical isochrones.

With 11 (IC 2391) and 15 (IC 2602) members in the Hipparcos catalogue respectively, they provide well determined parallax and proper motion estimates in the new reduction (Fig. 16). The distance moduli as derived for these two clusters from their parallax measurements are 5.80 ± 0.04 (IC 2391) and 5.86 ± 0.03 (IC 2602), the summary data are presented in Table 4. The distributions of the member stars are shown in Fig. 17. Figure 15 shows the HR diagrams of IC 2391 and IC 2602 in comparison with the diagram for the Pleiades. The positions of the sequences for the three young clusters coincide well, in the same way as the sequences for the older clusters UMa and Coma Ber (Sect. 6.2) and those of the much older Hyades and Praesepe clusters. With the improvement by a factor two in formal errors on the parallaxes for the Pleiades, IC 2391 and IC 2602 (with respect to the 1997 catalogue), this agreement is more substantial now than it was ever before. Just as for the Pleiades cluster, the parallax-based distance moduli for IC 2391 and IC 2602 are found to be systematically smaller than has been estimated from main sequence fitting, where values of 5.99 and 6.02 were reported by [Pinsonneault et al. \(1998\)](#) for IC 2391 and IC 2602 respectively. A distance modulus of 6.02 would imply a parallax of 6.25 mas, which, in view of Fig. 16 appears rather unlikely.

6.7. Blanco 1 (ζ Scl)

The cluster Blanco 1 has in the past been referred to as the ζ Scl cluster, after the bright star considered to be a cluster member ([Eggen 1970](#); [Perry et al. 1978](#); [de Epstein & Epstein 1985](#); [Westerlund et al. 1988](#)). The proper motion and parallax (at 6.49 ± 0.25 mas) measured by Hipparcos, however, clearly indicate that this star is not a cluster member. A suspicion that this

star might not be a cluster member was already earlier expressed by [Westerlund et al. \(1988\)](#). Also confirmed as a non-member is HD 225119 (HIP 291), based on both proper motion and parallax, while the remaining three B star members identified by [Westerlund et al. \(1988\)](#) are all confirmed. As this cluster can now no longer be associated with ζ Scl, it will be referred to as Blanco 1, after its discoverer ([Blanco 1949](#)). In previous studies of this system, the similarity between its member stars and those of the Pleiades cluster has been noticed repeatedly.

Based on the proper motions in the cluster area, 13 stars were ultimately selected as probable cluster members. A map of those stars can be seen in Fig. 17. The astrometric parameters derived from the Hipparcos data are presented in Table 4. The distribution of the parallaxes and their formal errors for the member stars is shown in Fig. 16.

6.8. NGC 2451

The cluster referred to here as NGC 2451 identifies with what has been called the Puppis Moving Group (PMG) by [Röser & Bastian \(1994\)](#), and by others as NGC 2451a. The reason for this is, that this cluster does not identify with the group of stars to which originally the name NGC 2451 was assigned. That group of stars are mainly in the background for the PMG, at about twice the distance, and only very few of those stars belong to the PMG. The PMG is of approximately the same age as IC 2391 and IC 2602. Membership determination for this cluster has in the past been somewhat problematic (see for example [Levato & Malaroda 1975a](#); [Maitzen & Catalano 1986](#)), and it has been suggested that there are in fact two clusters sharing nearly the same line-of-sight ([Röser & Bastian 1994](#)), although the existence of the second, more distant cluster, is still uncertain ([Baumgardt 1998](#)). An example of the problems surrounding this cluster is the list of spectral types for 24 stars in NGC 2451 by [Levato & Malaroda \(1975a\)](#). When incorporating the Hipparcos data, only 4 of these stars are confirmed as cluster members, 8 stars are definitely not members, while 12 stars are not included in the Hipparcos catalogue. Among the

non-members is the K5 IIa star HD 63032 (HIP 37819), which was still considered a certain member by [Levato & Malaroda \(1975a\)](#), but which has also been recognized as non-member by [Röser & Bastian \(1994\)](#). This star had been used in the past as an indicator of the age of NGC 2451. Its proper motion, at $(-10.2 \pm 0.18, 6.34 \pm 0.21)$ mas yr⁻¹ and parallax at 2.88 ± 0.19 mas well separates this star from the cluster. The non members in the field of NGC 2451 as contained in the Hipparcos catalogue do not show common parallax and proper motion values that would indicate membership of a second cluster behind NGC 2451. But with the second cluster at possibly twice the distance there may be very few if any actual members of it included in the Hipparcos catalogue. It should be noted that the PPM proper motions as used by [Röser & Bastian \(1994\)](#) are significantly offset in declination, by about 6 to 8 mas yr⁻¹ from the Hipparcos data. The formal internal errors of about 3 mas yr⁻¹ assigned to the PPM proper motions appear to be correct.

The Hipparcos catalogue contains data for 14 stars that can be identified as probable members of NGC 2451 as based on their proper motions and parallaxes. The combined parallax data for these stars is shown in Fig. 16. The mean parallax for the 14 selected stars is 5.45 ± 0.11 mas, equivalent to a distance modulus of 6.32 ± 0.04 . This is very close to the result obtained by fitting to theoretical isochrones by [Platais et al. \(2001\)](#), who give a distance modulus of 6.35 or 6.38, depending on the models used. However, it was noted that the model isochrones showed shapes different from that of the observation, which means that slightly different answers can be obtained when changing the weights of the stars contributing to the isochrone fit. The summary data for NGC 2451 are given in Table 4. The HR diagram for the cluster is shown in Fig. 15 together with the data for Blanco 1 and the Pleiades. The HR diagrams of all three clusters appear to coincide well when applying the astrometrically derived distance moduli.

7. Clusters between 250 and 500 pc

The astrometric parameters derived for clusters between 250 and 500 pc are generally less secure, as membership of individual stars can be doubtful, and generally few stars are contained in the Hipparcos catalogue. The clusters selected here are all well established, but do suffer, within the Hipparcos catalogue, from low numbers of (sometimes doubtful) members. A reasonably reliable set of astrometric parameters has been obtained for the following clusters: NGC 6475, NGC 7092, NGC 2516, NGC 2232, IC 4665, NGC 6633, Collinder 140, Trumpler 10, NGC 2422, NGC 3532 and NGC 2547. NGC 2264 is also described, but uncertainties in member identification do not allow for conclusions on its existence and astrometric parameters. The results for these clusters are presented in the sections below, including reference to other distance determinations for the same systems and comparisons with the results for nearby clusters of approximately the same age. All cluster solutions have followed the "combined abscissae" model as described in Sect. 4.4. A comparison was also made for NGC 6475 with the method used for the more nearby clusters as described in Sect. 4.3. The observed differences were well below the significance level.

7.1. NGC 6475 (M7)

NGC 6475 is a well populated cluster, which, with an estimated age of 200 to 260 Myr, is older than the Pleiades but younger than UMa and Coma Ber ([Patenaude 1978](#)). Membership has been determined in the past through both

proper motions ([Constantine et al. 1969](#)) and radial velocities by [Buscombe & Kennedy \(1968\)](#) and [Zentelis \(1983\)](#) for the brighter, and [Giesekeing \(1985\)](#) for the fainter potential members. A spectral study was done by [Abt \(1975\)](#) for the brightest members, and combined with *UBV* photometry by [Loden \(1984\)](#). Photometric studies in *ubv* were done by [Snowden \(1976\)](#), [Loden et al. \(1980\)](#) and [Nissen \(1988\)](#). Recent distance modulus estimates have varied from 7.17 ± 0.19 ([Nicolet 1981](#)) to 7.01 ([Nissen 1988](#)) and 7.08 ([Pinsonneault et al. 1998](#)), where the latter is an apparent distance modulus. The reddening of the cluster was determined by [Nissen \(1988\)](#) as $E(b - y) = 0.039 \pm 0.006$, equivalent to 0.053 ± 0.008 in $E(B - V)$. [Pinsonneault et al. \(1998\)](#) use a value of $E(B - V) = 0.06$.

Within the Hipparcos catalogue 17 to 22 stars can be identified as probable members based on their proper motions. They cover an area of about 3.5 degrees in diameter, which is similar, considering the difference in distance, to the area covered by, for example, the Pleiades cluster. For the solution made here 20 stars were finally selected. Solutions with slightly different selections did not produce significantly different results. In particular restricting the extent of the cluster to a 2 degrees diameter only changed the parallax from 3.71 ± 0.14 to 3.70 ± 0.14 . In all aspects the solution to the astrometric parameters of the cluster appears healthy, with only one rejected abscissa measurement out of 1328 in total used. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 18, in comparison with similar data for the Pleiades and Coma Ber.

7.2. NGC 7092 (M39)

A proper motion study by [Ebbighausen \(1940\)](#) and radial velocities by [Trumpler \(1928\)](#) established this group as a cluster with a radial velocity of -14 km s⁻¹, and served as the main membership criterion for the following decades. The individual data for the latter study do not appear to have been published, but some derived information was presented by [Eggen \(1951\)](#), who also determined a photometric parallax of 4.5 mas. Another early photometric study of M39 was presented by [Mävers \(1940\)](#), and covers an area of about 1.6 degrees diameter down to photographic magnitude 13. Based on a *UBV* photometric study, [Johnson \(1953\)](#) determined a distance modulus of 7.2 ± 0.2 , equivalent to a parallax of 3.6 mas. Around the same time, [Weaver \(1953\)](#) presented spectra and photoelectric magnitudes for some 30 potential cluster members. [Meadows \(1961\)](#) established that for the brightest members of M39 the rotational velocities are much like those of field stars of the same spectral types, a study continued and extended to radial velocities and a search for binary stars in the cluster by [Abt & Sanders \(1973\)](#). A few more radial velocities of M39 members were presented by [Geyer & Nelles \(1985\)](#). Using *H β* photometry calibrated to the [Crawford \(1973\)](#) relations, [Kilkenny et al. \(1975\)](#) derived individual distance moduli estimates for the brighter stars in the field, and determined, using *UBV* photometry and the zero-age main sequence by [Blaauw \(1963\)](#), a distance modulus of 6.87 ± 0.25 , as well as a reddening of $E(B - V) = 0.04 \pm 0.03$.

A new proper motion study by [McNamara & Sanders \(1977\)](#) seemed to indicate a rather low abundance of fainter stars in the cluster. From a total of 1710 stars studied, only 30 brighter stars were found to be probable members, and most of those are at $(B - V) < 0.3$. A distance modulus of 7.12 was determined by fitting to the [Johnson \(1957\)](#) calibration sequence, for which a reddening of 0.02 was assumed. The apparent sparsity of fainter




Fig. 16. The distribution of the parallaxes against their formal errors for the IC 2391 (*top left*), IC 2602 (*bottom left*), Blanco 1 (*top right*) and NGC 2451 (*bottom right*) members in the Hipparcos catalogue. The central vertical line represents the mean parallax of each cluster, as presented in Table 4. The black lines on either side show the 1σ error level, including the internal distance dispersion for the cluster. The outer grey lines similarly represent the 2σ level.

members may have been caused by a magnitude equation in the proper motions, as it was contradicted later by Platais (1984) in his proper motion study of the cluster (see also Platais 1994). This study was based on more homogeneous plate material. In it, 43 new (faint) candidate members were identified. The first distance-modulus determination that was at least indirectly based on parallax measurements (Nicolet 1981) provided a value of 7.46 ± 0.12 , equivalent to a parallax of 3.2 mas. A *UBV* photometric study by Mohan & Sagar (1985) based the distance modulus estimate of 7.4 ± 0.2 on a comparison with the zero-age main sequence as given by Schmidt-Kaler in (Voigt 1965), and estimated the age of the cluster to be between 200 and 400 Myr. The interstellar reddening has been determined for the cluster by Manteiga et al. (1991), based on IR photometry, at $E(B - V) = 0.01$. These authors put the age of M 39 to be between 240 and 480 Myr. Finally, Robichon et al. (1999) derived the parallax and distance modulus of M 39 from the first release of the Hipparcos data as 3.22 ± 0.29 mas and 7.46 ± 0.20 mag, respectively.

The new solution, based on 7 member stars, gives a parallax of 3.30 ± 0.19 and therefore a distance modulus of 7.41 ± 0.12 , based on the abscissa residuals for 903 field-of view transits, of which only one was rejected. The standard deviation for the astrometric solution is, at 1.03, close to the expected value. The proper motion of this cluster is relatively large, which eases the member selection. Based on its proper motion, the brightest star in the field, HIP 106346 (HD 205210) has been excluded from the solution. It appears not to be a cluster member, with also its parallax outside the 3σ area. Including this star in the solution brings the parallax down to 3.04 ± 0.17 mas (a distance modulus of 7.59 ± 0.12), which demonstrates how sensitive these solutions

can be to the selection of cluster members. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 20, in comparison with similar data for the Praesepe and Coma Ber.

7.3. NGC 2516

NGC 2516 has been described as a large and rich cluster. A study by Cox (1955) provided the first reliable magnitudes and colours in the region of NGC 2516. From these data, Cox derived a distance estimate of 400 ± 25 pc, and a reddening of $E(B - V) = 0.11$. A spectroscopic study of cluster members by Abt & Morgan (1969) showed a rich variety of peculiar stars to be present in the cluster. Dachs (1970) obtained *UBV* photometry for 70 stars in cluster, and derived a distance of 375 ± 20 pc (parallax of 2.7 mas) and a reddening of $E(B - V) = 0.09$. Dachs noted the presence of three evolved stars, and compared the cluster in age and content with the Pleiades, concluding that it is more than twice as rich. A similar study, but partially extended with measurements in RI and $H\beta$ was presented by Feinstein et al. (1973). In it, the distance of the cluster was estimated at 400 ± 40 pc, and the reddening at $E(B - V) = 0.12$. The age of the cluster was estimated at 60 Myr. Snowden (1975), using *ubv* photometry, estimated the age of the cluster more than twice higher, at 135 Myr, and re-emphasized its similarity with the Pleiades and IC 2602. The reddening was estimated at $E(b - y) = 0.088$. Based on Geneva photometry, Nicolet (1981) derived a distance modulus of 8.15 ± 0.10 , equivalent to a distance of 427 ± 22 pc, or a parallax of 2.3 mas. A study of the cluster in Walraven VBLUW photometry by Verschoor & van Genderen (1983) showed a reddening of the cluster by




Fig. 17. A tangential projection of the distribution on the sky for members of the clusters IC 2391 (*top left*), IC 2602 (*bottom left*), Blanco 1 (*top right*) and NGC 2451 (*bottom right*). The coordinates are in radians, relative to the assumed cluster centre.




Fig. 18. The HR-diagram positions of the stars used in the astrometric solutions to NGC 2516 and NGC 2422 (each compared with the Pleiades, open symbols), and NGC 6475 (compared with the Pleiades, open symbols, and Coma Ber, crosses).

$E(B - V) = 0.127 \pm 0.005$, with a spread among individual cluster members at 0.043 mag. The distance modulus of the cluster was derived as 7.85 ± 0.10 through a comparison with a ZAMS derived by The et al. (1980), which had been derived, through a transformation between photometric systems, from Johnson (1966). Based on Geneva photometric measurements for 67 stars, Cameron (1985) determined a distance modulus of 7.88 and a reddening of 0.12. A UBV study by Dachs & Kabus (1989) of 486 stars in the vicinity of the cluster indicated the presence of 362 cluster members in the sample. A distance modulus of 8.18 ± 0.38 and a reddening of 0.12 were derived. Spectroscopic observations for 36 bright stars in the cluster by González & Lapasset (2000) showed a radial velocity of $22.0 \pm 0.2 \text{ km s}^{-1}$. A CCD $UBVI$ photometric study by Sung et al. (2002) provided a distance modulus of 7.77 ± 0.11 and reddening of 0.112 ± 0.024 through comparisons with theoretical isochrones. The age was determined as 160 Myr.

Dachs & Kabus (1989) identified cluster members till at least $30'$ from the cluster centre. Within this range, 11 stars are found in the Hipparcos catalogue that very closely share the same proper motion (Fig. 19); a further three stars showed similar but not sufficiently coinciding proper motions, and were not used as cluster members. The cluster proper motion and parallax have been estimated by means of the 1166 abscissa residuals of the selected stars, giving a value for the parallax of $2.92 \pm 0.10 \text{ mas}$, equivalent to a distance modulus of 7.68 ± 0.07 . As for the Pleiades, a shorter distance than was expected from the pre-Hipparcos estimates is found. It may be worth noting too that by extending the search radius to 1.3 degrees, 8 more stars are found that closely share the proper motion and distribution of parallaxes of the 11 selected cluster members. Scaled to the distance of the Pleiades, the radius of 1.3 degrees in NGC 2516 is equivalent to a radius of 3.7 degrees. Members of the Pleiades cluster are known at least up to 4 to 5 degrees from the cluster centre (van Leeuwen et al. 1986). Including those stars in the cluster-parallax and proper motion determination gives a solution with very similar statistics as obtained with the initial selection, and a value for the parallax of $2.84 \pm 0.08 \text{ mas}$ instead (distance modulus at 7.73 ± 0.06). Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 18, in comparison with similar data for the Pleiades.

7.4. NGC 2232

The information on this cluster comes predominantly from Clariá (1972), who measured 43 stars in its vicinity in UBV and 22 in $H\beta$. From these data Clariá derived a distance for the cluster of 360 pc, equivalent to a distance modulus of 7.78 and a parallax of 2.8 mas. The eight brightest stars in the list of cluster members by Clariá are included in the Hipparcos catalogue. They do not cluster as well in proper motion as could be expected for a cluster at 360 pc, and two stars in particular, HD 45153 (HIP 30580) and HD 45975 (HIP 31011) stand out in both proper motion and parallax. These stars have been excluded from determining the astrometric parameters of the cluster. The combined abscissa residuals for the six remaining stars give a cluster parallax of $2.83 \pm 0.18 \text{ mas}$. Of the 522 residuals used, 2 were rejected. Including the two stars identified above as possible non-members puts the parallax at $3.27 \pm 0.18 \text{ mas}$, and increases the standard deviation of the solution from 0.96 to 1.04. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown




Fig. 19. Proper motion distribution of the stars identified as probable members of NGC 2516 in the Hipparcos catalogue.

in Fig. 21, in comparison with similar data for NGC 2451 and α Per.

7.5. IC 4665

Next to areas of the Pleiades and Praesepe clusters, the area of IC 4665 was one of the standard regions in the Johnson and Morgan UBV photometric system (Johnson 1954). A study by Hogg & Kron (1955) in UBV identified the cluster as somewhat younger than the Pleiades, at a parallax of about 2.3 mas (distance modulus of 8.2) and a reddening of $E(B - V) = 0.10$ to 0.18, based on a selection of 18 probable members. The membership selection was at least partly supported by the proper motion study of the cluster by Vasilevskis (1955), though this study was of limited accuracy. The cluster main sequence was shown very clearly in a photographic photometry study by McCarthy & O'Sullivan (1969), which incorporated further astrometric studies by Vasilevskis. Three studies in the early 70 s significantly refined the information available on IC 4665, through astrometric data (Sanders & van Altena 1972), photometry in $ubvy\beta$ (Crawford & Barnes 1972) and spectral classifications (Abt & Levato 1975). Crawford & Barnes (1972) derived a distance modulus of 7.5, and a reddening of $E(b - y) = 0.14 \pm 0.04$, where the error refers to the spread over individual cluster members. The proper motion of the cluster is quite small, and as a result there is increasing confusion between members and non-members for the fainter stars, which shows clearly from the HR diagrams of possible members by Sanders & van Altena (1972). Based on data obtained in the Geneva photometric system, Nicolet (1981) derived a distance modulus of 7.90 ± 0.22 , equivalent to a parallax of 2.6 mas. Geyer & Nelles (1985) determined the radial velocity of the cluster as -12.5 ± 4.7 , based on data for 5 stars. Prosser (1993) combined existing and new astrometric, photometric and spectroscopic data for the cluster and identified numerous new candidate members. The age of the cluster was estimated to be between 30 Myr and the age of the Pleiades. The radial velocity of the cluster was determined by Prosser & Giampapa (1994) at about -13 km s^{-1} .

The Hipparcos catalogue contains 6 or 7 stars that can be identified as probable members of the cluster. Four of these are




Fig. 20. The HR-diagram positions of the stars used in the astrometric solutions to NGC 3532 and NGC 7092 (each compared with Coma Ber, crosses, and Praesepe, open circles), and NGC 6633 (compared with the Praesepe, open circles).

contained in the astrometric study by Sanders & van Altena (1972), and identified as probable members in their discussion: S110 (HIP 86954), S115 (HIP 86960), S159 (HIP 87002) and S181 (HIP 87032). One star can be identified in (Vasilevskis 1955) as K 32 (HIP 86805), indicated as a possible member. Two more stars are just outside the areas covered by these studies, HIP 86993 and HIP 87592. The mean parallax as derived from 856 field transit abscissa residuals for these 7 stars is 2.81 ± 0.27 mas, equivalent to a distance modulus of 7.75 ± 0.21 . There were no rejected observations. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 22, in comparison with similar data for the Pleiades, NGC 2451 and α Per.

7.6. NGC 6633

A *UBV* photometric study of NGC 6633 by Hiltner et al. (1958) showed it to be similar in age to Praesepe, and subject to an interstellar reddening of $E(B - V) = 0.17$. They also derived a distance modulus of 7.50 ± 0.25 , equivalent to a parallax of 3.2 mas. A proper motion study covering 207 stars down to photographic magnitude 13.5 in an area of $50' \times 40'$ centred on the cluster was presented by Vasilevskis et al. (1958), and identified some 90 potential cluster members. A study by Sanders (1973) improved this still further, with 497 stars measured and 113 possible members. The problem faced by both these studies is the small size of the proper motions, which makes distinction from background stars difficult. The radial velocity for one of the giant members of the cluster was determined at -22.0 km s $^{-1}$ by Wallerstein et al. (1963), and later by Geyer & Nelles (1985) at -24.8 ± 1.0 . Photometric measurements in *ubvy* β for 37 stars were obtained by Schmidt (1976), who noted a slightly variable extinction with an average value of $E(b - y) = 0.124$, and determined a distance modulus of 7.71, using the calibration by Crawford (1973). A study by Levato & Abt (1977) determined spectral classifications for the brightest cluster members identified in the studies referred to above. Based on Geneva photometric measurements, Nicolet (1981) determined a distance

modulus of 7.47 ± 0.17 , while using *UBV* photometry by Hiltner et al. (1958), Cameron (1985) derived a distance modulus of 7.63 and a reddening of $E(B - V) = 0.17 \pm 0.007$, based on 39 members. Distance fitting was done with respect to the ZAMS as defined by Schmidt-Kaler in Voigt (1965). Based on the first release of the Hipparcos data, Robichon et al. (1999) determined a parallax of 2.70 ± 0.70 mas, using data for 4 stars.

In the new solution, the abscissa residuals of 917 field-of-view transits for 6 stars was used, from which a parallax of 2.67 ± 0.32 mas was derived, equivalent to a distance modulus of 7.87 ± 0.26 . Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 20, in comparison with similar data for Praesepe.

7.7. Collinder 140

A combined photometric and spectroscopic study of Collinder 140 was presented by Clariá & Rosenzweig (1978), who derived a distance modulus of 7.78 (equivalent to a distance of 360 pc), reddening of $E(B - V) = 0.05$, and age of 21 to 23 Myr. From a *UBV* photometric study, Williams (1978) placed the cluster at 420 ± 20 pc and at an age of 40 Myr, and very little reddening, at $E(B - V) = 0.01 \pm 0.01$. A spectroscopic study by Fitzgerald et al. (1980) derived a distance of 410 ± 30 pc, a reddening of $E(B - V) = 0.04$, and age of 20 ± 6 Myr. In his *UBV* photometric study of 38 clusters, Cameron (1985) determined a distance modulus of 7.96 (distance 391 pc) and reddening of $E(B - V) = 0.06$.

The distribution of proper motions and parallaxes in the region of Collinder 140 in the new Hipparcos catalogue shows the presence of 9 probable cluster members, for which the parallax, as derived from the 1413 individual field transits, is 2.66 ± 0.13 mas, equivalent to a distance modulus of 7.88 ± 0.11 . Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 21, in comparison with similar data for NGC 2451 and α Per.

Table 5. Summary data for the distant clusters as derived from combined-abscissae fitting.

Cluster	α, δ degr.	$\log(\text{Age})$ $E(B - V)$	$r(^{\circ})$ N_{stars}	ϖ $\sigma(\varpi)$	μ_{α^*} $\sigma(\mu_{\alpha^*})$	μ_{δ} $\sigma(\mu_{\delta})$	σ_{sol}	N_{absc} N_{rej}	dm $\sigma(\text{dm})$	ρ_1	ρ_2 ρ_3
NGC 6475	268.4	8.22	1.9	3.70	2.06	-4.98	1.04	1328	7.16	-4	1
C175-348	-34.8	0.06	20	0.14	0.17	0.10		1	0.08		-8
NGC 7092	322.9	8.57	0.3	3.30	-8.02	-20.36	1.03	903	7.41	-2	-6
C2130+482	48.4	0.02	7	0.19	0.20	0.19		1	0.12		-14
NGC 2516	119.4	8.08	0.8	2.92	-4.17	11.91	1.03	1166	7.68	0	-2
C0757-607	-60.7	0.12	11	0.10	0.11	0.11		6	0.07		-20
NGC 2232	96.9	7.49	0.8	2.84	-5.65	-2.56	0.96	522	7.73	-9	0
C0624-047	-4.7	0.01	6	0.18	0.16	0.14		2	0.14		4
IC 4665	266.8	7.63	1.8	2.81	-0.90	-8.27	1.04	856	7.75	-5	-16
C1743+057	5.6	0.19	7	0.27	0.24	0.16		0	0.21		-7
NGC 6633	276.5	8.76	0.9	2.67	1.01	-1.15	0.98	917	7.87	10	14
C1825+065	6.7	0.17	6	0.32	0.31	0.28		0	0.26		25
Coll 140	111.0	7.57	0.9	2.66	-7.50	4.02	1.01	1413	7.88	6	-7
C0722-321	-32.1	0.03	9	0.13	0.09	0.13		4	0.11		-3
Trumpler 10	131.7	7.38	1.4	2.58	-12.48	6.81	1.07	1532	7.94	8	6
C0646-423	-42.5	0.06	12	0.16	0.14	0.14		9	0.14		-9
NGC 2422	114.1	8.12	0.3	2.52	-6.72	0.82	1.05	879	7.99	-21	-6
C0734-143	-14.6	0.12	7	0.21	0.16	0.14		0	0.18		11
NGC 3532	166.5	8.45	0.2	2.43	-10.04	4.75	0.90	685	8.07	6	5
C1104-584	-58.7	0.02	6	0.24	0.24	0.21		0	0.22		33
NGC 2547	122.5	7.70	0.7	2.11	-9.34	4.18	0.99	1022	8.38	13	15
C0809-491	-49.2	0.05	8	0.17	0.18	0.16		2	0.17		0

Ages have been obtained from [Kharchenko et al. \(2005\)](#), reddening values come from various papers; the final two columns give the correlation coefficients between the proper motions and parallax determinations times 100 ($\rho_1 \equiv \rho(\varpi, \mu_{\alpha^*})$; $\rho_2 \equiv \rho(\varpi, \mu_{\delta})$; $\rho_3 \equiv \rho(\mu_{\alpha^*}, \mu_{\delta})$).




Fig. 21. The HR-diagram positions of the stars used in the astrometric solutions to Collinder 140, NGC 2232 and Trumpler 10 (each compared with α Per and NGC 2451, crosses).

7.8. Trumpler 10

Trumpler 10 is a not very well studied, fairly young cluster. [Lyngå \(1960\)](#) did a *UBV* photometric study in the region of the cluster, measuring some 29 stars, for which a further analysis was provided by [Lyngå \(1962\)](#), who derived an age estimate of 30 to 60 Myr. Spectroscopic types in the cluster were determined by [Levato & Malaroda \(1975b\)](#), who gave an estimate of the cluster distance of 440 ± 50 pc, equivalent to a distance modulus of 8.22 , and reddening at $E(B - V) = 0.02 \pm 0.01$. A proper

motion study by [Stock \(1984\)](#) seemed to throw some doubt on the existence of the cluster, but the proper motions in this study were not of sufficient accuracy to recognize it.

A search for potential cluster members in the region of Trumpler 10 identified 12 likely cluster members, with a mean parallax of 2.58 ± 0.16 mas, equivalent to a distance modulus of 7.94 ± 0.14 , close to the earlier ground-based estimate. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 21, in comparison with similar data for NGC 2451 and α Per.

7.9. NGC 2422 (M47)

The only two proper motion studies on this cluster are by van Schewick (1966) and Ishmukhamedov (1966), neither of which is easily accessible and played much of role in further investigations of this cluster. A *UBV* photometric study was presented by Smyth & Nandy (1962), and a spectroscopic study by Dworetsky (1975). Nicolet (1981) determined a distance modulus of 8.66 ± 0.16 using Geneva photometry. An extensive study of the cluster in *ubvy β* was carried out by Rojo Arellano et al. (1997), who determined a distance of 470.8 ± 4.8 (distance modulus 8.36) and a reddening of $E(b - y) = 0.089 \pm 0.026$. The age was estimated at 100 Myr. Analysis of the first release of the Hipparcos data by Robichon et al. (1999) showed a parallax of 2.01 ± 0.43 . A CCD-photometry study in *UBVRI* by Prisinzano et al. (2003) was used to study the luminosity and mass functions of the cluster.

The Hipparcos catalogue contains at least 7 stars that are probable cluster members. The abscissa residuals for the 879 field transits of these stars show a cluster parallax of 2.52 ± 0.21 mas, equivalent to a distance modulus of 7.99 ± 0.18 . There were no rejected observations. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 18, in comparison with similar data for the Pleiades.

7.10. NGC 3532

This rich open cluster was studied in photometry and astrometry by Koelbloed (1959), who derived a distance of 432 ± 40 pc (a distance modulus of 8.2), a very small amount of reddening ($E(B - V) = 0.01$), and age of 100 Myr. Further proper motions were obtained in an extensive study by King (1978), who identified 382 likely members in a field of about 1 degree square, down to photographic magnitude 11.0. A *UBV* photometric study of 183 stars in the region of the cluster by Fernandez & Salgado (1980) indicated a slightly larger distance of 490 ± 50 pc (a distance modulus of 8.45 ± 0.27) and age of 200 Myr. Gieseeking (1981) determined a radial velocity of 4.2 km s^{-1} for the cluster, and identified and measured a number of spectroscopic binaries contained in it. The larger distance and higher age seemed to be confirmed by *ubvy β* observations of 33 stars by Eggen (1981), who derived a distance modulus of 8.5 ± 0.25 , a reddening of $E(B - V) = 0.023 \pm 0.014$, and age of 350 Myr. A somewhat smaller distance modulus of 8.296 ± 0.096 was derived by Nicolet (1981) (distance 456 pc). Using photometric observations of 14 cluster stars in *UBV* and *ubvy β* , Johansson (1981) derived a distance modulus of 8.38 ± 0.51 , and a reddening of $E(B - V) = 0.10 \pm 0.04$. Further photometric observations for 68 stars, this time in *UBVRI*, were obtained by Wizinowich & Garrison (1982). This study showed some of the systematic differences that exist between different implementations of the same photometric system. Observations in *UBV* and DDO of 12 red giants in the cluster by Clariá & Lapasset (1988) showed a mean reddening of $E(B - V) = 0.07 \pm 0.02$. In their analysis of the first release of the Hipparcos data, Robichon et al. (1999) derived a parallax of 2.47 ± 0.39 mas for the cluster, equivalent to a distance modulus of 8.04 ± 0.35 .

In the new reduction of the Hipparcos data, 6 stars can be identified as very probable cluster members, having closely the same proper motion. Based on the 685 field-transit measurements for these stars, a cluster parallax of 2.43 ± 0.24 mas is derived, equivalent to a distance modulus of 8.07 ± 0.22 . Details of the solution are shown in Table 5. The HR-diagram positions




Fig. 22. The HR-diagram positions of the stars used in the astrometric solutions to NGC 2547 and IC 4665 (each compared with Pleiades, open symbols and α Per and NGC 2451, crosses).

of the stars used in the astrometric solution are shown in Fig. 20, in comparison with similar data for Praesepe and Coma Ber.

7.11. NGC 2547

Photometric and astrometric studies of NGC 2547 were presented by Fernie (1959, 1960), from which he derived an estimated distance of 360 pc (distance modulus 7.8) and reddening of $E(B - V) = 0.03$. Nicolet (1981) used Geneva photometry to derive a distance modulus of 8.22 ± 0.13 (distance 440 pc). Photoelectric data in *UBV* for 118 stars, and $H\beta$ for 23 stars in the region of NGC 2547 were obtained by Clariá (1982), from which was derived a mean distance of 450 pc (distance modulus 8.3), a reddening of $E(B - V) = 0.06 \pm 0.02$ and age of 57 Myr. Based on *UBV* data, Cameron (1985) derived a distance modulus of 8.1 and reddening of $E(B - V) = 0.05$. Analysis of the first release of the Hipparcos data by Robichon et al. (1999) showed a parallax of 2.31 ± 0.29 . A BVI CCD study by Naylor et al. (2002) gave a distance modulus between 8.00 and 8.15, and age between 20 and 35 Myr.

Eight stars were identified in the Hipparcos catalogue as probable cluster members based on their distribution of proper motions. The cluster parallax of 2.11 ± 0.17 (distance modulus 8.38 ± 0.17) was derived from the abscissa residuals of the 1022 field transits of these stars. There were 2 data points rejected. Details of the solution are shown in Table 5. The HR-diagram positions of the stars used in the astrometric solution are shown in Fig. 22, in comparison with similar data for the Pleiades, NGC 2451 and α Per.

7.12. NGC 2264

NGC 2264 is an extremely young cluster, which, probably for that reason, has been studied well. An extensive photometric survey in *UBV* was presented by Walker (1956), who identified the lower main sequence to lie above the expected locus when comparing his data with the “standard main sequence” of Johnson & Morgan (1953), to which the data were fitted for the B stars. A distance modulus of 9.50 was derived this way, based on which it was concluded that stars of spectral types later than A0 were situated above the main sequence. A proper motion study




Fig. 23. Proper motions for nine stars in the field of NGC 2264. Data points indicates with a larger central dot represent the five stars selected as probable members based on a combination of proper motions, parallaxes and apparent magnitudes.

Table 6. Data on stars selected in the NGC 2264 field.

HIP HD	id p	Hp $B - V$	ϖ σ_{ϖ}	$\mu_{\alpha,*}$ σ_{μ}	μ_{δ} σ_{μ}	g.o.f. rej
31 917		8.717	2.12	-2.57	-0.86	-1.44
47 662		-0.126	1.02	1.02	0.74	0
31 939	V37	8.100	1.49	-1.60	-3.47	0.17
47 731	0.94	-0.143	0.93	0.93	0.65	0
31 951		8.786	1.98	-2.96	-2.84	-1.17
47 751		-0.092	1.14	1.17	0.82	0
31 955	V58	7.904	1.03	-1.96	-4.42	0.12
47 777	0.98	-0.162	0.86	1.00	0.66	0
31 978	S Mon	4.555	3.55	-2.61	-1.61	-0.82
47 839		-0.233	0.50	0.56	0.39	1
32 030	V168	7.470	2.22	-3.16	-2.79	-0.18
47 961	0.98	-0.143	0.69	0.74	0.51	2
32 053	V206	8.942	3.90	-2.48	-3.04	0.00
48 055	0.94	-0.133	1.16	1.56	1.03	2
32 141		8.530	1.97	-4.73	-0.95	-0.95
48 232		0.035	0.95	0.90	0.63	0
32 245		8.373	3.94	-1.32	-3.01	1.61
48 427		0.062	0.95	1.01	0.70	2

by Vasilevskis et al. (1965) identified 140 stars with membership probabilities greater than 0.5 among 245 stars brighter than photographic magnitude 13, in a $40' \times 40'$ field centred on the cluster. A strongly variable reddening for the cluster stars was reported by Young (1978), possibly related to the molecular cloud associated with the cluster (Crutcher et al. 1978), though much less variation was indicated by the spectroscopic study of Barry et al. (1979) and the $UBVRI$ photometric study of Mendoza V. & Gomez (1980). Nicolet (1981) determined a distance modulus of 9.64 ± 0.67 , equivalent to a parallax of 1.2 mas. the radial velocity for the cluster was determined by Liu et al. (1989) as $24 \pm 8 \text{ km s}^{-1}$, based on observations of 12 early-type cluster members. Using a combination of photometric and spectroscopic data, Perez et al. (1987) derived a distance modulus of 9.88 ± 0.17 through a comparison with the main sequence as defined by Schmidt-Kaler in Schaifers et al. (1982).

The Hipparcos catalogue contains at least 4 stars classified by Vasilevskis et al. (1965) as highly probable members (see Table 6). In addition, the catalogue contains the brightest star in the field, S Mon, which has a proper motion very similar to the four stars mentioned above. A little outside the field studied by Vasilevskis et al. (1965), four more stars can be identified to share the same proper motion, and are within the same range of parallax values. Details on all these stars are provided in Table 6, while their proper motions are shown in Fig. 23. Taking the data for all 9 stars together, a parallax of $2.30 \pm 0.46 \text{ mas}$ is obtained, which is why this cluster became included in the initial selection of clusters within 500 pc. This solution shows, however, a relatively large standard deviation of 1.4, and also the combined positions of the selected stars in the HR diagram look suspicious. The stars that seem to contribute most to the large standard deviation are three in the south of the field and S Mon, for which parallaxes near 3.5 mas are found. Eliminating these, a parallax of $1.81 \pm 0.40 \text{ mas}$ is found (from the abscissae solution), and a standard deviation of 1.01. Of the 375 abscissae used, one was rejected in iterations. Given its uncertainty in the parallax determination as well as the final value, the data for this cluster have not been included in Table 5. They have been included in the discussion here to show the vulnerability of these solutions to more distant clusters to member selection criteria.

8. Discussion

One of the aims of this study is to provide for nearby clusters distance moduli that are determined completely independent from main-sequence fitting or other types of photometric calibration. Ultimately, when combined with homogeneous photometry for more extensive selections of cluster members, those data can provide a set of observational isochrones, to be compared both qualitatively and quantitatively with theoretical isochrones. A first, but very limited, combination with photometric data from the Hipparcos catalogue already shows some interesting relations as well as systematic differences. More extensive comparisons can be made when combining the cluster parallaxes with, for example, $ubvy\beta$ or Geneva photometry, which is available for many of the clusters discussed above (Mermilliod 1995).

8.1. Distance modulus and parallax comparisons

In this section the distance moduli as have been determined in various studies over the past 50 years are examined with respect to the new Hipparcos examinations. The comparisons are done as a function of cluster age, reflecting any systematic errors in the calibration sequences used. The ages used come from the catalogue by Kharchenko et al. (2005).

The calibration of the ZAMS by Johnson (1957) has made a large impact, direct as well as indirect, on distance calibrations of open clusters for about half a century. It was built on three assumptions, the distance of the Hyades, assumed to be 40 pc (compared to the current estimate of $46.45 \pm 0.50 \text{ pc}$), an area in the UBV colour-magnitude diagram ($0.55 < B - V < 0.85$) where the Pleiades and Hyades would share the same locus, and a connection between NGC 2362 and the Pleiades around $(B - V) = 0$, which was based on a theoretical evolutionary correction for the magnitudes based on studies by Tayler (1954, 1956). None of these assumptions could be verified at the time, as it is only by means of accurate parallaxes that such verification is made possible. Johnson estimated the uncertainty on the distance moduli determined this way to be of order 0.2 mag. The distance moduli derived by Johnson played an important role




Fig. 24. Differences in distance moduli between values derived by Johnson (1957) and the new Hipparcos reduction (*bottom graph*) and similarly (*middle graph*) for the data for the $ubvy\beta$ photometry (Crawford & Perry 1966; Crawford & Barnes 1969a, 1972, 1974; Crawford & Perry 1976; Crawford & Barnes 1969b), and (*top graph*) data from Cameron (1985). In the *bottom graph* the errors assigned to the distance moduli are mainly the uncertainty of 0.2 mag indicated by Johnson. In the other two graphs the error bars represent only the Hipparcos measurements.

in the early distance moduli determination through $ubvy\beta$ photometry (Crawford & Perry 1966; Crawford & Barnes 1969a, 1972). Figure 24 shows the observed differences between distance moduli for clusters based on these calibrations and the new Hipparcos determination. The distance of the Hyades as used by Johnson (1957) was underestimated by 15 per cent, resulting in errors of around 0.3 mag in the distance estimates for similar clusters, such as Praesepe. The assumption Johnson made about the coincidence of the Pleiades and the Hyades is not supported by the Hipparcos data. In the $ubvy\beta$ photometry it would not be supported either if systematic differences in c_1 are taken into account. The details of the connection between the Pleiades and the youngest open clusters, as implemented by Johnson, are also in need of adjustment.

The ZAMS definition by Johnson (1957) played also an important role in the ZAMS as presented by Schmidt-Kaler in Voigt (1965). This was used by Cameron (1985) for the distance modulus determination of 38 open clusters. As can be observed in Fig. 24, the same systematic differences are observed as for the Johnson and Crawford calibrations.




Fig. 25. Distance modulus differences. *Bottom:* between Becker & Fenkart (1971) and Hipparcos for 15 clusters. Error bars represent Hipparcos determinations only. *Middle:* between Nicolet (1981) and the new reduction for 13 clusters. The error bars represent the combined errors from the two determinations. *Top:* between Loktin & Matkin (1994) and the new reduction for 15 clusters. The error bars represent the Hipparcos determinations only.

The cluster distance determinations by Becker & Fenkart (1971) are at least partly still based on the initial calibrations by Johnson (1957), but with some modifications that affect mainly the young clusters. These modifications, based on incorporating the $(B - U)$ data, appear to have led to a systematic overestimate of the distances of clusters with ages less than about 100 Myr.

A comparison with the distance moduli as determined by Nicolet (1981) (Sect. 5) provides a test on whether, for example, stars in young clusters are of the same luminosity as field stars when found in the same photometric boxes, or whether there are systematic differences, possibly related to age. The comparison, shown in Fig. 25, seems to indicate a systematic overestimate of distance moduli as obtained through the photometric boxes for young clusters, by 0.1 to 0.4 mag, while the older clusters are not significantly offset. One may conclude that the random field stars with ground-based parallaxes as used by Nicolet more resemble the older than the younger cluster stars in luminosity at the same colours. However, the volume of stars with accurate parallax data now available could be used to check in much more detail what the actual luminosity variations of stars sharing photometric boxes are, before drawing any further conclusions.




Fig. 26. Distance modulus differences. Bottom: between Robichon et al. (1999) and Hipparcos for 17 clusters. Error bars represent the old Hipparcos determinations only. Top: between Kharchenko et al. (2005) and the new reduction for 19 clusters. The error bars represent the new Hipparcos reduction only.

Clear systematic differences can be seen in Fig. 25 for the comparison with the distance modulus determinations by Loktin & Matkin (1994). These determinations are based (Loktin & Matkin 1990) on theoretical isochrones by Mermilliod & Maeder (1986), Hejlesen (1980) and Vandenberg (1985), describing evolutionary models for different ranges of age and mass. The systematic differences that can be seen in Fig. 25 are likely to be a reflection of the offsets in those models, and the way these affect clusters of different ages.

An entirely different kind of comparison is that of the current determinations with those based on the first Hipparcos data release. Here the differences for the nearby clusters are small, though the errors have been reduced by about a factor two. For the more distant clusters the membership selection often becomes the critical parameter. This is most clearly demonstrated by IC 4765, for which a parallax based on 7 member stars was presented by Robichon et al. (1999), but for which there is no solution in the current paper. The more accurate proper motions in the area of this cluster in the new catalogue show only three possible members, all of which appear to be red giants. The general agreement for the nearby clusters between the current results and those derived using the 1997 catalogue is not entirely surprising. It relates to the size of the fields covered by most of these clusters compared to the radius within which correlations in the abscissa residuals were able to accumulate to correlations in astrometric parameters for neighbouring stars. Such accumulations have been shown to be only effective over distances of less than 1.5 degrees. For the Pleiades stars, spread over 8 degrees diameter on the sky, there were correlations for the brightest stars in the cluster centre, but all other stars remained largely unaffected.

A comparison with the catalogue of Kharchenko et al. (2005) was also made, but here conclusions are difficult to draw, as in particular for the range of clusters discussed here their methods of distance determination (based amongst others on results by

Robichon et al. 1999) varied. What is clear from the comparison is that the distance moduli by presented by Kharchenko et al. (2005) for the current selection of clusters are on average too large by about 0.15 mag.

In general, for many of the distance calibrations referred to above the same issue applied. Distance calibrations were presented as accurate, while not considering the inaccuracies and remaining noise levels of the actual calibration model itself. While a lot of attention was paid to exact reddening corrections, and sometimes metallicity variations, the most basic of errors as well as noise contribution, in the reference ZAMS, was generally not considered as a contribution to the final accuracies. Calibrations depended for a long time, directly or indirectly, on Johnson (1957), which itself was based on a distance estimate of the Hyades that was short by 15 per cent, an unverified assumption on the relation between the Pleiades and Hyades main sequences, and a very early estimate of the evolutionary luminosity effects on B stars. It is only now, when distances can be measured with completely independent methods, that the limitations of these assumptions are exposed and can be corrected for.

8.2. HR diagram comparisons

When examining the composite HR diagram of the clusters a few distinct groups appear, for which the distributions of stars largely coincide in colour–magnitude and colour–colour diagrams. Photometric data in the $ubvy\beta$ and Geneva systems was extracted from WEBDA¹ to further examine these groups. Additional photometry for UMa stars was obtained from Rufener (1988) and Hauck & Mermilliod (1998). In most cases field stars had to be removed from the selected data, but this was only done when a cluster main sequence could be clearly recognized. Clusters for which this was not possible have been left out. For the Hyades only stars were selected for which a kinematic parallax had been determined based on the new Hipparcos reduction, as described in Sect. 6.1. Thus, individual parallaxes were available for the selected stars. Directly measured individual parallaxes were used for the UMa stars.

The (provisional) groups contain the following clusters for which more extensive photometry is available:

1. the Hyades and Praesepe;
2. Coma Ber, UMa and NGC 7092;
3. Pleiades, Blanco 1 and NGC 2516;
4. α Per, NGC 2451, IC 4665, IC 2391, IC 2602 and NGC 6475.

The groups contain mostly clusters of similar age, though NGC 6475 appears a little out of place in group 4. For display, groups 1 and 2 have been taken together, and are shown in Fig. 27. A couple of observations can be made from this composite HR diagram:

- as was noted earlier, the Hyades and Praesepe data very closely overlap, and together form a very narrow sequence;
- the Coma Ber and UMa main sequences also overlap closely, and are sub luminous with respect to Hyades and Praesepe;
- NGC 7092 appears to follow Coma Ber and UMa for $0.1 < (B2 - V1) < 0.3$;
- increases in luminosity are observed for Hyades and Praesepe stars relative to Coma Ber and UMa for $(B2 - V1) < 0.2$;

¹ <http://www.univie.ac.at/webda/>




Fig. 27. The composite HR diagrams for groups 1 & 2 (left) and group 3 (right). Symbols on the left: Hyades, closed circles; Praesepe, open circles; UMa, closed triangles; Coma Ber, open triangles; NGC 7092, crosses. Symbols on the right: Pleiades, closed circles; Blanco 1, open circles; NGC 2516, crosses.

- increases in luminosity are observed for Coma Ber and UMa stars relative to NGC 7092 for $(B2 - V1) < 0.0$.

The third group contains three clusters which have been noted in the past as being very similar (Snowden 1975; Mermilliod et al. 2008). They share the same sequences in the HR diagram when their distance moduli are obtained from the Hipparcos parallaxes, as can be observed in Fig. 27. However, over the range $0.2 < (B2 - V1) < 0.65$ the combined sequence of the group-3 clusters is shifted by about 0.4 mag with respect to the group-1 sequence, and by about 0.25 mag with respect to the group-2 sequence. As current models providing theoretical isochrones appear not to accommodate for such shifts in luminosity, except through extreme assumptions on chemical composition, distance modulus estimates based on isochrones and those based on parallaxes remain systematically discrepant for clusters of the first and third groups. However, it should also be realized that isochrone calibration such as presented by (Pinsonneault et al. 2004) is based only on the position of the Hyades main sequence, and does not include an empirical calibration of the age-dependent luminosity effects. These are assumed to be all already fully covered by the theoretical models. The new result for the Pleiades, and its effective confirmation through NGC 2516 and Blanco 1, which both independently had been described to be very similar to the Pleiades, seem to show that this is

probably not the case. The issue of the Pleiades distance is still far from “largely settled”, as has been claimed by Pinsonneault et al. (2004) and An et al. (2007).

The clusters in group 4 are mostly young to very young, except for NGC 6475. In the region $0.15 < (B2 - V1) < 0.45$ the sequence fits well with the Coma Ber and UMa sequence, and is offset by about 0.15 mag from the group-1 sequence (Fig. 28). For values of $(B2 - V1) < -0.1$ the sequences of group 3 and 4 coincide. One complication with the group-4 sequence is the lack of well-studied actual clusters. The sequence for α Per shows more variation than is observed for Pleiades, Hyades and Praesepe, while the clusters IC 2639 and IC 2602 are relatively sparse.

8.3. The Strömgren c_0 index for F-type stars

The c_0 index (c_1 corrected for reddening) in the Strömgren photometry was shown by Crawford (1975) to be related to luminosity and surface gravity variations for F-type stars (see also Sect. 5.1). Variations in c_0 at a given value of β or $b - y$ are given by Δc_0 , i.e. c_0 relative to the calibration relation given in Crawford (1975) (see also Fig. 30). This is the relation described in Sect. 5.1 by Eq. (24). It is further evaluated in the Appendix, Sect. A. The difference in the Δc_0 values for the Hyades and




Fig. 28. The composite diagram for clusters in group 4 (see text).

Praesepe with respect to, for example, the Pleiades, has been attributed in the past to increased helium abundances (Nissen 1988; Dobson 1990) (see also Crawford & Barnes 1969a), although isochrone fitting by An et al. (2007) seems to suggest that this is not the case. It has been known for nearly half a century and has been referred to in the past as the Hyades anomaly. It is most clearly observed in the $(U - B)$ index, which is a measure of the Balmer jump for early-type stars. The significant difference in the Geneva $(U - B)$ index for the clusters in group 1 and group 3 can be seen in Fig. 29. The Strömgren c_0 index is closely related to the $(U - B)$ index.

A comparison between the c_0 versus $b - y$ diagrams for clusters in the Pleiades and Hyades groups by Nissen (1988) showed that the systematic difference in c_0 values between those clusters is about 0.033 to 0.04. Following Eq. (A.7), this would correspond to an expected difference of 0.3 to 0.4 in absolute magnitude, which is close to what is observed in the HR diagram for F stars in these clusters when using the Hipparcos parallax-based distance moduli. It is in this context also interesting to note that Nissen (1988) determined the difference between the true distance moduli, i.e. corrected for the c_0 effects, for the Pleiades and Hyades as 1.96 (which would imply a distance of 114 pc for the Pleiades), compared to 2.066 ± 0.038 determined by means of the Hipparcos data and 2.33 as derived from isochrone fitting (An et al. 2007). In a way, by calibrating their isochrones on the Hyades main sequence, An et al. (2007) have turned the “Hyades anomaly” into a “Pleiades anomaly”. An extensive discussion on how the Hyades anomaly may be related to the differences in parallax and isochrone-based cluster distance determinations was presented by Mermilliod et al. (1997).




Fig. 29. The $U - B$ versus $B2 - V2$ diagram in the Geneva photometry. The red dots represent data points in the Hyades and Praesepe clusters, the crosses for data from the Pleiades, Blanco 1 and NGC 2516. Some of the Pleiades points are offset because of localised differential reddening in the cluster. The difference in loci for the two groups of clusters is known as the Hyades anomaly.

A systematic difference of 0.033 to 0.04 in c_0 values would indicate, following Eq. (A.10), a lower surface gravity for the Hyades F stars by approximately 0.1, but that is also the noise level on that calibration, i.e. the same $\log g$ values can be observed for Δc_1 values that differ by 0.03 to 0.04. An explanation of the magnitude difference as due to metallicity variations would require an unrealistic, and also not observed, difference in the values for Δm_1 of around 0.15.

It is not clear why Stello & Nissen (2001) made no reference to the c_0 index differences, as discussed earlier by Nissen (1988), in their discussion on the Strömgren photometry and the Hipparcos parallax measurement for the Pleiades. From a purely observational point of view, a much smaller difference in absolute magnitudes between the Pleiades and Hyades stars (the result of a larger difference in distance moduli), as suggested by fits on theoretical isochrones, would be more difficult to explain, as this would contradict the general trend which can be clearly observed between the parameters ΔM_V and Δc_0 for F-type field stars (Fig. A.3). A similar conclusion of internal observational consistency for the Hipparcos cluster parallaxes was reached by Grenon (2002), based on comparisons of K stars in the Hyades, Pleiades and Praesepe clusters.

The relations used here show significant noise levels, and their applications do not provide exact results, they are merely an indication of trends and provide estimates of the effects differences in certain parameters may have. A comparison of F stars in the Hyades, Coma Ber and Pleiades clusters, for example, shows that the overall differences in c_1 are in the same direction and at




Fig. 30. Comparison of the c_1 (top) and m_1 (bottom) versus $H\beta$ relations for F stars in the Hyades (filled circles), Coma Ber (crosses) and the Pleiades (open circles). The calibration relation from Crawford (1975) is shown by filled triangles.

about the same amplitude as would be expected from the observed differences in the absolute magnitudes (Fig. 30), but it is difficult to assess whether there is not only a qualitative, but also a full quantitative agreement.

9. Space velocities

For general reference, the positions and space velocities for all 20 clusters and the individual UMA stars have been derived in galactic coordinates:

$$\begin{bmatrix} X \\ Y \\ Z \end{bmatrix} = R \cdot \begin{bmatrix} \cos l \cos b \\ \sin l \cos b \\ \sin b \end{bmatrix}, \quad (36)$$

and its derivative

$$\begin{bmatrix} u \\ v \\ w \end{bmatrix} = V_R \cdot \begin{bmatrix} \cos l \cos b \\ \sin l \cos b \\ \sin b \end{bmatrix} + \frac{\kappa \mu_{l*}}{\varpi} \begin{bmatrix} -\sin l \\ \cos l \\ 0 \end{bmatrix} - \frac{\kappa \mu_b}{\varpi} \begin{bmatrix} \cos l \sin b \\ \sin l \sin b \\ -\cos b \end{bmatrix}, \quad (37)$$

where $\kappa = 4.74047$ is the transformation factor for a proper motion of “1 mas yr⁻¹ at 1 kpc” to “km s⁻¹”. The accumulated results for the positions and space velocities are shown in Table 7. The space velocities are compared with those of the A and F stars within 125 pc (van Leeuwen 2007a) in Fig. 31.

The most striking of velocity coincidences is between the Hyades and Praesepe, which, within the accuracy of the measurements, have identical u and v velocities. Considering that these two clusters are also very similar in photometric aspects, it seems reasonable to assume a common origin, which would then also explain their very similar age and chemical composition.

There are no such striking coincidences between the UMA stars and Coma Ber, though both groups are somewhat anomalous in the distribution of velocities or positions. There is some similarity between the Pleiades and NGC 2516, but Blanco 1 does not seem to fit.




Fig. 31. Distribution of the velocities of 20 clusters and 8 UMA stars within the galactic plane, compared with the velocity distribution of A and F stars within 125 pc. The UMA stars form a small clump, and there are a few other coincidences in space velocity. The solar motion has not been subtracted.

10. Conclusions

The new reduction has eliminated problems in the Hipparcos astrometric data that were caused by peculiarities in the satellite dynamics and an occasional potential weakness in connectivity between the two fields of view. These were also issues raised by (Makarov 2002, 2003, 2006), but these effects can not be eliminated at the intermediate-data level, as had been shown already when the idea of such corrections was first presented by van Leeuwen (1999b). The new reduction dealt with these attitude problems there where they occurred (in the along-scan attitude reconstruction process) rather than with the indirect and diffuse effects they had on great-circle abscissa residuals. The attitude reconstruction is the only place in which they can be (and now have been) identified and corrected for. With also the connectivity issue having been addressed and tested in the new reduction, the current set of cluster parallax and proper motion values should not only be more precise, but also more accurate.

Differences between isochrone-fitting and parallax-based distance moduli for open clusters are not limited to the Pleiades, they are found to be systematic. The differences are observed between groups of clusters that show similar characteristics in for example the c_0 and $U - B$ indices for F-type stars, and could therefore be related to systematic surface gravity differences. A closer examination of field star data confirms that such variations are correlated to differences in absolute magnitudes, though not in a simple one-to-one relation. Differences are also observed between the Hyades and Praesepe group and Coma Ber and UMA, where none of the old “Hipparcos problems” ever applied, as concentrations of stars in these clusters are not dense enough. The notion that the parallax values for all these clusters have to be wrong because they do not fit the results obtained by traditional main sequence fitting, or the distance measurement for a single binary star in the cluster, such as expressed by Trimble et al. (2006), gives the impression that reduced data has to confirm theoretical models to be acceptable as results, rather than

Table 7. Positions and velocities of 20 clusters in the galactic reference frame.

Name	l	b	X	Y	Z	u	v	w
	degr.		pc			km s ⁻¹		
Hyades	179.1	-21.9	-43.1	0.7	-17.3	-41.1 ± 0.9	-19.2 ± 0.2	-1.4 ± 0.4
Coma Ber	221.8	83.8	-7.0	-6.3	86.2	-2.3 ± 0.1	-5.4 ± 0.1	-1.8 ± 1.0
Pleiades	166.4	-23.7	-107.0	25.9	-48.3	-6.7 ± 0.9	-25.0 ± 0.5	-12.8 ± 0.5
Praesepe	205.8	32.4	-138.4	-67.0	97.6	-41.5 ± 0.9	-19.8 ± 0.5	-9.7 ± 1.1
α Per	147.8	-6.1	-145.1	91.3	-18.2	-12.7 ± 0.9	-24.6 ± 0.7	-7.0 ± 0.2
IC 2391	270.4	-6.9	0.9	-143.9	-17.5	-22.7 ± 0.5	-18.5 ± 1.0	-6.3 ± 0.3
IC 2602	289.4	-5.0	49.2	-139.6	-13.0	-8.2 ± 0.4	-19.9 ± 0.9	-0.3 ± 0.2
Blanco 1	14.1	-79.3	37.4	9.4	-203.4	-17.7 ± 1.1	-6.7 ± 0.5	-7.1 ± 1.0
NGC 2451	252.5	-7.7	-54.8	-173.4	-24.6	-26.0 ± 0.6	-13.0 ± 1.0	-12.6 ± 0.3
NGC 6475	355.8	-4.5	268.7	-19.6	-21.0	-14.8 ± 1.0	-3.1 ± 0.3	-4.4 ± 0.3
NGC 7092	92.4	-2.2	-12.4	302.5	-11.9	28.7 ± 1.7	-6.9 ± 1.0	-13.2 ± 0.8
NGC 2516	273.7	-15.9	21.5	-328.7	-93.7	-18.8 ± 0.7	-23.9 ± 1.0	-3.4 ± 0.3
NGC 2232	214.4	-7.4	-288.0	-197.4	-45.6	-14.0 ± 0.8	-9.1 ± 0.6	-12.6 ± 0.7
IC 4665	30.6	16.8	293.1	173.5	103.0	-2.8 ± 1.1	-16.9 ± 1.2	-8.5 ± 0.7
NGC 6633	36.0	8.7	299.4	217.8	56.6	-22.3 ± 0.9	-17.5 ± 0.7	-6.9 ± 0.6
Coll 140	245.1	-7.8	-156.8	-337.9	-51.0	-22.2 ± 0.7	-18.1 ± 0.9	-12.2 ± 0.5
Trump 10	262.7	0.5	-49.2	-384.4	3.3	-27.1 ± 1.5	-21.8 ± 1.0	-9.8 ± 0.7
NGC 2422	231.0	3.0	-249.2	-308.1	21.0	-29.2 ± 0.8	-24.2 ± 0.9	-8.3 ± 0.9
NGC 3532	289.6	1.4	137.9	-387.6	10.2	-19.3 ± 2.1	-10.4 ± 1.2	0.9 ± 0.5
NGC 2547	264.4	-8.6	-45.4	-466.4	-71.0	-20.2 ± 1.6	-10.4 ± 1.0	-14.8 ± 1.1
HIP 51814	152.2	51.6	-14.6	7.7	20.8	12.5 ± 0.6	3.2 ± 0.3	-5.7 ± 0.8
HIP 53910	149.1	54.8	-12.1	7.2	20.0	13.5 ± 0.5	2.9 ± 0.3	-7.5 ± 0.8
HIP 58001	140.8	61.4	-9.5	7.7	22.4	15.6 ± 0.4	2.9 ± 0.3	-8.8 ± 0.9
HIP 59774	132.5	59.5	-8.5	9.2	21.3	14.9 ± 0.3	1.5 ± 0.4	-10.3 ± 0.9
HIP 62956	122.2	61.1	-6.5	10.3	22.2	14.1 ± 0.3	2.8 ± 0.4	-7.8 ± 0.9
HIP 63503	120.3	60.7	-6.3	10.8	22.2	13.1 ± 0.3	3.3 ± 0.4	-9.1 ± 0.9
HIP 64532	116.8	60.2	-5.7	11.2	21.9	14.1 ± 0.3	2.3 ± 0.5	-7.8 ± 0.9
HIP 65477	112.8	61.5	-4.6	11.0	22.0	14.1 ± 0.2	3.3 ± 0.4	-8.8 ± 0.9

The 8 HIP numbers are all UMa members, treated on an individual basis; the errors on the space velocities are in some cases uncertain as a result of uncertainty in the radial velocity accuracy.

that those data are judged upon the merits of the observations they have been derived from, and the quality of the data reductions. The Pleiades story over the past ten years has shown a number of examples where poor results, sometimes even based on flawed methods, were accepted (and quoted) without reservation, simply because they confirmed expectations. In this context it is worth while recalling a statement made by Crawford (1973) in his Survey Lecture at the IAU General Assembly in 1970: *In many cases, theory can help us observers quite a lot. We would be lost, or at least inefficient, without theory to guide us. However, we must be extremely careful not to force-fit or to let pre-conceived ideas mess us up. We are measuring observed parameters, and these we relate, or calibrate, to physical parameters.*

Acknowledgements. Discussions with various colleagues over the past couple of years have contributed to this paper, in particular with Robin Catchpole, Derek Jones, and Simon Hodgkin. Carme Jordi, Mike Irwin, Bob Carswell, Jan Lub, Anthony Brown and George Wallerstein read earlier drafts and provided much appreciated comments and suggestions.

Appendix A: Provisional (re-)calibration of Strömgren indices

The currently available volume of information available on F stars (parallaxes, photometric and spectroscopic data), allows for a much more detailed calibration of the Strömgren photometric indices with absolute magnitudes, metallicities, and spectroscopically determined effective temperatures and surface gravity values. Presented here are the provisional results of a more extensive study currently in preparation. It has been included here

in order to see how the observed systematic differences in colour indices and absolute magnitudes can be related to physical parameters as calibrated from nearby field stars.

The photometric data have been extracted from the *ubvy β* catalogue by Hauck & Mermilliod (1997, 1998). The spectroscopic data has been extracted from Cayrel de Strobel et al. (2001). For this exploratory exercise the data for multiple entries in the latter catalogue were simply averaged. The facilities of the Vizier² web page were used to extract samples of F stars, which were later further “cleaned” from luminosity class I and II stars and binaries. Only data with Hipparcos parallax determinations better than 10 per cent were kept, and all stars fainter than apparent magnitude about 10 were left out. This criterion is only approximate, as close to the limit a specific star may be fainter in one catalogue, and still just bright enough in another.

The first calibration done was a more detailed repeat of the luminosity calibrations of (Crawford 1975). This involved two linked calibrations, namely between absolute v magnitude (in what follows, v is always assumed to be corrected for the parallax-based distance modulus) and $b - y$, which determined dv values, and between the observed values of c_1 and $H\beta$ as a function of dv . The iterations use what is observed in the data, and was also implemented by Crawford, that stars occupying the lower boundary in the v versus $b - y$ diagram also occupy the lower envelope in the c_1 versus $H\beta$ diagram. Thus, the following relations were obtained, using only stars with Δc_1 values less than 0.02:

$$v_0(b - y) = (3.891 \pm 0.010)$$

² <http://webviz.u-strasbg.fr/viz-bin/VizieR>




Fig. A.1. The relation between $H\beta$ and c_1 in the Strömgren system for nearby F type stars with parallax errors below 10 per cent. The curved lines of black dots represent *from bottom to top* the calibration relation at Δv values of 1.2, 0.2, -0.8 and -1.8 mag. The crosses show the calibration relation determined and used by Crawford (1975).

$$+ (10.66 \pm 0.22) (b - y - 0.32), \quad (\text{A.1})$$

where the standard deviation was found to be 0.27 mag. The range in $b - y$ covered is 0.21 to 0.43. The values of Δc_1 were determined relative to the following calibrated relation, where $d\beta \equiv (H\beta - 2.64) \times 10$ ($d\beta$ thus ranges from -0.5 to $+0.8$), and $\Delta v \equiv v - v_0(b - y) + 0.75$:

$$\begin{aligned} c_1 = & (0.4370 \pm 0.0007) + (-0.049 \pm 0.010) \Delta v \\ & + (0.2205 \pm 0.0029) d\beta + (-0.032 \pm 0.003) \Delta v d\beta \\ & + (0.086 \pm 0.007) d\beta^2 + (-0.029 \pm 0.007) \Delta v d\beta^2 \\ & + (0.060 \pm 0.013) d\beta^3, \end{aligned} \quad (\text{A.2})$$

with a standard deviation of 0.029 mag. This relation is shown in Fig. A.1 for four different values of Δv superimposed on the actual data points. The lower curve, for $\Delta v = 1.2$, serves as the reference values $c_1(r)$ for calculating the Δc_1 values

$$c_1(r) \equiv 0.3782 + 0.1833 d\beta + 0.0512 d\beta^2 + 0.060 d\beta^3, \quad (\text{A.3})$$

and $\Delta c_1 \equiv c_1 - c_1(r)$.

For determining the Δm_1 values the reference function as presented by Crawford (1975) is used, which accurately fits a second-order polynomial

$$m_1(r) \equiv 0.1826 - 0.0550 d\beta + 0.0607 d\beta^2, \quad (\text{A.4})$$

where the standard deviation with respect to the discrete points defined by Crawford is 0.0005 mag (Fig. A.2). As was done above for Δc_1 , we define $\Delta m_1 \equiv m_1 - m_1(r)$. Contrary the situation for Δc_1 , there is no easily recognized relation between Δm_1 and the absolute magnitude differences Δv .

For completeness also the relation between $b - y$, Δc_1 , Δm_1 and $H\beta$ has been derived:

$$\begin{aligned} b - y = & (0.3376 \pm 0.0005) + (-0.1695 \pm 0.0013) d\beta \\ & + (0.0446 \pm 0.0021) d\beta^2 + (-0.0506 \pm 0.0048) \Delta c_1 \\ & + (-0.589 \pm 0.012) \Delta m_1 + (0.453 \pm 0.036) \Delta m_1 d\beta, \end{aligned} \quad (\text{A.5})$$

with a standard deviation of 0.014 mag. To compare with Crawford's original calibration, which was expressed in




Fig. A.2. The relation between $H\beta$ and m_1 for nearby F dwarfs with parallax accuracies better than 10 per cent. The line represents the definition of the lower envelope position by Crawford (1975).




Fig. A.3. Calibration of Δv . The four lines of black dots show (*from bottom to top*) the calibration relation for values of $\Delta c_1 = -0.02, 0.04, 0.10, 0.16$, and $\Delta m_1 = 0.015$.

$\delta\beta = 2.72 - \beta$, the following equivalent relation is obtained:

$$\begin{aligned} b - y = & (0.231 \pm 0.001) + (0.981 \pm 0.028) \delta\beta \\ & + (4.46 \pm 0.21) \delta\beta^2 + (-0.0506 \pm 0.0048) \Delta c_1 \\ & + (-0.227 \pm 0.035) \Delta m_1 + (-4.53 \pm 0.36) \Delta m_1 \delta\beta. \end{aligned} \quad (\text{A.6})$$

The main differences with respect to Crawford's original calibration are in the higher order terms, which are less well determined for the much smaller volume of data that was available to Crawford at the time of the calibration.

The calibration in terms of absolute magnitudes has been done for Δv as a function of $d\beta$, Δc_1 and Δm_1 :

$$\begin{aligned} \Delta v = & (-0.085 \pm 0.014) + (-0.053 \pm 0.029) d\beta \\ & + (-9.66 \pm 0.17) \Delta c_1 + (7.42 \pm 0.41) d\beta \Delta c_1 \\ & + (0.201 \pm 0.55) d\beta^2 + (2.43 \pm 0.31) \Delta m_1, \end{aligned} \quad (\text{A.7})$$

with a standard deviation of 0.34 mag. Important here is that we are now able to recognize the effect of variation in the metallicity index Δm_1 on the absolute magnitudes. The calibration relation is shown in Fig. A.3.




Fig. A.4. The relation between Δm_1 and Fe/H for 271 F stars. The calibration for $H\beta = 2.60, 2.65, 2.70$ is shown by the diagonal lines, with the lowest value of $H\beta$ as the *top* line.

Spectroscopic data for 271 F dwarfs with *ubv* β photometry were selected from Cayrel de Strobel et al. (2001). For each star also spectroscopically determined effective temperatures and surface gravity values are provided, allowing for a direct comparison between these physical parameters and the Strömrgren indices. The first of those relations is between $d\beta$ and the T_{eff} :

$$T_{\text{eff}} = (6294 \pm 4) + (861 \pm 16) d\beta \text{ K}, \quad (\text{A.8})$$

with a dispersion of 69 K. Thus, the stars examined here would range in temperature from 5862 ± 9 to 6983 ± 13 K.

The situation for the calibration of Δm_1 as a function of Fe/H is less simple due to the dependence on $H\beta$ and the uneven distribution and relatively small number of calibration points. The relation used in the final fit is as follows

$$\text{Fe/H} = (0.179 \pm 0.012) + (-0.119 \pm 0.038) d\beta \\ + (-11.21 \pm 0.38) \Delta m_1 + (-3.37 \pm 1.05) \Delta m_1 d\beta, \quad (\text{A.9})$$

with a standard deviation of 0.10. The data and the calibration are shown in Fig. A.4. These results are in general agreement with similar calibrations presented by Crawford (1975) and Gustafsson & Nissen (1972). The final calibration concerns the surface gravity $\log g$ as a function of Δc_1 . There is a generally well-defined relation:

$$\log g = (4.294 \pm 0.010) + (-2.91 \pm 0.14) \Delta c_1, \quad (\text{A.10})$$

with a standard deviation of 0.10. It is noted, however, that for the later-type stars more data points are found with significant disagreement between the $\log g$ and Δc_1 values. The data and the calibration are shown in Fig. A.5.

In general, when applying any of the above relations the standard deviation of the fit should be used as a measure of the uncertainty in the result obtained.

References

Abt, H. A. 1975, *PASP*, 87, 417
 Abt, H. A. 1978, *PASP*, 90, 692
 Abt, H. A., & Morgan, W. W. 1969, *AJ*, 74, 813
 Abt, H. A., & Sanders, W. L. 1973, *ApJ*, 186, 177
 Abt, H. A., & Levato, H. 1975, *PASP*, 87, 849
 An, D., Terndrup, D. M., Pinsonneault, M. H., et al. 2007, *ApJ*, 655, 233




Fig. A.5. The relation between Δc_1 and $\log g$ for 271 F stars of luminosity types III, IV and V. The diagonal line is the calibration curve.

Barry, D. C., Cromwell, R. H., & Schoolman, S. A. 1979, *ApJS*, 41, 119
 Baumgardt, H. 1998, *A&A*, 340, 402
 Becker, W., & Fenkart, R. 1971, *A&AS*, 4, 241
 Bertiau, F. C. 1958, *ApJ*, 128, 533
 Blaauw, A. 1963, *The Calibration of Luminosity Criteria* (University of Chicago Press, Chicago), 383
 Blanco, V. M. 1949, *PASP*, 61, 183
 Braes, L. L. E. 1961, *Monthly Notes of the Astronomical Society of South Africa*, 20, 7
 Braes, L. L. E. 1962, *Bull. Astron. Inst. Netherlands*, 16, 297
 Buscombe, W., & Kennedy, P. M. 1968, *MNRAS*, 139, 215
 Cameron, L. M. 1985, *A&A*, 147, 39
 Cayrel de Strobel, G., Soubiran, C., & Ralite, N. 2001, *A&A*, 373, 159
 Chen, B. 1999, *A&A*, 344, 494
 Clariá, J. J. 1972, *A&A*, 19, 303
 Clariá, J. J. 1982, *A&AS*, 47, 323
 Clariá, J. J., & Rosenzweig, P. 1978, *AJ*, 83, 278
 Clariá, J. J., & Lapasset, E. 1988, *MNRAS*, 235, 1129
 Constantine, S. M., Harris, B. J., & Nikoloff, I. 1969, *Proc. Astron. Soc. Aus.*, 1, 207
 Cox, A. N. 1955, *ApJ*, 121, 628
 Crawford, D. L. 1973, in *Problems of Calibration of Absolute Magnitudes and Temperature of Stars*, ed. B. Hauck, & B. E. Westerlund, *IAU Symp.*, 54, 93
 Crawford, D. L. 1975, *AJ*, 80, 955
 Crawford, D. L. 1978, *AJ*, 83, 48
 Crawford, D. L. 1979, *AJ*, 84, 1858
 Crawford, D. L., & Perry, C. L. 1966, *AJ*, 71, 206
 Crawford, D. L., & Barnes, J. V. 1969a, *AJ*, 74, 818
 Crawford, D. L., & Barnes, J. V. 1969b, *AJ*, 74, 407
 Crawford, D. L., & Barnes, J. V. 1972, *AJ*, 77, 862
 Crawford, D. L., & Barnes, J. V. 1974, *AJ*, 79, 687
 Crawford, D. L., & Perry, C. L. 1976, *AJ*, 81, 419
 Creze, M., Turon Lacarrieu, C., Golay, M., & Mandwewala, N. 1980, *A&A*, 85, 311
 Crutcher, R. M., Hartkopf, W. I., & Giguere, P. T. 1978, *ApJ*, 226, 839
 Dachs, J. 1970, *A&A*, 5, 312
 Dachs, J., & Kabus, H. 1989, *A&AS*, 78, 25
 Dambis, A. K. 1999, *Astron. Lett.*, 25, 7
 de Bruijne, J. H. J., Hoogerwerf, R., & de Zeeuw, P. T. 2001, *A&A*, 367, 111
 de Epstein, A. E. A., & Epstein, I. 1985, *AJ*, 90, 1211
 de Zeeuw, P. T., Hoogerwerf, R., de Bruijne, J. H. J., Brown, A. G. A., & Blaauw, A. 1999, *AJ*, 117, 354
 Dobson, A. K. 1990, *PASP*, 102, 88
 Domingo, A., & Figueras, F. 1999, *A&A*, 343, 446
 Dworetzky, M. M. 1975, *AJ*, 80, 131
 Ebbighausen, E. G. 1939, *ApJ*, 89, 431
 Ebbighausen, E. G. 1940, *ApJ*, 92, 434
 Eggen, O. J. 1951, *ApJ*, 113, 657
 Eggen, O. J. 1970, *ApJ*, 161, 159
 Eggen, O. J. 1981, *ApJ*, 246, 817
 ESA 1992, *The Hipparcos Input Catalogue*, SP No. 1136 (ESA)
 ESA 1997, *The Hipparcos and Tycho Catalogues*, SP No. 1200 (ESA)

- Feinstein, A., Marraco, H. G., & Mirabel, I. 1973, *A&AS*, 9, 233
- Fernandez, J. A., & Salgado, C. W. 1980, *A&AS*, 39, 11
- Fernie, J. D. 1959, *Monthly Notes of the Astron. Soc. S. Afr.*, 18, 57
- Fernie, J. D. 1960, *Monthly Notes of the Astron. Soc. S. Afr.*, 19, 120
- Fernie, J. D. 1965, *AJ*, 70, 575
- Fitzgerald, M. P., Miller, M., & Harris, G. L. H. 1980, *MNRAS*, 191, 95
- Fox Machado, L., Pérez Hernández, F., Suárez, J. C., Michel, E., & Lebreton, Y. 2006, *Mem. Soc. Astron. Ital.*, 77, 455
- Gatewood, G. 1995, *ApJ*, 445, 712
- Gatewood, G., & de Jonge, J. K. 1994, *ApJ*, 428, 166
- Gatewood, G., Castelaz, M., Han, I., et al. 1990, *ApJ*, 364, 114
- Gatewood, G., Kiewiet de Jonge, J., & Persinger, T. 1998, *AJ*, 116, 1501
- Gatewood, G., de Jonge, J. K., & Han, I. 2000, *ApJ*, 533, 938
- Geyer, E. H., & Nelles, B. 1985, *A&AS*, 62, 301
- Giannuzzi, M. A. 1995, *A&A*, 293, 360
- Gieseking, F. 1981, *A&A*, 99, 155
- Gieseking, F. 1985, *A&AS*, 61, 75
- González, J. F., & Lapasset, E. 2000, *AJ*, 119, 2296
- Grenon, M. 2002, *Highlights of Astronomy*, 12, 680
- Gustafsson, B., & Nissen, P. E. 1972, *A&A*, 19, 261
- Hauck, B., & Mermilliod, M. 1997, *VizieR Online Data Catalog*, 2215, 0
- Hauck, B., & Mermilliod, M. 1998, *A&AS*, 129, 431
- Hejlesen, P. M. 1980, *A&AS*, 39, 347
- Hill, G., & Perry, C. L. 1969, *AJ*, 74, 1011
- Hiltner, W. A., Iriarte, B., & Johnson, H. L. 1958, *ApJ*, 127, 539
- Hogg, A. R. 1960, *PASP*, 72, 85
- Hogg, A. R., & Kron, G. E. 1955, *AJ*, 60, 365
- Ishmukhamedov, K. Z. 1966, *Tsirkyulyar Tashkentskoj Astronomicheskoy Observatorii*, 346, 16
- Johansson, K. L. V. 1981, *A&AS*, 43, 421
- Johnson, H. L. 1953, *ApJ*, 117, 353
- Johnson, H. L. 1954, *ApJ*, 119, 181
- Johnson, H. L. 1957, *ApJ*, 126, 121
- Johnson, H. L. 1966, *ARA&A*, 4, 193
- Johnson, H. L., & Morgan, W. W. 1953, *ApJ*, 117, 313
- Johnson, H. L., Hoag, A. A., Iriarte, B., Mitchell, R. I., & Hallam, K. L. 1961, *Lowell Observatory Bulletin*, 5, 133
- Jordi, C., Masana, E., Figueras, F., & Torra, J. 1997, *A&AS*, 123, 83
- Jordi, C., Luri, X., Masana, E., et al. 2002, *Highlights of Astronomy*, 12, 684
- Kharchenko, N. V., Piskunov, A. E., Röser, S., Schilbach, E., & Scholz, R.-D. 2005, *A&A*, 438, 1163
- Kilkenny, D., Hilditch, C. E. J., Hilditch, R. W., Hill, P. W., & Lynas-Gray, A. E. 1975, *MNRAS*, 172, 5P
- King, D. S. 1978, *J. Proc. R. Soc. N.S.W.*, 111, 1–12 = Sydney Obs. Pap. No. 79, 111, 1
- Koelbloed, D. 1959, *Bull. Astron. Inst. Netherlands*, 14, 265
- Kovalevsky, J., Falin, J. L., Pieplu, J. L., et al. 1992, *A&A*, 258, 7
- Levato, H., & Malaroda, S. 1975a, *PASP*, 87, 823
- Levato, H., & Malaroda, S. 1975b, *PASP*, 87, 173
- Levato, H., & Abt, H. A. 1977, *PASP*, 89, 274
- Lindegren, L., Hoeg, E., van Leeuwen, F., et al. 1992, *A&A*, 258, 18
- Lindegren, L., Madsen, S., & Dravins, D. 2000, *A&A*, 356, 1119
- Liu, T., Janes, K. A., & Bania, T. M. 1989, *AJ*, 98, 626
- Loden, K., Lindblad, P. O., Schober, J., & Urban, A. 1980, *A&AS*, 41, 85
- Loden, L. O. 1984, *A&AS*, 58, 595
- Loktin, A. V., & Matkin, N. V. 1990, *SvA*, 34, 571
- Loktin, A. V., & Matkin, N. V. 1994, *A&A*, *Trans.*, 4, 153
- Lutz, T. E., & Kelker, D. E. 1973, *PASP*, 85, 573
- Lyngå, G. 1960, *Ark. Astron.*, 2, 379
- Lyngå, G. 1962, *Ark. Astron.*, 3, 65
- Madsen, S. 1999, in *Harmonizing Cosmic Distance Scales in a Post-HIPPARCOS Era*, ed. D. Egret & A. Heck, *Astron. Soc. Pac. Conf. Ser.*, 167, 78
- Madsen, S., Lindegren, L., & Dravins, D. 2000, in *Stellar Clusters and Associations: Convection, Rotation, and Dynamos*, ed. R. Pallavicini, G. Micela, & S. Sciortino, *ASP Conf. Ser.* 198, 137
- Madsen, S., Lindegren, L., & Dravins, D. 2001, in *Dynamics of Star Clusters and the Milky Way*, ed. S. Deiters, B. Fuchs, A. Just, R. Spurzem, & R. Wielen, *ASP Conf. Ser.*, 228, 506
- Madsen, S., Dravins, D., & Lindegren, L. 2002, *A&A*, 381, 446
- Maitzen, H. M., & Catalano, F. A. 1986, *A&AS*, 66, 37
- Makarov, V. 2002, *AJ*, 124, 3299
- Makarov, V. V. 2003, *AJ*, 126, 2408
- Makarov, V. V. 2006, *AJ*, 131, 2967
- Mamajek, E. E. 2006, *AJ*, 132, 2198
- Manteiga, M., Martinez-Roger, C., Morales, C., & Sabau, L. 1991, *A&AS*, 87, 419
- Mävers, F.-W. 1940, *Astron. Nachr.*, 270, 201
- McCarthy, M. F., & O'Sullivan, S. 1969, *Ric. Astron.*, 7, 483
- McNamara, B. J., & Sanders, W. L. 1977, *A&AS*, 30, 45
- Meadows, A. J. 1961, *ApJ*, 133, 907
- Mendoza V., E. E., & Gomez, T. 1980, *MNRAS*, 190, 623
- Mermilliod, J.-C. 1995, in *Information, On-Line Data in Astronomy*, ed. D. Egret, & M. A. Albrecht, 127
- Mermilliod, J.-C., & Maeder, A. 1986, *A&A*, 158, 45
- Mermilliod, J.-C., Turon, C., Robichon, N., Arenou, F., & Lebreton, Y. 1997, in *Hipparcos – Venice '97*, ed. R. M. Bonnet, E. Høg, P. L. Bernacca, L. Emiliani, A. Blaauw, C. Turon, J. Kovalevsky, L. Lindegren, H. Hassan, M. Bouffard, B. Strim, D. Heger, M. A. C. Perryman, & L. Woltjer, *ESA SP*, 402, 643
- Mermilliod, J.-C., Platais, I., James, D. J., Grenon, M., & Cargile, P. A. 2008, *A&A*, 485, 95
- Mitchell, R. I., & Johnson, H. L. 1957, *ApJ*, 125, 414
- Mohan, V., & Sagar, R. 1985, *MNRAS*, 213, 337
- Munari, U., Dallaporta, S., Siviero, A., et al. 2004, *A&A*, 418, L31
- Narayanan, V. K., & Gould, A. 1999, *ApJ*, 523, 328
- Naylor, T., Totten, E. J., Jeffries, R. D., et al. 2002, *MNRAS*, 335, 291
- Nicolet, B. 1981, *A&A*, 104, 185
- Nissen, P. E. 1970a, *A&A*, 8, 476
- Nissen, P. E. 1970b, *A&A*, 6, 138
- Nissen, P. E. 1988, *A&A*, 199, 146
- O'Mullane, W., Lammers, U., Bailer-Jones, C., et al. 2007, in *Astronomical Data Analysis Software and Systems XVI*, ed. R. A. Shaw, F. Hill, & D. J. Bell, *Astron. Soc. Pac. Conf. Ser.*, 376, 99
- Pan, X., Shao, M., & Kulkarni, S. R. 2004, *Nature*, 427, 326
- Patenaude, M. 1978, *A&A*, 66, 225
- Percival, S. M., Salaris, M., & Kilkenny, D. 2003, *A&A*, 400, 541
- Percival, S. M., Salaris, M., & Groenewegen, M. A. T. 2005, *A&A*, 429, 887
- Perez, M. R., The, P. S., & Westerlund, B. E. 1987, *PASP*, 99, 1050
- Perry, C. L., & Hill, G. 1969, *AJ*, 74, 899
- Perry, C. L., Walter, D. K., & Crawford, D. L. 1978, *PASP*, 90, 81
- Perryman, M. A. C., Lindegren, L., Kovalevsky, J., et al. 1997, *A&A*, 323, L49
- Perryman, M. A. C., Brown, A. G. A., Lebreton, Y., et al. 1998, *A&A*, 331, 81
- Petrie, R. M., & Heard, J. F. 1969, *Publications of the Dominion Astrophysical Observatory Victoria*, 13, 329
- Pinsonneault, M. H., Stauffer, J., Soderblom, D. R., King, J. R., & Hanson, R. B. 1998, *ApJ*, 504, 170
- Pinsonneault, M. H., Terndrup, D. M., Hanson, R. B., & Stauffer, J. R. 2004, *ApJ*, 600, 946
- Platais, I. 1994, *Bulletin d'Information du Centre de Données Stellaires*, 44, 9
- Platais, I., Kozhurina-Platais, V., Barnes, S., et al. 2001, *AJ*, 122, 1486
- Platais, I. K. 1984, *SvA Lett.*, 10, 84
- Prisinzano, L., Micela, G., Sciortino, S., & Favata, F. 2003, *A&A*, 404, 927
- Prosser, C. F. 1993, *AJ*, 105, 1441
- Prosser, C. F., & Giampapa, M. S. 1994, *AJ*, 108, 964
- Robichon, N., Arenou, F., Mermilliod, J. C., & Turon, C. 1999, *A&A*, 345, 471
- Rojo Arellano, E., Pena, J. H., & Gonzalez, D. 1997, *A&AS*, 123, 25
- Röser, S., & Bastian, U. 1994, *A&A*, 285, 875
- Rufener, F. 1988, *Catalogue of stars measured in the Geneva Observatory photometric system : 4 : 1988 (Sauverny: Observatoire de Geneve)*
- Sanders, W. L. 1973, *A&AS*, 9, 213
- Sanders, W. L., & van Altena, W. F. 1972, *A&A*, 17, 193
- Schaifers, K., Voigt, H. H., Landolt, H., Boernstein, R., & Hellwege, K. H. 1982, *A&A*, C: *Interstellar Matter, Galaxy, Universe (Landolt-Boernstein: Numerical Data and functional Relationships in Science and Technology. New Series, Berlin: Springer)*
- Schmidt, E. G. 1976, *PASP*, 88, 63
- Smyth, M. J., & Nandy, K. 1962, *Publications of the Royal Observatory of Edinburgh*, 3, 24
- Snowden, M. S. 1975, *PASP*, 87, 721
- Snowden, M. S. 1976, *PASP*, 88, 174
- Soderblom, D. R., King, J. R., Hanson, R. B., et al. 1998, *ApJ*, 504, 192
- Soderblom, D. R., Nelan, E., Benedict, G. F., et al. 2005, *AJ*, 129, 1616
- Southworth, J., Maxted, P. F. L., & Smalley, B. 2005, in *Transits of Venus: New Views of the Solar System and Galaxy*, ed. D. W. Kurtz, *IAU Colloq.*, 196, 361
- Stello, D., & Nissen, P. E. 2001, *A&A*, 374, 105
- Stock, J. 1984, *Rev. Mex. Astron. Astrofis.*, 9, 127
- Strömgren, B. 1966, *ARA&A*, 4, 433
- Sung, H., Bessel, M. S., Lee, B.-W., & Lee, S.-G. 2002, *AJ*, 123, 290
- Taylor, R. J. 1954, *ApJ*, 120, 332
- Taylor, R. J. 1956, *MNRAS*, 116, 25
- The, P. S., Bakker, R., & Antalova, A. 1980, *A&AS*, 41, 93
- Trimble, V., Aschwanden, M. J., & Hansen, C. J. 2006, *PASP*, 118, 947
- Trumpler, R. 1928, *PASP*, 40, 265
- Turon, C., Gomez, A., Crifo, F., et al. 1992, *A&A*, 258, 74

- Uggren, A. R., Weis, E. W., & Deluca, E. E. 1979, *AJ*, 84, 1586
- Valls-Gabaud, D. 2007, in *IAU Symposium*, ed. W. I. Hartkopf, E. F. Guinan, & P. Harmanec, *IAU Symposium*, Vol. 240, 281
- van Altena, W. F. 1966, *AJ*, 71, 482
- van der Marel, H. 1988, Ph.D. Thesis, Technische Universiteit Delft
- van der Marel, H., & Petersen, C. S. 1992, *A&A*, 258, 60
- van Leeuwen, F. 1980, in *Star Formation*, *IAU Symp.* 85, 157
- van Leeuwen, F. 1983, Ph.D. Thesis
- van Leeuwen, F. 1999a, *A&A*, 341, L71
- van Leeuwen, F. 1999b, in *Harmonizing cosmic distance scales in a post-Hipparcos era*, ed. D. Egret, & A. Heck, 167 (*PASPC*), 52
- van Leeuwen, F. 2005a, *A&A*, 439, 805
- van Leeuwen, F. 2005b, in *Transits of Venus: New Views of the Solar System and Galaxy*, *IAU Colloq.* 196, 347
- van Leeuwen, F. 2007a, *Hipparcos, the new reduction of the raw data* (Dordrecht: Springer)
- van Leeuwen, F. 2007b, *A&A*, 474, 653
- van Leeuwen, F. 2008, in *IAU Symp.*, 248, 82
- van Leeuwen, F., & Alphenaar, P. 1982, *The Messenger*, 28, 15
- van Leeuwen, F., & Evans, D. W. 1998, *A&A*, 323, 157
- van Leeuwen, F., Alphenaar, P., & Brand, J. 1986, *A&AS*, 65, 309
- van Leeuwen, F., Alphenaar, P., & Meys, J. J. M. 1987, *A&AS*, 67, 483
- van Leeuwen, F., & Fantino, E. 2003, *Space Sci. Rev.*, 108, 537
- van Schewick, H. 1966, *Veroeffentlichungen des Astronomisches Institute der Universitaet Bonn*, 74, 1
- Vandenberg, D. A. 1985, *ApJS*, 58, 711
- Vasilevskis, S. 1955, *AJ*, 60, 384
- Vasilevskis, S., Klemola, A., & Preston, G. 1958, *AJ*, 63, 387
- Vasilevskis, S., Sanders, W. L., & Balz, A. G. A. 1965, *AJ*, 70, 797
- Verschoor, J. N., & van Genderen, A. M. 1983, *A&AS*, 53, 419
- Voigt, H. H. 1965, *Landolt-Börnstein: Numerical Data and Functional Relationships in Science and Technology – New Series Gruppe/Group 6* (Landolt-Bornstein)
- Walker, M. F. 1956, *ApJS*, 2, 365
- Wallerstein, G., Westbrooke, W., & Hannibal, D. 1963, *PASP*, 75, 522
- Weaver, H. F. 1953, *ApJ*, 117, 366
- Westerlund, B. E., Lundgren, K., Pettersson, B., Garnier, R., & Breysacher, J. 1988, *A&AS*, 76, 101
- Whiteoak, J. B. 1961, *MNRAS*, 123, 245
- Williams, P. M. 1978, *MNRAS*, 183, 49
- Wizinowich, P., & Garrison, R. F. 1982, *AJ*, 87, 1390
- Young, A. 1978, *PASP*, 90, 144
- Zentelis, N. 1983, *A&AS*, 53, 445
- Zwahlen, N., North, P., Debernardi, Y., et al. 2004, *A&A*, 425, L45