

Erratum

A new method for the spectroscopic identification of stellar non-radial pulsation modes

I. The method and numerical tests

W. Zima

Instituut voor Sterrenkunde, Celestijnenlaan 200D, 3001 Leuven, Belgium
e-mail: zima@ster.kuleuven.be

A&A 455, 227–234 (2006), DOI: 10.1051/0004-6361:20064876

Key words. line: profiles – techniques: spectroscopic – stars: variables: general – errata, addenda

The formulae for χ_v^2 in Eq. (1) and for the errors in Eqs. (2) and (3) were defined and proposed by Daszynska-Daszkiewicz (private communication). A similar definition, but for complex amplitudes of photometric and radial velocity variations, has already been applied by Daszynska-Daszkiewicz et al. (2003, 2005) in their method of simultaneous determination of the degree ℓ and the non-adiabatic parameter f for non-radially pulsating stars.

I apologize for not mentioning this in the original paper.

References

- Zima, W. 2006, A&A, 455, 227
Daszyńska-Daszkiewicz, J., Dziembowski, W. A., & Pamyatnykh, A. A. 2003, A&A, 407, 999
Daszyńska-Daszkiewicz, J., Dziembowski, W. A., & Pamyatnykh, A. A. 2005, A&A, 441, 641