

## **Probing X-ray irradiation in the nucleus of NGC 1068 with observations of high-*J* lines of dense gas tracers (*Erratum*)**

J. P. Pérez-Beaupuits<sup>1</sup>, M. Spaans<sup>1</sup>, F. F. S. van der Tak<sup>2,1</sup>, S. Aalto<sup>3</sup>, S. García-Burillo<sup>4</sup>, A. Fuente<sup>4</sup>, and A. Usero<sup>4</sup>

<sup>1</sup> Kapteyn Astronomical Institute, University of Groningen, Landleven 12, 9747 AD Groningen, The Netherlands  
e-mail: [jp@astro.rug.nl](mailto:jp@astro.rug.nl)

<sup>2</sup> SRON Netherlands Institute for Space research, Landleven 12, 9747 AD Groningen, The Netherlands


<sup>3</sup> Onsala Rymdobservatorium, Chalmers Tekniska Högskola, 439 92 Onsala, Sweden

<sup>4</sup> Observatorio Astronómico Nacional, C/ Alfonso XII 3, 28014 Madrid, Spain

A&A 503, 459–466 (2009), DOI: [10.1051/0004-6361/200912350](https://doi.org/10.1051/0004-6361/200912350)

**Key words.** galaxies: ISM – galaxies: individual: NGC 1068 – galaxies: Seyfert – ISM: molecules: – radio lines: ISM –  
radio lines: galaxies – errata, addenda

An error occurred during the final processing of the publication of Fig. 1. The first spectrum of Fig. 1 has been published four times, replacing the three other spectra. The correct Fig. 1 is published on the next page.



**Fig. 1.** Molecular line emissions in NGC 1068. The velocity resolution was reduced to  $20 \text{ km s}^{-1}$  for all the lines. The spectra are centered with respect to the heliocentric systemic velocity  $v_{\text{sys}} = 1137 \text{ km s}^{-1}$ .