

LETTER TO THE EDITOR

Dust-temperature of an isolated star-forming cloud: *Herschel* observations of the Bok globule CB244[★]

A. Stutz^{1,2}, R. Launhardt¹, H. Linz¹, O. Krause¹, T. Henning¹, J. Kainulainen¹, M. Nielbock¹,
J. Steinacker^{3,1}, and P. André⁴

¹ Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany
e-mail: stutz@mpia.de

² Department of Astronomy and Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

³ LERMA & UMR 8112 du CNRS, Observatoire de Paris, 61 Av. de l'Observatoire, 75014 Paris, France

⁴ Laboratoire AIM, CEA/DSM-CNRS-Université Paris Diderot, IRFU/Service d'Astrophysique, C.E. Saclay, Orme des Merisiers, 91191 Gif-sur-Yvette, France

Received 29 March 2010 / Accepted 15 April 2010

ABSTRACT

We present *Herschel* observations of the isolated, low-mass star-forming Bok globule CB244. It contains two cold sources, a low-mass Class 0 protostar and a starless core, which is likely to be prestellar in nature, separated by 90'' (~18 000 AU). The *Herschel* data sample the peak of the *Planck* spectrum for these sources, and are therefore ideal for dust-temperature and column density modeling. With these data and a near-IR extinction map, the MIPS 70 μm mosaic, the SCUBA 850 μm map, and the IRAM 1.3 mm map, we model the dust-temperature and column density of CB 244 and present the first measured dust-temperature map of an entire star-forming molecular cloud. We find that the column-averaged dust-temperature near the protostar is ~17.7 K, while for the starless core it is ~10.6 K, and that the effect of external heating causes the cloud dust-temperature to rise to ~17 K where the hydrogen column density drops below 10^{21} cm^{-2} . The total hydrogen mass of CB 244 (assuming a distance of 200 pc) is $15 \pm 5 M_{\odot}$. The mass of the protostellar core is $1.6 \pm 0.1 M_{\odot}$ and the mass of the starless core is $5 \pm 2 M_{\odot}$, indicating that ~45% of the mass in the globule is participating in the star-formation process.

Key words. ISM: individual objects: CB244 – infrared: ISM – dust, extinction – ISM: clouds – submillimeter: ISM – stars: formation

1. Introduction

Bok globules are nearby, small, relatively isolated molecular clouds undergoing low-mass star-formation (e.g., Clemens & Barvainis 1988; Launhardt & Henning 1997). They tend to have only one or two star-forming cores that are embedded in a larger common cloud. These relatively simple characteristics make these objects ideal for studying the detailed processes taking place in low-mass star formation. Specifically, the temperature and density structure are fundamental physical parameters necessary to understand core fragmentation, collapse, and chemical evolution (e.g., Ward-Thompson et al. 2007; Stutz et al. 2008, 2009; Launhardt et al. 2010). The *Herschel* (Pilbratt et al. 2010) data cover the wavelength range which samples the peak of the Planck spectrum for cold sources (6–20 K). This wavelength regime is critical for accurate modeling of the temperature and density structure in the cold environments where stars are born (e.g., Shetty et al. 2009).

The *Herschel* guaranteed time key program “Early Phases of Star-formation” (EPoS; P.I. O. Krause) sample consists of the Photodetector Array Camera and Spectrometer (PACS;

Poglitsch et al. 2010) and the Spectral and Photometric Imaging REceiver (SPIRE; Griffin et al. 2010) imaging-mode observations of sites of both high- and low-mass star formation. The low-mass science demonstration phase portion of the sample is the focus of this contribution. The source CB 244 (L1262; Lynds 1962) is a Bok globule at a distance of ~200 pc (Hilton & Lahulla 1995), with an approximate extent of ~6', or about 0.5 pc. The CB 244 globule contains two submm peaks, one associated with a Class 0 protostar located at RA = 23^h25^m46.3^s, Dec = +74°17'39.1'', and one associated with a starless core located at RA = 23^h25^m27.1^s, Dec = +74°18'25.3''. The protostar drives a molecular outflow (e.g., Clemens et al. 1991), while the detection of the CB 244 starless core was first published as an additional source inside the globule by Launhardt (1996) and Shirley et al. (2000) and produces both an 8 μm (Tobin et al. 2010) and a 24 μm shadow. The YSO and starless core are separated by ~90'', and are therefore well resolved throughout the PACS and SPIRE bands. We use *Herschel* imaging to construct spatially resolved spectral energy distributions (SEDs) of the entire cloud that cover both sides of the peak of the SED. These data allow us to reconstruct the dust-temperature map and column density distribution of the Bok globule and hence the density profiles and mass distribution with unprecedented accuracy. They also reveal the role of

[★] *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.




Fig. 1. $8' \times 8'$ images of CB 244, shown at the wavelengths indicated in the *bottom right* of each panel. The images are shown on a log scale at stretches intended to highlight the structure of absorption or emission at each wavelength. The locations of the protostar and starless core, derived from the $8 \mu\text{m}$ image, are indicated with \times -symbols (note the offset between the \times 's and the starless core emission at $100\text{--}250 \mu\text{m}$ wavelengths). The panel contours are labeled with their corresponding wavelengths. The contour levels are: A_V extinction map levels = $\{5, 10, 15, 20, 25\}$ mags (inner dotted contours indicates the region inside which we place lower-limits on the extinction); SPIRE $350 \mu\text{m}$ = $\{0.5, 1.0, 1.5, 3.5\}$ mJy/ \square'' ; SCUBA $850 \mu\text{m}$ = $\{0.3, 0.75\}$ mJy/ \square'' ; IRAM 1.3 mm = $\{0.1, 0.3\}$ mJy/ \square'' . Approximate beam sizes are indicated as yellow circles.

external heating and shielding by the envelopes in the energy balance of such cores in isolated globules.

2. Observations and data processing

2.1. *Herschel* observations

The source CB 244 was observed with the PACS instrument on board the *Herschel* Space Observatory on 2009, December 30, during the science demonstration program. The globule CB 244 was observed at $100 \mu\text{m}$ and $160 \mu\text{m}$. We obtained two orthogonal scan maps with scan leg lengths of $9'$ using a scan speed of $20''/\text{s}$. The scan leg position angles guarantee an almost homogeneous coverage of CB 244. We produced highpass-filtered intensity maps from these data using the HIPE software package (Ott 2010), version 3.0, build 455. Besides the standard steps leading to level-1 calibrated data, a second-level deglitching as well as a correction for offsets in the detector sub-matrices were performed. Finally, the data were highpass-filtered, using a median window with a width of 271 data samples to remove the effects of bolometer temperature drifts during the course of the data acquisition. Furthermore, we masked emission structures, in this case both the YSO and starless core regions, before computing the running median. Masking bright sources minimizes over-subtraction of source emission in the highpass filtering step.

Finally, the data were projected onto a coordinate grid using the photProject routine inside HIPE. As a last step, the flux correction factors provided by the PACS ICC team were applied. We note that these data have relatively flat background values, indicating that the highpass reduction used in conjunction with the masking is a relatively robust scheme for avoiding artifacts and recovering extended emission in the PACS maps. In addition to these processing steps, we checked the pointing in the PACS $100 \mu\text{m}$ map against the MIPS $24 \mu\text{m}$ mosaic of the same region. Using point-sources detected in both images, we found a pointing offset of $\sim 1.25''$ in the $100 \mu\text{m}$ map relative to the $24 \mu\text{m}$ mosaic. Because the $160 \mu\text{m}$ PACS map contains no point-sources and was acquired at the same time as the $100 \mu\text{m}$ data, we blindly applied the same pointing correction. The final PACS 100 and $160 \mu\text{m}$ images are shown in Fig. 1.

Maps at 250 , 350 , and $500 \mu\text{m}$ were obtained with SPIRE on October 20, 2009. Two $9'$ scan legs were used to cover the source. Two repetitions resulted in 146 s of scanning time with the nominal speed of $30''/\text{s}$. The data were processed within HIPE with the standard photometer script up to level 1. During baseline removal, we masked out the high-emission area in the center of the field. For these data no cross-scan was obtained; therefore, the resulting maps still showed residual stripes along the scan direction. We used the Bendo et al. (2010) de-stripping scheme to mitigate this effect. In addition, we checked the




Fig. 2. Dust-temperature (color) and hydrogen column density (white contours) in CB 244. The line-of-sight-averaged mean dust-temperature was derived pixel by pixel from modified black-body SED fits to homogeneously beam-smoothed emission maps from *Herschel* PACS and SPIRE, *Spitzer* MIPS70, and ground-based extinction mapping and submillimeter dust-continuum maps at 0.8 and 1.3 mm (see text for details). Column density contours are at $\{0.1$ (thick) $0.3, 0.5, 1$ (thick), $2, 3.5, 5,$ and $7\} \times 10^{22}$ H/cm². The peak column density in the prestellar core just reaches 10^{23} H/cm². Two SEDs and the respective modified black-body fits are shown for the positions of the Class 0 protostar (“1”) and the prestellar core (“2”). The yellow dashed ellipses indicate the areas used to calculate source masses (see Sect. 4). The 37'' beam size is indicated by a grey circle. White areas represent regions where there is not enough signal-to-noise and/or coverage to get a reliable fit.

pointing in the SPIRE data by cross-correlating the images with the PACS 160 μm image; we found that the pointing offsets for all three wavelengths are on order of 2'' or smaller, and therefore we did not correct for this. The SPIRE 250, 350, and 500 μm images are shown in Fig. 1.

2.2. Other data

Near-IR data: We observed CB 244 in the near-IR K_S and H bands using Omega2000 at the Calar Alto Observatory. The $15' \times 15'$ field-of-view provided complete coverage of the globule. A standard near-IR data reduction scheme was adopted. The extracted photometric data were calibrated using the Two Micron All Sky Survey (2MASS; [Skrutskie et al. 2006](#)). Following the NICE method ([Lada et al. 1994](#)), the observed $(H - K_S)$ colors of stars were then related to visual extinction by $(H - K_S) = \langle (H - K_S)_0 \rangle + 0.063 \times A_V$, assuming the [Rieke & Lebofsky \(1985\)](#) extinction law, where $\langle (H - K_S)_0 \rangle$ is the mean intrinsic color of the stars determined from edges of the field. The A_V values for each star were used to create an extinction map. The resulting extinction map probes extinction values up to $A_V \approx 30$ mag, with a 3σ error of about 3 mag at $A_V = 0$ mag.

Spitzer data: The *Spitzer* observations presented here are from two programs: the MIPS observations are from program 53 (P.I. Rieke), while the IRAC observations are from program 58 (P.I. Lawrence). The MIPS observations were reduced using the data analysis tool (DAT; [Gordon et al. 2005](#)) according to steps outlined in [Stutz et al. \(2007\)](#). The IRAC frames were processed using the IRAC Pipeline v14.0, and mosaics were created from the basic calibrated data (BCD) frames using a custom IDL program (see [Gutermuth et al. 2008](#)).

(Sub)mm continuum data: The SCUBA 850 μm and the IRAM 1.3 mm maps are presented in [Launhardt et al. \(2010\)](#). The 850 μm and 1.3 mm maps cover both sources, but do not extend very far into the envelope. The maps have native beam sizes of 14.9'' (850 μm), and 10.9'' (1.3 mm). For further details see [Launhardt et al. \(2010\)](#). In Fig. 1 we show the 850 μm and 1.3 mm emission as contours.

3. Dust temperature and column density modeling

We used the PACS 100 and 160 μm , SPIRE 250, 350, and 500 μm data in conjunction with the NIR extinction map, MIPS 70 μm , SCUBA 850 μm , and IRAM 1.3 mm dust emission maps to model the line-of-sight-averaged temperature and column density of CB 244. The calibrated maps were all homogeneously convolved to the largest FWHM beam size of the data set, the SPIRE 500 μm 37'' beam, assuming Gaussian beam profiles, and projected onto a common $8'' \text{ pix}^{-1}$ grid. This approach has the drawback that all spatial structures smaller than this beam size are smoothed, but ensures that the fluxes at all wavelengths in a given image pixel are derived from the same area in the source. We corrected the filtering of large spatial-scale emission in the PACS data using the [Schlegel et al. \(1998\)](#) 100 μm IRAS maps and the ISO Serendipity Survey observations at 170 μm : we added a zero level of emission of 0.3 and 0.8 mJy/arcsec² to the 100 and 160 μm maps, respectively.

For each image pixel, an SED was extracted and fitted with a single-temperature modified black-body of the form $S_\nu = \Omega B_\nu(\nu, T_d) (1 - e^{-\tau(\nu)})$, where Ω is the solid angle of the emitting element, B_ν is the Planck function, T_d is the dust-temperature, and $\tau(\nu)$ is the optical depth at frequency ν . In a first iteration, we used a simple dust opacity model of the form $\kappa_\nu \propto \nu^\beta$ to fit for β , using χ^2 minimization. We found that $\beta = 1.9\text{--}2$ fitted the data best over the entire cloud. Note that compact structures were smoothed out; therefore this cannot be taken as evidence against grain growth in the centers of the two dense cores. In a second step, we then fixed the dust model to the [Ossenkopf & Henning \(1994\)](#) model for an MRN grain size distribution, with thin ice mantles and no coagulation, which has a $\kappa_{1.3\text{mm}} = 0.51 \text{ cm}^2 \text{ g}^{-1}$. The hydrogen-to-dust mass ratio was fixed to the canonical value of 110 (e.g., [Sodroski et al. 1997](#)). We then searched for the optimum T_d and hydrogen column density N_H values (the two free parameters) by calculating flux densities and A_V values and then comparing to the emission and extinction observations, using χ^2 minimization. The

resulting line-of-sight-averaged dust-temperature and column density maps are shown in Fig. 2 together with the SED fits at the central positions of the two submillimeter peaks. We found the best-fit column-averaged dust-temperatures for the protostar and prestellar core of ~ 17.7 K and ~ 10.6 K, respectively. Despite the fact that each individual image pixel was fitted independently and the maps at different wavelengths have different outer boundaries, both the temperature and the column density maps have very little noise and are very smooth, demonstrating the robustness of our fitting approach. Furthermore, we evaluated the effect of a possible filtered-out extended emission component in the chopped $850\ \mu\text{m}$ and $1.3\ \text{mm}$ maps. Adding in offsets of $60\ \text{mJy}/15''$ beam at $850\ \mu\text{m}$ and $12\ \text{mJy}/11''$ beam at $1.3\ \text{mm}$, corresponding to 20% of the peak surface brightness of the cold prestellar core, did not affect the fitted temperature and column density (less than a 2% change) for the prestellar core or the protostar. Hence, we conclude that possible missing extended emission in the (sub)mm maps does not affect our results.

4. Summary and conclusions

As shown in Fig. 2, the line-of-sight-averaged dust-temperature decreases constantly without any significant jump from $\sim 17 \pm 1$ K at the outer boundary of the cloud, where N_{H} drops below $10^{21}\ \text{cm}^{-2}$ and $A_{\text{V}} \approx 0.3$ mag, to ~ 13 K in the inner cloud at $N_{\text{H}} \approx 1\text{--}2 \times 10^{22}\ \text{cm}^{-2}$ and $A_{\text{V}} \approx 3\text{--}5$ mag. Note that due to the two major drawbacks of our simplified approach, beam-smoothing and line-of-sight-averaging, the local dust-temperatures at the source centers will be higher in the protostar and lower in the prestellar core. Nevertheless, the boundaries of the two sources can be derived from the dust-temperature map rather than from the smooth column density map. We find that $T_{\text{d}} = 13$ K marks the mean inner dust-temperature of the globule, from where the temperature towards the protostar rises steeply and that of the prestellar core decreases. We therefore adopt ellipses that follow the 13 K contour in the dust-temperature map as boundaries for the two sources (see Fig. 2). The mean radii of these ellipses are $50''$ (10000 AU) for the protostellar core and $76''$ (15000 AU) for the prestellar core. Integration of the column density map within these boundaries yields $M_{\text{H}} = 1.6 \pm 0.3\ M_{\odot}$ for the protostellar core and $M_{\text{H}} = 5 \pm 2\ M_{\odot}$ for the prestellar core, where the uncertainties are derived from an assumed ± 1 K uncertainty in the dust-temperature. The total mass of the globule (within the $N_{\text{H}} = 1 \times 10^{21}\ \text{cm}^{-2}$ contour) is $M_{\text{H}} = 15 \pm 5\ M_{\odot}$. We note that $M_{\text{gas}} = 1.36\ M_{\text{H}}$. These masses imply that $\sim 45\%$ of the mass of the globule is participating in the star-formation process.

Including the new *Herschel* data, we derive the following parameters for the protostar: $L_{\text{bol}} \sim 1.5\ L_{\odot}$, $L_{\text{submm}}/L_{\text{bol}} \sim 4\%$, $T_{\text{bol}} \sim 62$ K, and $M_{\text{env}}/L_{\text{bol}} \sim 1.45\ M_{\odot}/L_{\odot}$, (cf. Launhardt et al. 2010), confirming the Class 0 classification according to Chen et al. (1995) and Andre et al. (2000). Furthermore, the C^{18}O (2–1) FWHM line width for the prestellar core is $\sim 0.9\ \text{km s}^{-1}$ (Stutz et al., in prep.); using our fitted prestellar core mass and temperature, we find that the ratio of gravitational energy to thermal and turbulent energy is $E_{\text{grav}}/(E_{\text{therm}} + E_{\text{turb}}) \approx 0.9$, a marginally sub-critical value which is expected for prestellar cores. For comparison, Tobin et al. (2010) derive a mass of $\sim 3\text{--}4\ M_{\odot}$ for the prestellar core (integrated over a similar area and scaled to a distance of 200 pc) using the $8\ \mu\text{m}$ shadow (see Fig. 1), a good agreement given the uncertainties in both mass derivation methods. The

prominent and extended $3.6\ \mu\text{m}$ coreshine (see Fig. 1), originating from dust-grain scattering of the background radiation field, is an indication of grain growth (Steinacker et al. 2010) in the CB 244 cloud. These pieces of evidence together indicate that the CB 244 globule, and other globules like it, are excellent sources in which to study the earliest phases of low-mass star-formation. As a next step and in a follow-up paper, we will employ 3D-modeling to overcome the effects of line-of-sight averaging and beam-smoothing, and to reconstruct the full dust-temperature and density structure of CB 244 and the other sources in our sample. Measuring reliable temperatures and column densities with the *Herschel* data in a sample of prestellar and protostellar cores is a fundamental step towards revealing the initial conditions of low-mass star-formation.

Acknowledgements. The authors thank J. Tobin for helpful discussions and D. Johnstone for a critical and helpful referee report. PACS has been developed by a consortium of institutes led by MPE (Germany) and including UVIE (Austria); KU Leuven, CSL, IMEC (Belgium); CEA, LAM (France); MPIA (Germany); INAF- IFSI/OAA/OAP/OAT, LENS, SISSA (Italy); IAC (Spain). This development has been supported by the funding agencies BMVIT (Austria), ESA-PRODEX (Belgium), CEA/CNES (France), DLR (Germany), ASI/INAF (Italy), and CICYT/MCYT (Spain). SPIRE has been developed by a consortium of institutes led by Cardiff University (UK) and including Univ. Lethbridge (Canada); NAOC (China); CEA, LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain); Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, Univ. Sussex (UK); and Caltech, JPL, NHSC, Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); Stockholm Observatory (Sweden); STFC (UK); and NASA (USA).

References

- Andre, P., Ward-Thompson, D., & Barsony, M. 2000, *Protostars and Planets IV*, 59
- Bendo, G. J., et al. 2010, *A&A*, 518, L65
- Chen, H., Myers, P. C., Ladd, E. F., & Wood, D. O. S. 1995, *ApJ*, 445, 377
- Clemens, D. P., & Barvainis, R. 1988, *ApJS*, 68, 257
- Clemens, D. P., Yun, J. L., & Heyer, M. H. 1991, *ApJS*, 75, 877
- Gordon, K. D., Rieke, G. H., Engelbracht, C. W., et al. 2005, *PASP*, 117, 503
- Griffin, M. J., et al. 2010, *A&A*, 518, L3
- Gutermuth, R. A., Myers, P. C., Megeath, S. T., et al. 2008, *ApJ*, 674, 336
- Hilton, J., & Lahulla, J. F. 1995, *A&AS*, 113, 325
- Lada, C. J., Lada, E. A., Clemens, D. P., & Bally, J. 1994, *ApJ*, 429, 694
- Launhardt, R. 1996, Ph.D. Thesis, University of Jena
- Launhardt, R., & Henning, T. 1997, *A&A*, 326, 329
- Launhardt, R., et al. 2010, *ApJS*, 188, 139
- Lynds, B. T. 1962, *ApJS*, 7, 1
- Ossenkopf, V., & Henning, T. 1994, *A&A*, 291, 943
- Ott, S. 2010, in *Astronomical Data Analysis Software and Systems XIX*, ASP Conf. Ser., 00
- Pilbratt, G. L., et al. 2010, *A&A*, 518, L1
- Poglitsch, A., et al. 2010, *A&A*, 518, L2
- Rieke, G. H., & Lebofsky, M. J. 1985, *ApJ*, 288, 618
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Shetty, R., Kauffmann, J., Schnee, S., Goodman, A. A., & Ercolano, B. 2009, *ApJ*, 696, 2234
- Shirley, Y. L., Evans, II, N. J., Rawlings, J. M. C., & Gregersen, E. M. 2000, *ApJS*, 131, 249
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Sodroski, T. J., Odegard, N., Arendt, R. G., et al. 1997, *ApJ*, 480, 173
- Steinacker, J., Pagani, L., Bacmann, A., & Guieu, S. 2010, *A&A*, 511, A9
- Stutz, A. M., Bieging, J. H., Rieke, G. H., et al. 2007, *ApJ*, 665, 466
- Stutz, A. M., Rubin, M., Werner, M. W., et al. 2008, *ApJ*, 687, 389
- Stutz, A. M., Rieke, G. H., Bieging, J. H., et al. 2009, *ApJ*, 707, 137
- Tobin, J. J., Hartmann, L., Looney, L. W., & Chiang, H. 2010, *ApJ*, 712, 1010
- Ward-Thompson, D., André, P., Crutcher, R., et al. 2007, *Protostars and Planets V*, 33