LETTER TO THE EDITOR

Discovery of an extremely bright submillimeter galaxy at $z = 3.93$

J.-F. Lestrade¹, F. Combes¹, P. Salomé¹, A. Omont², F. Bertoldi³, P. André⁴, and N. Schneider⁴

¹ Observatoire de Paris, LERMA, CNRS, 61 Av. de l'Observatoire, 75014, Paris, France
 e-mail: jean-francois.lestrade@obspm.fr

² Institut d'Astrophysique de Paris, UMR 7095, CNRS, UPMC Univ. Paris 06, 98bis Boulevard Arago, 75014, Paris, France

³ Argelander Institute for Astronomy, University of Bonn, Auf dem Hügel 71, 53121 Bonn, Germany

⁴ Laboratoire AIM Paris-Saclay, CEA/IRFU/SAP - CNRS, Université Paris Diderot, 91191 Gif-sur-Yvette Cedex, France

Received 1 September 2010 / Accepted 13 October 2010

ABSTRACT

Serendipitously we have discovered a rare, bright submillimeter galaxy (SMG) with a flux density of 30 ± 2 mJy at $\lambda = 1.2$ mm, using MAMBO2 at the IRAM 30-m millimeter telescope. Although no optical counterpart is known for MM18423+5938, we were able to measure the redshift $z = 3.92960 \pm 0.00013$ from the detection of CO lines using the IRAM Eight MIXer Receiver (EMIR). In addition, by collecting all available photometric data in the far-infrared and radio to constrain its spectral energy distribution, we derive the FIR luminosity $4.8 \times 10^{14}/m L_{\odot}$ and mass $6.0 \times 10^9/m M_{\odot}$ for its dust, allowing for a magnification factor m caused by a probable gravitational lens. The corresponding star-formation rate is $8.3 \times 10^4/m M_{\odot}/\text{yr}$. The detection of three lines of the CO rotational ladder, and a significant upper limit for a fourth CO line, allow us to estimate an H_2 mass of between $1.9 \times 10^{11}/m M_{\odot}$ and $1.1 \times 10^{12}/m M_{\odot}$. The two lines $\text{C I}(^3P_1 - ^3P_0)$ and $\text{C I}(^3P_2 - ^3P_1)$ were clearly detected and yield a $[\text{C I}]/[\text{H}_2]$ number abundance of between 1.4×10^{-5} and 8.0×10^{-5} . Upper limits are presented for emission lines of HCN, HCO^+ , HNC, H_2O , and of other molecules. The moderate excitation of the CO lines is indicative of an extended starburst, and excludes the dominance of an AGN in heating this high- z SMG.

Key words. galaxies: evolution – galaxies: high-redshift – galaxies: ISM – infrared: galaxies – submillimeter: galaxies

1. Introduction

Deep blank-field millimeter and submillimeter surveys of small fields ($\sim 1 \text{ deg}^2$) have revealed many dusty, starburst submillimeter galaxies (SMGs) over the past decade with flux densities of a few to about ten mJy at $\lambda = 1.2$ mm (e.g. Bertoldi et al. 2007; Greve et al. 2008), and higher at $\lambda = 850 \mu\text{m}$ owing to dust emissivity (e.g. Smail et al. 1997). Recently, the South Pole Telescope survey, less deep but much larger in sky coverage (87 deg^2), has found 47 brighter SMGs with flux densities between 11 and 65 mJy at $\lambda = 1.4$ mm (Vieira et al. 2010). Whereas redshifts of SMGs are crucial to study their physical properties, most of these dust-obscured galaxies have very faint or no optical counterparts, making measurements of spectroscopic redshift extremely difficult or impossible (e.g. Smail et al. 2002).

These dust-enshrouded star-forming galaxies are expected to be at high redshifts and are identified with the most massive galaxies assembled during an energetic early phase of galaxy formation. Their abundance appears to peak at $z \sim 2.5$ (Chapman et al. 2005; Wardlow et al. 2010). Their star formation rate is prodigious at up to $10^3 M_{\odot} \text{ yr}^{-1}$, and the underlying starburst activity is believed to result from mergers (Blain et al. 2002).

Lestrade et al. (2009) discovered serendipitously a rare, bright point source, MM18423+5938, at $\lambda = 1.2$ mm (30 mJy) by mapping 50 separate fields totalling a sky area of 0.5 deg^2 with the MAMBO2 bolometer camera (Kreysa et al. at the IRAM 30-m millimeter telescope. Subsequently, some of us (PA and NS) searched but did not find local CO in the direction of the source suggesting not a young stellar object but an SMG instead, despite the Vieira et al.'s cumulative source count

that yields a chance as low as $\sim 7\%$ of finding a 30 mJy SMG. MM18423+5938 is detected at $70 \mu\text{m}$ but undetected at $24 \mu\text{m}$ in *MIPS/Spitzer* images. It is in neither the 2MASS catalogue, nor the NVSS VLA catalogue ($S_{1.4 \text{ GHz}} < 2.5$ mJy), and no optical and X-ray identifications are found in catalogues searched with NED (MM18423+593 is however outside the SDSS footprint). All these photometric data are summarized in Table 1.

We show in this Letter that MM18423+5938 is a bright, high-redshift SMG. We present in Sect. 2 our IRAM/EMIR spectroscopic observations of MM18423+5938 at millimeter wavelengths that yielded our detections of CO and C I, and upper limits for other molecular species. In Sect. 3, we model the data to infer the dust and gas content of MM18423+5938 and its general properties. To compute distances and luminosities, we adopt the Λ -CDM concordance cosmological model, $H_0 = 71 \text{ km s}^{-1}/\text{Mpc}$, $\Omega_M = 0.27$, and $\Omega_{\Lambda} = 0.73$ (Hinshaw et al. 2009).

2. Observations and data analysis

Lestrade et al. (2009) detected MM18423+5938 with a high but uncertain integrated flux density of 30–60 mJy, given that it was located close to the border of their MAMBO map. We reobserved MM18423+5938 with MAMBO at the IRAM 30-m telescope in the on-off wobbler-switching mode on 2010 January 16 using the map coordinates ($\alpha_{2000} = 18^{\text{h}}42^{\text{m}}22.5^{\text{s}} \pm 0.2^{\text{s}}$ and $\delta_{2000} = 59^{\circ}38'30'' \pm 2''$) and measured 30 ± 2 mJy at $\lambda = 1.2$ mm.

MM18423+5938 is located close to the border of the archived *MIPS/Spitzer* maps centered on the star GJ725AB

Table 1. Photometry available for MM18423+5938.

Band	S_ν (mJy)	Ref.
1.4 GHz	<2.5	NVSS
3 mm	$2^{+2.0}_{-1.5}$	this work
2 mm	9 ± 3	this work
1.2 mm	30 ± 2	this work
100 μ m	<600	IRAS
60 μ m	<100	IRAS
70 μ m	31 ± 4	<i>Spitzer/MIPS</i>
24 μ m	<0.6	<i>Spitzer/MIPS</i>
<i>BVR</i>	<i>B</i> mag > 21	NED (DSS)

(AOR 4199424). We determined the flux densities of the star GJ725AB and MM18423+5938 at 70 μ m and 24 μ m by means of aperture photometry, scaling with the stellar photospheric flux densities of GJ725AB predicted by the NextGen stellar atmospheric model (Allard et al. 2001). We note a difference of $9''$, i.e. at the 3σ level, between the positions of the source in our MAMBO map (Lestrade et al. 2009) and the archived 70 μ m *MIPS Spitzer* map, which is not understood.

To measure the redshift of MM18423+5938, we used the strategy of observation developed by Weiss et al. (2009) with the multi-band heterodyne receiver Eight Mixer Receiver (EMIR¹) at the IRAM 30-m telescope. The 3 mm setup (E090) of EMIR provides 7.43 GHz of instantaneous, dual linear polarization bandwidth. The entire frequency range from 77.7 to 115.8 GHz in the 3 mm band can be searched with six tunings spaced to provide 0.5 GHz overlap. This range corresponds to $0 < z < 0.48$ and $1 < z < 10$ for the CO lines between ($J = 1-0$) and ($J = 8-7$). Observations were conducted from 2010 July 29th to August 2nd with precipitable water vapor comprised between 3 and 7 mm and with standard system temperatures of 110 K for the E090 setup. Data were processed with 16 units of the Wide band Line Multiple Autocorrelator (WILMA) providing a spectral resolution of 2 MHz for the E090 setup. The observations were conducted in wobbler-switching mode, with a switching frequency of 1 Hz and an azimuthal wobbler throw of $100''$. Pointing and focus offsets were determined once every two hours and found to be stable. Calibration was done every 6 min using the standard hot/cold load absorber. The data were reduced with the CLASS software.

We started to scan the whole 3 mm band by integrating data for ~ 2 h for each tuning. We discovered unambiguously a line at 93.52 GHz after 20 min of integration during our third tuning on the second night, and continued to integrate dual polarisation data for 1.5 h in total to reach an rms noise level of $T_{\text{mb}} = 0.8$ mK in 60 km s^{-1} channels (Fig. 1 lefthand top panel). At this stage, we successively assumed that this line could be CO(1–0), CO(2–1), ... to calculate for each of these assumptions the corresponding redshift and predict the frequencies of the higher J transitions accessible in the 2 mm band of EMIR (setup E150 from 127 to 176 GHz). We then tuned to these frequencies with the E150 setup and swiftly detected a line at 140.26 GHz that corresponds to CO(6–5) at $z = 3.92960 \pm 0.00013$, in addition to the line at 93.52 GHz for CO(4–3). This identification of the CO transitions and determination of the redshift of MM18423+5938 were carried out during the same night of July 30/31 in ~ 6 h. The rest of the allocated time (15 h) was used to search for CO(7–6) (detected), CO(9–8) (undetected, consequently CO(10–9) was not searched), and for other species, C I (two lines detected),

Fig. 1. The three CO rotational lines detected, and the two C I lines. The vertical scale is T_{mb} in mK.

and HCN & HNC(5–4), LiH(1–0) & HCO+(5–4), H_2O_o , H_2O_p , $^{13}\text{CO}(5-4)$, CS(8–7), and CS(9–8) (all undetected but interesting upper limits are discussed below, see Table 3). Two CO lines (5–4 and 8–7) unfortunately are in the atmospheric O_2 and H_2O lines opacity domains and could not be observed. The spectra were of high quality, stable, and flat. Their mean levels measure the continuum flux densities at 2 mm and 3 mm (Table 1) owing to the excellent weather conditions (precipitable water vapor ~ 4 mm).

Figure 1 displays the three CO lines detected, along with the two C I lines. Spectra were smoothed to a resolution of $30-50 \text{ km s}^{-1}$. Gaussian models were fitted to the lines and the results are reported in Table 2. Line widths found are normal *albeit* small suggesting that we are observing a galaxy seen rather face-on.

3. Results

3.1. Dust emission

The photometric data are collected in Table 1. To model these data in Fig. 2, we choose to use the well-established Milky Way dust model of Desert et al. (1990) as a template. This model consists of three main components, PAH and both very small and large grains, and the emissivity slope is assumed to be $\beta = 2$. The large dust grains are dominant in mass and their temperature is estimated to be $T_d = 45$ K, constrained by the Rayleigh-Jeans part of the emission. The very small grains are made of two temperature components, ~ 80 and ~ 130 K, constrained by the 70 μ m *Spitzer* measurement and the robust relationship between $L'_{\text{CO}(3-2)}$ and L_{FIR} found by Iono et al. (2009). A single temperature component for the small grains in the model was tried but the resulting L_{FIR} is significantly inconsistent with this relationship. The total mass of the dust required by this model is $M_{\text{dust}} = 6.0 \times 10^9 / m M_\odot$, and for a gas-to-dust mass ratio of 150, we infer that $M_{\text{gas}} = 9.2 \times 10^{11} / m M_\odot$. We suspect there is a gravitational lens along the line of sight, with an amplification factor m , or these values would be implausibly at least one order of magnitude larger than for rare hyperluminous objects. We derive the total FIR luminosity $L_{\text{FIR}} = 4.8 \times 10^{14} / m L_\odot$ and the star-formation rate $SFR = 8.3 \times 10^4 / m M_\odot/\text{yr}$, by applying the relation of Kennicutt (1998).

¹ <http://www.iram.es/IRAMES/mainWiki/EmirforAstronomers>

Table 2. Observed line parameters.

Line	ν_{obs} [GHz]	T_{mb} [mK]	S_{ν} [mJy]	ΔV_{FWHM} [km s $^{-1}$]	I [Jy km s $^{-1}$]	V^* [km s $^{-1}$]	$L'/m/10^{10}$ [K km s $^{-1}$ pc 2]
CO(4–3)	93.5249	5.3 ± 0.6	$26.7 \pm 3.$	175 ± 20	4.95 ± 0.5	0 ± 8	$19.7 \pm 2.$
CO(6–5)	140.2695	6.4 ± 1	$32.8 \pm 5.$	189 ± 19	6.6 ± 0.6	-7 ± 9	$11.5 \pm 1.$
CO(7–6)	163.6342	4.2 ± 0.9	21.5 ± 4.6	172 ± 27	3.9 ± 0.5	11 ± 11	5.0 ± 0.6
C I($^3P_1 - ^3P_0$)	99.8378	1.9 ± 0.6	$9.6 \pm 3.$	225 ± 55	2.3 ± 0.5	-50 ± 24	8.0 ± 1.7
C I($^3P_2 - ^3P_1$)	164.1802	$4.2 \pm 1.$	$21.5 \pm 5.$	184 ± 49	4.2 ± 0.8	8 ± 19	$5.3 \pm 1.$

Notes. Quoted errors are statistical errors from Gaussian fits. The systematic calibration uncertainty is 10%. ^(*) The velocity is given relative to $z = 3.929605$. **Fig. 2.** Available photometric data for MM18423+5938 (Table 1). These are superimposed upon our dust emission model (full red curve) based on the Milky Way dust model by Desert et al. (1990) adapted for our source at $z = 3.93$. It consists of three components; the large grains at $T_d = 45$ K, containing most of the mass (dotted line), the very small grains at hotter temperature (dot-dash), and the PAH (dashed line). **Fig. 3.** CO line flux density (points) or upper limits (arrow) measured at the IRAM-30 m, with the best-fit LVG model (dash line) computed with $n_{\text{H}_2} = 10^3 \text{ cm}^{-3}$, $T_k = 45 \text{ K}$, and $N_{\text{CO}}/\Delta V = 3 \times 10^{18} \text{ cm}^{-2}/\text{km s}^{-1}$.

3.2. CO lines

The CO SED of MM18423+5938 in Fig. 3 indicates moderate line excitation, peaking only at $J = 5$; CO SEDs peak at $J = 6$ or 7 in local starbursts such as M82 and NGC253 (Weiss et al. 2007), peak at higher J in AGN-dominated sources, e.g. $J = 10$ in APM0827, and reach a plateau for $J > 8$ in Mrk231 (van der Werf et al. 2010). We ran several LVG models, to constrain the H_2 volumic density and the kinetic temperature (cf. Combes et al. 1999). The moderate excitation implies a regime of low temperature and/or density. The gas kinetic temperature is taken to be equal to the dust temperature ($T_d = 45 \text{ K}$). When models are run with higher T_k , they all imply $n(\text{H}_2)$ lower than 10^3 cm^{-3} , which is not realistic for CO(7–6)-emitting clouds. Our estimate is therefore $T_k = 45 \text{ K}$, $n(\text{H}_2) = 10^3 \text{ cm}^{-3}$, and a column density per velocity interval

$N_{\text{CO}}/\Delta V = 3 \times 10^{18} \text{ cm}^{-2}/\text{km s}^{-1}$, as adopted to model the data in Fig. 3. From the CO(1–0) line derived from our LVG model, we infer an H_2 mass of $1.9 \times 10^{11}/m M_\odot$ with the conversion factor $M(\text{H}_2)/L'_{\text{CO}} = 0.8 M_\odot/(\text{K km s}^{-1} \text{ pc}^2)$ adopted for ULIRGs by Solomon et al. (1997). This latter value yields a lower limit, while the standard conversion ratio for the Milky Way, 5.75 times higher, yields the upper limit $M(\text{H}_2) = 1.1 \times 10^{12}/m M_\odot$. The true mass must lie between these two values, and indicates large amounts of molecular gas even if allowing for an amplification m . Assuming a typical intrinsic extension of 3 kpc (e.g. Tacconi et al. 2006), the surface filling factor of the molecular component is 0.3 for $m = 1$, and 0.03 for $m = 10$. The $L_{\text{FIR}}/L'_{\text{CO}}$ of $2 \times 10^3 L_\odot/\text{K km s}^{-1} \text{ pc}^2$ is consistent with ratios of other luminous infrared galaxies within the scatter (Iono et al. 2009).

3.3. Atomic carbon lines

The two lines C I($^3P_1 - ^3P_0$) and C I($^3P_2 - ^3P_1$) were clearly detected. They have comparable central velocity and line width (Table 2), implying that they originate from the same region in the source. The relation between the integrated C I($^3P_1 - ^3P_0$) brightness temperature and the beam averaged C I column density with the usual assumption of the optically thin limit is given by

$$N_{\text{CI}} = \frac{8\pi k \nu_{10}^2}{hc^3 A_{10}} Q(T_{\text{ex}}) \frac{1}{3} e^{T_1/T_{\text{ex}}} \int T_{\text{mb}} dv,$$

where $Q(T_{\text{ex}}) = 1 + 3e^{-T_1/T_{\text{ex}}} + 5e^{-T_2/T_{\text{ex}}}$ is the C I partition function, and $T_1 = 23.6 \text{ K}$ and $T_2 = 62.5 \text{ K}$ are the energies above the ground state. When dealing with high- z sources, we can use the definition of the line luminosity (e.g. Solomon et al. 1997) and derive the C I mass via (cf Weiss et al. 2003, 2005)

$$M_{\text{CI}} = 1.902 \times 10^{-4} Q(T_{\text{ex}}) e^{23.6/T_{\text{ex}}} L'_{\text{CI}(\text{^3P}_1 - \text{^3P}_0)} [M_\odot].$$

The mass estimated from the higher-excitation line is expressed in an analogous way, and we can then deduce that

$$\frac{L'_{\text{CI}(\text{^3P}_2 - \text{^3P}_1)}}{L'_{\text{CI}(\text{^3P}_1 - \text{^3P}_0)}} = 2.087 e^{-38.9/T_{\text{ex}}},$$

where line luminosities are given in Table 2. The derived T_{ex} is 33.9 K. The mass of atomic carbon thus amounts to $M_{\text{CI}} = 1.0 \times 10^8/m M_\odot$. Given our lower and upper limits to the H_2 mass from the CO lines, the $[\text{C I}]/[\text{H}_2]$ number abundance is between 1.4×10^{-5} and 8.0×10^{-5} . This is somewhat higher than the average $[\text{C I}]/[\text{H}_2]$ number abundance found in comparable star-forming objects (e.g. Barvainis et al. 2007; Pety et al. 2004; Weiss et al. 2003, 2005; Riechers et al. 2009; Danielson et al. 2010). In these latter estimates, although the observed $L_{\text{CI}}/L_{\text{CO}}$ values are comparable, the $[\text{C I}]/[\text{H}_2]$ number abundances are

Table 3. Line upper limits.

Line	ν_{obs} [GHz]	$S_{\nu}(3\sigma)$ [mJy]	$L'/10^{10}$ [K km s ⁻¹ pc ²]
CO(9–8)	210.344	<10.	<1.4
¹³ CO(4–3)	89.412	<10.	<8.1
HCN& HNC(5–4)	90.	<7.	<5.5
LiH(1–0) & HCO+(5–4)	90.	<7.	<5.5
H ₂ O _o (1,1,0–1,0,1)	112.978	<10.	<5.0
H ₂ O _p (2,1,1–2,0,2)	152.554	<12.	<3.3
CS(8–7)	79.488	<11.	<11.2
CS(9–8)	89.420	<7.	<5.6

somewhat dissimilar because of the various CO-to-H₂ conversion factors adopted by these authors. Abundance lower than 1.8×10^{-5} has been reported (Casey et al. 2010). The contribution of the atomic carbon to the cooling is low, $L_{\text{C I}}/L_{\text{FIR}} = 2.5 \times 10^{-6}$, comparable to that of nearby star-forming galaxies (Gerin & Phillips 2000).

3.4. Other lines

We searched for high-density tracers, such as HCN, HNC, and HCO⁺, in particular their lowest level available, i.e. $J = 5-4$. The upper limits found (Table 3) confirm that, on average, the H₂ density is not high, as found by our LVG models of the CO line excitation. The HCN luminosity is higher than one third of the CO luminosity in local AGN-dominated objects (Imanishi et al. 2004), and we note that our 3σ upper limit ($L'_{\text{HCN}(5-4)}/L'_{\text{CO}(5-4)} \sim 0.28$) is close to this limit. However, observation of HCN(1–0) and CO(1–0) are needed to conclude.

We also searched for H₂O emission, using the first ortho and para lines in their ground states, i.e. H₂O($J_{K_a K_c}=1_{10} \rightarrow 1_{01}$) and H₂O($J_{K_a K_c}=2_{11} \rightarrow 2_{02}$) but only obtained interesting upper limits. Assuming that these water lines are optically thick, these upper limits yield a filling factor of dense clumps lower than 30% that of CO clouds.

At $z = 2.3$, a tentative detection of the H₂O($J_{K_a K_c}=2_{11} \rightarrow 2_{02}$) line was reported in IRAS F10214 (Encrenaz et al. 1993; Casoli et al. 1994), while at $z = 0.685$, the fundamental transition of ortho-water, H₂O($J_{K_a K_c}=1_{10} \rightarrow 1_{01}$), was detected in absorption towards B0218+357 (Combes & Wiklind 1997). Water lines were searched for other starburst galaxies at high z (Riechers et al. 2006; Wagg et al. 2006), and significant upper limits set. A search for H₂O($J_{K_a K_c}=1_{10} \rightarrow 1_{01}$) emission toward the $z = 3.91$ quasar APM 08279+5255 provided a surface-filling factor lower than 12% of the CO one (Wagg et al. 2006; Weiss et al. 2007).

Finally, other interesting molecules were undetected in the observed bands, although with insufficient sensitivity: ¹³CO(4–3) (limiting the ¹³CO/¹²CO emission ratio to <1/2), CS(8–7), LiH (1–0) lines, several lines of formaldehyde H₂CO, and SiO, which is a tracer of shocks. Their lowest transitions observed are H₂CO(7(2,5)–8(0,8)), H₂CO(6(1,6)–5(1,5)), SiO(9–8) and upper limits are $L' < 20 \times 10^{10}$ K km s⁻¹ pc².

4. Discussion and conclusion

We have found that the brightest SMG in the North, MM18423+5938, is at a redshift $z = 3.92960 \pm 0.00013$. This source is part of a most interesting population of similar objects recently found by the Herschel Atlas survey (Negrello et al. 2010, submitted). From our modelled SED, the FIR luminosity $4.8 \times 10^{14}/m L_{\odot}$ and mass $6.0 \times 10^9/m M_{\odot}$ for the dust of MM18423+5938, and the implied star-formation rate of

$8.3 \times 10^4/m M_{\odot}/\text{yr}$, lead to suspect a gravitational lens along the line of sight with an amplification factor m as found for SMMJ2135-0102 (Swinbank et al. 2010). From our modelled CO SED, we have found that M(H₂) is between 1.9 and $10.8 \times 10^{11}/m M_{\odot}$ depending on the conversion ratio, indicating large amounts of molecular gas. The CO line excitation is moderate which indicates that there is both no strong heating by a central AGN and a starburst that is not too extreme. The average density of the molecular medium is low, of the order of 10^3 cm^{-3} , and the gas kinetic temperature is assumed to be 45 K in our model. The atomic carbon lines, assumed optically thin, are excited to $T_{\text{ex}} = 33.9\text{K}$ and yield $M_{\text{C I}} = 1.0 \times 10^8/m M_{\odot}$, which corresponds to a [C I]/[H₂] number abundance of between 1.4×10^{-5} and 8.0×10^{-5} when the limits on the H₂ mass derived from the CO lines are used. In this high- z SMG, the C I-to-CO luminosity ratio is consistent with those of other high- z galaxies.

The moderate CO line excitation found excludes a dominant AGN in MM18423+5938, unlike Mrk231 where CO is excited up to $J = 13$ (van der Werf et al. 2010). This moderate excitation favors an extended gas disk (typically 3 kpc), rather than a compact nuclear starburst (300 pc) and consequently a high CO-to-H₂ conversion ratio. This high-redshift SMG, with a star formation efficiency of $L_{\text{FIR}}/L'_{\text{CO}} = 2400$, is comparable to the lower- z submillimeter galaxies studied by Greve et al. (2005).

Acknowledgements. Based on observations carried out with the IRAM 30 m telescope. IRAM is supported by INSU/CNRS (France), M.P.G. (Germany) and I.G.N. (Spain). The authors are grateful to the IRAM staff for their support, and to the referee for helpful comments.

References

- Allard, F., Hauschildt, P. H., Alexander, D. R., et al. 2001, ApJ, 556, 357
- Bertoldi, F., Carilli, C., Aravena, M., et al. 2007, ApJSS, 172, 132
- Blain, A. W., Smail, I., Ivison, R. J., et al. 2002, PhR, 369, 111
- Casey, C. M., Chapman, S. C., Daddi, E., et al. 2009, MNRAS, 400, 670
- Casoli, F., Gerin, M., Encrenaz, P. J., & Combes, F. 1994, A&A, 287, 716
- Chapman, S. C., Blain, A. W., Smail, I., & Ivison, R. J. 2005, ApJ, 622, 772
- Combes, W. T. 1997, ApJ, 486, L79
- Combes, F., Maoli, R., & Omont, A. 1999, A&A, 345, 369
- Danielson, A., Swinbank, A., Smail, I., et al. 2010, MNRAS [arXiv:1008.3183]
- Desert, F.-X., Boulanger, F., & Puget, J.-L. 1990, A&A, 237, 215
- Encrenaz, P., Combes, F., Casoli, F., et al. 1993, A&A, 273, L19
- Gerin, M., & Phillips, T. G. 2000, ApJ, 537, 644
- Greve, T. R., Bertoldi, F., Smail, I., et al. 2005, MNRAS, 359, 1165
- Greve, T. R., Pope, A., Scott, D., et al. 2008, MNRAS, 389, 1489
- Hinshaw, G., Weiland, J. L., Hill, R. S., et al. 2009, ApJS, 180, 225
- Imanishi, M., Nakanishi, K., Kuno, N., & Kohno, K. 2004, AJ, 128, 2037
- Iono, D., Wilson, C. D., Yun, M. S., et al. 2009, ApJ, 695, 1537
- Kennicutt, R. 1998, ARA&A, 38, 189
- Kreysa, E., Gemuend, H. P., Gromke, J., et al. 1998, SPIE, 3357, 319
- Lestrade, J.-F., Wyatt, M. C., Bertoldi, F., et al. 2009, A&A, 506, 145
- Pety, J., Beelen, A., Cox, P., et al. 2004, A&A, 428, L21
- Riechers, D. A., Weiss, A., Walter, F., et al. 2006, ApJ, 649, 635
- Riechers, D. A., Walter, F., Bertoldi, F., et al. 2009, ApJ, 703, 1398
- Riechers, D. A., Capak, P. L., Carilli, C., et al. 2010, ApJ, 720, L131
- Smail, I., Ivison, R. J., & Blain, A. W. 1997, ApJ, 490, L5
- Smail, I., Ivison, R. J., Blain, A. W., & Kneib, J. P. 2002, MNRAS, 331, 495
- Swinbank, A. M., Smail, I., Longmore, S., et al. 2010, Nature, 464, 733
- Solomon, P. M., Downes, D., Radford, S. J. E., & Barrett, J. W. 1997, ApJ, 478, 144
- Tacconi, L. J., Neri, R., Chapman, S. C., et al. 2006, ApJ, 640, 228
- van der Werf, P., Isaak, K., Meijerink, R., et al. 2010, A&A, 518, L42
- Vieira, J. D., Crawford, T. M., Switzer, E. R., et al. 2010, ApJ, 719, 763
- Wagg, J., Wilner, D. J., Neri, R., Downes, D., & Wiklind, T. 2006, ApJ, 651, 46
- Wardlow, J. L., Smail, I., Coppin, K. E. K., et al. 2010, MNRAS, submitted [arXiv:1006.2137]
- Weiss, A., Henkel, C., Downes, D., & Walter, F. 2003, A&A, 409, L41
- Weiss, A., Downes, D., Henkel, C., & Walter, F. 2005, A&A, 429, L25
- Weiss, A., Downes, D., Neri, R., et al. 2007, A&A, 467, 955
- Weiss, A., Ivison, R. J., Downes, D., et al. 2009, ApJ, 705, L45