The direct oxygen abundances of metal-rich galaxies derived from electron temperature (Corrigendum)

Y. C. Liang^{1,2}, F. Hammer², S. Y. Yin^{1,3,4}, H. Flores², M. Rodrigues², and Y. B. Yang^{2,1}

¹ National Astronomical Observatories, Chinese Academy of Sciences, 20A Datun Road, Chaoyang District, Beijing 100012, PR China

e-mail: ycliang@nao.cas.cn

² GEPI, Observatoire de Paris-Meudon, 92195 Meudon, France

³ Department of Physics, Hebei Normal University, Shijiazhuang 050016, PR China

⁴ Department of Physics, College of Technology, Harbin University, Haerbin 150086, PR China

A&A 473, 411–421 (2007), DOI: [10.1051/0004-6361:20077436](https://doi.org/10.1051/0004-6361:20077436)

Key words. galaxies: abundances – galaxies: stellar content – galaxies: ISM – galaxies: spiral – galaxies: starburst – errata, addenda

The published Eq. (8) in our paper (Liang et al. 2007, A&A, 473, 411) should be replaced by the formula below:

$$\log R_{23} = 73.1323 - 32.7809x + 4.80642x^2 - 0.229631x^3, \quad (8)$$

where x refers to $12 + \log(\text{O}/\text{H})$.

This does not affect our result, shown by the solid line in Fig. 7b, since it was computed with Eq. (1) here actually.

Acknowledgements. We thank Vallery Stanishev for pointing out this problem.