

# Kinetics of the $\text{OCN}^-$ and $\text{HOCN}$ formation from the $\text{HNCO} + \text{H}_2\text{O}$ thermal reaction in interstellar ice analogs

P. Theule<sup>1</sup>, F. Duvernay<sup>1</sup>, A. Ilmane<sup>1</sup>, T. Hasegawa<sup>2</sup>, O. Morata<sup>2</sup>, S. Coussan<sup>1</sup>, G. Danger<sup>1</sup>, and T. Chiavassa<sup>1</sup>

<sup>1</sup> Université de Provence, laboratoire de Physique des Interactions Ioniques et Moléculaires, Centre de St-Jérôme, Avenue Escadrille Normandie-Niémen, 13397 Marseille, France  
e-mail: patrice.theule@univ-provence.fr

<sup>2</sup> Academia Sinica Institute of Astronomy and Astrophysics Astronomy-Mathematics Building, NTU, Taipei, ROC

Received 3 November 2010 / Accepted 21 February 2011

## ABSTRACT

**Aims.** We study in the laboratory the kinetics of the low-temperature  $\text{OCN}^-$  and  $\text{HOCN}$  formation from the purely thermal reaction of solid  $\text{HNCO}$  and  $\text{H}_2\text{O}$ . The cyanate ion  $\text{OCN}^-$  is an intermediate in the isomerization process of isocyanic acid  $\text{HNCO}$  into cyanic acid  $\text{HOCN}$  in water ice.

**Methods.** We study the reaction, isomerization and desorption kinetics of the  $\text{HNCO}/\text{OCN}^-/\text{HOCN}$  system using Fourier transform infrared spectroscopy.

**Results.** Activation energies of  $26 \pm 2 \text{ kJ mol}^{-1}$  (3127 K) and  $36 \pm 1 \text{ kJ mol}^{-1}$  (4330 K) are found for the  $\text{HNCO} + \text{H}_2\text{O} \rightarrow \text{OCN}^- + \text{H}_3\text{O}^+$  and  $\text{OCN}^- + \text{H}_3\text{O}^+ \rightarrow \text{HOCN} + \text{H}_2\text{O}$  reactions respectively. Desorption energies of  $37 \pm 3 \text{ kJ mol}^{-1}$  (4450 K) and  $40 \pm 3 \text{ kJ mol}^{-1}$  (4811 K) are measured for  $\text{HNCO}$  and  $\text{OCN}^-$ , respectively.

**Conclusions.** The present experiment has the important implication that the  $\text{H}_2\text{O} + \text{HNCO}$  reaction alone cannot account for the observed abundances of solid  $\text{OCN}^-$  in astronomical IR sources.

**Key words.** astrochemistry – ISM: molecules – molecular processes – molecular data

## 1. Introduction

The spectra of several protostellar objects exhibit a strong absorption feature at 4.62 microns ( $2165 \text{ cm}^{-1}$ ), called the XCN band, which is commonly attributed to the solid-phase cyanate ion  $\text{OCN}^-$  (Grim & Greenberg 1987; Lowenthal et al. 2000; Novozamsky et al. 2001; van Broekhuizen et al. 2005). A systematic study of this feature with the VLT-ISAAC spectrometer shows large source-to-source abundance variations (van Broekhuizen et al. 2005). However, this feature seems absent from the limited sample of starless dark molecular clouds investigated by Knez et al. (2005). Thus, the  $\text{OCN}^-$  band is often seen as a manifestation of thermal or non-thermal energetic processes of the protostellar environment.

The formation of the solid-state cyanate ion  $\text{OCN}^-$  can be explained by different processes: proton bombardment (Moore et al. 1983), electron bombardment (Bennett et al. 2010), vacuum ultraviolet photolysis of  $\text{NH}_3:\text{CO}:\text{H}_2\text{O}$  ices where the photochemical formation of isocyanic acid  $\text{HNCO}$  is followed by a proton transfer to ammonia  $\text{NH}_3$  (d'Hendecourt et al. 1986; Grim et al. 1989), or directly from an acid-base reaction of  $\text{HNCO}$  with solid  $\text{NH}_3$  or  $\text{H}_2\text{O}$  (Demyk et al. 1998; Raunier et al. 2003b; van Broekhuizen et al. 2004). A good review of all the experiments carried out on  $\text{OCN}^-$  and of the processes that lead to its formation is given in Table 1 of van Broekhuizen et al. (2005) and a quantitative comparison of these different processes is given in van Broekhuizen et al. (2004). Based on UV fluence and  $\text{NH}_3$  abundance arguments, the authors conclude that the UV photoprocessing of  $\text{NH}_3:\text{CO}:\text{H}_2\text{O}$  ices may not account for the observed  $\text{OCN}^-$  abundances, and that  $\text{OCN}^-$  is most

likely formed thermally from  $\text{HNCO}$ . The VLT studies of low-mass young stellar objects (YSOs) (van Broekhuizen et al. 2005) show that if  $\text{OCN}^-$  abundances allow quantitatively for a photochemical mechanism, the large variations in abundances are more difficult to explain by photochemistry; thus the thermal formation, with either  $\text{NH}_3$  or  $\text{H}_2\text{O}$ , should be seriously considered. However, the validity of these mechanisms to reproduce or not the observed abundances of  $\text{OCN}^-$  depends on their respective kinetics, and no kinetic studies have been available so far.

Experimental and theoretical low-temperature studies on the acid-base reaction of  $\text{HNCO}$  with solid  $\text{NH}_3$  have confirmed the spontaneous character and the effectiveness of its formation from  $\text{HNCO}$  (Raunier et al. 2003a,b, 2004; van Broekhuizen et al. 2004). Moreover, the thermal deprotonation of  $\text{HNCO}$  by  $\text{H}_2\text{O}$  is observed (Raunier et al. 2003b), though it is less effective than the deprotonation of  $\text{HNCO}$  by  $\text{NH}_3$ , which indicates a higher activation energy. However, whatever their activation energies are, these reactions are limited by the diffusion and the abundances of available reactants. If we consider an average temperature of 30–40 K for YSOs ices, the mobility of  $\text{H}_2\text{O}$ ,  $\text{NH}_3$  or  $\text{HNCO}$  is negligible, therefore the most likely reaction to occur is statistically set by the most abundant reactants species, i.e.  $\text{H}_2\text{O}$ , because the  $\text{NH}_3$  abundance with respect to  $\text{H}_2\text{O}$  is typically less than 5% (van Broekhuizen et al. 2005). Thus one can think that the dominant reaction may be the thermal reaction of  $\text{H}_2\text{O}$  with  $\text{HNCO}$ , and that the  $\text{OCN}^-$  abundance is mostly set by the kinetics of the  $\text{H}_2\text{O} + \text{HNCO}$  reaction, assuming the presence of the solid-state  $\text{HNCO}$  reactant within interstellar ices.

However,  $\text{HNCO}$  has never been observed in the solid phase of the ISM. Yet, gas phase isocyanic acid  $\text{HNCO}$  was one of the

first interstellar molecules identified in the interstellar medium (ISM) (Snyder & Buhl 1972). HNCO is also the simplest species containing all four most abundant elements, H, N, C and O, and four isomers have this elemental composition: isocyanic acid HNCO, cyanic acid HOCN, fulminic acid HCNO and isofulminic acid HONC. HNCO is the most stable isomer, and *ab initio* calculations give the energies of their ground electronic singlet states relative to HNCO to be 24.7, 70.7 and 84.1 kcal mol<sup>-1</sup>, respectively (Mladenović & Lewerenz 2008). However, it is only recently that fulminic acid HCNO has been detected and the HNCO/HCNO abundance ratio is observed to change by one order of magnitude depending on the object (Marcelino et al. 2009). Cyanic acid HOCN has also been tentatively identified in SgrB2 (Brünken et al. 2009) and the HOCN/HNCO abundance ratio estimated at 0.5%. What can be deduced from this ratio is strongly dependent on the chemical pathways that lead to the formation of the two isomers and on the isomerization pathways from one isomer to another. The chemistry related to HNCO and its isomers is not known and there has been much speculation about HNCO formation. For some authors, HNCO is formed only in the gas phase (Turner et al. 1999; Iglesias 1977), for other authors, HNCO is formed on grain surfaces and then desorbed (Hasegawa et al. 1992; Garrod et al. 2008; Tideswell et al. 2010). For the latter authors, HNCO is a transient species that is rapidly processed to OCN<sup>-</sup> on grain surface during the collapse phase, and then, during the hot core stage, these species are desorbed and subsequently destroyed in the gas-phase returning to HNCO (Tideswell et al. 2010). HNCO is also proposed to be formed in the gas-phase in post-shock gas because HNCO abundance is enhanced in the presence of shocks and correlates well the SiO abundances (Zinchenko et al. 2000).

In this paper we address the solid-state chemistry of the HNCO/OCN<sup>-</sup>/HOCN/H<sub>2</sub>O system on the basis of laboratory experiments using Fourier-transform infrared (FTIR) spectroscopy on interstellar ice analogs. Without speculating on HNCO formation mechanisms in the interstellar medium, we study the kinetics of the conversion of HNCO into OCN<sup>-</sup> and then into HOCN in a pure H<sub>2</sub>O environment. The H<sub>2</sub>O + HNCO reaction gives the salt H<sub>3</sub>O<sup>+</sup>OCN<sup>-</sup> with an activation energy  $E_1 = 26 \pm 2$  kJ mol<sup>-1</sup> (3127 K) (1 kJ mol<sup>-1</sup> = 120.272 K). Then, the proton is transferred from H<sub>3</sub>O<sup>+</sup> to OCN<sup>-</sup> to give the isocyanic acid HOCN, with an activation energy  $E_2 = 36 \pm 1$  kJ mol<sup>-1</sup> (4330 K). We show that OCN<sup>-</sup> follows a kinetic law that can be written as  $dx(\text{OCN}^-)/dt = k_1 \times x(\text{HNCO})^2 - k_2 \times x(\text{OCN}^-)^2$ . We also study the desorption kinetics of HNCO and OCN<sup>-</sup> to derive their desorption energies,  $E_{\text{des}}(\text{HNCO}) = 37 \pm 3$  kJ mol<sup>-1</sup> (4450 K) and  $E_{\text{des}}(\text{OCN}^-) = 40 \pm 3$  kJ mol<sup>-1</sup> (4811 K). This laboratory work on the isolated system HNCO:H<sub>2</sub>O provides the reference input data to a gas-grain chemistry model that we will use to investigate the observed solid-state OCN<sup>-</sup> abundances in a complete interstellar ice.

## 2. Experimental

The HNCO monomer is synthesized from the thermal decomposition of the cyanuric acid polymer (Aldrich Chemical Co., 98%) at 650°C under a primary vacuum (Raunier et al. 2003b). HNCO and H<sub>2</sub>O are inserted and outgassed in two separated vacuum lines using standard manometric techniques, and then they are co-deposited. The two gases are sprayed onto a highly polished copper surface maintained at 10 K using closed-cycle helium cryostats (ARS Cryo, model DE-204 SB) in a vacuum chamber at 10<sup>-8</sup> mbar. The temperature is controlled between 10 K and 300 K using a heating resistance and a Lakeshore Model 336

temperature controller. The IR spectra are recorded at different temperatures in the reflection mode using a Vertex 70 spectrometer with a MCT detector. A typical spectrum has a 1 cm<sup>-1</sup> resolution and is averaged over one hundred interferograms.

The concentration ratio of the different mixtures is first set at room temperature before deposition using leak valves to adjust fluxes and then derived at 10 K after deposition from the IR spectra by integrating the bands to estimate the column density of HNCO, according to their band strengths. For H<sub>2</sub>O, we use the values of  $3.1 \times 10^{-17}$  cm molecule<sup>-1</sup> for the band strength of the libration mode at 760 cm<sup>-1</sup> and  $2.0 \times 10^{-16}$  cm molecule<sup>-1</sup> for the band strength of the OH stretching mode at 3280 cm<sup>-1</sup> (d'Hendecourt & Allamandola 1986). For HNCO, we adopt the value  $7.2 \times 10^{-17}$  cm molecule<sup>-1</sup> for the band strength of the  $\nu_{\text{as}}\text{NCO}$  mode at 2251 cm<sup>-1</sup>, which is the value given for a H<sub>2</sub>O-rich ice (van Broekhuizen et al. 2004). For OCN<sup>-</sup> we use the value  $1.3 \times 10^{-16}$  cm molecule<sup>-1</sup> for the band strength of the  $\nu_{\text{as}}\text{NCO}$  mode (van Broekhuizen et al. 2004). We also use these bands to monitor the evolution of the HNCO, OCN<sup>-</sup> abundances because they are consumed or produced during the reactions to estimate the different kinetical parameters. However, the calculated concentration ratio depends on the band strengths provided by the literature. These values depend on the nature, composition, and temperature of the ice in which they are found, and this dependence is a major source of uncertainties when evaluating the column densities of frozen molecules. A 20% uncertainty on the band strengths and consequently on calculated column densities is probably sufficiently conservative. Moreover, some bands are overlapping in the HNCO:H<sub>2</sub>O mixture, especially the H<sub>2</sub>O libration mode and the  $\delta\text{HNC}$  mode of HNCO, which adds uncertainty to the column density determination.


In a typical kinetic experiment, the temperature is set to a fixed value using the temperature controller and measured using a DTGS 670 silicon diode with a 0.1 K uncertainty. Infrared spectra are recorded at this fixed temperature at fixed intervals of time.

The activation energy of desorption and the order of desorption for HNCO, OCN<sup>-</sup> and HOCN can be obtained from temperature program desorption (TPD) experiments followed either by mass spectrometry of the desorbed gas phase species or infrared spectroscopy of the remaining solid-state species on their respective characteristic bands. The latter technique has the advantage to distinguish between the desorption of HNCO, OCN<sup>-</sup> and HOCN, whereas their fragmentation is identical in a quadrupole mass spectrometer. The IR solid-phase signal from the remaining material correlates with the desorbing gas phase material measured with a quadrupole mass spectrometer within the measurement uncertainty (Bossa et al. 2009). Thus we will follow the TPD experiments using IR spectroscopy of the solid-state species. The TPD linear variation of the temperature with time (temperature ramp,  $T = T_0 + \beta \times t$ ) is performed using the temperature controller.

## 3. Results

### 3.1. Formation of H<sub>3</sub>O<sup>+</sup>OCN<sup>-</sup> and HOCN from a HNCO:H<sub>2</sub>O ice mixture

We start with a HNCO:H<sub>2</sub>O ice mixture where H<sub>2</sub>O is in excess with respect to HNCO. Figure 2 shows the thermal evolution of a HNCO:H<sub>2</sub>O ice mixture in a 1:5 ratio deposited between 10 and 14 K. A typical ice sample has a thickness of few 100 monolayers. At 10 K the spectrum is dominated by the strong water absorption bands at 3280, 1628, and 760 cm<sup>-1</sup>, assigned to the



**Fig. 1.** Infrared spectrum of a HNCO:H<sub>2</sub>O mixture in a 1:5 concentration ratio at  $T = 120$  K. A small amount of OCN<sup>-</sup> is formed during the deposition.

**Table 1.** Positions and integrated band strengths of a HNCO:H<sub>2</sub>O mixture at 10 K.


HNCO:H <sub>2</sub> O ice mixture at 15 K			
Position (cm <sup>-1</sup> )	$A$ (cm molecule <sup>-1</sup> )	mode	molecule
3560, 3373, 3233	–	$\nu$ NH	HNCO
3280	$2.1 \times 10^{-16a}$	$\nu$ OH	H <sub>2</sub> O
2900	weak <sup>c</sup>	$\nu$ OH	H <sub>3</sub> O <sup>+</sup>
2295	–	$\nu_{as}$ NCO	HOCN
2251	$7.2 \times 10^{-17b}$	$\nu_{as}$ NCO	HNCO
2170	$1.3 \times 10^{-16b}$	$\nu_{as}$ NCO	OCN <sup>-</sup>
1750	– <sup>c</sup>	$\delta$ OH	H <sub>3</sub> O <sup>+</sup>
1660	$3.1 \times 10^{-17}$	$\delta$ OH	H <sub>2</sub> O
1205	– <sup>c</sup>	$\nu$ OH	H <sub>3</sub> O <sup>+</sup>
858	–	$\delta$ HNC	HNCO
760	$3.1 \times 10^{-17a}$	$\nu_L$	H <sub>2</sub> O

**References.** <sup>(a)</sup> d’Hendecourt & Allamandola (1986); <sup>(b)</sup> van Broekhuizen et al. (2004); <sup>(c)</sup> Falk & Guiguere (1957).

OH stretching mode, OH bending mode, and libration mode, respectively. Isocyanic acid HNCO is characterized by strong bands located at 3373, 3233, 2251, and 858 cm<sup>-1</sup>, but only the 2251 cm<sup>-1</sup> band, assigned to the NCO antisymmetric stretch vibration, is not blended by the water bands, which is why it will be used to monitor the HNCO evolution. The 2251 cm<sup>-1</sup> band has actually two components: one component at 2260 cm<sup>-1</sup> corresponding to HNCO in a H<sub>2</sub>O environment (Raunier et al. 2003b), and a component 2234 cm<sup>-1</sup> corresponding to HNCO in a HNCO environment. The proportion of the two components in the 2251 cm<sup>-1</sup> band depends on the ice sample deposition, and accordingly the position of the band may be shifted from one deposition to another by few wavenumbers. The position and shape of this band are also slightly changing while HNCO is reorganizing around 120 K. Figure 1 shows a typical HNCO:H<sub>2</sub>O mixture spectrum just after the temperature is brought to 120 K. Although the HNCO + H<sub>2</sub>O reaction is not possible in the gas phase at room temperature, during the deposition step when the gases are thermalizing from 300 K to 14 K on the cold surface, a little amount of OCN<sup>-</sup> is formed, as can be seen from its band at 2170 cm<sup>-1</sup> as seen in Fig. 1. The amount of initially formed OCN<sup>-</sup> varies from one deposition to another.

All the observed bands are listed in Table 1 along with their corresponding assignments and band strengths.

When the temperature is set to a fixed value between 100 K and 130 K, the OCN<sup>-</sup> band at 2170 cm<sup>-1</sup> is progressively growing while the bands corresponding to HNCO and H<sub>2</sub>O decrease as can be seen in Fig. 2. The OCN<sup>-</sup> feature keeps growing until it reaches a stationary state corresponding to this temperature. This



**Fig. 2.** Infrared spectrum **a)** and difference spectrum (spectrum at time  $t$  minus spectrum at  $t = 0$ ) **b)** of a HNCO:H<sub>2</sub>O mixture at  $T = 120$  K at  $t = 0$  s, 3300 s, 21 300 s and 53 700 s. OCN<sup>-</sup> has a band at 2170 cm<sup>-1</sup> and HOCN at 2295 cm<sup>-1</sup>. The HNCO band has two components: one component at 2260 cm<sup>-1</sup> corresponding to HNCO in a H<sub>2</sub>O environment and one component at 2234 cm<sup>-1</sup> corresponding to HNCO in a HNCO environment. The variation of the second component is the result of a rearrangement that can give a positive band at 2251 cm<sup>-1</sup> in some spectra. The bands at 2349 cm<sup>-1</sup> correspond to fluctuations of the gas phase CO<sub>2</sub>.


indicates that the H<sub>3</sub>O<sup>+</sup>OCN<sup>-</sup> species is formed thermally by the reaction involving HNCO and H<sub>2</sub>O. Owing to their low band strengths it is impossible to see the H<sub>3</sub>O<sup>+</sup> features at 1205, 1750 and 2900 cm<sup>-1</sup> (Falk & Guiguere 1957), but because H<sub>2</sub>O is the only other species, the counter ion of OCN<sup>-</sup> is necessarily H<sub>3</sub>O<sup>+</sup>, and the H<sub>3</sub>O<sup>+</sup>OCN<sup>-</sup> species is formed from a thermally induced proton transfer from HNCO to H<sub>2</sub>O. It is important to know whether the reaction HNCO + H<sub>2</sub>O that gives OCN<sup>-</sup> + H<sub>3</sub>O<sup>+</sup> is a total reaction (HNCO + H<sub>2</sub>O → OCN<sup>-</sup> + H<sub>3</sub>O<sup>+</sup>) or an equilibrium (HNCO + H<sub>2</sub>O ⇌ OCN<sup>-</sup> + H<sub>3</sub>O<sup>+</sup>). In the former case all HNCO is consumed to give OCN<sup>-</sup> after a certain time; in the latter case the reverse reaction occurs where OCN<sup>-</sup> gives HNCO, and an equilibrium depending on the temperature and Gibbs energies is reached. In the former case the HNCO/OCN<sup>-</sup> ratio is only defined by the kinetics, in the latter case the HNCO/OCN<sup>-</sup> ratio is defined by the thermodynamics.

We can see from Fig. 2 that the reaction is not total, and thus, the system should be in an equilibrium. If this is true, we should be able to derive an equilibrium constant from the final abundances of the reactants and products, the equilibrium constant being dependent on the temperature. To test this latter assumption, after waiting for a sufficiently long time for a stationary state to be achieved at a fixed temperature of 125 K,


the temperature is lowered by 5 K to displace the equilibrium constant. The system does not respond to the temperature variation, the abundance of each species remains the same as at 125 K, and no displacement of the equilibrium constant is observed. Thus, we conclude that the system is not in equilibrium either. At that point we see that we cannot push the analogy between liquid solutions and solid solutions further. Some molecules are in configurations that enable the total reaction, some molecules are not, like in clusters of pure HNCO for example. Only a limited set of HNCO molecules are in a favorable configuration, one HNCO molecule making a hydrogen bond with the electron lone pair of one H<sub>2</sub>O molecule and being activated enough (Raunier et al. 2003b). We conclude that for the population of molecules that are statistically in a favorable configuration, the reaction is total, because the reverse reaction is not observed.

Figure 2 shows that in addition to the the OCN<sup>-</sup> band at 2170 cm<sup>-1</sup>, a new band appears at 2295 cm<sup>-1</sup>. We assign this band to the  $\nu_{\text{as}}\text{NCO}$  mode of cyanic acid HOCN (Raunier et al. 2003b), since this transition has been observed at 2280 cm<sup>-1</sup> in a rare gas matrix environment at 13 K (Teles et al. 1989). The +15 cm<sup>-1</sup> difference observed for HOCN is consistent with the +8 cm<sup>-1</sup> frequency difference observed for the  $\nu_{\text{as}}\text{NCO}$  band between an Argon matrix at 10 K, 2259 cm<sup>-1</sup>, and a solid HNCO:H<sub>2</sub>O ice mixture at 120 K, 2251 cm<sup>-1</sup>. Not all experiments show a HOCN band. This can be explained because the yield of conversion of HNCO into OCN<sup>-</sup> is a few percents of the total amount of HNCO, the speed of conversion of this low amount of OCN<sup>-</sup> into HOCN must be very slow and HOCN should not be observed. The amount of HOCN formed during the experiment must therefore come from the initial amount of OCN<sup>-</sup>, formed at  $t = 0$  during the deposition when the gas is thermalizing on the surface from 300 K to 10 K. Not all experiments have enough OCN<sup>-</sup> formed initially at the deposition to form an observable amount of HOCN during the experiment.

In the first reaction a proton is transferred at a rate  $k_1$  from the HNCO nitrogen atom to the H<sub>2</sub>O oxygen atom to form the H<sub>3</sub>O<sup>+</sup>OCN<sup>-</sup> species. In the second reaction, a proton is transferred at rate  $k_2$  from the H<sub>3</sub>O<sup>+</sup> oxygen atom to the OCN<sup>-</sup> oxygen atom (via an assisted proton transfer through a chain of H<sub>2</sub>O molecules) to form HOCN as expressed in system of Eqs. (1):



Indeed, to derive the rates of formation ( $k_1$ ) and destruction ( $k_2$ ) of OCN<sup>-</sup>, we need to include the desorption of each species in the total reaction network, as expressed in the system of Eqs. (2):



To extract the reaction rates  $k_1$  and  $k_2$  from our experimental kinetic data we need to derive the  $k_{\text{des}}(\text{HNCO})$ ,  $k_{\text{des}}(\text{OCN}^-)$  and  $k_{\text{des}}(\text{HOCN})$  desorption rates. As for the desorption of OCN<sup>-</sup>, we do not know whether the proton is attaching to the nitrogen or the oxygen before desorbing, and it desorbs either as HNCO or HOCN.

### 3.2. HNCO and OCN<sup>-</sup> desorption

Figure 4 shows the decrease of the solid phase material along with the temperature ramp during a TPD experiment.

The desorption curves obtained from the IR signal are fitted to the expression Bossa et al. (2009)

$$\frac{C(T)}{C(T = T_0)} = \exp(-\nu_1 \times (T - T_0)/\beta \times \exp(-E_{\text{des}}/(R \times T))), \quad (3)$$


where  $C(T)$  is the column density of the solid-phase molecules at temperature  $T$ .  $T_0$  and  $\beta$  are the initial temperature and the heating rate respectively,  $\nu_1$  the first-order pre-exponential term,  $E_{\text{des}}$  the desorption energy and  $R$  the ideal gas constant.

The desorption curves depend on the HNCO:H<sub>2</sub>O concentration ratio. If the HNCO:H<sub>2</sub>O concentration ratio is high, as seen in Fig. 4, the desorption curves shows two inflexion points corresponding to two desorption events. The first one corresponds to the desorption of pure HNCO, and the second one corresponds to HNCO embedded in water ice, desorbing along with the H<sub>2</sub>O mantle. Indeed, the second inflexion point gives the desorption energies of H<sub>2</sub>O,  $E_{\text{des}}(\text{H}_2\text{O}) \approx 44$  kJ mol<sup>-1</sup>. If the HNCO:H<sub>2</sub>O concentration ratio is low, the desorption curves gives the desorption energies of H<sub>2</sub>O.

Averaging the desorption energies obtained from different temperature programmed experiments, at different temperature ramp rate and HNCO:H<sub>2</sub>O concentration ratios, considering pure desorption and not desorption along with water, we obtain the following desorption energies:  $E_{\text{des}}(\text{HNCO}) = 37 \pm 3$  kJ mol<sup>-1</sup>, and  $E_{\text{des}}(\text{OCN}^-) = 40 \pm 3$  kJ mol<sup>-1</sup>. HOCN is produced in an insufficient amount and mainly desorbs along with H<sub>2</sub>O, so that it has not been possible to measure its desorption energy. We were unable to derive the pre-exponential parameter of the desorption kinetics, but it is possible to use a value of  $\nu_0 = 2.9 \times 10^{12}$  s<sup>-1</sup> using the expression for the characteristic desorption frequency given in Hasegawa et al. (1992). These TPD experiments also tell us about the residence time of the species. If the temperature is higher than 130 K, the residence time will be shorter than any chemical reaction timescale.

### 3.3. Kinetics studies on the HNCO:OCN<sup>-</sup>:HOCN:H<sub>2</sub>O system

We see that in two consecutive steps process isocyanic acid HNCO is isomerized into cyanic acid HOCN: a proton is first dragged from the nitrogen atom to a water molecule, and second a proton from another water molecule is attached to the oxygen atom of OCN<sup>-</sup>. This reaction is purely thermal, and does not necessitate an earlier photochemical formation of radicals. To derive a kinetic law for the formation of OCN<sup>-</sup> from HNCO and the subsequent formation of HOCN from OCN<sup>-</sup>, we need to carry out experiments at a temperature high enough that the formation can be measured in a realistic time scale of a few hours. This corresponds to a temperature above 100 K. However, as can be seen in Fig. 4, for a temperature above 130 K, the desorption of HNCO, OCN<sup>-</sup>, and HOCN needs to be taken into account to solve system 2, because they are desorbing at a timescale close to the chemical reactions involved in this system. Measuring kinetics below 130 K allows us to conveniently simplify system 2 into system 1. However, measures at 130 K are very convenient because the kinetics are fast and the HOCN band is clearly visible. In this case the desorption equations need to be taken into account. Above 130 K the desorption process is dominant.



**Fig. 3.** Time evolution of the  $\text{HNCO}$  (circles),  $\text{OCN}^-$  (crosses) and  $\text{HO CN}$  (squares) normalized abundances recorded by FTIR spectroscopy at 120 K for a  $\text{HNCO}:\text{H}_2\text{O}$  mixture in a 1:3 concentration ratio. Fitting the experimental data to the analytical expressions of evolution for each species (dashed lines) gives the reaction rates at  $T = 120$  K.

### 3.3.1. The temperature dependence of the reaction rates


To simplify the system of Eqs. (2) into the system of Eqs. (1) we will first only consider the chemical reactions while neglecting the desorption of the species as long as the temperature is lower than, or equal to, approximately 120 K. The general form of the kinetic law for the first reaction can be written as

$$\begin{cases} \frac{dx(\text{HNCO})}{dt} = -k_1(T) \times x(\text{H}_2\text{O})^1 \times x(\text{HNCO})^1 \\ \frac{dx(\text{OCN}^-)}{dt} = k_1(T) \times x(\text{H}_2\text{O})^1 \times x(\text{HNCO})^1 \\ \quad - k_2(T) \times x(\text{OCN}^-)^1 \times x(\text{H}_3\text{O}^+)^1 \\ \frac{dx(\text{HO CN})}{dt} = k_2(T) \times x(\text{OCN}^-)^1 \times x(\text{H}_3\text{O}^+)^1, \end{cases} \quad (4)$$

where  $x(\text{H}_2\text{O})$ ,  $x(\text{HNCO})$ ,  $x(\text{OCN}^-)$  and  $x(\text{HO CN})$  are  $\text{H}_2\text{O}$ ,  $\text{HNCO}$ ,  $\text{OCN}^-$  and  $\text{HO CN}$  molar fractions, respectively. For example we have  $x(\text{HNCO}) = \frac{n(\text{HNCO})}{n(\text{HNCO}) + n(\text{H}_2\text{O})}$ . When  $\text{H}_2\text{O}$  is much more abundant than any other species,  $\text{H}_2\text{O}$  is a solvent in a solid solution and molar fractions tend to concentrations.

To determine the temperature dependence of  $k_1(T)$  and  $k_2(T)$  we record the time evolution of the  $\text{HNCO}:\text{OCN}^-:\text{HO CN}:\text{H}_2\text{O}$  system at a fixed temperature and for different  $\text{HNCO}:\text{H}_2\text{O}$  concentration ratios. Figure 3 shows the evolution of the  $\text{HNCO}$ ,  $\text{OCN}^-$ , and  $\text{HO CN}$  integrated abundances at  $T = 120$  K for a 1:3 concentration ratio, which is equivalent to  $(\text{HNCO}) = 0.25$ . To monitor the time evolution of  $\text{HNCO}$ , we record the integrated absorbance of the  $2260 \text{ cm}^{-1}$  band on a difference spectrum, i.e. spectrum at time  $t$  minus spectrum at  $t = 0$ . This band corresponds to  $\text{HNCO}$  in a  $\text{H}_2\text{O}$  environment at 120 K that is consumed during the reaction. The  $\text{OCN}^-$  evolution is monitored using the  $2170 \text{ cm}^{-1}$  band and  $\text{HO CN}$  the  $2295 \text{ cm}^{-1}$  band. Moreover, because of the two components in the  $\text{HNCO}$  band, the changes in the  $\text{HNCO}$  band at  $2260 \text{ cm}^{-1}$  are difficult to monitor, which is why integrating the  $\text{OCN}^-$   $2170 \text{ cm}^{-1}$  band gives more accurate results on the  $k_1$  rate and is thus preferable.

We separate the  $\text{H}_2\text{O}$  molecules population into two kinds. The first kind,  $x(\text{H}_2\text{O}^{\text{reaction}})$ , are  $\text{H}_2\text{O}$  molecules that are making a favorable hydrogen bond with the  $\text{HNCO}$  nitrogen atom, in a configuration that the reaction is possible (Raunier et al. 2003b). The second kind,  $x(\text{H}_2\text{O}^{\text{catalysis}})$ , are  $\text{H}_2\text{O}$  molecules either far from  $\text{HNCO}$  or close to  $\text{HNCO}$ , but they do not form a hydrogen bond with the  $\text{HNCO}$  nitrogen atom. Thus the  $x(\text{H}_2\text{O}^{\text{catalysis}})$  molecules cannot react with  $\text{HNCO}$ , but they have a catalytic effect by decreasing the activation barrier of the proton transfer by long-range interactions (Raunier et al. 2003b). This population does not enter into the kinetic laws,



**Fig. 4.** Desorption rate curves from the integrated absorption bands of pure  $\text{HNCO}$  (full squares) and  $\text{HNCO}$  for a  $\text{HNCO}:\text{H}_2\text{O}$  mixture where  $\text{H}_2\text{O}$  is in excess (crosses) and where  $\text{H}_2\text{O}$  is not in excess (full circles). The dotted lines are the results from the fits using a first-order desorption rate (Eq. (3)). Inset:  $\text{OCN}^-$  (triangles) desorption curve in a  $\text{HNCO}:\text{H}_2\text{O}$  mixture where  $\text{H}_2\text{O}$  is not in a 1:5 concentration ratio. Only the first part of the desorption curve is fitted, corresponding to the desorption of  $\text{OCN}^-$  alone.

but is indirectly taken into account in the value of the activation barrier. This ice catalytic effect is shown by the fact that the  $\text{HNCO} + \text{H}_2\text{O}$  reaction does not occur at room temperature in the gas phase if we mix the two reactants in the same volume at a few mbars pressure. Thus we can write  $x(\text{H}_2\text{O}^{\text{reaction}}) = x(\text{HNCO})$ , since one  $\text{HNCO}$  molecule can react with only one  $\text{H}_2\text{O}^{\text{reaction}}$ . This is equivalent to write  $\frac{dx(\text{HNCO})}{dt} = k_1(T) \times x(\text{HNCO})^2$ . In the same way, because one molecule of  $\text{OCN}^-$  reacts one at one with one molecule of  $\text{H}_3\text{O}^+$ , which is making a favorable hydrogen bond with its oxygen atom, we can write  $x(\text{OCN}^-) = x(\text{H}_3\text{O}^+)$  and give an order 2 to this reaction.

Assuming such an order two for the two reactions, we can rewrite system 4 as

$$\begin{cases} \frac{dx(\text{HNCO})}{dt} = -k_1(T) \times x(\text{HNCO})^2 \\ \frac{dx(\text{OCN}^-)}{dt} = k_1(T) \times x(\text{HNCO})^2 - k_2(T) \times x(\text{OCN}^-)^2 \\ \frac{dx(\text{HO CN})}{dt} = k_2(T) \times x(\text{OCN}^-)^2. \end{cases} \quad (5)$$

We can then derive the analytical expressions for each species. Because  $\text{HO CN}$  originates in the  $\text{OCN}^-$  formed at  $t = 0$ ,  $k_1$  is not taken into account in the kinetics of  $x(\text{HO CN})$ , which is only determined by the  $k_2$  rate. The initial amount of  $\text{OCN}^-$  ( $t = 0$ ) does need to be considered for the expressions of  $x(\text{HNCO}(t))$  and  $x(\text{OCN}^-(t))$  because we are working on a difference spectrum, where the initial amount  $\text{OCN}^-(t = 0)$  does not appear. Moreover  $k_2$  does not need to be taken into account in the kinetics of  $x(\text{OCN}^-)$ . The contribution of the second term in the expression of the kinetics of  $x(\text{OCN}^-)$  is a small correction that can be neglected, because  $x(\text{OCN}^-)^2 \ll x(\text{HNCO})^2$ . Indeed, experimentally, the  $\text{OCN}^-$  band keeps growing along with time showing no influence of  $k_2$  as an intermediate species should, and its evolution is perfectly fitted taking only  $k_1$  into account. With these approximations we can write the analytical expressions of the species as


$$\begin{cases} x(\text{HNCO})(t) = \frac{x(\text{HNCO})(t=0)}{1 + x(\text{HNCO})(t=0) \times k_1 \times t} \\ x(\text{OCN}^-)(t) = \frac{x(\text{HNCO})(t=0)^2 \times k_1 \times t}{1 + x(\text{HNCO})(t=0) \times k_1 \times t} \\ x(\text{HO CN})(t) = \frac{x(\text{OCN}^-)(t=0) \times k_2 \times t}{1 + x(\text{OCN}^-)(t=0) \times k_2 \times t}. \end{cases} \quad (6)$$

Fitting the experimental data against these analytical expressions at a fixed temperature, as shown in Fig. 3, allows us to extract  $k_1$

**Table 2.** Experimental reaction rates  $k_1$  ( $\text{HNCO} + \text{H}_2\text{O} \rightarrow \text{OCN}^- + \text{H}_3\text{O}^+$  reaction) and  $k_2$  ( $\text{OCN}^- + \text{H}_3\text{O}^+ \rightarrow \text{HOCN} + \text{H}_2\text{O}$  reaction) as a function of temperature.

Reaction rates		
Temperature (K)	$k_1$ ( $\text{s}^{-1}$ )	$k_2$ ( $\text{s}^{-1}$ )
110	$0.20 \times 10^{-3}$	.
115	$0.59 \times 10^{-3}$	$2 \times 10^{-3}$
120	$0.12 \times 10^{-2}$	$9 \times 10^{-3}$
120	$0.15 \times 10^{-2}$	$2 \times 10^{-2}$
120	$0.26 \times 10^{-2}$	.
120	$0.13 \times 10^{-2}$	$2 \times 10^{-2}$
125	$0.65 \times 10^{-2}$	$4 \times 10^{-2}$
130	$0.13 \times 10^{-1}$	.
130	$0.16 \times 10^{-1}$	.
130	$0.11 \times 10^{-1}$	.

**Notes.** Not all experiments have enough  $\text{OCN}^-$  formed initially at the deposition to form an observable amount of  $\text{HOCN}$  during the experiment and measure  $k_2$ .



**Fig. 5.**  $\ln(k_1)$  (crosses) and  $\ln(k_2)$  (circles) as a function of the inverse of the temperature,  $k_1$  and  $k_2$  being the rates for the  $\text{HNCO} + \text{H}_2\text{O} \rightarrow \text{OCN}^- + \text{H}_3\text{O}^+$  and the  $\text{OCN}^- + \text{H}_3\text{O}^+ \rightarrow \text{HOCN} + \text{H}_2\text{O}$  reactions respectively. The uncertainty on a point is given by the dispersion of the measurements for the same temperature around the averaged value. The dashed lines represent the fit of the experimental points against an Arrhenius law.

and  $k_2$  for this temperature. We record the kinetics of the reaction between 110 K and 130 K as seen in Table 2. The rate at  $T = 130$  K is obtained adding a desorption term to system 4, since the molecules at the surface are desorbing while the molecules in the mantle are reacting. The accuracy on each reaction rate is estimated by reproducing individual identical experiments at both 120 K and 130 K. That our analytical expressions correctly fit the experimental data supports our assumptions on the partial orders.

We can now determine the temperature dependence of  $k_1(T)$  by fitting the reaction rates at different temperatures against an Arrhenius law  $k_1(T) = \nu_1 \times \exp\left(\frac{-E_1}{RT}\right)$ , as shown in Fig. 5.

We find an activation energy  $E_1 = 26 \pm 2 \text{ kJ mol}^{-1}$  and a pre-exponential factor  $\nu_1 = 3_{-2}^{+5} \times 10^8 \text{ s}^{-1}$ . This latter value is on the same order of magnitude than typical pre-exponential factors for second-order reactions in liquid solutions. However, the uncertainty on the derived pre-exponential factor is high. The value we derive for the activation energy is lower than the  $42 \text{ kJ mol}^{-1}$  calculated in Raunier et al. (2003b). This can be explained because this value has been derived for a  $\text{HNCO}$  molecule adsorbed on a  $\text{H}_2\text{O}$  surface. The experimentally derived value corresponds to bulk  $\text{H}_2\text{O}$  ice, which corresponds to double the solvation effect and thus lowers the barrier height even more. Given the uncertainties on the reaction rates, it is impossible to determine

a possible temperature dependence of the pre-exponential factor as in a modified Arrhenius law. Consequently we can write the reaction rate for the  $\text{HNCO} + \text{H}_2\text{O} \rightarrow \text{OCN}^- + \text{H}_3\text{O}^+$  reaction as  $k_1(T) = 3_{-2}^{+5} \times 10^8 \times \exp\left(\frac{-26 \pm 2 \text{ kJ mol}^{-1}}{RT}\right) \text{ s}^{-1}$ .

As for the  $\text{OCN}^- + \text{H}_3\text{O}^+ \rightarrow \text{HOCN} + \text{H}_2\text{O}$  reaction, we find an activation energy  $E_2 = 36 \pm 1 \text{ kJ mol}^{-1}$ , and a pre-exponential factor of  $\nu_2 = 4_{-2}^{+4} \times 10^{13} \text{ s}^{-1}$ . Accordingly we can write the reaction rate for the  $\text{OCN}^- + \text{H}_3\text{O}^+ \rightleftharpoons \text{HOCN} + \text{H}_2\text{O}$  reaction as  $k_2(T) = 4 \times 10^{13} \exp\left(\frac{-36 \pm 1 \text{ kJ mol}^{-1}}{RT}\right) \text{ s}^{-1}$ .

## 4. Discussion

We showed that the  $\text{HNCO} + \text{H}_2\text{O}$  acid-base reaction converts  $\text{HNCO}$  into the  $\text{H}_3\text{O}^+\text{OCN}^-$  species. This salt then transforms into  $\text{HOCN}$  and  $\text{H}_2\text{O}$ , the overall process being the isomerization of the isocyanic acid  $\text{HNCO}$  into cyanic acid  $\text{HOCN}$  within ice. We derive the expressions for the kinetic laws and reaction rates of formation and destruction of  $\text{OCN}^-$  from the  $\text{HNCO} + \text{H}_2\text{O}$  and  $\text{OCN}^- + \text{H}_3\text{O}^+$  reactions respectively. Whether or not this purely thermal mechanism is relevant to  $\text{OCN}^-$  formation in the interstellar environment is a question, first, of the kinetics of this reaction at a particular temperature and other competitive reactions such as photodissociations or hydrogenations, and second, of the physical environment the ice is located in. Indeed, in a protostellar environment, the grains are subjected to different temperatures, photon fluxes and density conditions depending on their location in the YSO, and  $\text{OCN}^-$  can be formed, desorbed, or destroyed accordingly. We will now look at the observed  $\text{OCN}^-$  abundances in different objects and then discuss the production of  $\text{OCN}^-$  from the  $\text{HNCO} + \text{H}_2\text{O}$  reaction, first in a dense molecular cloud and then in a YSO.

### 4.1. Observed $\text{OCN}^-$ abundances in the IR spectra of young stellar objects

Table 3 shows  $\text{OCN}^-$  abundances observed for several background stars and low-mass YSOs. We can see first that the  $\text{OCN}^-$  band is absent in the three background stars Elias 13, Elias 16, and EC 118 spectra (Knez et al. 2005). These objects give the column densities of solid-state material in dense molecular clouds before the star formation process begins. Thus, when the star starts collapsing, there is no  $\text{OCN}^-$ . On the other hand, all low-mass YSOs exhibit a  $\text{OCN}^-$  feature, with varying column densities and abundances with respect to  $\text{H}_2\text{O}$ , from one object to another. These differences in  $\text{OCN}^-$  abundances could be explained by (i) a difference of elapsed time, i.e. the older a YSO is, the more  $\text{OCN}^-$  is formed; (ii) a difference in environmental conditions (temperature, radiation flux, density). To test the first hypothesis, we can compare the  $\text{OCN}^-$  abundances of Class I objects with Class II or Class III objects. Within Class I objects, we can compare abundances for objects of a decreasing spectral index  $\alpha$ , which is defined as the derivative of the spectral energy distribution (SED) between 2 and  $14 \mu\text{m}$ . We do not observe any obvious trend between the  $\text{OCN}^-$  abundances and the class of an YSO or its spectral index. This might rule out the first hypothesis, and the variations of  $\text{OCN}^-$  abundances could be explained by a difference in environmental conditions: differences of masses, luminosities, of portion of the ice warmed. However, there are so few data that it is impossible to draw a reliable conclusion at that point.

**Table 3.** Column densities of  $\text{OCN}^-$  and  $\text{H}_2\text{O}$ , and abundance of  $\text{OCN}^-$  in background stars (top of the table) and low-mass YSO (bottom of the table).


Source	Spectral index <sup>a</sup>	Class	$N(\text{OCN}^-)^b$ $\times 10^{16} \text{ mol cm}^{-2}$	$N(\text{H}_2\text{O})^a$ $\times 10^{16} \text{ mol cm}^{-2}$	$[\text{OCN}^-]^c$ %
Elias 13	...	...	...	100	0
Elias 16	...	...	<2.3	250	0
EC 118 (CK2)	...	...	...	3.5	0
HH46	1.70	I	4.90(38)	779(77)	0.63(8)
HH100 IRS	0.8	I	0.80(4)	245(24)	0.33(8)
EC 90A	-0.09	I	0.20(12)	169(16)	0.12(5)
EC 90B	-0.09	I	1.12(24)	169(16)	0.86(12)
IRS 51	-0.15	I/II	1.46(26)	192	0.76(12)
RNO 91	0.03	II/III	1.74(20)	425(36)	0.41(6)

**Notes.** The values into parentheses are uncertainties. <sup>(a)</sup> Boogert et al. (2008); <sup>(b)</sup> van Broekhuizen et al. (2005); <sup>(c)</sup> as a percentage of the abundance of  $\text{H}_2\text{O}$ .

#### 4.2. Modeling $\text{OCN}^-$ abundances in a dense molecular cloud

$\text{HNCO}$ ,  $\text{H}_2\text{O}$ ,  $\text{OCN}^-$  and HOCN can be produced and destroyed in the gas and in the solid phase by different types of reactions, or go from one phase to the other by accretion and desorption. To take into account all these competitive reactions in a dense molecular cloud, we use a rate equations gas-grain model Hasegawa et al. (1992) along with the usual physical parameters used in this reference, hydrogen density  $N(\text{H}) = n(\text{H}) + n(\text{H}_2) = 2 \times 10^{-4} \text{ cm}^{-3}$ , dust to gas mass ratio 1%, density  $3 \text{ g cm}^{-3}$ , surface sites of adsorption  $10^6$ , a high visual extinction  $A_V = 500$ , and the initial gas-phase conditions with all hydrogen into its molecular form at the initial time. We use the most recent reactions database for the gas phase (OSU database) and the solid phase (Garrod et al. 2008). The  $\text{HNCO} + \text{H}_2\text{O}$  and  $\text{H}_3\text{O}^+ + \text{OCN}^-$  reactions we measured in this work are added to this reaction network with their measured kinetic parameters, along with the desorption energies of each species. The latter reaction is the only one in the reactions network that can produce  $\text{OCN}^-$  and for which kinetics parameters are known. Solid  $\text{HNCO}$  can be produced in the gas phase from various reactions and then be accreted on the grains, or it can be formed from surface reactions, essentially from the reaction  $\text{NH} + \text{CO}$ , which has no activation energy barrier according to Garrod et al. (2008) reaction network. Because we calculate a very low quantity of the  $\text{HNCO}$  parent molecule in the solid-state,  $1 \times 10^{-14} \text{ cm}^{-3}$  after  $10^5$  years, and because the  $\text{OCN}^-$  that can be converted from this quantity is even lower, we cannot expect  $\text{OCN}^-$  to be produced. Indeed, using the derived expression for the temperature dependence of the  $k_1$  reaction rate,  $k_1(T) = 3_{-2}^{+5} \times 10^8 \exp \frac{-26 \pm 2 \text{ kJ mol}^{-1}}{RT} \text{ s}^{-1}$ , we can calculate that  $1 \times 10^{-144} \text{ s}^{-1} \leq k_1(10 \text{ K}) \leq 3 \times 10^{-133} \text{ s}^{-1}$ , which makes this reaction impossible to occur at such a low temperature, which is typical of a molecular cloud. No  $\text{OCN}^-$  is produced in a dense molecular cloud, which fits with the observations of background stars.

We now investigate which reactions and which parameters govern the  $\text{OCN}^-$  production in a homogeneous object such as a molecular cloud. Figure 6 shows that the abundance of solid  $\text{HNCO}$  is decreasing after 100 years, which could prevent any  $\text{OCN}^-$  formation from  $\text{HNCO}$  at higher temperature. This decrease cannot be explained by evaporation, which is almost zero at 10 K, or by photodissociation reactions caused by the high visual extinction and as shown by the absence of any photoproduct. To investigate why no  $\text{OCN}^-$  is formed from the  $\text{HNCO} + \text{H}_2\text{O}$  reaction the and why the amount of  $\text{HNCO}$  decreases, at  $3 \times 10^5$  years, we arbitrarily impose the



**Fig. 6.** Time evolution of the abundances with respect to solid-state water of  $\text{HNCO}$  (solid line),  $\text{HNCHO}$  (dotted line) and  $\text{NH}_2\text{CHO}$  (dashed line) in a dense molecular cloud where solid  $\text{HNCO}$  density is arbitrarily fixed to be 5% of solid  $\text{H}_2\text{O}$  density at  $3 \times 10^5$  years (arrow). No  $\text{OCN}^-$  is produced from the  $\text{HNCO} + \text{H}_2\text{O}$  thermal reaction at 10 K and  $\text{OCN}^-$  density is zero all the time. Almost all  $\text{HNCO}$  is consumed in hydrogenation reactions toward  $\text{HNCHO}$  and  $\text{NH}_2\text{CHO}$ , whose increases in density are directly correlated to  $\text{HNCO}$  decrease. In addition to add 5% of  $\text{HNCO}$  compared to the solid  $\text{H}_2\text{O}$  density, H and  $\text{H}_2$  are reduced by 6 orders of magnitude (dot-dashed line): the  $\text{HNCO}$  abundance is only slightly decreasing. *Inset:* zoom of the evolution of the different abundances after  $t = 3 \times 10^5$  years.

solid  $\text{HNCO}$  density to be 5% of the solid  $\text{H}_2\text{O}$  density, which is on the order of the amount necessary to produce observed quantities of  $\text{OCN}^-$  (van Broekhuizen et al. 2004), and monitor which products this injected quantity is directed to. As seen in Fig. 6, where the arrow shows the arbitrary input of  $\text{HNCO}$  in the model, none of this quantity is directed into  $\text{OCN}^-$ . However, all abundances of the products from the reactions of  $\text{HNCO}$  with H atom or  $\text{NH}$ ,  $\text{NH}_2$ ,  $\text{OH}$ ,  $\text{CH}_3$ ,  $\text{CH}_3\text{O}$ ,  $\text{CH}_2\text{OH}$  or  $\text{HCO}$  radicals, are correlated to the sudden addition of solid  $\text{HNCO}$ . The two hydrogenation reactions  $\text{HNCO} + \text{H} \rightarrow \text{HNCHO}$  and  $\text{HNCO} + \text{H} \rightarrow \text{NH}_2\text{CO}$  are especially important, with rates on the order of  $10^{12} \text{ s}^{-1}$  (Garrod et al. 2008). Indeed, their two products  $\text{HNCHO}$  and  $\text{NH}_2\text{CO}$  are closely correlated with  $\text{HNCO}$ .

To confirm the predominance of the hydrogenation reactions, at  $3 \times 10^5$  years, in addition to impose the solid  $\text{HNCO}$  density to be 5% of the solid  $\text{H}_2\text{O}$  density, we arbitrarily decrease the quantity of H and  $\text{H}_2$  by 6 orders of magnitude. When we do this, solid  $\text{HNCO}$  is almost stable, slightly decreasing, and this slight

decrease is nevertheless directed into reactions with hydrogen, as shown in Fig. 6. We can therefore conclude that the hydrogenation reactions are dominating the disappearance of HNCO, and that the reactions of formation of OCN<sup>-</sup> must counterbalance them to explain the observed quantities of OCN<sup>-</sup>.

The HNCO+H<sub>2</sub>O reaction has an activation energy that is too high to compete with the hydrogenation reactions. However, the HNCO + NH<sub>3</sub> reaction can also produce OCN<sup>-</sup> and is known to have a very low activation energy (Raunier et al. 2003a). There are currently no experimental data on the kinetics of the HNCO:NH<sub>3</sub>:H<sub>2</sub>O system, so we will arbitrarily appoint a low activation energy to this reaction, between 0 and 5 kJ mol<sup>-1</sup>. With 0 kJ mol<sup>-1</sup> it is possible to produce OCN<sup>-</sup> at 10 K from the reaction with NH<sub>3</sub>, although with a very low abundance, as is shown by the correlated formation of NH<sub>4</sub><sup>+</sup>. It is impossible to produce OCN<sup>-</sup> with a 5 kJ mol<sup>-1</sup> activation energy at 10 K, but it is possible at 40 K. This shows the importance of measuring the value of the HNCO + NH<sub>3</sub> reaction activation energy and to experimentally investigate the HNCO hydrogenation reactions.

### 4.3. Modeling OCN<sup>-</sup> abundances in a young stellar object

We then investigated if there is in an YSO a particular region or time where OCN<sup>-</sup> can be formed from the reaction HNCO+H<sub>2</sub>O. Using the measured expression for  $k_1(T)$  at  $T = 80$  K, which is close to the limit temperature to encounter water ice and HNCO on grains, we have  $5 \times 10^{-11} \leq k_1(80 \text{ K}) \leq 2 \times 10^{-7} \text{ s}^{-1}$ , with an average reaction rate of  $k_1(80 \text{ K}) \approx 0.1 \text{ yr}^{-1}$ , which may be enough to counterbalance hydrogenation reactions. Indeed, if the surface temperature is warmer, the residence time of H on grains is shorter and hydrogenation reactions may not be dominant anymore. For this investigation, we choose to use a very simple disk model (Aikawa & Herbst 1999). This model corresponds to a class I/II object and is stationary. We define a grid along the  $R$  radial direction and  $Z$  transverse direction of the disk. For each point of the grid, the code is run for  $10^6$  years, which is the extreme limit of validity of this model. The initial conditions for all molecules except solid HNCO are fixed by running the model from the diffuse medium to the dense molecular cloud medium for  $3 \times 10^5$  years. At the initial time there is no OCN<sup>-</sup> as discussed in the previous paragraph. The abundance of solid HNCO is fixed to be 5% of the abundance of H<sub>2</sub>O, which is the quantity necessary to reproduce observed OCN<sup>-</sup> abundances (van Broekhuizen et al. 2005) using thermal reactions. After  $10^6$  years, OCN<sup>-</sup> cannot be formed from the HNCO + H<sub>2</sub>O reaction alone. Once again HNCO is directed into its hydrogenation products. This disagrees with existing solid OCN<sup>-</sup> observations. This means that alternative reactions with very low activation energies, such as the HNCO + NH<sub>3</sub> reaction, can counterbalance the hydrogenation reactions and produce OCN<sup>-</sup> from HNCO, or that OCN<sup>-</sup> is produced from another route than from HNCO. Additional experimental work is needed on both the HNCO + NH<sub>3</sub> reaction and the HNCO hydrogenation reactions to be able to discuss quantitatively the different ways of OCN<sup>-</sup> formation.

## 5. Conclusion

We studied the kinetics of the conversion of isocyanic acid HNCO into OCN<sup>-</sup> and then of OCN<sup>-</sup> into cyanic acid HOCN

in a H<sub>2</sub>O environment. The H<sub>2</sub>O + HNCO reaction gives the H<sub>3</sub>O<sup>+</sup>OCN<sup>-</sup> species with a kinetic rate  $k_1(T) = \nu_1 \times \exp(-E_a/RT)$ ,  $\nu_1 = 3^{+5}_2 \times 10^8 \text{ s}^{-1}$ ,  $E_1 = 26 \pm 2 \text{ kJ mol}^{-1}$  (3127 K). The H<sub>3</sub>O<sup>+</sup> to OCN<sup>-</sup> reaction has a kinetic rate  $k_2(T) = \nu_2 \times \exp(-E_2/RT)$ ,  $\nu_2 = 4^{+4}_2 \times 10^{-13} \text{ s}^{-1}$ ,  $E_2 = 36 \pm 1 \text{ kJ mol}^{-1}$  (4330 K). We show that OCN<sup>-</sup> follows a kinetic law that can be written as  $dx(\text{OCN}^-)/dt = k_1 \times x(\text{HNCO})^2 - k_2 \times x(\text{OCN}^-)^2$ . The desorption energies of HNCO and OCN<sup>-</sup> are measured to be  $E_{\text{des}}(\text{HNCO}) = 37 \pm 3 \text{ kJ mol}^{-1}$  (4450 K) and  $E_{\text{des}}(\text{OCN}^-) = 40 \pm 3 \text{ kJ mol}^{-1}$  (4811 K), respectively. An attempt to incorporate the derived kinetics laws into an existing gas-grain model was unsuccessful in explaining the observed abundances of OCN<sup>-</sup> in YSO ices. The model shows that hydrogenation reactions present in reactions databases are dominating the chemistry of HNCO, preventing the formation of OCN<sup>-</sup> from the HNCO + H<sub>2</sub>O reaction only. Experimental data on the hydrogenation of HNCO are therefore needed. Moreover, other reactions may be responsible of the OCN<sup>-</sup> formation, such as the HNCO + NH<sub>3</sub> reaction. We need similar kinetics measurements on these reactions to compare them with hydrogenation reactions.

*Acknowledgements.* This work has been funded by the French national program Physique Chimie du Milieu Interstellaire (PCMI) and the Centre National d'Études Spatiales (CNES). P. Theulé particularly thanks the Academia Sinica Institute for Astronomy and Astrophysics for hosting him during several stays.

## References

- Aikawa, Y., & Herbst, E. 1999, A&A, 351, 233  
 Bennett, C. J., Jones, B., Knox, E., et al. 2010, ApJ, 723, 641  
 Boogert, A. C. A., Pontoppidan, K. M., Knez, C., et al. 2008, ApJ, 678, 985  
 Bossa, J. B., Theule, P., Duvernay, F., & Chiavassa, T. 2009, ApJ, 707, 1524  
 Brünken, S., Gottlieb, C. A., McCarthy, M. C., & Thaddeus, P. 2009, ApJ, 697, 880  
 Demyk, K., Dartois, E., d'Hendecourt, L. B., et al. 1998, A&A, 339, 353  
 d'Hendecourt, L. B., & Allamandola, L. J. 1986, A&AS, 64, 453  
 d'Hendecourt, L. B., Allamandola, L. J., Grim, R. J. A., & Greenberg, J. M. 1986, A&A, 158, 119  
 Falk, M., & Guigere, P. A. 1957, Can. J. Chem, 35, 1195  
 Garrod, R. T., Weaver, S. L. W., & Herbst, E. 2008, ApJ, 682, 283  
 Grim, R. J. A., & Greenberg, J. M. 1987, ApJ, 321, L91  
 Grim, R. J. A., Greenberg, J. M., de Groot, M. S., et al. 1989, A&AS, 78, 161  
 Hasegawa, T., Herbst, E., & Leung, C. M. 1992, A&AS, 82, 167  
 Iglesias, E. 1977, ApJ, 218, 697  
 Knez, C., Boogert, A. C. A., Pontoppidan, K. M., et al. 2005, ApJ, 635, L145  
 Lowenthal, M. S., Khanna, R. K., & Moore, M. H. 2000, BAAS, 32, 1108  
 Marcelino, N., Cernicharo, J., Tercero, B., & Roueff, E. 2009, ApJ, 690, L27  
 Mladenović, M., & Lewerenz, M. 2008, Chem. Phys., 343, 129  
 Moore, M. H., Donn, B., Khanna, R., & A'Hearn, M. F. 1983, Icarus, 54, 388  
 Novozamsky, J. H., Schutte, W. A., & Keane, J. V. 2001, A&A, 379, 588  
 OSU astrochemistry database, network 2009  
 Raunier, S., Chiavassa, T., Marinelli, F., Allouche, A., & Aycard, J.-P. 2003a, Chem. Phys. Lett., 368, 594  
 Raunier, S. S., Chiavassa, T., Allouche, A., Marinelli, F., & Aycard, J.-P. 2003b, Chem. Phys., 288, 197  
 Raunier, S., Chiavassa, T., Marinelli, F., & Aycard, J.-P. 2004, Chem. Phys., 302, 259  
 Snyder, L. E., & Buhl, D. 1972, ApJ, 177, 619  
 Teles, J. H., Maier, G., Hess, B. A., et al. 1989, Chemische Berichte, 122, 753  
 Tideswell, D. M., Fuller, G. A., Millar, T. J., & Markwick, A. J. 2010, A&A, 510, A85  
 Turner, B. E., Terzieva, R., & Herbst, E. 1999, ApJ, 518, 699  
 van Broekhuizen, F. A., Keane, J. V., & Schutte, W. A. 2004, A&A, 415, 425  
 van Broekhuizen, F. A., Pontoppidan, K. M., Fraser, H. J., & van Dishoeck, E. F. 2005, A&A, 441, 249  
 Zinchenko, I., Henkel, C., & Mao, R. Q. 2000, A&A, 361, 1079