Amplitudes and lifetimes of solar-like oscillations observed by CoRoT

Red-giant versus main-sequence stars

(Corrigendum)

F. Baudin¹, C. Barban², K. Belkacem^{3,2}, S. Hekker^{4,5,6}, T. Morel³, R. Samadi², O. Benomar¹, M.-J. Goupil²,
 F. Carrier⁴, J. Ballot⁷, S. Deheuvels², J. De Ridder⁴, A. P. Hatzes⁸, T. Kallinger^{9,10}, and W. W. Weiss⁹

¹ Institut d'Astrophysique Spatiale, CNRS, Université Paris XI, 91405 Orsay Cedex, France
 e-mail: frederic.baudin@ias.u-psud.fr

² LESIA, Université Pierre et Marie Curie, Université Denis Diderot, Obs. de Paris, 92195 Meudon Cedex, France

³ Institut d'Astrophysique et de Géophysique, Université de Liège, Allée du 6 Août 17, 4000 Liège, Belgium

⁴ Instituut voor Sterrenkunde, KU Leuven, Celestijnenlaan 200D, 3001 Leuven, Belgium

⁵ School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

⁶ Astronomical Institute "Anton Pannekoek", University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

⁷ Institut de Recherche en Astrophysique et Planétologie, CNRS, Université de Toulouse, 31400 Toulouse, France

⁸ Thüringer Landessternwarte, 07778 Tautenburg, Germany

⁹ University of Vienna, Institute for Astronomy, Türkenschanzstrasse 17, 1180 Vienna, Austria

¹⁰ Department of Physics and Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, BC V6T 1Z1, Canada

A&A, 529, A84 (2011), DOI: [10.1051/0004-6361/201014037](https://doi.org/10.1051/0004-6361/201014037)

Key words. stars: oscillations – methods: data analysis – asteroseismology – errata, addenda

In Sects. 4.3 and 4.4, uncertainties were not properly taken into account in the computation of the scaling relation between mode widths (Sect. 4.3) and effective temperature for main-sequence stars (MS), and of the scaling relation between mode amplitude and effective temperature, again for main-sequence stars (other scaling relations are correct). The correct results are

$$\Gamma \propto T_{\text{eff}}^s \quad \text{with} \quad \begin{cases} s \approx -0.3 \pm 0.9 & \text{for RG (unchanged)} \\ s \approx 16 \pm 2 & \text{for MS (corrected)} \end{cases}$$

and:

$$A_{\text{max}}^I \propto \left(\frac{L}{M}\right)^s \quad \text{with} \quad \begin{cases} s = 0.92 \pm 0.14 & \text{for RG (unchanged)} \\ s = 0.42 \pm 0.14 & \text{for MS (corrected)}. \end{cases}$$

The corrected values are only slightly changed and well inside former uncertainties. Moreover, the corrected uncertainties are smaller than the former ones. Figure 7 is also slightly changed (see Fig. 1). The conclusions remain unchanged with these more precise results.

Acknowledgements. This wrong computation was pointed out by T. Appourchaux who performed a thorough check of the former published results.

Fig. 1. Measured mode linewidths versus T_{eff} for the red giants ($T_{\text{eff}} < 5000$ K) and for the main-sequence stars ($T_{\text{eff}} > 5000$ K). The dot-dash line indicates the fit to the red-giant width and the triple-dot-dash line the fit to main-sequence stars.