

# The *Herschel*<sup>\*</sup> view of the on-going star formation in the Vela-C molecular cloud

T. Giannini<sup>1</sup>, D. Elia<sup>2</sup>, D. Lorenzetti<sup>1</sup>, S. Molinari<sup>2</sup>, F. Motte<sup>2</sup>, E. Schisano<sup>2</sup>, S. Pezzuto<sup>2</sup>, M. Pestalozzi<sup>2</sup>,  
A. M. Di Giorgio<sup>2</sup>, P. André<sup>3</sup>, T. Hill<sup>3</sup>, M. Benedettini<sup>2</sup>, S. Bontemps<sup>4</sup>, J. Di Francesco<sup>5,6</sup>,  
C. Fallscheer<sup>5,6</sup>, M. Hennemann<sup>3</sup>, J. Kirk<sup>7</sup>, V. Minier<sup>3</sup>, Q. Nguyễn Lương<sup>8</sup>, D. Polychroni<sup>2</sup>, K. L. J. Rygl<sup>2</sup>,  
P. Saraceno<sup>2</sup>, N. Schneider<sup>3</sup>, L. Spinoglio<sup>2</sup>, L. Testi<sup>9</sup>, D. Ward-Thompson<sup>10</sup>, and G. J. White<sup>11,12</sup>

<sup>1</sup> INAF – Osservatorio Astronomico di Roma, via Frascati 33, 00040 Monte Porzio Catone, Italy  
e-mail: giannini@mporzio.astro.it

<sup>2</sup> INAF – Istituto di Astrofisica e Planetologia Spaziale, via Fosso del Cavaliere 100, 00133 Rome, Italy

<sup>3</sup> Laboratoire AIM, CEA/IRFU CNRS/INSU Université Paris Diderot, CEA-Saclay, 91191 Gif-sur-Yvette Cedex, France

<sup>4</sup> Université de Bordeaux, OASU, Bordeaux, France

<sup>5</sup> University of Victoria, Department of Physics and Astronomy, PO Box 3055, STN CSC, Victoria, BC, V8W 3P6, Canada

<sup>6</sup> National Research Council of Canada, Herzberg Institute of Astrophysics, 5071 West Saanich Rd., Victoria, BC, V9E 2E7, Canada

<sup>7</sup> School of Physics and Astronomy, Cardiff University, Queens Buildings, The Parade, Cardiff, CF243AA, UK

<sup>8</sup> Laboratoire AIM Paris-Saclay, CEA/IRFU – CNRS/INSU – Université Paris Diderot, Service d'Astrophysique, Bât. 709, CEA-Saclay, 91191 Gif-sur-Yvette Cedex, France

<sup>9</sup> ESO, Karl Schwarzschild str. 2, 85748 Garching bei Munchen, Germany

<sup>10</sup> School of Physics and Astronomy, Cardiff University, Queens Buildings, The Parade, Cardiff, CF243AA, UK

<sup>11</sup> Department of Physics and Astronomy, The Open University, Milton Keynes, UK

<sup>12</sup> Department of Physics and Astronomy, McMaster University, Hamilton, ON, L8S 4M1, Canada

Received 2 August 2011 / Accepted 30 January 2012

## ABSTRACT

**Aims.** As part of the *Herschel* guaranteed time key programme “HOBYS”, we present the PACS and SPIRE photometric survey of the star-forming region Vela-C, one of the nearest sites of low-to-high-mass star formation in the Galactic plane. Our main objectives are to take a census of the cold sources and to derive their mass distribution down to a few solar masses.

**Methods.** Vela-C was observed with PACS and SPIRE in parallel mode at five wavelengths between 70  $\mu\text{m}$  and 500  $\mu\text{m}$  over an area of about 3 square degrees. A photometric catalogue was extracted from the detections in each of the five bands, using a threshold of  $5\sigma$  over the local background. Out of this catalogue we selected a robust sub-sample of 268 sources, of which  $\sim 75\%$  are cloud clumps (diameter between 0.05 pc and 0.13 pc) and 25% are cores (diameter between 0.025 pc and 0.05 pc). Their spectral energy distributions (SEDs) were fitted with a modified black body function. We classify 48 sources as protostellar, based on their detection at 70  $\mu\text{m}$  or at shorter wavelengths, and 218 as starless, because of non-detections at 70  $\mu\text{m}$ . For two other sources, we do not provide a secure classification, but suggest they are Class 0 protostars.

**Results.** From the SED fitting we derived key physical parameters (i.e. mass, temperature, bolometric luminosity). Protostellar sources are in general warmer ( $\langle T \rangle = 12.8$  K) and more compact ( $\langle \text{diameter} \rangle = 0.040$  pc) than starless sources ( $\langle T \rangle = 10.3$  K,  $\langle \text{diameter} \rangle = 0.067$  pc). Both these findings can be ascribed to the presence of an internal source(s) of moderate heating, which also causes a temperature gradient and hence a more peaked intensity distribution. Moreover, the reduced dimensions of protostellar sources may indicate that they will not fragment further. A virial analysis of the starless sources gives an upper limit of 90% probability for the sources to be gravitationally bound and therefore prestellar in nature. A luminosity vs. mass diagram of the two populations shows that protostellar sources are in the early accretion phase, while prestellar sources populate a region of the diagram where mass accretion has not started yet. We fitted a power law  $N(\log M) \propto M^{-1.1 \pm 0.2}$  to the linear portion of the mass distribution of prestellar sources. This is in between that typical of CO clumps and those of cores in nearby star-forming regions. We interpret this as a result of the inhomogeneity of our sample, which is composed of comparable fractions of clumps and cores.

**Key words.** ISM: individual objects: Vela-C – ISM: clouds – stars: formation – submillimeter: ISM – circumstellar matter


## 1. Introduction

Most of the stellar content in our Galaxy forms in cold ( $T \sim 10$ – $30$  K) and dense ( $n > 10^3$   $\text{cm}^{-3}$ ) cores within the giant molecular clouds (GMC) of the Galactic disc (e.g., Blitz 1991). The main properties of the stellar population, such as the efficiency

of the mass conversion into stars and the shape of the initial mass function (IMF) may very well be closely related to the physical properties and mass distribution of the progenitor structures in the parental cloud (e.g. Mc Kee & Ostriker 2007).

Being one of the nearest GMCs in the Galactic disc, the Vela molecular ridge (VMR, Murphy & May 1991,  $l \approx 260^\circ$ – $275^\circ$ ,  $b \approx \pm 3^\circ$ ) represents an ideal observational target. It is composed of four molecular clouds at a distance between 700 pc (clouds A, C, D) and 2000 pc (cloud B, Liseau et al. 1992),

\* *Herschel* is an ESA space observatory with science instruments provided by European-led Principal Investigator consortia and with important participation from NASA.



**Fig. 1.** Composite 3-colour image of Vela-C: PACS  $70\ \mu\text{m}$  (blue), PACS  $160\ \mu\text{m}$  (green), and SPIRE  $500\ \mu\text{m}$  (red). The blueish regions near the centre and towards the bottom right of the map are the HII regions RCW 36 and RCW 34, respectively. Red and blue diamonds represent the locations of prestellar and protostellar sources, respectively, belonging to the sample of 268 objects used for the SED analysis. Bigger diamonds indicate sources with  $M \geq 20 M_{\odot}$ , while yellow diamonds indicate a few sources inside RCW 34 (see text). The top left segment corresponds to a length of 5 pc in the sky for a distance to Vela-C of 700 pc.

and harbours protostars with masses up to  $10 M_{\odot}$ , both isolated and clustered (Massi et al. 2000, 2003). Vela-C is the most massive component and hosts the youngest stellar population (Yamaguchi et al. 1999); this latter has been investigated by combining near-infrared data ( $J$ ,  $H$ ,  $K$ -band images) with far-infrared surveys (MSX, IRAS), and around thirty isolated protostars and seven embedded young clusters associated with  $\text{C}^{18}\text{O}$  clumps have been found (Liseau et al. 1992; Lorenzetti et al. 1993; Massi et al. 2003; Baba et al. 2006). A bright HII region, RCW 36, has been associated by Massi et al. (2003) with an early-type star (spectral type O5-B0). A second HII region, RCW 34, was originally associated to Vela-C by several authors (Herbst 1975; Murphy & May 1991), but recent observations favour a much longer distance ( $d = 2.5$  kpc, Bik et al. 2010). Vela-C has been imaged at  $250\ \mu\text{m}$ ,  $350\ \mu\text{m}$  and  $500\ \mu\text{m}$  with the Balloon-borne Large Aperture Submillimeter Telescope (BLAST, Pascale et al. 2008) that provided the first census of the compact dust emission in a range of evolutionary stages and lifetimes (Netterfield et al. 2009).

As a target of the *Herschel* guaranteed time key programme HOBYS (“*Herschel* imaging survey of OB Young Stellar objects”, Motte et al. 2010), Vela-C has been observed with the PACS (Poglitsch et al. 2010) and SPIRE (Griffin et al. 2010) cameras between  $70\ \mu\text{m}$  and  $500\ \mu\text{m}$ . The extended emission in the form of filaments and ridges has been presented by Hill et al. (2011), who identified five sub-regions with different column densities. Here we will focus on the determination of the physical parameters of the compact sources, their evolutionary stage and their mass distribution. The latter has been investigated in several star-forming regions through different tracers (e.g. Testi & Sargent 1998; Motte et al. 1998, 2001; Kramer et al. 1998; Reid & Wilson 2006; Enoch et al. 2008; Rathborne et al. 2009; Könyves et al. 2010; Ikeda & Kitamura 2011). In this paper, we intend to enlarge the statistics by providing the mass distribution over a wide range, going from subsolar values up to tens of solar masses.

This paper is organised as follows. In Sect. 2 we describe the observations, the data reduction procedures and the photometric results. In Sect. 3 we describe the spectral energy distribution (SED) analysis along with the fitted physical parameters. In


Sect. 4 we discuss the evolutionary stages of the sources along with their mass distribution. Our results are then summarised in Sect. 5.

## 2. Observations, data reduction and results

Vela-C was observed on May 18, 2010 with the *Herschel* parallel mode, i.e., using simultaneously PACS at  $70/160\ \mu\text{m}$  and SPIRE at  $250/350/500\ \mu\text{m}$ . A common area of  $\sim 3$  square degrees was covered by both instruments around the position  $\alpha_{(J2000.0)} = 08^{\text{h}}59^{\text{m}}55^{\text{s}}$ ,  $\delta_{(J2000.0)} = -43^{\circ}53'00''$ . The field was observed in two orthogonal directions at the scan speed of  $20''/\text{s}$ . The data reduction strategy is described in detail in Traficante et al. (2011): here we summarise only the fundamental steps. From archival data to the Level 1 stage, we used scripts prepared in the *Herschel* interactive processing environment (HIPE, Ott 2010), partially customised compared to the standard pipeline. The obtained time-ordered data (TODs) of each bolometric detector were then processed further by means of dedicated IDL routines and finally maps were created using the FORTRAN code ROMAGAL. Images of Vela-C at all wavelengths have been presented by Hill et al. (2011), here we show in Fig. 1 the composite three-colour image at  $70/160/500\ \mu\text{m}$  for the readers’ convenience.

### 2.1. Source detection and photometry

The detection and photometry of compact sources were carried out using the Curvature Threshold Extractor package (CuTeX, Molinari et al. 2011) applying the same strategy as Molinari et al. (2010). Briefly, in each band, sources were identified as peaks in the second-derivative images of the original HOBYS maps, then an elliptical Gaussian fit was performed to provide 1) the total flux, integrated down to the zero intensity level, 2) the observed full width at half maximum (FWHM, defined as the geometrical mean of the major and minor ellipse axes), and 3) the peak intensity. The latter, when divided by the local rms noise, allows us to obtain an *a posteriori* estimate of the signal-to-noise ratio (S/N). We filtered out all sources with  $S/N < 5$ . Following Elia et al. (2010), entries at different wavelengths in the PACS/SPIRE



**Fig. 2.** Percentage of recovered artificial sources as a function of their fluxes in the five *Herschel* bands.

catalogue were attributed to the same source based on simple positional criteria. In practice, we associated two sources detected in two different bands if their mutual angular distance does not exceed the radius of the *Herschel* half-power beam-width (HPBW<sup>1</sup>) at the longer wavelength. Around 15% of the entries in the catalogue present multiple associations at decreasing wavelength; as a general rule, the closest counterpart was kept without attempting to divide the flux at the longer wavelength among the sources. We assigned to each source the celestial coordinates of the counterpart at the shortest wavelength, where the spatial resolution is higher. Finally, to exclude artefacts, we discarded sources with an axes ratio >2 and a position angle randomly changing with wavelength by more than 20°.

## 2.2. Catalogue statistics

The complete *Herschel* catalogue of Vela-C will be published in the near future as part of the HOBYS data products. Here we limit ourselves to give a statistical summary of the detected sources in Table 1. In the second column we list the number of entries in each of the five bands: this number increases with increasing wavelength up to  $\lambda = 250 \mu\text{m}$ , then decreases at longer wavelengths. This behaviour can be explained in the light of the sensitivity limits given in the third column: since the number of entries in the first three bands increases despite the loss of sensitivity, this increasing is a real effect reflecting the intrinsic properties of the cloud. Conversely, in the last two SPIRE bands, the poorer sensitivity limits, together with the loss of angular resolution, are likely responsible for the reduced number of the detected sources.


The flux completeness limits (fourth column) are estimated at the 90% of confidence level, estimated by recovering artificial Gaussian sources randomly spread over the map, whose fluxes and diameters cover the same ranges found for the observed sources. In Fig. 2, we show the percentage of the recovered artificial sources as a function of the flux for each *Herschel* band. For comparison, the BLAST completeness limits at 250  $\mu\text{m}$  and 350  $\mu\text{m}$  (estimated in Fig. 3 of Netterfield et al. 2009) are more than a factor of 15 larger than the SPIRE limits at the same wavelengths.

<sup>1</sup> The values of the *Herschel* HPBW are 5.0'' at 70  $\mu\text{m}$ , 11.4'' at 160  $\mu\text{m}$ , 17.8'' at 250  $\mu\text{m}$ , 25.0'' at 350  $\mu\text{m}$ , and 35.7'' at 500  $\mu\text{m}$ .

**Table 1.** Statistics of the 5  $\sigma$  PACS/SPIRE catalogue.

Band	#	Sensitivity limit (Jy)	90% Completeness limit <sup>a</sup> (Jy)
Total entries <sup>b</sup>	1686		–
70 $\mu\text{m}$	658	0.04	0.21
160 $\mu\text{m}$	871	0.09	0.67
250 $\mu\text{m}$	966	0.11	1.05 (17)
350 $\mu\text{m}$	697	0.35	1.32 (22)
500 $\mu\text{m}$	416	0.46	1.95

**Notes.** (a) For comparison we report in parentheses the BLAST completeness limit estimated from Netterfield et al. (2009) (their Fig. 3). (b) Number of sources detected in at least one band.



**Fig. 3.** Examples of modified black body fits to the SEDs of two sources (undetected and detected at 70  $\mu\text{m}$ , respectively). Crosses and dots mark the data points before and after the correction applied at longer wavelengths to minimize the effects of the poorer spatial resolution (see text). Values of the main fitted parameters are reported.

## 2.3. Source diameters

In each of the five bands, the physical  $\text{FWHM}_{\text{dec}}(\lambda)$  was derived by deconvolving the observed FWHM with the HPBW at the same wavelength:

$$\text{FWHM}_{\text{dec}}(\lambda) = \sqrt{\text{FWHM}_{\lambda}^2 - \text{HPBW}_{\lambda}^2}. \quad (1)$$

Noticeably, the  $\text{FWHM}_{\text{dec}}(\lambda)$  increases with  $\lambda$ , with mean values of 7.5'', 13.4'', 22.7'', 28.6'', 41.0'' in the five *Herschel* bands, respectively. Since we are most interested in tracing the cold dust emission (namely that at temperature  $\lesssim 25$  K), which is likely poorly related to the 70  $\mu\text{m}$  flux, we define as angular source diameter ( $\theta$ ) the  $\text{FWHM}_{\text{dec}}$  measured at 160  $\mu\text{m}$ , where we detect a considerable number of sources with a good angular resolution (see also Motte et al. 2010). For a small fraction of sources that remained undetected at 160  $\mu\text{m}$ ,  $\theta$  is the  $\text{FWHM}_{\text{dec}}$  measured at 250  $\mu\text{m}$ .

We consider as spatially resolved the detections with  $\theta$  larger than  $\sim 60\%$  of the angular resolution at 160  $\mu\text{m}$  (corresponding to a diameter  $D \sim 0.025$  pc at the Vela-C distance), as also adopted in the Aquila Rift and Polaris by André et al. (2010). Noticeably, around  $\sim 25\%$  of our sources have a diameter between 0.025 pc and 0.05 pc (“cores”), while most sources are larger (“clumps”,  $0.5 \leq D \leq 0.13$  pc); in the following analysis, we will not distinguish any longer between the two categories, and will refer to both of them with the general term “source”.



### 3. Analysis

#### 3.1. Flux scaling

The physical parameters of the detected sources were derived by fitting their SEDs with a modified black body curve. Two examples of the observed SEDs are shown in Fig. 3. These illustrate a trend seen in many of the SEDs, namely that the *observed* fluxes (depicted with crosses) flatten or even rise longwards  $\lambda \geq 250 \mu\text{m}$ . This effect is a direct consequence of the increase of the  $\text{FWHM}_{\text{dec}}(\lambda)$  with wavelength (Sect. 2.2), which in turn implies increasing areas over which the emission is integrated. In practice, this flattening indicates that a single-temperature, modified black body fit is not completely adequate to model the observed photometric points. However, since the small amount of available data prevents us from using a multiple-temperature, modified black body model, we restricted the fit only to the innermost portions of the sources, whose spatial scales are defined by the source angular diameter  $\theta$ . To estimate the emission coming from these restricted solid angles, we followed the flux scaling method adopted by Motte et al. (2010) that is described in detail by Nguyễn Lương et al. (2011). This method is based on the idea that for quasi-spherical, self-gravitating sources, the radial density law is  $\propto r^{-2}$  (with  $r \sim 0.1\text{--}1$  pc), and thus  $M(<r) \propto r$  (Shu 1977). This relationship implies that fluxes scale linearly with the source radius, provided that the emission is optically thin and the temperature gradient is weak ( $T(r) \propto r^{-q}$ , with  $q \sim 0$ , see Motte & André 2001 for more details). All these conditions have been empirically verified in our case by taking the photometry at different apertures for some isolated sources, and finding a linear increase of fluxes with increasing radii. Hence, fluxes at  $250 \mu\text{m}$ ,  $350 \mu\text{m}$ , and  $500 \mu\text{m}$ , were reduced by the ratio  $\text{FWHM}_{\text{dec}}(\lambda)/\theta$ . The average values of these factors are 1.9 at  $250 \mu\text{m}$ , 2.0 at  $350 \mu\text{m}$ , and 3.2 at  $500 \mu\text{m}$ , respectively. In Fig. 3, dots represent fluxes after the above correction.

#### 3.2. Selection criteria

Once the SPIRE fluxes were rescaled, we further filtered the catalogue to select a robust sub-sample for SED analysis. The final list was composed of sources fulfilling the following criteria, i.e., sources

- (1) with detections in at least three adjacent bands between  $160 \mu\text{m}$  and  $500 \mu\text{m}$ , to contain the peak of the emission of the cold dust;
- (2) without a dip in the SED between three adjacent wavelengths;
- (3) not peaking at  $500 \mu\text{m}$ . Note that conditions (2) and (3) exclude most of the artefacts or detections affected by incorrect positional associations or bad photometry;
- (4) spatially resolved at  $160 \mu\text{m}$ ;
- (5) not presenting multiple associations at  $\lambda \geq 160 \mu\text{m}$ , to alleviate confusion problems;
- (6) not belonging to the RCW34 region, to exclude sources that could be more distant than  $700$  pc (see Sect. 1). These sources, identified by eye in the *Herschel* maps, are depicted in yellow in Fig. 1.

The first three criteria are the most stringent ones, since they lead to a selection of only 388 sources out of 1686 present in the catalogue. Applying criteria (4) to (6), we obtained a final sample of 268 sources to be fitted. Their locations are depicted in Fig. 1 with different colours indicating their evolutionary stage (see below).

#### 3.3. SED fitting

Since cold dust emission is likely unrelated to the photometry measured at  $70 \mu\text{m}$ , we fitted the SEDs longward  $\lambda \geq 160 \mu\text{m}$  and checked a posteriori whether or not the modified black body curve also fits the  $70 \mu\text{m}$  point. The assumed relation is

$$F_\nu = (1 - e^{-\tau_\nu}) B_\nu(T_d) \Omega, \quad (2)$$

where  $F_\nu$  is the observed flux at the frequency  $\nu$ ,  $\Omega$  is the source area (in sr),  $B_\nu(T_d)$  is the black body function at the dust temperature  $T_d$ , and  $\tau_\nu$  is the optical depth, parametrised as  $\tau_\nu = (\nu/\nu_0)^\beta$ , where  $\nu_0 = c/\lambda_0$  is the frequency at which  $\tau = 1$ , and  $\beta = 2$ , as predicted by simple dust emission models (Hildebrand 1983). A linear least-squares fit to the scaled fluxes (with their photometric uncertainties) was performed by comparing the observations with a database of models where  $T_d$  and  $\lambda_0$  are allowed to vary in the following ranges:  $8 \text{ K} \leq T_d \leq 40 \text{ K}$ ;  $10 \mu\text{m} \leq \lambda_0 \leq 500 \mu\text{m}$ . We applied the PACS colour corrections, amounting to a maximum of 4% of the flux, to the model fluxes before comparing them with the data (Müller et al 2011). As a check, we also verified that the fitted values of  $\lambda_0$  never exceed  $160 \mu\text{m}$  (see Sect. 4.1 and Fig. 5); this ensures that the emission is optically thin in the SPIRE wavelength range, namely where flux scaling is performed (see the discussion in Sect. 3.1).

Finally, masses were derived from the dust emission using the relation

$$M = (d^2 \Omega / k_{300}) \tau_{300}, \quad (3)$$

where  $k_{300} = 0.1 \text{ cm}^2 \text{ g}^{-1}$  is the opacity per unit gas mass computed at  $\lambda = 300 \mu\text{m}$  assuming a gas-to-dust ratio of 100 (Beckwith et al. 1990), and  $d = 700$  pc. (The details of the derivation of Eq. (3) from (2) will be given in Pezzuto et al., in prep.).

#### 3.4. Separating starless from protostellar sources

Since Vela-C was not observed by *Spitzer* at  $24 \mu\text{m}$ , which would have revealed embedded sources with adequate sensitivity, we used the *Herschel* data themselves, and where available, other mid- and far-IR surveys to distinguish starless and protostellar sources. This distinction was primarily made on the basis of the presence of a  $70 \mu\text{m}$  counterpart (see also Bontemps et al. 2010). Indeed, a tight correlation between  $70 \mu\text{m}$  fluxes and internal luminosities of protostars has been demonstrated (Dunham et al. 2008), and, furthermore, external heating by the interstellar radiation field cannot produce a  $70 \mu\text{m}$  emission from a source detectable with *Herschel*. Hence, we consider as starless all sources without a  $70 \mu\text{m}$  counterpart (e.g. Fig. 3, left panel). Conversely, protostellar are *i*) sources whose  $70 \mu\text{m}$  flux lies above the best modified black body fit by more than  $3\sigma$  (e.g. Fig. 3, right panel), or, *ii*) sources detected in the mid-IR at  $\lambda < 70 \mu\text{m}$  by other surveys (IRAS/MSX/Akari). Out of 268 objects, 218 are starless and 48 are protostellar. Two other sources, not detected at  $\lambda < 70 \mu\text{m}$  by other surveys, were fitted with a single modified black body curve from  $70 \mu\text{m}$  to  $500 \mu\text{m}$ . The temperature of both sources is  $\sim 25$  K, i.e. warmer than starless sources (Sect. 3.4). On this basis we surmise that these sources are embedded Class 0 protostars (André et al. 1993).

#### 3.5. Physical parameters


In this section, we discuss the statistics of the key physical parameters measured directly from the observations or derived

from SEDs fits. These quantities are listed in Table 2 and plotted in Fig. 4.

Starless sources exhibit a very narrow temperature distribution (top left panel), strongly peaked at  $T_d \sim 10$  K. The high cutoff is at  $T_d \sim 15$  K, while the low one coincides with the minimum temperature of 8 K considered in the fit. On average, protostellar sources are warmer than starless sources; their temperature distribution spreads almost uniformly between 9 K and 15 K, though a few sources are found with  $T_d$  up to 25 K. The overall temperature distribution (not shown in Fig. 4) resembles that found from BLAST data of the whole Vela region (Netterfield et al. 2009). This latter distribution, however, does not show the sharp  $\sim 10$  K peak, but rather has a shallower shape peaking at  $\sim 12$  K. This marginal difference may be caused by both the better angular resolution of *Herschel*, which allows deeper penetration into colder regions, and to the inclusion of warmer regions (e.g., Vela-D, Olmi et al. 2009) in the Netterfield et al. sample. It is also interesting to compare the compact source distribution with the average temperature of the surrounding cloud. The latter was derived by Hill et al. (2011) by fitting a modified black body to the *Herschel* SED of each pixel. In the parts of the cloud where no compact sources are detected, the temperature is fairly constant at  $T_d \sim 14$  K (black line in Fig. 4). Practically all the starless sources are colder than the ambient medium, a result expected for dense structures heated only by external radiation. Interestingly, the same result is found for most of the protostellar sources. This finding reasonably indicates that the external envelope we are tracing is moderately heated by the internal protostar(s), which is therefore likely to be in the very early phases of the evolution.

The mass distribution (top right panel of Fig. 4) shows a nearly monothonic decrease from lower to higher masses. There are no significant differences between the average masses of starless and protostellar sources ( $\sim 5 M_\odot$ , Table 2); this suggests that the envelope remains substantially unperturbed by the internal gravitational collapse and, again, that the protostellar sources are relatively unevolved, because little mass accretion is taking place. Noticeably, 35 sources (7 of which are protostellar and 28 are starless) have  $10 \leq M/M_\odot \leq 20$  and 8 (1 of which is protostellar and 7 are starless) have  $20 \leq M/M_\odot \leq 55$ , whose locations are depicted with big crosses in Fig. 1. Apart from a few exceptions, these massive objects are fairly compact (with diameters between 0.04 pc and 0.07 pc), hence they are potentially able to form massive stars. For comparison, Hill et al. (2011) found in Vela-C nine objects with masses between  $13 M_\odot$  and  $70 M_\odot$ . We consider the agreement between the two determinations satisfactory, given the different source extraction methods (Hill et al. 2011, took the bias created by bright rimmed effects and source multiplicity into account in the photometry) and the different prescriptions under which the SED fitting has been performed (for example, Hill et al. did not apply flux scaling).

The source diameter distributions are presented in the bottom left panel of Fig. 4. For starless sources, it spreads from the resolution limit of 0.025 pc up to 0.13 pc, peaking around 0.05 pc. These dimensions, as anticipated in Sect. 2, span from those of progenitors of single stars up to those typical of large clumps, which may give rise to the formation of protostellar clusters. Conversely, protostellar sources are significantly more compact; this suggests that no further fragmentation will occur and that the majority of the protostellar sources likely host single stellar objects or stellar systems with a small number of members. The small sizes of protostellar sources can be alternatively (and additionally) explained by considering that the internal source(s) of heating likely causes a temperature gradient and hence a more



**Fig. 4.** Histograms of the main physical parameters of the sources in Vela-C (top left: temperature; top right: mass; bottom left: diameter; bottom right: *Herschel* luminosity). Starless and protostellar sources are separately highlighted in red and blue, respectively. The black line in the top left panel represents the temperature of the surrounding cloud, taken from Hill et al. (2011), while that in the bottom left panel highlights the spatial resolution limit.

peaked intensity distribution that is fitted with a compact spatial extent.

Finally, in the bottom right panel of Fig. 4, we plot the distributions of luminosity in the *Herschel* bands ( $L_{\text{FIR}}$ ). This quantity has been computed by interpolating the corrected fluxes between all available wavelengths between  $70 \mu\text{m}$  and  $500 \mu\text{m}$ . For starless sources,  $L_{\text{FIR}}$  is a good approximation of bolometric luminosity; typically, it ranges between  $0.05 L_\odot$  and  $0.5 L_\odot$ , with a small percentage of objects at higher luminosities not exceeding  $5 L_\odot$ . For protostellar sources,  $L_{\text{FIR}}$  has to be considered as a lower limit to the bolometric luminosity because of the lack of near- and mid-infrared data points in the SEDs.

## 4. Discussion

### 4.1. Virial analysis of starless sources

Starless sources are defined as *prestellar* if they are gravitationally bound (e.g., André et al. 2000; Di Francesco et al. 2007), hence they can potentially form one or more stars. In principle, spectroscopic observations would be required to derive the virial masses and to securely establish the sources' dynamical states. Since such observations are not available for Vela-C, we have assumed thermal pressure support and neglected the internal turbulence. In this simplified view, the virial mass can be surrogated by the critical Bonnor-Ebert mass:

$$M_{\text{BE}} \approx 2.4 R_{\text{BE}}^2 / G. \quad (4)$$


Here  $a$  is the sound speed at the source temperature<sup>2</sup>,  $G$  is the gravitational constant, and  $R_{\text{BE}}$  the Bonnor-Ebert radius (in pc).

<sup>2</sup>  $a = \sqrt{k_B T_d / \mu}$ , where  $k_B$  is the Boltzmann constant,  $T_d$  is the fitted temperature, and  $\mu = 2.33 m(H)$  is the mean molecular weight, with  $m(H)$  the atomic hydrogen mass.

**Table 2.** Statistics of the main physical parameters of the sources in Vela-C.

Parameter	Starless (218)			Protostellar (48)		
	median	average	min-max	median	average	min-max
$M (M_{\odot})$	3.3	5.5	0.13–55.8	2.7	4.8	0.15–29.1
$T_d$ (K)	10.0	10.3	8.0–15.2	11.4	12.8	9.0–24.2
$D$ (pc)	0.064	0.067	0.025–0.13	0.040	0.040	0.025–0.07
$L_{\text{FIR}} (L_{\odot})$	0.17	0.22	0.04–4.8	0.6	8.0	0.08–138

**Notes.** We give median, average, minimum, and maximum values.



**Fig. 5.** Mass vs. diameter diagram for the sources in Vela-C. Red filled circles and red open circles are prestellar and unbound sources, respectively. Blue filled circles are protostellar sources. Dotted lines are the loci of sources with  $M = 0.5 M_{\text{BE}}$ , for  $T = 8$  K and  $T = 20$  K, while the dashed vertical line represents the spatial resolution limit. We also show the curve representing the relation between mass and diameter in the case  $\lambda_0 = 160 \mu\text{m}$ . This relation shifts towards higher masses with increasing  $\lambda_0$ ; hence, since all our sources are located below the reported curve, they have optically thin emission at  $\lambda > 160 \mu\text{m}$ .

We assumed  $R_{\text{BE}} = D/2$ . Starless sources with  $M/M_{\text{BE}} \geq 0.5$  were selected as being gravitationally bound (Pound & Blitz 1993) and candidate prestellar. By this definition, we found very many prestellar sources (206 objects), i.e.,  $\sim 94\%$  of the starless ones, remarkably higher than the percentage (69%) found in the Aquila Rift cloud (Könyves et al. 2010; André et al. 2010). Moreover, this percentage maintains high (193 objects, i.e., 88%) even if we demand that bound sources have  $M$  at least equal to  $1 M_{\text{BE}}$ .

To check this result further, we plot all sources in a mass vs. diameter diagram (Fig. 5). First we note that, as a consequence of condition (4) in the selection criteria of Sect. 3.2, all points lie to the right of the spatial resolution limit of 0.025 pc, shown with a vertical dashed line. Second, we plot the relation between mass and diameter as a function of the second variable in Eq. (3), namely the wavelength  $\lambda_0$  at which the optical depth  $\tau = 1$ . This relation shifts towards higher masses with increasing  $\lambda_0$ , therefore all the sources located below (above) the curve for a given value of  $\lambda_0$  have optically thin (thick) emission at  $\lambda > \lambda_0$ . As an example, we show this relation for the value  $\lambda_0 = 160 \mu\text{m}$ , which is the shortest wavelength that we include in the SEDs fits.

Because all our sources are located below this curve, their emission is optically thin at wavelengths longer than  $160 \mu\text{m}$ .

Consistently with our virial classification, most of the prestellar sources in Fig. 5 lie above or between the lines of  $0.5 M_{\text{BE}}$ , computed at 8 K and 20 K. We note, however, that a relevant fraction of these objects have a mass up to a factor of ten larger than the Bonnor-Ebert mass at 20 K. Reasonably, these objects are dynamically unstable, if turbulence and magnetic field supports against gravity are neglected.


All sources classified as unbound are located below the Bonnor-Ebert curve at  $T = 8$  K. Noticeably, most of them cluster around the value  $M \sim 0.2 M_{\odot}$ . This corresponds to the mass detection limit, estimated from the flux sensitivity limits of Table 1 and by assuming average values of  $T$  and  $\lambda_0$  in Eqs. (2) and (3). Because this mass limit decreases with increasing temperature, we deduce that we are able to efficiently probe only the unbound sources particularly warm. Indeed, for the sub-sample of this category (12 objects) we obtain  $\langle T_d \rangle = 13.8$  K, namely higher than the value of  $\langle T_d \rangle = 10.3$  K, referring to the whole sample of starless sources (see Table 2). Noticeably, our inability to detect cold, unbound sources, may explain the very high percentage of prestellar sources with respect to the unbound ones (see above). Moreover, this fraction could even more decrease if the most massive prestellar cores were gravitationally unstable, as already noticed above.

Finally, to show the results of Table 2 more clearly, we also plot in the same diagram the positions of protostellar sources. As already noted in Sect. 3.5, they cluster in Fig. 5 at diameters smaller than starless sources, and close to our spatial resolution limit. Since no significant differences in mass (as well as in temperature values) are recognisable with respect to starless sources, this result should reflect an increase in average density.

#### 4.2. Luminosity vs. mass diagram

Another interesting perspective to be examined is the relation between the bolometric luminosities ( $L_{\text{bol}}$ ) and the envelope masses of the sources ( $M$ ). To make this comparison, we took as  $L_{\text{bol}}$  the  $L_{\text{FIR}}$  values, even if these adequately represent bolometric luminosities only for starless sources. Previous studies (e.g. Saraceno et al. 1996; André et al. 2000; Molinari et al. 2008; Elia et al. 2010; Bontemps et al. 2010; Henneman et al. 2010) have illustrated that the  $L_{\text{bol}}$  vs.  $M$  plot is a meaningful tool for characterising the evolutionary status of cores and clumps, since tracks representing model predictions can be directly compared with the locations of the observed sources. In such a diagram, a collapsing core is expected to initially follow an almost vertical path (i.e., mass constant with time, and luminosity strongly increasing with accretion), then subsequently protostellar outflow activity increasingly disperses the envelope while the luminosity remains almost constant, which results in a horizontal track.






**Fig. 6.**  $L_{\text{bol}}-M$  diagram for prestellar and protostellar sources in Vela-C (red and blue circles, respectively). Grey solid lines represent the evolutionary tracks for low-mass objects adopted by Molinari et al. (2008), for initial values of 0.5, 1, 5, and  $10 M_{\odot}$ , respectively. An arrow indicates the evolution direction, while green lines delimit the region of transition between Class 0 and Class I sources (André et al. 2000). The red dotted line represents the best-fitting power law ( $L_{\text{bol}} \propto M^{0.42 \pm 0.04}$ ) for the distribution of the prestellar sources.

In Fig. 6, the  $L_{\text{bol}}$ ,  $M$  pairs of the Vela-C protostellar and prestellar sources are plotted, along with the evolutionary tracks for low-mass objects adopted by Molinari et al. (2008), based on the simplified assumption of a central protostar accreting mass from the envelope at a constant rate  $\dot{M} = 10^{-5} M_{\odot} \text{ yr}^{-1}$ . More evidently than in previously shown diagrams, a significant distinction between the two populations, although not a complete segregation, is seen. In the same plot, we delimit with green lines the transition region between Class 0 and Class I objects (André et al. 2000), which roughly separates the vertical portions of the tracks, where Class 0 objects are expected to be found, from the horizontal ones, where Class I objects are located. Noticeably, the large majority of the protostellar sources populate the region corresponding to Class 0 objects. This is in contrast with our previous results based on SED fitting, according to which 48 sources are young protostars (although not much evolved) and just two sources are identified as candidate Class 0 sources. This discrepancy can be partly reconciled however if considering that our underestimate of the bolometric luminosities of protostellar sources reflects underestimates of the actual ages. Moreover, the André et al. (2000) limit separating Class 0 and I could not be completely adequate in Vela, as already found in Taurus by Motte & André (2001).

Also, the distribution of the prestellar sources is quite flat and homogeneously concentrated in a region of the diagram corresponding to a very early stage where no mass accretion has started. A tentative power-law fit gives a dependence  $L_{\text{bol}} \propto M^{(0.42 \pm 0.04)}$ . This slope is shallower than that found by Brand et al. (2001) for their sample of protostellar sources, namely 0.85. A similar scaling relation has been found for the CO luminosity of clumps: from the theoretical point of view,  $L_{\text{CO}} \propto M_{\text{vir}}^{-1}$  at the virial equilibrium (Scoville & Sanders 1987; see also Wolfire et al. 1993, and references therein); this slope has been found by Rengarajan (1984), while other authors find lower values (e.g. Solomon et al. 1987; Yonekura et al. 1997; Miyazaki & Tsuboi 2000). A direct comparison with our result, however, cannot be performed because of the different classes of sources considered, or different quantities compared.



**Fig. 7.** Source mass distribution in Vela-C. The error bars correspond to  $\sqrt{N}$  statistical uncertainties. Continuous red line represents the fit to the linear portion of the mass distribution (namely that consistent with a single straight line within the errorbars) for all prestellar sources. A similar fit performed on the restricted sample of prestellar sources with diameters less than 0.08 pc is represented with a grey dashed line. The derived slopes are reported as well. Mass distribution of CO clumps (Kramer et al. 1998), the single-star IMF (Kroupa 2001) and the multiple-system IMF (Chabrier 2005) are shown for comparison with an orange, a green, and a blue line, respectively. The dotted vertical line represents the mass completeness limit of prestellar sources.

#### 4.3. Source mass distribution

Figure 7 shows the mass distribution of prestellar sources in Vela-C. We estimate a mass completeness limit of  $\sim 4 M_{\odot}$ , derived by assuming the average temperature of prestellar sources of Table 2 and the flux 90% completeness limit at  $160 \mu\text{m}$  (see Table 1). Note that this value is significantly better than the BLAST completeness limit ( $\sim 14 M_{\odot}$  for sources colder than 14 K). We fitted a power law  $N(\log M) \propto M^{\gamma}$  to the linear portion of the distribution, finding a slope  $\gamma = -1.1 \pm 0.2$ . The uncertainty was determined by considering both the statistical error of the data and the variation of the slope with the histogram binning, which was varied from 0.1 to 0.3 in  $\log(M/M_{\odot})$ . Noticeably, no variations are found in the slope taking into account only the 193 objects for which  $M \geq 1M_{\text{BE}}$ , since their mass distribution differs from that of the whole sample only in the mass range below the completeness limit, where the fit is not performed.

The value of  $\gamma$  we find in Vela-C is shallower than that found by Netterfield et al. ( $\gamma = -1.9 \pm 0.2$ , for sources with  $T_{\text{d}} < 14 \text{ K}$ ). If we limit the fit to the BLAST mass completeness limit of  $14 M_{\odot}$ , however, we obtain a slope  $\gamma = -1.9 \pm 0.2$ , which reconciles our high-mass end distribution with that obtained with BLAST. More important, and thanks to *Herschel's* massive increase in sensitivity, which significantly extends the mass range, a possible change of slope is recognizable at  $M \gtrsim 10 M_{\odot}$ . This is close to the value of  $\sim 9 M_{\odot}$  where a steepening was probed, for example, in Orion A (Ikeda et al. 2007). Recent theoretical models by Padoan & Nordlund (2011) predict this change of slope, even if at lower masses ( $M \sim 3-5 M_{\odot}$ ). Unfortunately, no firm conclusion can be drawn on the base of our data because of the poor statistics in the high-mass bins.

Compared with other literature values, our  $\gamma$  value of  $-1.1$  is between the mass distribution slope of CO clumps ( $\gamma \sim -0.7$ ,

Kramer et al. 1998) and that typical of prestellar cores, as for example that measured in the Aquila Rift ( $\gamma = -1.45 \pm 0.2$ ) on a sample of sources with  $2 \lesssim M/M_{\odot} \lesssim 10$  and diameter typically less than 0.08 pc (Könyves et al. 2010). Hence, our slope reflects the heterogenous nature of our sample, which is composed of objects with a variety of diameters typical of both clumps and cores. Indeed, if we include in the mass distribution fit only sources with diameter less than 0.08 pc (142 objects), we obtain a slope  $\gamma = -1.4 \pm 0.2$  (grey dotted line in Fig. 7), which well reconciles with the slope of the Aquila Rift. Such a slope is also consistent with the IMF slope for  $1.0 \lesssim M/M_{\odot}$  (for single-star IMF  $\gamma = -1.3 \pm 0.7$ , Kroupa 2001, while for multiple-systems IMF  $\gamma = -1.35 \pm 0.3$ , Chabrier 2005), likely indicating that fragmentation will proceed less efficiently in these small objects.

Finally, a substantial agreement within the error bars is provided by the comparison with surveys covering a mass range largely overlapping that of Vela-C. For example,  $\gamma \sim 1.3 \pm 0.2$  for  $0.8 < M/M_{\odot} < 6$  in Perseus, Serpens and Ophiucus (Enoch et al. 2008) and  $\gamma = -1.3 \pm 0.1$  for  $3 \lesssim M/M_{\odot} \lesssim 60$  in Orion A (Ikeda et al. 2007).

## 5. Conclusions

We have reported on the *Herschel* observations of the Vela-C star-forming region over an area of  $\sim 3$  square degrees. From our analysis we obtained the following results:

- From a  $5\sigma$  level catalogue of cold compact sources in five *Herschel* bands between  $70\mu\text{m}$  and  $500\mu\text{m}$ , we have selected a robust sub-sample of 268 sources. Their physical diameters indicate that our sample is mainly composed of cloud clumps ( $0.05 \text{ pc} \lesssim D \lesssim 0.13 \text{ pc}$ ), together with a  $\sim 25\%$  of cores ( $0.025 \text{ pc} \lesssim D \lesssim 0.05 \text{ pc}$ ).
- Based on the detection of a  $70\mu\text{m}$  flux, we identified 218 starless and 48 protostellar sources. For two other sources we do not give a secure classification, but suggest them to be candidate Class 0 protostars.
- Source physical parameters have been derived from modified black body fits to the SEDs. Both starless and protostellar sources are on average colder than the surrounding medium. This indicates that the radiation from interstellar field and/or from embedded protostars is inefficient in penetrating the cold dust in depth.
- Protostellar sources are on average slightly warmer and more compact than starless sources. Both these findings can be ascribed to the presence of an internal source(s) of moderate heating, which also causes a temperature gradient and a more peaked intensity distribution. Moreover, the reduced dimensions of protostellar sources may indicate that they will not fragment further.
- No significant differences are found between the masses of the two groups. These range from sub-solar up to tens of solar masses. In particular, we found eight objects with  $M > 20 M_{\odot}$ , which are potential candidate progenitors of high-mass stars.
- More than 90% of the starless sources are prestellar (i.e. bound) if a virial analysis is applied. This percentage, however, should be considered as an upper limit, both because our sensitivity does not allow us to efficiently probe the coldest unbound sources and because several of the massive prestellar sources could be gravitationally unstable.
- A luminosity vs. mass diagram for the two populations of prestellar and protostellar sources was constructed. Prestellar

sources cluster in a well-defined region of the diagram that corresponds to a very early stage in which no mass accretion is expected to have started. A tentative power-law fit to the observed distribution gives a dependence  $L_{\text{bol}} \propto M^{(0.42 \pm 0.04)}$ . Conversely, protostellar sources populate the diagram region that corresponds to the early accretion phase.

- The mass distribution of the prestellar sources with  $M \geq 4 M_{\odot}$  shows a slope of  $-1.1 \pm 0.2$ . This is between that typical of CO clumps and those of cores in closeby star-forming regions, maybe reflecting the heterogeneous nature of our sample, which is a mixture of cores and clumps. We signal a possible a change of slope in the mass distribution for  $M \gtrsim 10 M_{\odot}$ , even if the very big errorbars in the higher mass bins prevent us to draw firm conclusions on its reliability.

*Acknowledgements.* SPIRE has been developed by a consortium of institutes led by Cardiff Univ. (UK) and including Univ.Lethbridge (Canada); NAOC (China); CEA, LAM (France); IFSI, Univ. Padua (Italy); IAC (Spain); Stockholm Observatory (Sweden); Imperial College London, RAL, UCL-MSSL, UKATC, Univ. Sussex (UK); Caltech, JPL, NHSC, Univ. Colorado (USA). This development has been supported by national funding agencies: CSA (Canada); NAOC (China); CEA, CNES, CNRS (France); ASI (Italy); MCINN (Spain); SNSB (Sweden); STFC and UKSA (UK); and NASA (USA). PACS has been developed by a consortium of institutes led by MPE (Germany) and including UVIE (Austria); KU Leuven, CSL, IMEC (Belgium); CEA, LAM (France); MPIA (Germany); INAF-IFSI/OAA/OAP/OAT, LENS, SISSA (Italy); IAC (Spain). This development has been supported by the funding agencies BMVIT (Austria), ESA-PRODEX (Belgium), CEA/CNES (France), DLR (Germany), ASI/INAF (Italy), and CICYT/MCYT (Spain). Data processing and maps production has been possible thanks to ASI generous support via contracts I/005/07/0-1, I/005/011/0 and I/038/080/0.

## References

- André, P., Ward-Thompson, D., & Barsony, M. 1993, *ApJ*, 406, 122  
 André, P., Ward-Thompson, D., & Barsony, M. 2000, in *Protostars & Planets IV*, ed. V. Mannings, A. P. Boss, S. S. Russell (Tucson: University of Arizona Press), 59  
 André, P., Men'shchikov, A., Bontemps, S., et al. 2010, *A&A*, 518, L102  
 Baba, D., Sato, S., Nagashima, C., et al. 2006, *AJ*, 132, 1692  
 Beckwit, S. V. W., Sargent, A. I., Chini, R. S., et al. 1990, *AJ*, 99, 924  
 Bik, A., Puga, E., Waters L. B. F. M., et al. 2010, *ApJ*, 713, 883  
 Blitz, L. 1991, in *The Physics of Star Formation and Early Stellar Evolution*, ed. C. J. Lada, & N. D. Kylafis (Dordrecht: Kluwer), 3  
 Bontemps, S., André, P., Könyves, V., et al. 2010, *A&A*, 518, L85  
 Brand J., Cesaroni, R., Palla, F., et al. 2001, *A&A*, 370, 230  
 Chabrier, G. 2005, *ASSL*, 327, 41  
 Di Francesco, J., Evans, II, N. J., Caselli, P., et al. 2007, in *Protostars & Planets V*, 17  
 Dunham, M. M., Crapsi, A., Evans II, N. J., et al. 2008, *ApJ*, 179, 249  
 Elia, D., Schisano, E., Molinari, S., et al. 2010, *A&A*, 518, L97  
 Enoch, M. L., Evans II, N. J., Sargent, A. et al. 2008, *ApJ*, 684, 1240  
 Herbst, W. 1975, *AJ*, 80, 683  
 Griffin, M. J., Abergel, A., Abreu, A., et al. 2010, *A&A*, 518, L3  
 Ikeda, N., & Kitamura, Y. 2011, *ApJ*, 732, 101  
 Ikeda, N., Sunada, K., & Kitamura Y. 2007, *ApJ*, 665, 1194  
 Hennemann, M., Motte, F., Bontemps, S., et al. 2010, *A&A*, 518, L84  
 Hildebrand, R. H. 1983, *QJRAS*, 24, 267  
 Hill, T., Motte, F., Didelon, P., et al. 2011, *A&A*, 533, A94  
 Könyves, V., André, P., Men'shchikov, A., et al. 2010, *A&A*, 518, L106  
 Kramer, C., Stutzki, J., Robrig, R., et al. 1998, *A&A*, 329, 249  
 Kroupa, P. 2001, *MNRAS*, 322, 231  
 Liseau, R., Lorenzetti, D., Nisini, B., et al. 1992, *A&A*, 265, 577  
 Lorenzetti, D., Spinoglio, L., & Liseau, R. 1993, *A&A*, 275, 489  
 Massi, F., Lorenzetti, D., Giannini, T., et al. 2000, *A&A*, 353, 589  
 Massi, F., Lorenzetti, D., & Giannini, T. 2003, *A&A*, 399, 147  
 Mc Kee, C. F., & Ostriker, E. C. 2007, *ARA&A*, 45, 565  
 Miyazaki, A., & Tsuboi, M. 2000, *ApJ*, 536, 357  
 Molinari, S., Pezzuto S., Cesaroni, R., et al. 2008, *A&A*, 481, 345  
 Molinari, S., Swinyard, B., Bally, J., et al. 2010, *A&A*, 518, L100  
 Molinari, S., Faustini, F., Schisano, E., et al. 2011, *A&A*, 530, A133  
 Motte, F., & André, P. 2001, *A&A*, 365, 440  
 Motte, F., André, P., & Neri, R. 1998, *A&A*, 336, 150  
 Motte, F., André, P., Ward-Thompson, D., et al. 2001, *A&A*, 372, 41



- Motte, F., Zavagno, A., Bontemps, S., et al. 2010, *A&A*, 518, L77
- Müller, T., Okumura, K., & Klaas, U. 2011, PACS technical note PICC-ME-IN-038, <http://herschel.esac.esa.int//twiki/pub/PacsCalibrationWeb/cc-report-v1.pdf>
- Murphy, D. C., & May, J. 1991, *A&A*, 247, 202
- Netterfield, C. B., Ade, P. A. R., Bock, J. J., et al. 2009, *ApJ*, 707, 1824
- Nguyễn Lương, Q., Motte, F., Hennemann, M., et al. 2011, *A&A*, 535, A76
- Olmí, L., Ade, P. A. R., Anglés-Alcázar, D., et al. 2009, *ApJ*, 707, 1836
- Ott, S. 2010, in *Astronomical Data Analysis Software and System XIX*, ed. M. O. Y. Mizumoto, K.-I. Morita, ASP Conf. Ser., 434, 139
- Padoan, P., & Nordlund, Å. 2011, *ApJ*, 741, L22
- Pascale, E., Ade, P. A. R., Bock, J. J., et al. 2008, *ApJ*, 681, 400
- Poglitsch, A., Waelkens, C., Geis, N., et al. 2010, *A&A*, 518, L2
- Pound, M. W., & Blitz, L. 1993, *ApJ*, 418, 328
- Rathborne, J. M., Lada, C. J., Muench, A. A., et al. 2009, *ApJ*, 699, 742
- Reid, M. A., & Wilson, C. D. 2006, *ApJ*, 644, 990
- Rengarajan, T. N. 1984, *ApJ*, 287, 671
- Saraceno, P., André, Ph., Ceccarelli, C., et al. 1996, *A&A*, 309, 827
- Scoville, N. Z., & Sanders, D. B. 1987, in *Interstellar processes*, Proc. Symp., Grand Teton National Park, WY, July 1–7, 1986 (Dordrecht: D. Reidel Publishing Co.), 21
- Shu, F. H. 1977, *ApJ*, 214, 488
- Solomon P. M., Rivolo, A. R., Barrett, J., et al. 1987, *ApJ*, 319, 730
- Testi, L., & Sargent, A. I. 1998, *ApJ*, 508, L91
- Traficante, A., Calzoletti, L., Veneziani, M., et al. 2011, *MNRAS*, 416, 2932
- Wolfire, M. G., Hollenbach, D., & Tielens, A. G. G. M. 1993, *ApJ*, 402, 195
- Yamaguchi, N., Mizuno, N., Saito, H., et al. 1999, *PASJ*, 51, 775
- Yonekura, Y., Dobashi, K., Mizuno, A., et al. 1997, *ApJS*, 110, 2