

## The Na-O anticorrelation in horizontal branch stars

### II. NGC 1851<sup>★</sup> (*Corrigendum*)

R. G. Gratton<sup>1</sup>, S. Lucatello<sup>1</sup>, E. Carretta<sup>2</sup>, A. Bragaglia<sup>2</sup>, V. D'Orazi<sup>1</sup>, Y. Momany<sup>1,3</sup>, A. Sollima<sup>1</sup>,  
M. Salaris<sup>4</sup>, and S. Cassisi<sup>5</sup>

<sup>1</sup> INAF – Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, 35122 Padova, Italy  
e-mail: [raffaele.gratton@oapd.inaf.it](mailto:raffaele.gratton@oapd.inaf.it)

<sup>2</sup> INAF – Osservatorio Astronomico di Bologna, via Ranzani 1, 40127 Bologna, Italy

<sup>3</sup> European Southern Observatory, Alonso de Cordova 3107, Vitacura, Santiago, Chile

<sup>4</sup> Astrophysics Research Institute, Liverpool John Moores University, UK

<sup>5</sup> INAF – Osservatorio Astronomico di Teramo, via Collurania, Teramo, Italy

A&A 539, A19 (2012), DOI: [10.1051/0004-6361/201118491](https://doi.org/10.1051/0004-6361/201118491)

**Key words.** stars: evolution – stars: abundances – stars: Population II – globular clusters: individual: NGC 1851 – globular clusters: general – errata, addenda

The Al abundances in NGC 1851 RHB stars given in Table 3 of this paper should be corrected upward by 0.2 dex because we realized they were obtained by assuming an inconsistent, too high, solar abundance.

---

\* Based on observations collected at ESO telescopes under program 386.D-0086.