Chemistry of massive young stellar objects with a disk-like structure[★]

K. Isokoski¹, S. Bottinelli^{2,3}, and E. F. van Dishoeck^{4,5}

¹ Raymond and Beverly Sackler Laboratory for Astrophysics, Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

e-mail: isokoski@strw.leidenuniv.nl

² Université de Toulouse, UPS-OMP, IRAP, 31400 Toulouse, France

³ CNRS, IRAP, 9 Av. Colonel Roche, BP 44346, 31028 Toulouse Cedex 4, France

⁴ Leiden Observatory, Leiden University, PO Box 9513, 2300 RA Leiden, The Netherlands

⁵ Max-Planck Institut für Extraterrestrische Physik (MPE), Giessenbachstr. 1, 85748 Garching, Germany

Received 19 December 2012 / Accepted 21 March 2013

ABSTRACT

Aims. Our goal is to take an inventory of complex molecules in three well-known high-mass protostars for which disks or toroids have been claimed and to study the similarities and differences with a sample of massive young stellar objects (YSOs) without evidence of such flattened disk-like structures. With a disk-like geometry, UV radiation can escape more readily and potentially affect the ice and gas chemistry on hot-core scales.

Methods. A partial submillimeter line survey, targeting CH₃OH, H₂CO, C₂H₅OH, HCOOCH₃, CH₃OCH₃, CH₃CN, HNC, NH₂CHO, C₂H₅CN, CH₂CO, HCOOH, CH₃CHO, and CH₃CCH, was made toward three massive YSOs with disk-like structures, IRAS 20126+4104, IRAS 18089-1732, and G31.41+0.31. Rotation temperatures and column densities were determined by the rotation diagram method, as well as by independent spectral modeling. The molecular abundances were compared with previous observations of massive YSOs without evidence of any disk structure, targeting the same molecules with the same settings and using the same analysis method.

Results. Consistent with previous studies, different complex organic species have different characteristic rotation temperatures and can be classified either as warm (>100 K) or cold (<100 K). The excitation temperatures and abundance ratios are similar from source to source and no significant difference can be established between the two source types. Acetone, CH₃COCH₃, is detected for the first time in G31.41+0.31 and IRAS 18089-1732. Temperatures and abundances derived from the two analysis methods generally agree within factors of a few.

Conclusions. The lack of chemical differentiation between massive YSOs with and without observed disks suggest either that the chemical complexity is already fully established in the ices in the cold prestellar phase or that the material experiences similar physical conditions and UV exposure through outflow cavities during the short embedded lifetime.

Key words. astrochemistry – line: identification – methods: observational – stars: formation – ISM: abundances – ISM: molecules

1. Introduction

Millimeter lines from complex organic molecules are widely associated with high-mass star forming regions and indeed form one of the signposts of the deeply embedded phase of star formation (e.g., Blake et al. 1987; Hatchell et al. 1998; Gibb et al. 2000; Fontani et al. 2007; Requena-Torres et al. 2008; Belloche et al. 2009; Zernickel et al. 2012). Many studies of the chemistry in such regions have been carried out, either through complete spectral surveys of individual sources or by targeting individual molecules in a larger number of sources (see Herbst & van Dishoeck 2009; Caselli & Ceccarelli 2012, for reviews). In spite of all this work, only few systematic studies of the abundances of commonly observed complex molecules have been performed across a sample of massive young stellar objects (YSOs), to search for similarities or differences depending on physical structure and evolutionary state of the object. Intercomparison of published data sets is often complicated by the use of different telescopes with different beams, different frequency ranges and different analysis techniques.

Chemical abundances depend on the physical structure of the source such as temperature, density and their evolution with time, as well as the amount of UV radiation impinging on the gas and dust. In contrast with the case of solar-mass stars, the physical structures and mechanisms for forming massive ($M > 8 M_{\odot}$) stars are still poorly understood. Indeed, theoretically, the powerful UV radiation pressure from a high-mass protostellar object (HMPO) should prevent further accretion and so inhibit the formation of more massive stars (Zinnecker & Yorke 2007). However, a number of recent studies have claimed the presence of disk- or toroid-like “equatorial” structures surrounding a handful of high-mass protostars (Cesaroni et al. 2007). These data support theories in which high-mass star formation proceeds in a similar way as that of low-mass stars via a disk accretion phase in which high accretion rates and non spherically symmetric structures overcome the problem of radiation pressure. The best evidence so far is that for a ~5000 AU disk in Keplerian rotation around IRAS 20126+4104, claimed by Cesaroni et al. (1999) on the basis of the presence of a velocity gradient in the CH₃CN emission perpendicular to the direction of the outflow, as predicted by the disk-accretion paradigm. Surprisingly, even in the best case of IRAS 20126+4104, the

[★] Appendices are available in electronic form at <http://www.aanda.org>

Fig. 1. Illustration of a protostar with a spherical structure (*left*) and a protostar with a flattened disk-like structure (*right*) with enhanced UV photons illuminating the walls of the outflow cavity.

detailed chemistry of these (potential) disks has not yet been studied.

Most chemical models invoke grain surface chemistry to create different generations of complex organic molecules (Tielens & Charnley 1997). Hydrogenation of solid O, C, N and CO during the cold ($T_d < 20$ K) prestellar phase leads to ample production of CH_3OH and other hydrogenated species like H_2O , CH_4 and NH_3 (Tielens & Hagen 1982). Exposure to UV radiation results in photodissociation of these simple ices, with the fragments becoming mobile as the cloud core heats up during the protostellar phase. First generation complex molecules result from the subsequent recombination of the photofragments, and will eventually evaporate once the grain temperature rises above the ice sublimation temperature of ~ 100 K (Garrod & Herbst 2006; Garrod et al. 2008). Good examples are $\text{C}_2\text{H}_5\text{OH}$, HCOOCH_3 and CH_3OCH_3 resulting from mild UV processing of CH_3OH ice (Öberg et al. 2009). Finally, a hot core gas phase chemistry between evaporated molecules can drive further complexity in second generation species (e.g., Millar et al. 1991; Charnley et al. 1992, 1995).

One of the most obvious consequences of an equatorial rather than spherical structure is that UV radiation can more easily escape the central source and illuminate the surface layers of the surrounding disk or toroid as well as the larger scale envelope (Bruderer et al. 2009, 2010) (Fig. 1). This can trigger enhanced formation of complex organic molecules in the ices relative to methanol. Another effect of UV radiation is that it drives increased photodissociation of gaseous N_2 and CO. The resulting atomic N and C would then be available for grain surface chemistry potentially leading to enhanced abundances of species like HNC and NH_2CHO .

To investigate the effects of disk-like structures on the chemistry, we present here a single-dish survey using the *James Clerk Maxwell Telescope* (JCMT) of three HMPOs for which large equatorial structures (size > 2000 AU) have been inferred, namely IRAS 20126+4104, IRAS 18089–1732, and G31.41+0.31. The results are compared with those of a recent survey of a sample of HMPOs by Bisschop et al. (2007, hereafter BIS07), targeting many of the same molecules and settings. The use of the same telescope and analysis method allows a meaningful comparison between the two samples of sources. BIS07 found that the O-rich complex molecules are closely correlated with the grain surface product CH_3OH supporting the above general chemical scenario. N-rich organic molecules do not appear to be correlated with O-rich ones, but overall, the relative abundances of the various species are found to be remarkably constant within one chemical family. One of the main questions is whether this similarity in abundance ratios also holds for sources with disk-like structures. Although our

observations do not spatially resolve these structures, they are sensitive enough and span a large enough energy range to determine abundances on scales of $\sim 1''$ and thereby set the scene for future high-angular resolution observations with interferometers like the Atacama Large Millimeter/submillimeter Array (ALMA). Moreover, current interferometers resolve out part of the emission, which is why single-dish observations are still meaningful.

This paper is organized as follows. In Sect. 2, the observed sources and frequency settings are introduced and the details of the observations are presented. Section 3 focuses on the data analysis methods. Specifically, two techniques are used to determine excitation temperatures and column densities: the rotation diagram method employed by BIS07 and spectral modeling tools in which the observed spectra are simulated directly. Section 4 presents the results from both analysis methods and discusses their advantages and disadvantages. Finally, Sect. 5 compares our results with those of BIS07 to draw conclusions on similarities and differences in chemical abundances between sources with and without large equatorial structures.

2. Observations

2.1. Observed sources

Table 1 gives the coordinates, luminosity L , distance d , galactocentric radius R_{GC} , velocity of the local standard of rest V_{LSR} and the typical line width for the observed sources. The selected sources are massive YSOs, for which strong evidence exists for a circumstellar disk structure. All sources are expected to harbor a *hot core*: a compact, dense ($\geq 10^7 \text{ cm}^{-3}$) and warm (≥ 100 K) region with complex chemistry triggered by the grain mantle evaporation (Kurtz et al. 2000). CH_3CN emission from ≥ 100 K gas is present in all sources. Moreover, CH_3OH 7_K-6_K transitions (338.5 GHz) with main beam temperatures of ≥ 1 K have been observed for these sources. Sources also needed to be visible from the JCMT¹.

2.1.1. IRAS 20126+4104

IRAS 20126+4104 (hereafter IRAS 20126) is a luminous ($\sim 10^4 L_\odot$) YSO located relatively nearby at a distance of 1.64 kpc (Moscadelli et al. 2011). It was first identified in the IRAS point source catalog by the IR colors typical of ultracompact HII regions and by H_2O maser emission characteristic of high-mass star formation (Comoretto et al. 1990). IRAS 20126 features a ~ 0.25 -pc scale inner jet traced by H_2O maser spots in the SE-NW direction with decreasing velocity gradient (Tofani et al. 1995). Source and masers are embedded inside a dense, parsec-scale molecular clump (Estalella et al. 1993; Cesaroni et al. 1999). The inner jet feeds into a larger scale bipolar outflow with the two having reversed velocity lobes (Wilking et al. 1990; Cesaroni et al. 1999). The reversal is likely to be due to precession of the inner jet caused by a companion separated by a distance of $\sim 0.5''$ (850 AU) (Hofner et al. 1999; Shepherd et al. 2000; Cesaroni et al. 2005; Sridharan et al. 2005). A rotating, flattened, Keplerian disk structure has been detected perpendicular to the inner jet. Observations of CH_3CN transitions show a

¹ The *James Clerk Maxwell Telescope* is operated by the Joint Astronomy Centre, on behalf of the Particle Physics and Astronomy Research council of the United Kingdom, the Netherlands Organization for Scientific Research and the National Research Council of Canada. The project ID is m09bn10.

Table 1. Coordinates, luminosity, distance, galactocentric radius, velocity of the local standard of rest, $^{12}\text{C}/^{13}\text{C}$ isotope ratio and the typical line width for the observed sources.

Sources	$\alpha(2000)$	$\delta(2000)$	L [$10^5 L_\odot$]	d [kpc]	R_{GC}^* [kpc]	V_{LSR} [km s^{-1}]	$^{12}\text{C}/^{13}\text{C}$	ΔV [km s^{-1}]
IRAS 20126+4104	20:14:26.04	+41:13:32.5	0.13 ^a	1.64 ^b	8.3	−3.5	70	6 ^c
IRAS 18089-1732	18:11:51.40	−17:31:28.5	0.32 ^d	2.34 ^e	6.2	33.8	54	5 ^c
G31.41+0.31	18:47:34.33	−01:12:46.5	2.6 ^f	7.9 ^g	4.5	97.0	41	6–10 ^h

Notes. ^(*) The galactocentric radii were calculated using distances d in this table and a IAU recommended distance from the galactic center $R_0 = 8.5$ kpc. The $^{12}\text{C}/^{13}\text{C}$ isotope ratios are calculated from Eq. (11).

References. ^(a) Cesaroni et al. (1997), ^(b) Moscadelli et al. (2011), ^(c) Leurini et al. (2007), ^(d) Sridharan et al. (2002), ^(e) Xu et al. (2011), ^(f) Cesaroni et al. (1994b), ^(g) Churchwell et al. (1990), ^(h) Fontani et al. (2007).

Keplerian circumstellar disk (radius ~ 1000 AU) with a velocity gradient perpendicular to the jet and a hot core with a diameter of ~ 0.0082 pc and a temperature of ~ 200 K at a geometric center of the outflow (Cesaroni et al. 1997, 1999; Zhang et al. 1998). Direct near-infrared (NIR) observations show a disk structure as a dark line (Sridharan et al. 2005). The disk shows a temperature and density gradient and is going through infall of material with a rate of $\sim 2 \times 10^{-3} M_\odot \text{ yr}^{-1}$ as expected for a protostar of this mass and luminosity (Cesaroni et al. 2005). Recent modeling of the Spectral Energy Distribution (SED) of IRAS 20126 shows indeed a better fit when a disk is included (Johnston et al. 2011).

2.1.2. IRAS 18089-1732

IRAS 18089-1732 (hereafter IRAS 18089) is a luminous ($\sim 10^{4.5} L_\odot$, Sridharan et al. 2002) YSO located at a distance of 2.34 kpc (Xu et al. 2011). It was identified based on CS detections of bright IRAS point sources with colors similar to ultracompact HII regions and the absence of significant free-free emission (Sridharan et al. 2002), and with H_2O and CH_3OH maser emission (Beuther et al. 2002). The CO line profile shows a wing structure characteristic of an outflow, although no clear outflow structure could be resolved from the CO maps (Beuther et al. 2002). A collimated outflow in the Northern direction is however seen in SiO emission on scales of $5''$ (Beuther et al. 2004). IRAS 18089 has a highly non-circular dust core of ~ 3000 AU diameter ($\sim 1''$), with optically thick CH_3CN at ~ 350 K (Beuther et al. 2005). HCOOCH_3 was found to be optically thin, with emission confined to the core, and showing a velocity gradient perpendicular to the outflow indicative of a rotating disk (Beuther et al. 2004). Also hot NH_3 shows a velocity gradient perpendicular to the outflow, although no Keplerian rotation was found, possibly due to infall and/or self gravitation (Beuther & Walsh 2008). Several hot-core molecules (HCOOCH_3 , CH_3CN , CH_3OCH_3 , HNCO , NH_2CHO , CH_3OH , $\text{C}_2\text{H}_5\text{OH}$) were mapped by Beuther et al. (2005) but no column densities or abundances were reported.

2.1.3. G31.41+0.31

G31.41+0.31 (hereafter G31) is a luminous ($2.6 \times 10^5 L_\odot$, Cesaroni et al. 1994b) YSO at a distance of 7.9 kpc (Churchwell et al. 1990). Preliminary evidence for a rotating disk with a perpendicular bipolar outflow was reported by Cesaroni et al. (1994a,b) and Olmi et al. (1996) showing a velocity gradient across the core in the NE-SW direction, similar to previously detected OH masers (Gaume & Mutel 1987). High-angular resolution CH_3CN observations by Beltrán et al. (2005) could not

detect Keplerian rotation typical for less luminous stars, nevertheless a toroidal structure undergoing gravitational collapse and fast accretion ($\sim 3 \times 10^{-2} M_\odot \text{ yr}^{-1}$) onto the central object was found. The G31 hot molecular core (HMC) is part of a complex region in which multiple stellar sources are detected (Benjamin et al. 2003); indeed, it is separated from an ultracompact (UC) HII region by only $\sim 5''$ and overlaps with a diffuse halo of free-free emission, possibly associated with the UC HII region itself (Cesaroni et al. 1998). More recent interferometric observations confirm the velocity gradient in the NH_3 (4,4) inversion transition and in CH_3CN data (Cesaroni et al. 2010, 2011). Line profiles look like a rotating toroid with infall motion. Several complex hot-core molecules have been observed in G31, including glycolaldehyde CH_2OHCHO (Beltrán et al. 2009), but again no abundance ratios have been presented.

2.2. Observational details

The observations were performed at the JCMT on Mauna Kea, Hawaii, between August 2007 and September 2009. The observations of the 338 GHz region covering CH_3OH ($7_K \rightarrow 6_K$) transitions were taken from JCMT archive. The front ends consisted of the facility receivers A3 (230 GHz region) and HARP-B (340 GHz region). The back-end was the digital autocorrelation spectrometer (ACSIS), covering 400 and 250 MHz of instantaneous bandwidth for A3 and HARP-B, respectively, with a channel width of 50 kHz. The noise level for both receivers was $T_{\text{rms}} \sim 20$ mK on a T_A^* scale when binned to 0.5 km s^{-1} . The integration time was ~ 1 h and 1.8 h for A3 and HARP-B, respectively. The spectra were scaled from the observed antenna temperature, T_A^* , to main-beam temperature, T_{MB} , using main beam efficiencies η_{MB} of 0.69 and 0.63 at 230 GHz and 345 GHz, respectively. We adopt a T_A^* calibration error of 20%.

The HPBW (half-power beam width, θ_B) for the 230 and 345 GHz band observations are $20\text{--}21''$ and $14''$, respectively. Emission from a volume with a source diameter $\theta_S \leq \theta_B$ undergoes beam dilution described by the beam-filling factor, η_{BF} :

$$\eta_{\text{BF}} = \frac{\theta_S^2}{\theta_S^2 + \theta_B^2}. \quad (1)$$

Table 2 gives the observed frequency settings and the targeted molecular lines. The settings were taken from BIS07 and were chosen to cover at least one strong line for the target molecules as well as lines of other interesting species. Strong lines of target molecules were chosen due to their high main-beam temperatures and minimum line-confusion in line surveys of Orion-KL by Sutton et al. (1985) and Schilke et al. (1997) at 230 GHz and

Table 2. Observed frequency settings and molecular lines.

Molecule	Freq. [GHz]	E_{up} [K]	$\mu^2 S$ [D ²]	Transition	Additional molecules
CH ₃ CN	239.1195	144.77	811.86	13 _K –12 _K	CH ₃ ¹³ CN, HCOOCH ₃ , CH ₃ OCH ₃
HNCO	219.7983	58.02	28.112	10 _{0,10,11} –9 _{0,9,10}	H ₂ ¹³ CO, C ₂ H ₅ CN
	240.8809	112.53	30.431	11 _{1,11,12} –10 _{1,10,11}	CH ₃ OCH ₃ , CH ₃ OH, HN ¹³ CO
HCOOCH ₃	222.3453	37.89	42.100	8 _{5,4} –7 _{4,3}	CH ₃ OCH ₃ , NH ₂ CHO
	225.2568	125.50	33.070	18 _{6,12} –17 _{6,12}	H ₂ CO, CH ₃ OCH ₃ , ¹³ CH ₃ OH
H ₂ CO	364.2752	158.42	52.165	5 _{3,3} –4 _{3,2}	C ₂ H ₅ OH
CH ₃ CN	331.0716	151.11	513.924	18 _K –17 _K	HCOOCH ₃ , HNCO, CH ₃ ¹³ CN
HNCO	352.8979	187.25	43.387	16 _{1,15,17} –15 _{1,14,16}	C ₂ H ₅ CN, C ₂ H ₅ OH, HCOOCH ₃
NH ₂ CHO	345.1826	151.59	664.219	17 _{0,17} –16 _{0,16}	HCOOCH ₃ , C ₂ H ₅ OH, ¹³ CH ₃ OH
HCOOCH ₃	354.6084	293.39	87.321	33 _{2,32} –32 _{2,31}	NH ₂ CHO, C ₂ H ₅ CN

345 GHz, respectively. In order to determine rotation temperatures, we covered at least two transitions for a given species with $E_{\text{up}} < 100$ K and $E_{\text{up}} > 100$ K each. BIS07 used the single pixel receiver B3, the predecessor of HARP-B, together with DAS (Digital Autocorrelation Spectrometer) as the back-end, covering a larger instantaneous bandwidth of 500 MHz. Only the central receptor of HARP-B array is analyzed here as no significant off source emission was detected in the complex molecules.

Data reduction and line fitting were done using the CLASS software package². Line assignments were done by comparison of observed frequencies corrected for source velocity with the JPL³, CDMS⁴ and NIST⁵ catalogs (Pickett et al. 1998; Müller et al. 2005). The line assignment/detection was based on Gaussian fitting with the following criteria: (i) the fitted line position had to be within ± 1 MHz of the catalog frequency; (ii) the FWHM had to be consistent with those given in Table 1; and (iii) a $S/N > 3$ is required on the peak intensity. Section A.1 in the Appendix describes in more detail the error analysis. In general, our errors on the integrated intensities are conservative and suggest a lower S/N than that on the peak intensity or obtained using more traditional error estimates.

3. Data analysis

3.1. Rotation diagrams

Rotation temperatures and column densities were obtained through the rotation diagram (RTD) method (Goldsmith & Langer 1999), when 3 or more lines are detected over a sufficiently large energy range. Integrated main-beam temperatures, $\int T_{\text{MB}} dV$, can be related to the column density in the upper energy level by:

$$\frac{N_{\text{up}}}{g_{\text{up}}} = \frac{3k \int T_{\text{MB}} dV}{8\pi^3 \nu \mu^2 S}, \quad (2)$$

where N_{up} is the column density in the upper level, g_{up} is the degeneracy of the upper level, k is Boltzmann's constant, ν is the transition frequency, μ is the dipole moment and S is the line

strength. The total *beam-averaged* column density N_{T} in cm^{-2} can then be computed from:

$$\frac{N_{\text{up}}}{g_{\text{up}}} = \frac{N_{\text{T}}}{Q(T_{\text{rot}})} e^{-E_{\text{up}}/T_{\text{rot}}}, \quad (3)$$

where $Q(T_{\text{rot}})$ is the rotational partition function, and E_{up} is the upper level energy in K.

Blended transitions of a given species with similar E_{up} ($\Delta E_{\text{up}} < 30$ K) were assigned intensities according to their Einstein coefficients (A) and upper level degeneracies (g_{up}):

$$\int T_{\text{MB}} dV(i) = \int T_{\text{MB}} dV \times \frac{A^i g_{\text{up}}^i}{\sum_i A^i g_{\text{up}}^i}, \quad (4)$$

where the summation is over all the contributing transitions. Blended transitions with different E_{up} , or with contamination from transitions belonging to other species, were excluded from the RTD fit.

The beam averaged column density, N_{T} , is converted to the *source-averaged* column density N_{S} using the beam-filling factor η_{BF} :

$$N_{\text{S}} = \frac{N_{\text{T}}}{\eta_{\text{BF}}}. \quad (5)$$

The standard RTD method assumes that the lines are optically thin. Lines with strong optical depth, determined from the arguments in Sect. 4.1 as well as the models discussed in Sect. 3.2, were excluded from the fit. For CH₃OH, also low- E_{up} lines arising from a cold extended component (see Sect. 4.4.1) were excluded.

Differential beam dilution is taken into account by multiplying the line intensities in the 230 GHz range by $\eta_{\text{BF}}(340 \text{ GHz})/\eta_{\text{BF}}(230 \text{ GHz})$. For warm and cold molecules, beam dilution is derived assuming source diameters $\theta_{T=100 \text{ K}}$ (see Eq. (10) below) and $14''$, respectively (see Eq. (1)). All emission is thus assumed to be contained within the smallest beam size. Same approach was used in BIS07.

The vibrational partition function was ignored assuming that all emission arises from the ground vibrational state so that the vibrational partition function can be set to unity. This approximation can cause an underestimation of the derived column densities for larger molecules, even at temperatures of 100–200 K. Indeed, Widicus Weaver et al. (2005) show that the error can be up to a factor of 2 for temperatures up to 300 K. Since this approximation affects all complex molecules (albeit at different levels), the overall error in abundance ratios will be less than a factor of 2 and well within the other uncertainties.

² CLASS is part of the GILDAS software package developed by IRAM.

³ <http://spec.jpl.nasa.gov/ftp/pub/catalog/catform.html>

⁴ <http://www.ph1.uni-koeln.de/vorhersagen>

⁵ <http://physics.nist.gov/PhysRefData/Micro/Html/contents.html>

3.2. Spectral modeling

The alternative method for analysing the emission is to model the observed spectra directly. For this purpose, we used the so-called “Weeds” extension of the CLASS software package⁶, developed to facilitate the analysis of large millimeter and sub-millimeter spectral surveys (Maret et al. 2011). In this model, the excitation of a species is assumed to be in LTE (Local Thermodynamic Equilibrium) at a temperature T_{ex} . The brightness temperature, T_{B} , of a given species as a function of the rest frequency ν is then given by:

$$T_{\text{B}}(\nu) = \eta_{\text{BF}} \left[J_{\nu}(T_{\text{ex}}) - J_{\nu}(T_{\text{bg}}) \right] (1 - e^{-\tau(\nu)}), \quad (6)$$

where η_{BF} is the beam filling factor (see Eq. (1)), J_{ν} is the radiation field such that:

$$J_{\nu}(T) = \frac{h\nu/k}{e^{h\nu/kT} - 1}$$

and T_{bg} is the temperature of the background emission. The HPBW θ_{B} is calculated within the Weeds model as $\theta_{\text{B}} = 1.22c/\nu D$, where c is the speed of light and D is the diameter of the telescope⁷. The opacity $\tau(\nu)$ is:

$$\tau(\nu) = \frac{c^2}{8\pi\nu^2} \frac{N_{\text{T}}}{Q(T_{\text{ex}})} \sum_i A^i g_{\text{up}}^i e^{-E_{\text{up}}^i/kT_{\text{ex}}} (e^{h\nu_0^i/kT_{\text{ex}}} - 1) \phi^i \quad (7)$$

where the summation is over each line i of the considered species. ν_0^i is the rest frequency of the line and ϕ^i is the profile function of the line, given by:

$$\phi^i = \frac{1}{\Delta\nu \sqrt{2\pi}} e^{-(\nu - \nu_0^i)^2 / 2\Delta\nu^2}, \quad (8)$$

where $\Delta\nu$ is the line width in frequency units at $1/e$ height. $\Delta\nu$ can be expressed as a function of the line FWHM in velocity units ΔV by:

$$\Delta\nu = \frac{\nu_0^i}{c \sqrt{8 \ln 2}} \Delta V. \quad (9)$$

The input parameters in the model are the column density N_{S} in cm^{-2} , excitation temperature T_{ex} in K, the source diameter θ_{S} in arcseconds, offset velocity from the source LSR (Local Standard of Rest) in km s^{-1} and the line FWHM in km s^{-1} . All parameters excluding the source diameter are optimized manually to obtain the best agreement with the observed spectra. The source diameter for emission from cold species is allowed to vary over a larger area; generally $14''$ is used. The emission from hot-core molecules targeted in our work is assumed to originate from the central region with $T_{\text{dust}} \geq 100$ K. The source diameter for the warm emission is calculated using a relation derived from dust modeling of a large range of sources (BIS07):

$$R_{T=100 \text{ K}} \approx 2.3 \times 10^{14} \left(\sqrt{\frac{L}{L_{\odot}}} \right) \text{ cm}, \quad (10)$$

where $L/L_{\odot}$ is the luminosity of the source relative to the solar luminosity. Table 3 gives the calculated $R_{T=100 \text{ K}}$ radii and diameters for the observed sources.

⁶ CLASS is part of the GILDAS software package developed by IRAM.

⁷ For JCMT with 15-m antenna diameter, the equation gives a θ_{B} of $21.9''$ and $14.8''$ for the 230 GHz band and 345 GHz bands, respectively.

Table 3. Source radii and angular diameters for $T=100$ K.

Sources	$R_{T=100\text{K}}$ [AU]	$\theta_{\text{S}, T=100 \text{ K}}$ [']
IRAS 20126+4104	1753	2.2
IRAS 18089-1732	2750	2.4
G31.41+0.31	7840	2.0

In the analysis for individual molecules, the initial values for T_{ex} were based on the T_{rot} from the RTD analysis in case of optically thin species. For optically thick species the value of T_{ex} from the ^{13}C -isotope was used. If T_{ex} for the isotopologue could not be obtained, the initial temperature was guessed. The T_{ex} value was then optimized visually based on the relative intensity of the emission lines. The simulated emission was not allowed to exceed the emission of optically thin lines in the spectrum in any of the observed spectral ranges. Coinciding and blended transitions, which together contribute to an optically thick line, are excluded in the analysis.

Due to the visual optimization and the possibility of overlapping lines (particularly in the line-rich source G31), the resulting T_{ex} values are only a rough estimate (± 50 – 100 K) and do not differ significantly from those from the RTD method. The column density for each species was constrained by optically thin, unblended lines, where available.

For the specific case of CH_3OH , which requires two temperature components for a proper fit, the analysis was also done with the CASSIS analysis package⁸. CASSIS has the advantage that it can properly model the emission from overlapping optically thick lines, as well as from nested regions of emission. For CH_3OH , warm emission from the compact inner region may be absorbed by the surrounding colder gas, which can influence the derived hot core abundances.

4. Results

4.1. General results and comparison between sources

The observed sources, IRAS 20126, IRAS 18089 and G31, differ from each other significantly in the observed chemical richness in the JCMT single-dish data. Figure 2 presents two frequency ranges with lines from several observed species. For G31, strong lines of all complex organic species are detected, whereas for IRAS 20126 many targeted lines are below the detection limit. Many complex molecules are also found in IRAS 18089 but with weaker lines than for G31. Among the serendipitous discoveries acetone, CH_3COCH_3 , is possibly seen in G31 and IRAS 18089 for the first time (see Sect. 4.5). Integrated line intensities for all detected lines and selected upper limits are given in Tables A.1–A.14. The rotation diagrams for the detected species are shown in Figs. B.1–B.13 whereas the optimized parameters in the Weeds model for each molecule and source are available online in Tables C.1–C.3.

4.2. Optical depth determinations

To assess the importance of optical depths effects, the ratio of lines of different isotopologues are compared. The expected

⁸ CASSIS has been developed by IRAP-UPS/CNRS (<http://cassis.irap.omp.eu>).

Fig. 2. Spectral ranges 238.83–239.26 GHz and 339.94–340.18 GHz with lines from several targeted species for the three sources.

Table 4. Isotopologue line intensity ratios in the observed sources.

Species	IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
$\text{CH}_3\text{OH}/^{13}\text{CH}_3\text{OH}$	>6	>18	6
$\text{CH}_3\text{CN}/\text{CH}_3^{13}\text{CN}$	>65	4.7	4.5

Notes. Lower limits are those for which the ^{13}C -isotopologue was not detected.

$^{12}\text{C}/^{13}\text{C}$ isotope ratio depends on the galactocentric radius, R_{GC} , according to Eq. (11) (Wilson & Rood 1994).

$$^{12}\text{C}/^{13}\text{C} = (7.5 \pm 1.9) R_{\text{GC}} [\text{kpc}] + (7.6 \pm 12.9). \quad (11)$$

The isotope ratios derived for our sources using Eq. (11) are given in Table 1. The galactocentric radii were calculated trigonometrically from the galactic coordinates, using the IAU value for the distance to the Galactic center $R_0 = 8.5$ kpc (Kerr & Lynden-Bell 1986).

Table 4 lists the observed isotopologue intensity ratios for the most abundant species in our sources. The $\text{CH}_3\text{OH}/^{13}\text{CH}_3\text{OH}$ ratios are derived from a low-energy transition $2_{2,0,+0}-3_{1,3,+0}$ with $E_{\text{up}} = 44.6$ K. No high E_{up} transitions were reliably detected for $^{13}\text{CH}_3\text{OH}$ in our sources in our standard settings and the low E_{up} ratios are therefore taken to apply to cold methanol. For G31.41+0.31, one additional observation was carried out to cover a transition with $E_{\text{up}} > 100$ K. A $\text{CH}_3\text{OH}/^{13}\text{CH}_3\text{OH}$ intensity ratio of 4.0 is derived for a transition with $E_{\text{up}} \approx 210$ K ($13_{1,12,-0}-13_{0,13,+0}$). This indicates that also warm methanol is optically thick in this source.

The $\text{CH}_3\text{CN}/\text{CH}_3^{13}\text{CN}$ ratios are derived from the 13_3-12_3 line intensities for G31 and IRAS 18089 and indicate that CH_3CN is optically thick in these two sources, but not in IRAS 20126. For H_2CO and HNCO , isotopologue lines are detected but for different transitions than the main isotopologues. Thus, a model is needed to infer the optical depths. Fits to each of the isotopologues independently at a fixed temperature of 150 K using the $R_{T=100\text{ K}}$ source size gives column density ratios that are significantly smaller than the overall isotope ratios, suggesting that these species are also optically thick for G31 and IRAS 18089. In practice, we have excluded the optically thick lines (as indicated by the Weeds model) from the RTD fits for all species.

4.3. Temperatures

Table 5 summarizes the derived temperatures from the RTD fit for the various species. As also found by BIS07, molecules can be classified into *cold* (<100 K) and *warm* (>100 K), and our categorization is similar to theirs. The Weeds analysis is consistent with this grouping. There is however variation in temperatures within the groups, with warm species having rotation temperatures from 70 to 300 K, and cold species from 40 to 100 K. Some variation is seen in rotation temperatures of individual species between different sources; the rotation temperatures are generally higher in G31 than in IRAS 18089, while IRAS 20126 has the lowest of the three.

The T_{rot} value for CH_3OH is ~ 300 K for G31 and IRAS 18089 and ~ 100 K for IRAS 20126. For G31, several lines with $E_{\text{up}} > 400$ K are detected, which makes the RTD fit more robust. For IRAS 18089 and IRAS 20126 the accuracy of the RTD fits suffers from the small range of E_{up} in the detected transitions. In addition to optically thick lines, low- E_{up} lines have been excluded from the fit. These lines are underestimated by the RTD fit and probably originate from a colder extended region also seen in the ^{13}C lines (Fig. 3). The T_{rot} from the RTD analysis therefore represents the warm CH_3OH alone. See Sect. 4.4.1 for a more detailed discussion on the CH_3OH emission.

For CH_3CN , the T_{rot} values range from ~ 200 and ~ 350 K for the three sources. The $\text{CH}_3^{13}\text{CN}$ rotation diagram gives a value of T_{rot} of only 53 K, however (Fig. 3). This illustrates the large uncertainties at high temperatures for optically thick species and the possibility of a cold component in addition to the warm one.

Contrary to the general trend, the T_{rot} value for H_2CO is somewhat higher (204 K) in IRAS 18089 than in G31 (157 K). The discrepancy could be influenced by the small number of lines used for the analysis. All fitted lines are those belonging to the para- H_2CO , and the T_{rot} fits are thus not affected by fluxes from different spin states. A single transition of

Table 5. Temperatures (K) derived from the RTD analysis and the Weeds or CASSIS (CH₃OH) model.

	IRAS 20126+4104		IRAS 18089-1732		G31.41+0.31	
	RTD	Model	RTD	Model	RTD	Model
warm species						
H ₂ CO	123 ± 21	150	204 ± 82	(150)	157 ± 44	(150)
CH ₃ OH	122 ± 17	300, 14 ± 1	291 ± 37	300, 15 ± 2	323 ± 34	200, 14 ± 2
C ₂ H ₅ OH	–	(100)	85 ± 18	150	120 ± 15	100
HNCO	–	(200)	92 ± 25	200	111 ± 32	200
NH ₂ CHO	–	300	72 ± 28	100	94 ± 50	300
CH ₃ CN	217 ± 352	(200)	[346 ± 106]	(200)	[311 ± 68]	(300)
C ₂ H ₅ CN	–	(80)	84 ± 33	80	105 ± 23	80
HCOOCH ₃	–	(200)	118 ± 20	200	174 ± 11	300
CH ₃ OCH ₃	–	(100)	66 ± 11	100	90 ± 6	100
cold species						
CH ₂ CO	–	(50)	71 ± 11	50	97 ± 101	50
CH ₃ CHO	–	(50)	–	(50)	–	50
HCOOH	–	(40)	–	(40)	–	40
CH ₃ CCH	40 ± 10	35	46 ± 12	40	67 ± 14	80

Notes. The species are classified *warm* and *cold* according to BIS07. Square bracketed values are T_{rot} values for optically thick species and round bracketed values are T_{ex} values assumed based on temperatures derived from the other sources in this study. – Means not enough lines were detected for a rotation diagram. Typical uncertainties in the Weeds excitation temperatures are ± 50 K.

Fig. 3. Rotation diagrams for ¹³CH₃OH (*left panel*) and CH₃¹³CN (*right panel*) in G31. Triangles represent blended lines and are not included from the fit.

H₂¹³CO (3_{1,2}–2_{1,1}) was covered, and no information on the excitation temperature can therefore be obtained from the minor isotopologue.

The T_{rot} values of the other complex species, HNCO, C₂H₅OH, C₂H₅CN, NH₂CHO and CH₃OCH₃ are around 100 K and are slightly higher in G31 than in IRAS 18089. The T_{rot} of HCOOCH₃ stands out in both sources, in G31 being closer to 200 K. No lines belonging to these species were detected in IRAS 20126.

The species classified as cold by BIS07, CH₂CO and CH₃CCH, indeed show cold rotation temperatures in all sources. Not enough lines of CH₃CHO or HCOOH, which are also classified as cold in BIS07, are observed in our sources for making rotation diagrams.

4.4. Column densities

Table 7 presents the column densities derived from the RTD analysis, Weeds or CASSIS model, and from the ¹³C-isotopologues for the optically thick species. Following BIS07, the column densities for warm molecules are given as source-averaged values (see Eq. (5)). The emission from cold molecules extends over a larger volume and the values are given

as beam-averaged column densities. Typical uncertainty of the column densities derived from the RTD analysis is $\sim 40\%$.

4.4.1. CH₃OH

An accurate determination of the CH₃OH column density is essential for comparing the abundance ratios of complex organic species. For hot-core molecules, it is particularly important to quantify the warm CH₃OH emission. The column densities of CH₃OH in BIS07 were determined by the RTD method excluding the optically thick lines. The same is done in our analysis. Our rotation diagrams however also show emission from low- E_{up} transitions, which are strongly underestimated by the RTD fit on the warm lines, providing further evidence for the presence of a colder component. We have therefore also excluded these transitions. The fit to the higher E_{up} , optically thin lines should give the warm CH₃OH column density obtained in a similar way as BIS07.

CH₃OH emission was also simulated using a two-component CASSIS model. A single-component model is not able to simultaneously reproduce the warm and cold lines without overestimating the lines from intermediate energy levels. Indeed, a better agreement is obtained using a two-component model, consisting of a warm compact component and a cold extended component. Table 6 shows the best model parameters obtained from the fitting. The warm components are fixed to $\theta_{T=100\text{ K}}$ while the cold component is allowed to extend beyond the beam diameter. The warm CH₃OH column densities derived from the CASSIS fit are in agreement with the values derived from the RTD analysis. The best fits plotted onto the CH₃OH 7_K–6_K transitions (338.5 GHz) are shown in Fig. 4.

Several ¹³CH₃OH lines are detected in G31. However, only low- E_{up} lines are reliably detected since the high- E_{up} lines are very weak or blended. Assigning these lines to the cold component and assuming $T_{\text{rot}} = 20$ K (Öberg et al. 2011; Requena-Torres et al. 2008), the beam-averaged

Table 6. CASSIS model parameters for CH₃OH in the observed sources.

Source	Component ^a	N_S [cm ⁻²]	T_{ex} [K]	$FWHM$ [km s ⁻¹]	θ_S [$''$]	V_{LSR} [km s ⁻¹]	χ^2
IRAS 20126+4104	comp.	$(1.1 \pm 0.1)\text{E}+17$	300 ^c	8.15 ± 0.60	2.2 ^b	-4.05 ± 0.70	1.9
	ext.	$(2.2^{+0.9}_{-0.7})\text{E}+15$	14 ± 1	$7.00^{+0.60}_{-0.75}$	$\gg 14.0$	$-4.25^{+0.80}_{-0.25}$	
IRAS 18089-1732	comp.	$(2.0^{+0.1}_{-0.2})\text{E}+17$	300 ^c	$5.45^{+0.40}_{-0.5}$	2.4 ^b	33.4 ± 0.2	4.4
	ext.	$(2.4^{+1.4}_{-0.5})\text{E}+15$	15 ± 2	$3.50^{+0.50}_{-0.55}$	$\gg 14.0$	$32.3^{+0.4}_{-0.2}$	
G31.41+0.31	comp.	$(1.0 \pm 0.2)\text{E}+18$	200 ^c	6.35 ± 0.35	2.0 ^b	97.2 ± 0.2	7.4
	ext.	$(1.2 \pm 0.6)\text{E}+16$	14 ± 2	3.95 ± 0.40	$\gg 14.0$	$97.2^{+0.2}_{-0.3}$	

Notes. ^(a) “comp.” for warm, compact component; “ext.” for colder, more extended component. ^(b) Fixed to $\theta_{S,T=100\text{ K}}$. ^(c) Fixed.

Fig. 4. CASSIS two-component models for CH₃OH emission in the 338 GHz region covering the 7_K–6_K transitions with $K = 0$ –6.

¹³CH₃OH column density is $2.1 \times 10^{15} \text{ cm}^{-2}$, corresponding to a CH₃OH column density of $1.7 \times 10^{17} \text{ cm}^{-2}$ for the 14 $''$ volume. This is more than an order of magnitude higher than found in the Weeds model for the cold component, supporting the optically thick interpretation of the cold component. Similarly, for the case of IRAS 18089, the detected ¹³CH₃OH lines may indicate a high optical depth.

As mentioned before, the ¹³CH₃OH line with $E_{\text{up}} = 211 \text{ K}$, covered in additional observations for G31, reveals optical thickness in the warm component as well. The CH₃OH column density derived from this line assuming temperature of 300 K is $2.7 \times 10^{18} \text{ cm}^{-2}$ and the CASSIS fit as well as the RTD analysis underestimate the warm column density by a factor of ~ 2.5 . In principle, the CASSIS analysis could be made consistent with the ¹³CH₃OH value by letting the warm source size vary as well. However, since such information is not known for other molecules, as well as for consistency with BIS07, we have chosen to keep the warm source size fixed at the 100 K radius.

In summary, it seems plausible that at least in our sources, methanol emission arises from two temperature components, a warm ($T_{\text{ex}} \approx 300 \text{ K}$) compact component and a cold ($T_{\text{ex}} \approx 20 \text{ K}$) significantly more extended component. At least for G31, the CH₃OH emission is optically thick in both components.

4.4.2. Other molecules

Overall, the column densities of the various species follow the same pattern in all sources, and hence seem to be well correlated

relative to each other. CH₃OH is the most abundant complex molecule in all sources. The other species have in general half to one order of magnitude lower column densities.

The CH₃CN emission is optically thick for G31 and IRAS 18089. Column densities from the RTD analysis are therefore underestimated. Due to a lack of sufficient optically thin lines, also the Weeds analysis underestimates the column densities. Values derived from the ¹³C-isotopologues are thus more reliable, even though some fraction may arise from a colder component. Indeed, the column density for CH₃CN derived from the 13-isotopologue is an order of magnitude larger than that from the RTD of the main isotopologue alone. The same procedure was used by BIS07.

For H₂CO, the best estimates come from the RTD analysis on the optically thin para-H₂CO lines. The derived H₂CO column densities assuming T_{rot} from Table 5 and a statistical ortho-to-para ratio of 3 are given in Table 7. The column densities derived from the ¹³C-isotopologue of ortho-H₂CO are still larger than from RTD analysis using optically thin lines, which indicates either larger ortho-to-para ratio in these sources (ratio of ~ 3 to 5 has been predicted for cold clouds by Kahane et al. 1984) or non LTE excitation. BIS07 derived the H₂CO column densities from the ¹³C-isotopologue.

The column density derived for HCOOCH₃ from the RTD diagram is significantly smaller than that derived from the Weeds model. The HCOOCH₃ emission is stronger in the 230 GHz beam (20–21 $''$) than in the 345 GHz beam (14 $''$) probably indicating significant extended emission. The RTD analysis was performed on the entire dataset, while only lines in the 230 GHz band was used in the Weeds modeling.

5. Discussion

5.1. Comparison to massive YSOs without a disk structure

In order to see the effect of a flattened circumstellar structure on the chemistry around the YSO, we compare the temperatures and abundances of the complex organic molecules in sources studied in this work and those studied in BIS07. In addition to our sources (IRAS 20126, IRAS 18089 and G31), AFGL 2591, NGC 7538 IRS1 and G24.78 from the BIS07 sample are now known to have disk structures as well (van der Tak et al. 2006; Wang et al. 2012; Pestalozzi et al. 2004, 2009; Knez et al. 2009; Beltrán et al. 2005). A disk-like structure may be present also for other sources but current evidence is not as strong as for the above sources. For the comparison with the BIS07 sample, we use the results primarily from the RTD method to exclude method based differences.

Table 7. Column densities for the targeted species from the RTD analysis, Weeds or CASSIS (CH₃OH) models and those derived from ¹³C-isotopologues.

Species	IRAS 20126+4104			IRAS 18089-1732			G31.41+0.31		
	RTD	Model	$N(^{12}\text{C})^a$	RTD	Model	$N(^{12}\text{C})^a$	RTD	Model	$N(^{12}\text{C})^a$
H ₂ CO ^b	8.4E+15	6.0E+15	<4.7E+16	1.9E+16	1.5E+16	6.3E+16	6.5E+16	6.0E+16	9.6E+16
CH ₃ OH	2.9E+16	1.1E+17	<4.0E+17 ^c	2.2E+17	2.0E+17	2.0E+18 ^c	1.2E+18	1.0E+18	2.7E+18 ^d
C ₂ H ₅ OH	–	<0.3E+16	–	3.9E+16	2.5E+16	–	1.5E+17	1.0E+17	–
HNCO	–	1.0E+15	<7.0E+15	2.5E+15	0.4E+16	<3.8E+16	3.6E+15	0.6E+16	<8.2E+16
NH ₂ CHO	–	0.3E+15	–	8.4E+14	0.5E+15	–	2.1E+15	0.4E+16	–
CH ₃ CN	1.5E+15	1.3E+15	<1.4E+15	>4.7E+15	>3.5E+15	2.2E+16	>1.6E+16	>2.0E+16	1.6E+17
C ₂ H ₅ CN	–	<0.5E+15	–	3.1E+15	0.4E+16	–	2.0E+16	2.0E+16	–
HCOOCH ₃	–	<0.2E+16	–	2.8E+16	3.0E+16	–	1.7E+17*	1.0E+18	–
CH ₃ OCH ₃	–	<0.1E+17	–	1.3E+17	1.0E+17	–	5.4E+17	5.0E+17	–
CH ₂ CO	–	<2.5E+13	–	7.9E+13	7.5E+13	–	–	3.3E+14	–
CH ₃ CHO	–	<0.5E+14	–	–	<0.5E+14	–	–	5.0E+14	–
HCOOH	–	<2.5E+13	–	–	<2.5E+13	–	3.2E+15	1.0E+15	–
CH ₃ CCH	2.7E+14	3.5E+14	–	5.2E+14	0.9E+15	–	1.0E+15	1.4E+15	–

Notes. Column densities for the warm molecules are source-averaged, while those for cold molecules are beam-averaged. Column densities where optical depth has a significant effect are labeled as lower limits. ^(*) Rotation diagram fit on HCOOCH₃ on all lines, while Weeds model on the 230 GHz lines only. ^(a) $N(^{12}\text{C})$ obtained from the $N(^{13}\text{C})$ adopting a ¹²C/¹³C ratio equal to 70, 54 and 41 for IRAS 20126, IRAS 18089 and G31.41, respectively. ^(b) Column density from ¹³C-isotopologue from ortho-H₂¹³CO, corrected using the statistical ortho to para ratio of 3:1. ^(c) Calculated from the line intensity of transition 2_{2,0,+0}–3_{1,3,+0} at 345.084 GHz with $E_{\text{up}} = 44.6$ K, assuming a T_{rot} of 300 K. ^(d) Calculated from the line intensity of transition 13_{1,12,-0}–13_{0,13,+0} at 341.132 GHz with $E_{\text{up}} = 206$ K, assuming a T_{rot} of 300 K.

Fig. 5. Source-averaged column densities for the targeted warm species in $\theta_{\text{T}}=100$ K volume. Column densities from the RTD analysis are marked in black bars. The red and blue bars show column densities from the Weeds or CASSIS (CH₃OH) models and from ¹³C-isotopologue, respectively. Arrows indicate upper limits.

5.1.1. Rotation temperatures

Figure 7 shows the rotation temperatures of the complex species for the observed sources together with those from BIS07. As in BIS07, we find that the complex molecules can be divided into warm and cold species based on their rotation temperatures. The rotation temperatures obtained for the molecules in our sources agree with the division; CH₃OH, H₂CO, C₂H₅OH, HCOOCH₃, CH₃OCH₃, HNCO, NH₂CHO, CH₃CN and C₂H₅CN show rotation temperatures generally ≥ 100 K while CH₂CO, CH₃CHO, HCOOH and CH₃CCH have rotation temperatures < 100 K.

The rotation temperatures of the cold molecules show a very small scatter (± 25 K) from source to source, with or without a disk structure, ignoring the one outlier for HCOOH. Of the

warm species, C₂H₅OH has a consistent rotation temperature of 100–150 K from source to source. The RTD analysis for these species is reliable due to low optical depths and lack of anomalous excitation.

The other warm species show more scatter in the derived rotation temperatures. This is particularly the case for CH₃OH and CH₃CN, which have rotation temperatures ranging from 100 to 350 K. No systematic difference is seen between the two source types. Moreover, the HCOOCH₃ rotation temperatures for AFGL 2591 and NGC 7538 IRS1 are similar to the diskless sources, disproving any difference between the two source types. The scatter in the rotation temperatures could indicate that they exist in environments with different

Fig. 6. Beam-averaged column densities for the targeted cold species. Column densities from the RTD analysis are marked in black bars while the red bars show column densities from the Weeds model. Arrows indicate upper limits.

temperatures, but it may also be caused by optical depth effects (H₂CO, HNCO, CH₃OH and CH₃CN) and (anomalous) radiative excitation. The latter has been previously seen for, e.g., NH₂CHO and HCOOCH₃ (Nummelin et al. 2000) as well as HNCO (Churchwell et al. 1986). For species with significant radiative excitation, e.g., HCOOCH₃, the scatter of the data points in the RTD plots is higher and results in additional uncertainty since the linear fits do not properly capture this scatter.

5.1.2. Column densities

Figures 8 and 9 show the column densities for the targeted species in our sources and those from BIS07 for warm and cold molecules, respectively. For the warm species the column densities within the sources vary by 1–2 orders of magnitude. There is also a significant variation between sources. G31 is chemically richest of the sources, with highest column densities of all the targeted species, compared to other sources. IRAS 20126 is among the chemically poorest sources. The pattern of column densities is remarkably similar, however, and the disk sources do not stand out.

5.1.3. Abundance ratios

Table 8 gives the abundance ratios for the targeted species with respect to CH₃OH for oxygen-bearing species and HNCO for nitrogen-bearing species. Column densities from the RTD analysis are primarily used. In cases where column densities are not available from the RTD analysis (mainly for IRAS 20126) the values are taken from the Weeds analysis.

The resulting abundance ratios are presented in Fig. 10. Values from the RTD method are shown in black bars and can thus be directly compared to BIS07, whereas those from the Weeds model and ¹³C-isotopologue are in red and blue bars. In general, the abundance ratios in different sources have larger variations within the source types than between them. For example, H₂CO has both lowest and highest abundance ratio for two disk candidates, G31 and IRAS 20126, respectively. The H₂CO results depend somewhat on the analysis method used.

C₂H₅OH, HCOOCH₃ and CH₃OCH₃ abundances peak for G31 and IRAS 18089, and are generally lower for other sources. CH₃OCH₃ also peaks for some diskless sources. The N-bearing species NH₂CHO has a large variation in the abundance ratio

with respect to HNCO. Among our sources, NH₂CHO peaks for G31, the source with largest abundance of O-bearing species, and the least clear disk structure.

Figure 11 shows the abundance ratios of N-bearing species with respect to CH₃OH. Again, a lot of variation is seen within each source type, with no specific trend found for sources with a disk-like structure.

We stress that the absolute inferred column densities and abundance ratios are uncertain by a factor of a few up to an order of magnitude, as already indicated by the different analysis techniques. An independent assessment of the accuracy of the results can be made by comparison with the inferred column densities and abundances of Zernickel et al. (2012) for NGC 6334 I (=NGC 6334 IRS), who observed a completely different set of lines of the same molecules with *Herschel*-HIFI and the SMA in beams ranging from 2–40''. Comparison with the results of BIS07 shows good agreement within a factor of a few for the abundances of several species (CH₃OH, CH₃OCH₃, CH₃OCHO, C₂H₅OH, C₂H₅CN) whereas others differ by an order of magnitude (H₂CO, CH₃CN). The species that show the largest discrepancy are those with highly optically thick lines and without a large set of isotopologue lines. Thus, a combination of differences in adopted source sizes and optical depth effects can account for the discrepancies. Because our approach is the same for all sources, the relative values from source to source should be more reliable.

5.2. Chemical and physical implications

The main goal of this study is to investigate similarities and differences between sources with and without a disk-like structures. The presence of a flattened accretion disk should allow the UV radiation from the central object to escape more readily and then impinge on the gas and dust in the outer walls of a flared disk or outflow cavity (Bruderer et al. 2010). Increased UV radiation could manifest itself as enhancement in the complex organic species produced through UV photodissociation from CH₃OH in the solid state, such as CH₃OCH₃, C₂H₅OH and CH₃OCHO (Öberg et al. 2009). In this scenario, CH₃OH is photodissociated into various radicals such as CH₃, CH₃O and CH₂OH which can become mobile at higher ice temperatures (20–40 K) and form the observed complex organic molecules (Garrod & Herbst 2006). Higher temperatures favor diffusion of larger radicals resulting in the formation of larger complex organic molecules compared with small molecules like H₂CO. Another related parameter that plays a role is the CO content of ices, with H₂CO, CHO- and COOH-containing species enriched in cold, CO-rich CH₃OH ices in which CO has not evaporated. Indeed Öberg et al. (2011) show that the increased abundance of HCOOCH₃ in low-mass YSOs compared with the high-mass sample of BIS07 may be due to this effect. Although only limited laboratory experiments exist on N-containing molecules, the abundance of N-bearing complex organic molecules could be enhanced due to photodissociation of N₂ or NH₃.

Our results show that there is no consistent enhancement of complex organic species produced through CH₃OH photochemistry, nor of N-bearing species, that can be attributed to the presence of a flattened disk structure on the scales probed by our data. The variations within source types are larger than between the source types. One explanation for this lack of differentiation could be that all high-mass sources have a source structure through which UV radiation can impact the larger surroundings. In particular, just like low-mass sources, all high-mass YSOs are expected to have outflows and cavities through

Fig. 7. Rotation temperatures for selected species in massive YSOs with (open stars) and without (asterisks) observed disk structure.

Fig. 8. Source-averaged column densities for warm molecules. Column densities from the RTD analysis are marked in black bars. Sources without disk structure are marked with an asterisk. The red and blue bars show column densities from the Weeds or CASSIS (CH_3OH) models and from ^{13}C -isotopologue, respectively. Upper limits are indicated with arrows.

which UV radiation can escape and affect the chemistry, whether or not they have a large disk-like structure. This scenario can be tested with future high angular resolution data on $<1''$ scale with ALMA which should then reveal the emission from complex molecules coating the walls of the outflow cavities.

An alternative explanation is that enhanced temperature in the YSO environment is not needed for the production of complex organic molecules and that they are already formed in the prestellar stage through UV- and cosmic-ray processing of cold ices, using just the internal UV field produced by interaction of cosmic rays with H_2 (Prasad & Tarafdar 1983) rather than that of the star. This scenario has gained support over the last few years

with the detection of HCOOCH_3 and CH_3OCH_3 in cold low-mass cores away from the YSO where the molecules are either released by shocks (Arce et al. 2008; Öberg et al. 2010) or by photodesorption (Bacmann et al. 2012) although the precise formation mechanisms are still unclear. Gas-phase processes may also contribute in some cases. Again, high spatial resolution and sensitivity such as ALMA will be key to testing this scenario.

6. Summary and conclusions

We have carried out a partial submillimeter line survey, targeting several complex organic species toward selected massive

Fig. 9. Beam-averaged column densities for cold molecules in sources observed in this study and those from BIS07. Sources without disk structure are marked with an asterisk. Column densities from the RTD analysis are marked in black bars while the red bars show column densities from the Weeds model. Upper limits are indicated with arrows.

Table 8. Abundance ratios of complex species in the observed sources and those in other chemically rich environments.

Source	H ₂ CO	C ₂ H ₅ OH	NH ₂ CHO	HCOOCH ₃	CH ₃ OCH ₃	Refs.
Massive protostellar objects with a disk structure						
IRAS 20126+4104	0.29 ^p	<0.10	<0.30	<0.34	<0.34	this work
IRAS 18089-1732	0.09 ^p	0.18	0.34	0.13	0.60	this work
G31.41+0.31	0.06 ^p	0.12	0.59	0.14	0.46	this work
AFGL 2591	0.28 ^o	<0.02	<1.00	0.51	<0.16	1
NGC 7538 IRS1	0.21 ^o	0.05	0.25	0.12	<0.13	1
G24.78	0.23 ^o	0.03	0.14	0.11	0.43	1
Massive protostellar objects with no detected disk structure						
G75.78	0.20 ^o	<0.02	0.05	0.06	0.21	1
W 33A	0.27 ^o	0.02	0.32	0.13	0.14	1
NGC 6334 IRS1	0.13 ^o	0.02	0.17	0.12	0.60	1
W 3(H ₂ O)	0.18 ^o	0.01	0.27	0.05	0.15	1
Low-mass outflows						
B1-b core	1.2	<1.0	–	2.3	<0.8	2
SMM4-W	0.6	<0.4	–	3.5	1.1	3
L1157 outflow	–	0.7	–	1.8	–	4, 5
Low-mass protostars						
SMM1	6.6	<3.4	–	10	5.3	3
SMM4	2.2	<0.6	–	<1.0	0.8	3
NGC1333 IRAS 4A env.	–	–	–	56	<22	6, 7, 8
NGC1333 IRAS 4B env.	–	–	–	26	<19	6, 7, 8
IRAS 16293 env.	4	–	–	30	20	8, 9, 10
Low-mass hot corinos						
IRAS 16293 A	<0.02	1.4	–	0.6	0.6	11, 12
IRAS 16293 B	0.6	–	–	0.8	1.6	11, 12
NGC 1333 IRAS 2A	–	–	–	–	2	13, 14

Notes. The abundances are column densities with respect to CH₃OH for oxygen-bearing species and with respect to HNCO for nitrogen-bearing species. The CH₃OH and HNCO column densities are taken from the RTD analysis when available. ^(p) Para-H₂CO. ^(o) Ortho-H₂CO.

References. (1) Bisschop et al. (2007); (2) Öberg et al. (2010); (3) Öberg et al. (2011); (4) Bachiller & Perez Gutierrez (1997); (5) Arce et al. (2008); (6) Maret et al. (2005); (7) Bottinelli et al. (2007); (8) Herbst & van Dishoeck (2009); (9) van Dishoeck et al. (1995); (10) Cazaux et al. (2003); (11) Kuan et al. (2004); (12) Bisschop et al. (2008); (13) Huang et al. (2005); (14) Jørgensen et al. (2005).

Fig. 10. Abundance ratios of complex species with respect to CH_3OH for oxygen-bearing species and HNCO for nitrogen-bearing species. The black bars indicate abundance ratios calculated with the CH_3OH and HNCO column density derived from the RTD analysis similar to BIS07. The red and blue bars indicate abundance ratios where CH_3OH and HNCO column densities are derived from CASSIS or Weeds (HNCO) model, respectively. Upper limits are marked with arrows.

Fig. 11. Abundance ratios of N-bearing species HNCO , CH_3CN , $\text{C}_2\text{H}_5\text{CN}$ and NH_2CHO with respect to CH_3OH (from RTD analysis). Arrows indicate upper limits.

YSOs with strong evidence of circumstellar accretion disks, IRAS 20126+4104, IRAS 18089-1732 and G31.41+0.31. This is the first time that molecular abundances are reported for these well-known sources. The analysis is performed using both the rotation diagram method and spectral modeling. The inferred rotation temperatures and molecular abundances are compared to sources without reported disk structures analyzed using the same techniques. The molecules can be divided into two different groups based on their rotation temperatures, independent of source type. In particular, the cold (<100 K) species have remarkably constant rotation temperatures from source to source. The warm (>100 K) species exhibit more scatter possibly due to optical depth effects, non-LTE conditions and radiative excitation. The column densities peak for the same sources, with G31.41+0.31 being chemically the richest of the studied sources.

The relative abundances of species follow the same pattern, and no chemical differentiation could be established between the two source types within the (considerable) uncertainties. The lack of chemical differentiation between massive YSOs with and without observed disk-like structure suggests similarity in the physical conditions and level of UV exposure for all sources, for example through outflow cavities. Alternatively, the complex molecules may already be formed in the cold prestellar stage under similar conditions. Ultimately and indirectly, these results imply that the mechanism of the formation of massive stars does not differ significantly from source to source. Future high angular and high sensitivity observations of optically thin lines will be key to distinguishing the different scenarios and derive accurate abundance ratios that can be directly compared with chemical models.

Acknowledgements. Astrochemistry in Leiden is supported by the Netherlands Research School for Astronomy (NOVA), by a Spinoza grant from the Netherlands Organisation for Scientific Research (NWO), and by the European Community's Seventh Framework Programme FP7/2007-2013 under grant agreement 238258 (LASSIE) and 291141 (CHEMPPLAN).

References

- Arce, H. G., Santiago-García, J., Jørgensen, J. K., Tafalla, M., & Bachiller, R. 2008, *ApJ*, 681, L21
- Bachiller, R., & Perez Gutierrez, M. 1997, *ApJ*, 487, L93
- Bacmann, A., Taquet, V., Faure, A., Kahane, C., & Ceccarelli, C. 2012, *A&A*, 541, L12
- Belloche, A., Garrod, R. T., Müller, H. S. P., et al. 2009, *A&A*, 499, 215
- Beltrán, M. T., Cesaroni, R., Neri, R., et al. 2005, *A&A*, 435, 901
- Beltrán, M. T., Codella, C., Viti, S., Neri, R., & Cesaroni, R. 2009, *ApJ*, 690, L93
- Benjamin, R. A., Churchwell, E., Babler, B. L., et al. 2003, *PASP*, 115, 953
- Blake, G. A., & Walsh, A. J. 2008, *ApJ*, 673, L55
- Beuther, H., Schilke, P., Sridharan, T. K., et al. 2002, *A&A*, 383, 892
- Beuther, H., Hunter, T. R., Zhang, Q., et al. 2004, *ApJ*, 616, L23
- Beuther, H., Zhang, Q., Sridharan, T. K., & Chen, Y. 2005, *ApJ*, 628, 800
- Bisschop, S. E., Jørgensen, J. K., van Dishoeck, E. F., & de Wachter, E. B. M. 2007, *A&A*, 465, 913 (BIS07)
- Bisschop, S. E., Jørgensen, J. K., Bourke, T. L., Bottinelli, S., & van Dishoeck, E. F. 2008, *A&A*, 488, 959
- Blake, G. A., Sutton, E. C., Masson, C. R., & Phillips, T. G. 1987, *ApJ*, 315, 621
- Bottinelli, S., Ceccarelli, C., Williams, J. P., & Lefloch, B. 2007, *A&A*, 463, 601
- Bruderer, S., Benz, A. O., Doty, S. D., van Dishoeck, E. F., & Bourke, T. L. 2009, *ApJ*, 700, 872
- Bruderer, S., Benz, A. O., Stäuber, P., & Doty, S. D. 2010, *ApJ*, 720, 1432
- Caselli, P., & Ceccarelli, C. 2012, *A&ARv*, 20, 56
- Cazaux, S., Tielens, A. G. G. M., Ceccarelli, C., et al. 2003, *ApJ*, 593, L51
- Cesaroni, R., Churchwell, E., Hofner, P., Walmsley, C. M., & Kurtz, S. 1994a, *A&A*, 288, 903
- Cesaroni, R., Olmi, L., Walmsley, C. M., Churchwell, E., & Hofner, P. 1994b, *ApJ*, 435, L137
- Cesaroni, R., Felli, M., Testi, L., Walmsley, C. M., & Olmi, L. 1997, *A&A*, 325, 725
- Cesaroni, R., Hofner, P., Walmsley, C. M., & Churchwell, E. 1998, *A&A*, 331, 709
- Cesaroni, R., Felli, M., Jenness, T., et al. 1999, *A&A*, 345, 949
- Cesaroni, R., Neri, R., Olmi, L., et al. 2005, *A&A*, 434, 1039
- Cesaroni, R., Galli, D., Lodato, G., Walmsley, C. M., & Zhang, Q. 2007, *Protostars and Planets V*, 197
- Cesaroni, R., Hofner, P., Araya, E., & Kurtz, S. 2010, *A&A*, 509, A50
- Cesaroni, R., Beltrán, M. T., Zhang, Q., Beuther, H., & Fallscheer, C. 2011, *A&A*, 533, A73
- Charnley, S. B., Tielens, A. G. G. M., & Millar, T. J. 1992, *ApJ*, 399, L71
- Charnley, S. B., Kress, M. E., Tielens, A. G. G. M., & Millar, T. J. 1995, *ApJ*, 448, 232
- Churchwell, E., Wood, D., Myers, P. C., & Myers, R. V. 1986, *ApJ*, 305, 405
- Churchwell, E., Walmsley, C. M., & Cesaroni, R. 1990, *A&AS*, 83, 119
- Comoretto, G., Palagi, F., Cesaroni, R., et al. 1990, *A&AS*, 84, 179
- Estalella, R., Mauersberger, R., Torrelles, J. M., et al. 1993, *ApJ*, 419, 698
- Fontani, F., Pascucci, I., Caselli, P., et al. 2007, *A&A*, 470, 639
- Garrod, R. T., & Herbst, E. 2006, *A&A*, 457, 927
- Garrod, R. T., Weaver, S. L. W., & Herbst, E. 2008, *ApJ*, 682, 283
- Gaume, R. A., & Mutel, R. L. 1987, *ApJS*, 65, 193
- Gibb, E., Nummelin, A., Irvine, W. M., Whittet, D. C. B., & Bergman, P. 2000, *ApJ*, 545, 309
- Goldsmith, P. F., & Langer, W. D. 1999, *ApJ*, 517, 209
- Hatchell, J., Thompson, M. A., Millar, T. J., & MacDonald, G. H. 1998, *A&AS*, 133, 29
- Herbst, E., & van Dishoeck, E. F. 2009, *ARA&A*, 47, 427
- Hofner, P., Cesaroni, R., Rodríguez, L. F., & Martí, J. 1999, *A&A*, 345, L43
- Huang, H.-C., Kuan, Y.-J., Charnley, S. B., et al. 2005, *Adv. Space Res.*, 36, 146
- Johnston, K. G., Keto, E., Robitaille, T. P., & Wood, K. 2011, *MNRAS*, 415, 2953
- Jørgensen, J. K., Bourke, T. L., Myers, P. C., et al. 2005, *ApJ*, 632, 973
- Kahane, C., Lucas, R., Frerking, M. A., Langer, W. D., & Encrenaz, P. 1984, *A&A*, 137, 211
- Kerr, F. J., & Lynden-Bell, D. 1986, *MNRAS*, 221, 1023
- Knez, C., Lacy, J. H., Evans, II, N. J., van Dishoeck, E. F., & Richter, M. J. 2009, *ApJ*, 696, 471
- Kuan, Y.-J., Huang, H.-C., Charnley, S. B., et al. 2004, *ApJ*, 616, L27
- Kurtz, S., Cesaroni, R., Churchwell, E., Hofner, P., & Walmsley, C. M. 2000, *Protostars and Planets IV*, 299
- Leurini, S., Schilke, P., Wyrowski, F., & Menten, K. M. 2007, *A&A*, 466, 215
- Maret, S., Ceccarelli, C., Tielens, A. G. G. M., et al. 2005, *A&A*, 442, 527
- Maret, S., Hily-Blant, P., Pety, J., Bardeau, S., & Reynier, E. 2011, *A&A*, 526, A47
- Millar, T. J., Herbst, E., & Charnley, S. B. 1991, *ApJ*, 369, 147
- Moscadelli, L., Cesaroni, R., Rioja, M. J., Dodson, R., & Reid, M. J. 2011, *A&A*, 526, A66
- Müller, H. S. P., Schlöder, F., Stutzki, J., & Winnewisser, G. 2005, *J. Mol. Struct.*, 742, 215
- Nummelin, A., Bergman, P., Hjalmarson, Å., et al. 2000, *ApJS*, 128, 213
- Öberg, K. I., Garrod, R. T., van Dishoeck, E. F., & Linnartz, H. 2009, *A&A*, 504, 891
- Öberg, K. I., Bottinelli, S., Jørgensen, J. K., & van Dishoeck, E. F. 2010, *ApJ*, 716, 825
- Öberg, K. I., van der Marel, N., Kristensen, L. E., & van Dishoeck, E. F. 2011, *ApJ*, 740, 14
- Olmi, L., Cesaroni, R., & Walmsley, C. M. 1996, *A&A*, 307, 599
- Pestalozzi, M. R., Elitzur, M., Conway, J. E., & Booth, R. S. 2004, *ApJ*, 603, L113
- Pestalozzi, M. R., Elitzur, M., & Conway, J. E. 2009, *A&A*, 501, 999
- Pickett, H. M., Poynter, R. L., Cohen, E. A., et al. 1998, *J. Quant. Spec. Radiat. Transf.*, 60, 883
- Prasad, S. S., & Tarafdar, S. P. 1983, *ApJ*, 267, 603
- Requena-Torres, M. A., Martín-Pintado, J., Martín, S., & Morris, M. R. 2008, *ApJ*, 672, 352
- Schilke, P., Groesbeck, T. D., Blake, G. A., & Phillips, T. G. 1997, *ApJS*, 108, 301
- Shepherd, D. S., Yu, K. C., Bally, J., & Testi, L. 2000, *ApJ*, 535, 833
- Sridharan, T. K., Beuther, H., Schilke, P., Menten, K. M., & Wyrowski, F. 2002, *ApJ*, 566, 931
- Sridharan, T. K., Williams, S. J., & Fuller, G. A. 2005, *ApJ*, 631, L73
- Sutton, E. C., Blake, G. A., Masson, C. R., & Phillips, T. G. 1985, *ApJS*, 58, 341
- Tielens, A. G. G. M., & Charnley, S. B. 1997, *Origins of Life and Evolution of the Biosphere*, 27, 23
- Tielens, A. G. G. M., & Hagen, W. 1982, *A&A*, 114, 245
- Tofani, G., Felli, M., Taylor, G. B., & Hunter, T. R. 1995, *A&AS*, 112, 299
- van der Tak, F. F. S., Walmsley, C. M., Herpin, F., & Ceccarelli, C. 2006, *A&A*, 447, 1011
- van Dishoeck, E. F., Blake, G. A., Jansen, D. J., & Groesbeck, T. D. 1995, *ApJ*, 447, 760
- Wang, K.-S., van der Tak, F. F. S., & Hogerheijde, M. R. 2012, *A&A*, 543, A22
- Widicus Weaver, S. L., Butler, R. A. H., Drouin, B. J., et al. 2005, *ApJS*, 158, 188
- Wilking, B. A., Blackwell, J. H., & Mundy, L. G. 1990, *AJ*, 100, 758
- Xu, Y., Moscadelli, L., Reid, M. J., et al. 2011, *ApJ*, 733, 25
- Zernickel, A., Schilke, P., Schmiedeke, A., et al. 2012, *A&A*, 546, A87
- Zhang, Q., Hunter, T. R., & Sridharan, T. K. 1998, *ApJ*, 505, L151
- Zinnecker, H., & Yorke, H. W. 2007, *ARA&A*, 45, 481

Appendix A: Detected lines per species for all sources

The line assignment and detection is based on Gaussian fitting with the following criteria: (i) the fitted line position has to be within ± 1 MHz of the catalog frequency; (ii) the FWHM is consistent with the those given in Table 1; and (iii) the peak intensity has at least a $S/N = 3$. The errors on the integrated intensities are computed as follows.

The integrated main beam temperatures, $\int T_{\text{MB}} dV$, were obtained by Gaussian fits to the lines (Eq. (A.1)).

$$\int T_{\text{MB}} dV = c^{st} T_0 \Delta V \quad (\text{A.1})$$

with

$$c^{st} = \frac{\sqrt{\pi/\ln 2}}{2}$$

where T_0 is the peak intensity and ΔV is the FWHM of the line. The error, $d \int T_{\text{MB}} dV$, is calculated from Eq. (A.2).

$$d \int T_{\text{MB}} dV = c^{st} [\Delta V d(T_0) + T_0 d(\Delta V)] \quad (\text{A.2})$$

with

$$d(T_0) = \sqrt{\text{rms}^2 + (\text{cal} \times T_0)^2 + \sigma_{T_0}^2}$$

and

$$d(\Delta V) = \sigma_{\Delta V}$$

where rms is the root mean square amplitude of the noise in the spectral bin δv , cal is the calibration uncertainty of the telescope, and σ 's are the statistical errors on T_0 and ΔV from the Gaussian fits.

The errors on the integrated intensities derived from Eq. (A.2) include all statistical errors from the Gaussian fit.

$$d \int T_{\text{MB}} dV = c^{st} \Delta V \sqrt{\text{rms}^2 + (\text{cal} \times T_0)^2}. \quad (\text{A.3})$$

For undetected transitions the upper limits were determined as 3σ limits (Eq. (A.4)) using:

$$\sigma = 1.2 \sqrt{\delta v \Delta V} \text{rms}, \quad (\text{A.4})$$

where 1.2 is the coefficient related to the calibration uncertainty of 20%.

CLASS was used to determine the Gaussian fits and the uncertainties in the individual parameters. The formal errors on the integrated intensities derived from Eq. (A.3) in some cases yield a $S/N < 2.5$. This is caused by: (i) a conservative estimate of the statistical error on the FWHM parameter in the Gaussian fitter of CLASS; (ii) all statistical errors are included into our error calculation. Considering the higher S/N on the peak intensity, a more traditional error estimate without the statistical errors from the Gaussian fit (Eq. (A.3)) would result in a $S/N > 3$ for the integrated intensity as well. All weak line fits were confirmed by visual inspection.

Table A.1. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s^{-1}) for H_2CO and its isotopic species.

Frequency [GHz]	Transition	E_{up} [K]	A [s^{-1}]	Sources	IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
H_2CO							
218.222 ^p	3 _{0,3} –2 _{0,2}	21	2.8(–4)	2.82 (0.65)	–	–	>7.14 (1.71)
218.476 ^p	3 _{2,2} –2 _{2,1}	68	1.6(–4)	–	–	–	4.15 (1.03)
363.946 ^p	5 _{2,4} –4 _{2,3}	100	1.2(–3)	4.65 (1.08)	9.92 (2.15)	–	>20.29 (3.31)
364.103 ^p	5 _{4,1/2} –4 _{4,0/1}	241	5.0(–4)	1.90 (0.58)	4.25 (1.07)	–	10.97 (2.88)
364.275 ^o	5 _{3,3} –4 _{3,2}	143/158	8.9(–4)	>5.08 (1.50)	>12.62 (3.34)	–	>21.56 (5.08)
364.289 ^o	5 _{3,2} –4 _{3,1}	143/158	8.9(–4)	>5.78 (1.71)	>10.68 (2.83)	–	>20.46 (4.85)
H_2^{13}CO							
219.909 ^o	3 _{1,2} –2 _{1,1}	33	2.6(–4)	<0.34	1.08 (0.60)	–	1.60 (0.55)

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. > Means lower limit (optically thick line). – Means frequency not observed. ^(p) Para- H_2CO . ^(o) Ortho- H_2CO .

Table A.2. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for CH₃OH and its isotopic species.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources		
				IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
CH₃OH						
218.440	4 _{2,0} –3 _{1,0}	45	6.9(–5)	–	–	10.10 (2.22) [∴]
219.984	25 _{3,0} –24 _{4,0}	802	3.0(–5)	<1.33	<0.24	<0.45 (0.24) [∴]
219.994	23 _{5,0} –22 _{6,0}	776	2.6(–5)	<1.33	<0.24	<1.74 (0.56) [∴]
240.739	26 _{3,-0} –26 _{2,+0}	864	1.1(–4)	<0.20	<0.40 (0.32)	<0.78 (0.47) [∴]
240.818	5 _{1,2} –4 _{1,2}	834	8.5(–5)	<0.20	<0.23	<3.58 (0.88) [∴]
240.861	5 _{-4,2} –4 _{-4,2}	779	3.2(–5)	<0.20	<0.27	<0.81 (0.37) [∴]
240.870	5 _{0,2} –4 _{0,2}	769	8.9(–5)	<0.20	<0.90 (0.71) [∴]	<3.26 (1.06) [∴]
240.916	5 _{3,-2/+2} –4 _{3,-2/+2}	693	5.7(–5)	<0.20	<1.17 (0.46) [∴]	1.43 (0.45)
240.932	5 _{4,-2/+2} –4 _{4,-2/+2}	649	3.2(–5)	<0.20	<0.57 (0.20)	~0.42 (0.21)
240.937	5 _{-2,2} –4 _{-2,2}	680	7.4(–5)	<0.20	<1.03 (0.69) [∴]	~0.49 (0.24)
240.939	5 _{0,+2} –4 _{0,+2}	543	1.2(–4)	<0.20	<1.03 (0.69) [∴]	2.54 (0.86)
240.948	5 _{3,2} –4 _{3,2}	656	5.6(–5)	<0.20	~0.24 (0.12) [*]	~0.74 (0.30) [*]
240.952	5 _{2,2} –4 _{2,2}	621	7.5(–5)	<0.20	~0.24 (0.12) [*]	~0.74 (0.30) [*]
240.959	5 _{-1,2} –4 _{-1,2}	567	8.5(–5)	<0.20	1.18 (0.42) [∴]	1.74 (0.61) [∴]
240.961	5 _{1,+1} –4 _{1,+1}	360	8.5(–5)	<0.20	1.18 (0.42) [∴]	1.74 (0.61) [∴]
241.043	22 _{-6,0} –23 _{-5,0}	776	3.4(–5)	–	~0.35 (0.48)	<0.38 (0.54) [∴]
330.794	8 _{-3,0} –9 _{-2,0}	146	8.0(–5)	~0.28 (0.57)	3.23 (1.12)	8.06 (2.43)
331.220	16 _{-1,0} –15 _{-2,0}	321	7.8(–5)	<0.39 ^c	2.66 (0.79)	>3.19 (0.91) ^b
338.124	7 _{0,0} –6 _{0,0}	78	2.5(–4)	—	–	30.57 (11.11) [∴]
338.345	7 _{-1,0} –6 _{-1,0}	70	2.5(–4)	7.29 (2.05) [∴]	10.34 (2.58) [∴]	38.03 (9.06) [∴]
338.405	7 _{6,0} –6 _{6,0}	244	6.7(–5)	<8.90 (2.54) [∴]	<12.95 (3.14) [∴]	<44.10 (10.87) [∴]
338.409	7 _{0,+0} –6 _{0,+0}	65	2.5(–4)	8.90 (2.54) [∴]	12.95 (3.14) [∴]	44.10 (10.87) [∴]
338.431	7 _{-6,0} –6 _{-6,0}	254	6.7(–5)	~0.72 (0.14)	~2.23 (1.06)	8.49 (2.90)
338.442	7 _{6,-0/+0} –6 _{6,-0/+0}	259	6.7(–5)	~1.20 (0.79) [∴]	~2.29 (1.27)	8.80 (3.44)
338.457	7 _{-5,0} –6 _{-5,0}	189	1.2(–4)	1.80 (0.66) [∴]	~2.55 (1.44)	10.12 (3.74)
338.475	7 _{5,0} –6 _{5,0}	201	1.2(–4)	~1.43 (0.94) [∴]	~1.89 (1.21)	9.22 (3.46)
338.486	7 _{5,-0/+0} –6 _{5,-0/+0}	203	1.2(–4)	~2.86 (0.94) [∴]	>3.40 (3.00)	>13.31 (5.28)
338.504	7 _{-4,0} –6 _{-4,0}	153	1.7(–4)	~1.78 (0.85)	~3.39 (1.88)	>12.14 (4.99)
338.5126	7 _{4/4,-0/+0} –6 _{4/4,-0/+0}	145	1.7(–4)	>2.61 (0.94) [∴]	>5.87 (1.81) [∴]	>26.88 (9.59) [∴]
338.5129	7 _{2,-0} –6 _{2,-0}	103	2.3(–4)	>2.61 (0.94) [∴]	>5.87 (1.81) [∴]	>26.88 (9.59) [∴]
338.530	7 _{4,0} –6 _{4,0}	161	1.7(–4)	~1.34 (0.59)	~3.69 (2.22)	11.33 (4.40)
338.541	7 _{3,+0/-0} –6 _{3,+0/-0}	115	2.1(–4)	>3.75 (1.24)	>8.71 (2.71)	>29.00 (8.01)
338.560	7 _{-3,0} –6 _{-3,0}	128	2.1(–4)	~2.14 (0.95)	>4.50 (1.87)	14.94 (4.70)
338.583	7 _{3,0} –6 _{3,0}	113	2.1(–4)	2.74 (1.17)	>4.02 (1.61)	18.35 (5.14)
338.615	7 _{1,0} –6 _{1,0}	86	2.5(–4)	4.48 (1.33) [∴]	9.01 (2.57) [∴]	28.49 (6.87) [∴]
338.640	7 _{2,+0} –6 _{2,+0}	103	2.3(–4)	>2.99 (1.07)	>6.46 (2.36)	21.12 (5.79)
338.722	7 _{2,0} –6 _{2,0}	87	2.3(–4)	2.60 (0.70) [∴]	5.24 (1.36) [∴]	17.99 (8.33) [∴]
338.723	7 _{-2,0} –6 _{-2,0}	91	2.3(–4)	2.60 (0.70) [∴]	5.24 (1.36) [∴]	17.99 (8.33) [∴]
340.141	2 _{2,+0} –3 _{1,+0}	45	4.1(–5)	1.56 (0.71) [∴]	3.31 (1.04) [∴]	12.35 (2.83) [∴]
342.730 ^a	13 _{1,12,-0} –13 _{0,13,+0}	147	2.4(–5)	–	–	14.19 (3.54)
344.971	12 _{7,1} –11 _{6,1}	762	1.3(–4)	<0.12	–	~2.27 (1.07)
364.159	9 _{3,1} –9 _{2,1}	522	6.5(–5)	~0.20 (0.41)	2.97 (1.10) [∴]	2.66 (0.77)
¹³CH₃OH						
222.468	21 _{1,20,0} –21 _{0,21,0}	544	3.6(–5)	<0.23	<0.32	<0.59
225.404	16 _{2,14,0} –15 _{3,12,0}	331	2.0(–5)	<0.19	<0.45 (0.35) [∴]	<3.49 (1.55) [∴]
338.760	13 _{0,13,+0} –12 _{1,12,+0}	206	2.2(–4)	<0.20	<0.39	~5.55 (2.36)
341.132 ^a	13 _{1,12,-0} –13 _{0,13,+0}	211	2.4(–5)	–	–	3.55 (2.22)
345.084	2 _{2,0,+0} –3 _{1,3,+0}	45	2.9(–5)	<0.29	<0.24	<1.69 (0.77) [∴]
345.133	4 _{0,4,0} –3 _{-1,3,0}	36	8.2(–5)	<0.29	1.81 (0.70)	4.06 (1.25)
354.446	4 _{1,3,0} –3 _{0,3,0}	44	1.3(–4)	<0.18	1.68 (0.65)	6.32 (1.59)
363.924	14 _{3,12,1} –14 _{2,13,1}	655	4.7(–5)	<0.24	<2.06 (0.94) [∴]	<1.40 (0.92) [∴]

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. ^(a) From additional observations on G31 only. ^(b) Line close to the edge of the frequency band. ^(c) Used to constrain the RTD fit. > Means lower limit (optically thick line). < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. ∴ Means blended frequency. ∴ Cold ($E_{\text{up}} < 100$ K) CH₃OH line (treated as a blend with cold component). ^(*) Calculated flux (see Eq. (4)).

Table A.3. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for C₂H₅OH.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources		
				IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
C ₂ H ₅ OH						
218.461	5 _{3,2,2} –4 _{2,3,2}	24	6.6(–5)	–	–	<2.60 (0.52):
218.554	21 _{5,16,2} –21 _{4,17,2}	226	6.2(–5)	—	–	~0.77 (0.65)
222.217	20 _{5,15,2} –20 _{4,16,2}	208	6.5(–5)	<0.22	~0.56 (0.59)	<1.29 (0.26):
222.419	20 _{4,17,1} –19 _{5,15,0}	256	1.1(–5)	<0.22	<0.34 (0.30):	<1.18 (0.24):
222.519	26 _{4,23,2} –26 _{3,24,2}	316	6.7(–5)	<0.22	<0.27	~0.26 (0.24)
225.105	13 _{10,3/4,0} –12 _{10,2/3,0}	255	4.2(–5)	<0.19	<0.25	~0.66 (0.42)
225.108	13 _{11,2/3,0} –12 _{11,1/2,0}	280	2.9(–5)	<0.19	<0.25	~0.40 (0.42)
225.110	13 _{6,8/7,1} –12 _{6,7/6,1}	181	8.0(–5)	<0.19	<0.25	1.12 (0.42)
225.112	13 _{9,5/4,0} –12 _{9,4/3,0}	231	5.3(–5)	<0.19	<0.25	~0.74 (0.42)
225.116	13 _{12,1/2,0} –12 _{12,0/1,0}	308	1.5(–5)	<0.25	<0.25	<0.55 (0.43):
225.131	13 _{8,5/6,0} –12 _{8,4/5,0}	211	6.4(–5)	<0.19	<0.25	~1.16 (1.12)
225.171	13 _{7,6/7,0} –12 _{7,5/6,0}	192	7.3(–5)	<0.19	<0.25	~1.86 (0.84)
225.210	19 _{5,14,2} –19 _{4,15,2}	191	6.7(–5)	<0.19	<0.25	~1.98 (1.05)
225.229	17 _{2,15,2} –16 _{3,14,2}	137	2.5(–5)	<0.19	~0.37 (0.24)	<2.76 (0.55):
225.249	13 _{6,8/7,0} –12 _{6,7/8,0}	176	8.1(–5)	<0.19	~0.40 (0.30)	~2.34 (0.93)
225.279	13 _{5,9,1} –12 _{5,8,1}	168	8.7(–5)	<0.19	<0.25	~1.61 (0.77)*
225.283	13 _{5,8,1} –12 _{5,7,1}	168	8.7(–5)	<0.19	<0.25	~1.61 (0.77)*
225.400	13 _{5,9,0} –12 _{5,8,0}	163	8.7(–5)	<0.19	~0.13 (0.15)	0.74 (0.32)*
225.404	13 _{5,8,0} –12 _{5,7,0}	163	8.7(–5)	<0.19	~0.13 (0.15)	0.74 (0.32)*
225.457	13 _{3,11,1} –12 _{3,10,1}	148	9.7(–5)	<0.19	<0.25	~0.95 (0.48)
238.841	21 _{2,19,0} –21 _{1,21,1}	258	4.8(–5)	—	<0.22	<2.28 (0.46):
239.020	28 _{4,25,2} –27 _{5,22,2}	362	2.9(–5)	<0.21	<0.22	<4.79 (0.96):
239.186	16 _{0,16,1} –15 _{1,14,0}	171	4.4(–5)	<0.21	<0.22	<0.92 (0.18):
240.654	4 _{2,2,1} –3 _{1,2,0}	75	4.1(–5)	–	<0.22	–
240.782	15 _{2,13,0} –14 _{1,13,1}	163	2.9(–5)	<0.2	<0.23	<3.38 (0.68):
240.839	14 _{1,13,0} –13 _{0,13,1}	147	5.1(–5)	<0.20	~0.20 (0.33)	~0.85 (0.52)
330.985	6 _{3,4,1} –5 _{2,4,0}	90	1.2(–4)	<0.31	–	~1.46 (0.84)
331.027	13 _{5,8,1} –13 _{4,10,0}	168	1.1(–4)	<0.29	<0.29	<1.84 (0.37):
331.079	43 _{7,37,2} –43 _{6,38,2}	860	2.2(–4)	<0.29	~0.36 (0.39)	~0.15 (0.22)
331.095	27 _{7,20,2} –27 _{6,21,2}	380	2.1(–4)	<0.29	<0.29	~0.33 (0.55)
338.088	25 _{1,24,1} –24 _{2,22,0}	332	5.9(–5)	–	–	~3.64 (1.99)
338.099	18 _{7,11,2} –18 _{6,12,2}	205	2.1(–4)	–	–	~2.05 (1.54)*
338.110	18 _{7,12,2} –18 _{6,13,2}	205	2.1(–4)	–	–	~2.05 (1.54)*
338.163	10 _{2,8,1} –9 _{1,8,0}	113	8.6(–5)	–	–	2.05 (0.98)
338.412	17 _{7,10,2} –17 _{6,11,2}	190	2.1(–4)	<4.41 (1.31):	<13.78 (3.29):	<21.70 (5.13):
338.417	17 _{7,11,2} –17 _{6,12,2}	190	2.1(–4)	<4.41 (1.31):	<0.71	<21.70 (5.13):
338.672	16 _{7,9,2} –16 _{6,10,2}	176	2.0(–4)	<0.32	<0.83	1.63 (0.85)*
338.674	16 _{7,10,2} –16 _{6,11,2}	176	2.0(–4)	<0.32	<0.83	1.63 (0.85)*
339.979	9 _{4,6,2} –8 _{3,5,2}	58	2.2(–4)	<0.15	~0.55 (0.47)	3.23 (0.93)
345.174	7 _{7,0/1,0} –6 _{6,0/1,1}	140	2.5(–4)	<0.26	~1.16 (0.70)	~2.28 (1.56)
345.229	21 _{1,21,0} –20 _{1,20,0}	242	3.7(–4)	<0.29	~0.58 (0.37)	<5.52 (1.10):
345.295	21 _{1,21,1} –20 _{1,20,1}	246	3.7(–4)	<0.29	~1.03 (1.00)	~2.91 (1.31)
345.312	22 _{3,19,1} –21 _{4,17,0}	286	5.0(–5)	<0.29	<0.19	<0.39
345.333	21 _{0,21,0} –20 _{0,20,0}	242	3.7(–4)	<0.29	<6.54 (1.31):	<17.35 (3.47):
345.408	21 _{0,21,1} –20 _{0,20,1}	246	3.7(–4)	<0.29	~0.75 (0.47)	~2.47 (1.13)
352.858	21 _{1,20,2} –20 _{2,19,2}	196	1.2(–4)	<0.18	~0.51 (0.43)	~1.56 (0.78)
353.034	12 _{3,9,2} –11 _{2,10,2}	77	1.9(–4)	<0.16	~0.61 (0.59)	3.12 (1.00)
354.758	20 _{3,17,1} –19 _{3,16,1}	249	4.0(–4)	<0.18	~0.59 (0.52)	3.96 (1.05)
363.968	21 _{7,15/14,1} –20 _{7,14/13,1}	314	3.9(–4)	<0.22	~0.64 (0.44)	2.66 (1.02)
364.001	21 _{8,14/13,0} –20 _{8,13/12,0}	327	3.7(–4)	<0.22	~0.70 (0.48)	2.28 (0.80)
364.233	21 _{7,15/14,0} –20 _{7,14/13,0}	309	3.9(–4)	<0.24	~0.64 (0.56)	~1.94 (1.48)

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line. * Calculated flux (see Eq. (4)).

Table A.4. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for HNCO and its isotopic species.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
HNCO						
219.657	10 _{3,8/7,9/11/10} -9 _{3,7/6,9/10/8}	447	1.4(-4)	<1.33	<0.73 (0.70):	<1.52 (0.63):
219.736	10 _{2,9/8,9/11/10} -9 _{2,8/7,9/10/8}	231	1.4(-4)	<0.28	<0.84 (0.75):	~0.64 (0.91)
219.798	10 _{0,10,9/11/10} -9 _{0,9,9/10/8}	58	1.5(-4)	<0.28	1.31 (0.50)	1.22 (0.55)
240.876	11 _{1,11,10/12/11} -10 _{1,10,10/11/9}	113	2.0(-4)	<0.20	<0.82 (0.54)	<3.10 (0.93):
330.849	15 _{1,14,14/16/15} -14 _{1,13,14/15/13}	170	5.2(-4)	~0.32 (0.31)	<1.53 (0.71):	2.01 (0.71)
352.898	16 _{1,15,15/17/16} -15 _{1,14,15/16/14}	187	6.3(-4)	<0.18	~1.84 (1.00)	<3.12 (1.02):
HN ¹³ CO						
219.664	10 _{3,8/7,10} -9 _{3,7/6,10}	47	1.4(-4)	<1.32	<0.21	<0.41
219.740	10 _{2,9,11/9/10} -9 _{2,8,10/8/9}	31	1.4(-4)	<0.28	<0.24	<0.26
219.744	10 _{2,8,11/9/10} -9 _{2,7,10/8/9}	31	1.4(-4)	<0.28	<0.24	<0.26
219.804	10 _{0,10,10/9/8} -9 _{0,9,10/9/8}	58	1.5(-4)	<0.28	<0.24	<0.63
240.881	11 _{1,11,12/11/10} -10 _{10,11,11/10/9}	113	2.0(-4)	<0.20	<0.23	<0.98
330.860	15 _{1,14,16/14/15} -14 _{1,13,15/13/14}	170	5.2(-4)	<0.31	<0.32	<0.31

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line.

Table A.5. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for NH₂CHO.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
NH ₂ CHO						
218.460	10 _{10,1,9} -9 _{9,1,8}	61	7.5(-4)	–	–	<2.45 (0.49):
339.716	16 _{16,8,8/9} -15 _{15,8,7/8}	329	2.2(-3)	<0.15	<0.17	<4.04 (0.81):
339.781	16 _{16,7,10/9} -15 _{15,7,9/8}	284	2.3(-3)	~0.74 (0.42)	~0.29 (0.29)	<2.66 (1.20):
339.904	16 _{16,6,11/10} -15 _{15,6,10/9}	246	2.5(-3)	<0.89 (0.39):	~0.48 (0.30)	~1.41 (0.76)
340.135	16 _{16,5,12} -15 _{15,5,11}	213	2.6(-3)	<0.78 (0.16):	<1.40 (0.28):	<1.11 (0.22):
340.139	16 _{16,5,11} -15 _{15,5,10}	213	2.6(-3)	<0.78 (0.16):	<1.40 (0.28):	<1.11 (0.22):
345.183	17 _{17,0,17} -16 _{16,0,16}	151	3.0(-3)	<0.20	1.16 (0.53)	2.42 (0.94)
345.327	16 _{16,1,15} -15 _{15,1,14}	145	3.0(-3)	<0.43	<1.87 (0.37):	<5.46 (1.09):

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line.

Table A.6. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for CH₃CN and its isotopic species.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources	IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
CH ₃ CN							
238.844	13 ₈ -12 ₈	537	5.0(-4)	–	<0.22	<0.17	
238.913	13 ₇ -12 ₇	430	5.8(-4)	<0.21	~0.37 (0.27)	1.82 (0.62)	
238.972	13 ₆ -12 ₆	337	6.4(-4)	~0.33 (0.38)	~0.84 (0.37)	3.22 (1.11)	
239.023	13 ₅ -12 ₅	259	6.9(-4)	~0.93 (0.49)	<6.43 (1.29):	3.19 (0.88)	
239.064	13 ₄ -12 ₄	195	7.4(-4)	~0.51 (0.38)	1.17 (0.45)	5.25 (1.44)	
239.096	13 ₃ -12 ₃	145	7.7(-4)	1.04 (0.40)	1.64 (0.55)	5.21 (1.37)	
239.120	13 ₂ -12 ₂	109	7.9(-4)	~0.63 (0.38)	1.80 (0.61)	5.71 (1.37)	
239.133	13 ₁ -12 ₁	87	8.1(-4)	0.73 (0.26)*	1.72 (0.52)*	4.56 (1.19)	
239.138	13 ₀ -12 ₀	80	8.1(-4)	0.73 (0.26)*	1.72 (0.52)*	4.60 (1.15)	
330.843	18 ₆ -17 ₆	408	1.9(-3)	~0.31 (0.26)	4.63 (1.28)	7.44 (2.20)	
330.913	18 ₅ -17 ₅	329	2.0(-3)	–	3.59 (1.13)	5.91 (1.87)	
330.970	18 ₄ -17 ₄	265	2.1(-3)	–	2.81 (0.78)	6.70 (2.07)	
331.014	18 ₃ -17 ₃	215	2.1(-3)	–	4.66 (1.30)	8.26 (2.38)	
331.046	18 ₂ -17 ₂	180	2.2(-3)	–	4.51 (1.31)	10.24 (2.87)	
331.065	18 ₁ -17 ₁	158	2.2(-3)	–	4.01 (1.09)*	6.25 (1.96)	
331.072	18 ₀ -17 ₀	151	2.2(-3)	–	4.01 (1.09)*	7.86 (2.30)	
CH ₃ ¹³ CN							
238.855	13 ₆ -12 ₆	337	9.2(-4)	–	<0.22	<10.23 (2.58):	
238.905	13 ₅ -12 ₅	259	1.0(-3)	<0.21	<0.22	<1.00 (0.48):	
238.946	13 ₄ -12 ₄	195	1.1(-3)	<0.21	<0.22	2.21 (0.70)	
238.978	13 ₃ -12 ₃	145	1.1(-3)	<0.21	~0.35 (0.31)	1.51 (0.58)	
239.001	13 ₂ -12 ₂	109	1.2(-3)	<0.21	<0.22	~0.62 (0.35)	
239.015	13 ₁ -12 ₁	87	1.2(-3)	<0.21	<0.22	4.61 (0.97)*	
239.020	13 ₀ -12 ₀	80	1.2(-3)	<0.21	<0.22	4.61 (0.97)*	
330.806	18 ₄ -17 ₄	265	3.0(-3)	<0.31	<0.32	<8.08 (2.52):	
330.851	18 ₃ -17 ₃	215	3.1(-3)	<0.31	<2.66 (0.53):	<4.12 (1.23):	
330.882	18 ₂ -17 ₂	179	3.1(-3)	<0.31	<0.32	~0.69 (0.49)	
330.901	18 ₁ -17 ₁	158	3.2(-3)	<0.31	<1.49 (0.30):	2.37 (0.93)	
330.908	18 ₀ -17 ₀	151	3.2(-3)	<0.31	<1.49 (0.30):	3.48 (1.12)	

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line. * Calculated flux (see Eq. (4)).

Table A.7. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for C₂H₅CN.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources	IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
C ₂ H ₅ CN							
218.390	24 _{3,21} -23 _{3,20}	140	8.7(-4)	<0.18	–	~1.03 (0.45)	
219.903	12 _{3,10} -11 _{2,9}	44	3.0(-5)	<0.28	<0.24	~0.18 (0.13)	
225.236	25 _{4,21} -24 _{4,20}	158	9.4(-4)	<0.00	~0.53 (0.36)	2.53 (0.85)	
225.307	12 _{3,9} -11 _{2,10}	44	3.2(-5)	<0.19	<0.25	<1.46 (0.57):	
225.317	23 _{2,22} -22 _{1,21}	122	4.8(-5)	<0.19	<0.25	<1.79 (0.65):	
240.699	16 _{9,7/8} -17 _{8,10/9}	148	7.9(-6)	<0.20	<0.48 (0.10):	<0.89 (0.46):	
240.861	28 _{1,28} -27 _{0,27}	169	1.0(-4)	<0.20	<0.27 (0.05):	~0.78 (0.37)	
338.143	37 _{3,34} -36 _{3,33}	317	3.3(-3)	–	–	5.74 (2.44)	
339.895	39 _{2,38} -38 _{2,37}	334	3.3(-3)	<0.15	~0.38 (0.68)	3.45 (1.05)	
339.968	38 _{2,36} -37 _{2,35}	327	3.3(-3)	<0.15	~0.50 (0.32)	2.20 (0.69)	
340.149	39 _{1,38} -38 _{1,37}	334	3.3(-3)	<0.15	~0.26 (0.17)	2.20 (0.82)	
352.992	21 _{2,19} -20 _{1,20}	105	1.4(-5)	<0.13	<0.24	<0.19	
353.089	23 _{3,20} -22 _{2,21}	129	5.8(-5)	<0.16	<0.24	~0.58 (0.29)	
354.477	40 _{3,38} -39 _{3,37}	361	3.8(-3)	<0.18	<3.38 (0.68):	1.33 (0.65)	

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line.

Table A.8. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for HCOOCH₃.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
HCOOCH ₃						
218.281	17 _{3,14,2} –16 _{3,13,2}	100	1.5(–4)	<0.18	–	1.38 (0.42)
218.298	17 _{3,14,0} –16 _{3,13,0}	100	1.5(–4)	<0.18	–	1.00 (0.430)
219.584	18 _{13,5/6,3} –17 _{13,4/5,3}	401	7.7(–5)	<1.33	–	~0.17 (0.15)
219.592	28 _{9,19,2} –28 _{8,20,2}	295	1.6(–5)	<1.33	–	~0.30 (0.22)
219.623	18 _{12,6/7,3} –17 _{12,5/6,3}	384	8.9(–5)	<1.33	–	~0.66 (0.32)
219.642	18 _{13,6,4} –17 _{13,5,4}	401	7.7(–5)	<1.33	<0.21	~0.55 (0.43)
219.696	18 _{11,8/7,3} –17 _{11,7/6,3}	369	1.0(–4)	<0.28	<0.21	0.69 (0.25)
219.705	18 _{4,15,3} –17 _{4,14,3}	300	1.5(–4)	<0.28	<0.21	0.77 (0.27)
219.764	18 _{9,9,5} –17 _{9,8,5}	342	1.2(–4)	<0.28	<0.21	0.61 (0.24)
219.822	18 _{10,9/8,3} –17 _{10,8/7,3}	355	1.1(–4)	<0.28	<0.21	<1.10 (0.55):
222.149	18 _{6,12,3} –17 _{6,11,3}	312	1.5(–4)	<0.22	1.25 (0.25):	<3.05 (0.83):
222.177	18 _{6,12,5} –17 _{6,11,5}	312	1.5(–4)	<0.18	<0.20	~1.14 (0.67)
222.421	18 _{8,10,2} –17 _{8,9,2}	144	1.3(–4)	<0.22	<1.32 (0.26):	1.62 (0.45)
222.438	18 _{8,11,0} –17 _{8,10,0}	144	1.3(–4)	<0.22	0.35 (0.16)*	1.07 (0.39)*
222.440	18 _{8,10,0} –17 _{8,9,0}	144	1.3(–4)	<0.22	0.35 (0.16)*	1.07 (0.39)*
222.442	18 _{8,11,1} –17 _{8,10,1}	144	1.3(–4)	<0.22	0.35 (0.16)*	1.07 (0.39)*
225.372	20 _{21,9,3} –19 _{21,8,3}	307	1.7(–4)	<0.19	<0.31	1.38 (0.56)
225.449	20 _{11,9,3} –19 _{11,8,3}	307	1.7(–4)	<0.19	<0.18	<1.30 (0.55):
238.927	20 _{3,18,1} –19 _{2,17,2}	128	2.1(–5)	<0.21	<0.22	~0.40 (0.23)
238.933	20 _{3,18,0} –19 _{2,17,0}	128	2.1(–5)	<0.21	<0.22	~0.47 (0.20)
238.947	19 _{3,16,5} –18 _{3,15,5}	309	2.0(–4)	<0.21	~0.29 (0.26)	<0.88 (0.38):
239.111	7 _{6,2,4} –6 _{5,2,4}	227	2.7(–5)	<0.21	<0.22	<0.15 (0.17)
241.059	30 _{3,27,0} –30 _{3,28,0}	281	5.8(–6)	–	<0.23	~0.33 (0.31)
241.068	30 _{3,27,0} –30 _{2,28,0}	281	1.1(–5)	–	<0.23	~0.76 (0.53)
330.941	30 _{2/1,29,4/5} –29 _{2/1,28,4/5}	443	5.5(–4)	<0.31	~0.73 (0.38)	~0.75 (0.83)
331.021	25 _{5,21,0} –24 _{4,20,0}	210	3.5(–5)	<0.34	<0.40	<8.29 (1.66):
331.036	25 _{5,21,1} –24 _{4,20,2}	210	3.5(–5)	<0.32	<0.34	<10.01 (3.05):
331.121	27 _{12,16,4} –26 _{12,15,4}	505	4.5(–4)	<0.29	–	<0.74 (0.69)
331.149	28 _{4,25,1} –27 _{4,24,1}	248	5.3(–4)	<0.29	0.69 (0.31)	2.60 (1.39)
331.160	28 _{4,25,0} –27 _{4,24,0}	248	5.3(–4)	<0.29	<0.99 (0.42):	<2.24 (1.15):
331.161	27 _{11,17/16,3} –26 _{11,16/15,3}	490	4.6(–4)	<0.29	<0.99 (0.42):	<2.24 (1.15):
338.338	27 _{8,19,2} –26 _{8,18,2}	267	5.4(–4)	<0.26	<0.39	6.26 (2.37)
338.356	27 _{8,19,0} –26 _{8,18,0}	267	5.4(–4)	<0.23	<0.44	5.09 (2.08)
338.393	28 _{5,24,3} –27 _{5,23,3}	443	5.7(–4)	<0.23	<0.43	<3.80 (2.05):
338.396	27 _{7,21,1} –26 _{7,20,1}	258	5.5(–4)	<0.23	<0.45	<3.80 (2.05):
338.414	27 _{7,21,0} –26 _{7,20,0}	258	5.5(–4)	<8.06 (2.04):	<0.45	<43.32 (9.59):
339.882	29 _{3,26,3} –28 _{3,25,3}	450	5.8(–4)	<0.22	<0.79 (0.16):	~0.73 (0.33)
340.044	29 _{4,26,4} –28 _{4,25,4}	450	5.8(–4)	<0.15	<0.12	<3.75 (0.75):
340.115	27 _{7,20,5} –26 _{7,19,5}	444	5.5(–4)	<0.15	~0.30 (0.31):	0.77 (0.32)
345.068	28 _{14,14,2} –27 _{14,13,2}	370	4.7(–4)	<0.26	0.28 (0.11)	<3.93 (1.62):
345.069	28 _{14,15/14,0} –27 _{14,14/13,0}	370	4.7(–4)	<0.26	0.56 (0.22)	<3.93 (1.62):
345.073	16 _{6,11,0} –15 _{5,10,0}	104	3.9(–5)	<0.26	<0.19	<5.62 (1.29):
345.085	19 _{13,6,2} –19 _{12,7,2}	224	3.5(–5)	–	~0.37 (0.20)	<1.73 (0.35):
345.091	28 _{14,15,1} –27 _{14,14,1}	370	4.7(–4)	<0.26	<0.50 (0.32):	<4.14 (1.70):
345.148	28 _{6,23,3} –27 _{6,22,3}	452	6.0(–4)	<0.26	~0.55 (0.34)	~0.59 (0.54)
345.163	11 _{8,3,5} –10 _{7,3,5}	269	7.1(–5)	<0.26	<0.19	~0.50 (0.47)
345.230	18 _{13,5,2} –18 _{12,6,2}	213	3.2(–5)	<0.25	~0.59 (0.40)	<8.18 (3.04):
345.242	18 _{13,6,1} –18 _{12,7,1}	213	3.2(–5)	<0.25	<0.85 (0.83):	<8.18 (3.04):
345.248	28 _{10,19,4} –27 _{10,18,4}	493	5.5(–4)	<0.26	<0.19	<0.34
352.817	33 _{0/1/0/1,33,3} –32 _{1/1/0/0,32,3}	479	1.1(–4)	<0.18	~0.38 (0.28)	<2.68 (0.76):
352.841	33 _{1/0,33,4/5} –32 _{1/0,32,4/5}	479	7.7(–4)	<0.18	<0.50 (0.42):	1.72 (0.47)
352.912	31 _{2,29,2} –30 _{3,28,1}	286	8.5(–5)	<0.18	<0.26	0.30 (0.08)
352.918	31 _{3,29,1} –30 _{3,28,1}	286	6.6(–4)	<0.18	<1.10 (0.36):	2.32 (0.62)*
352.922	31 _{2,29,2} –30 _{2,28,2}	286	6.6(–4)	<0.18	<1.10 (0.36):	2.32 (0.62)*
352.926	31 _{3,29,0} –30 _{3,28,0}	286	6.6(–4)	<0.18	<1.10 (0.36):	2.32 (0.62)*
352.930	31 _{2,29,0} –30 _{2,28,0}	286	6.6(–4)	<0.18	<1.10 (0.36):	2.32 (0.62)*

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line. * Calculated flux (see Eq. (4)).

Table A.9. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for HCOOCH₃ continued.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
HCOOCH ₃						
354.427	29 _{16,13/14,3} –28 _{16,12/13,3}	614	4.7(–4)	<0.18	<0.23	~1.04 (0.64)
354.477	29 _{15,15,4} –28 _{15,14,4}	593	5.0(–4)	<0.13	<0.23	<0.40 (0.58):
354.574	12 _{8,5/4,3} –11 _{7,4/5,3}	276	7.4(–5)	<0.18	<0.23	<1.44 (0.78):
354.605	29 _{15,15/14,3} –28 _{15,14/13,3}	593	5.0(–4)	<0.18	<0.23	1.96 (0.46)
354.608	33 _{1/0,33,1/2} –32 _{1/0,32,1/2}	293	7.2(–4)	<0.18	1.11 (0.42)*	3.14 (0.49)*
354.608	33 _{0/1,33,2/1} –32 _{1/0,32,1/2}	293	6.8(–4)	<0.18	1.05 (0.40)*	2.75 (0.47)*
354.629	28 _{7,21,5} –27 _{7,20,5}	461	6.4(–4)	<0.18	<0.23	~1.20 (0.68)
354.742	12 _{8,5,1} –11 _{7,5,1}	88	7.3(–5)	<0.18	<0.23	1.19 (0.44)
354.759	12 _{8,4,2} –11 _{7,4,2}	88	7.3(–5)	<0.16	<0.17	<3.23 (0.65):
354.806	12 _{8,5/4,0} –11 _{7,4/5,0}	88	7.3(–5)	<0.18	<0.23	3.90 (1.18)
354.839	29 _{14,15/16,3} –28 _{14,14/15,3}	574	5.2(–4)	<0.15	<0.36 (0.46):	<4.91 (1.40):
364.297	33 _{1/2,32,2/1} –32 _{1,31,2}	308	7.3(–4)	<0.24	<6.81 (1.36):	<3.67 (1.16):
364.302	33 _{2/1,32,0} –32 _{2/1,31,0}	308	7.3(–4)	0.80 (0.91)	<6.81 (1.36):	<5.00 (1.42):

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line. * Calculated flux (see Eq. (4)).

Table A.10. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for CH₃OCH₃.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
CH ₃ OCH ₃						
218.490	23 _{3,21,2/3} –23 _{2,22,2/3}	264	3.4(–5)	<0.18	–	<1.19 (0.75):
218.492	23 _{3,21,1} –23 _{2,22,1}	264	3.4(–5)	<0.18	–	<1.19 (0.75):
218.495	23 _{3,21,0} –23 _{2,22,0}	264	3.4(–5)	<0.18	–	<1.19 (0.75):
222.239	4 _{3,2,2} –3 _{2,1,2}	22	2.8(–5)	<0.22	0.82 (0.56)	~0.93 (0.43)
222.248	4 _{3,2,3/1} –3 _{2,1,3/1}	22	4.2(–5)	<0.22	~0.43 (0.20)	<1.15 (0.35):
222.255	4 _{3,2,0} –3 _{2,1,0}	22	4.9(–5)	<0.22	~0.62 (0.21)	1.11 (0.98)
222.259	4 _{3,1,1} –3 _{2,1,1}	22	1.6(–5)	<0.22	~0.30 (0.21)*	2.37 (0.74)*
222.260	4 _{3,1,2} –3 _{2,1,2}	22	2.1(–5)	<0.22	~0.11 (0.21)*	0.83 (0.26)*
222.326	25 _{3,23,0} –24 _{4,20,0}	308	7.9(–6)	<0.22	<0.27	~0.42 (0.19)*
222.327	25 _{3,22,1} –24 _{4,20,1}	308	7.9(–6)	<0.22	<0.27	~1.12 (0.52)*
222.329	25 _{3,22/23,2/3} –24 _{4,20/20,2/3}	308	7.9(–6)	<0.22	<0.27	~0.42 (0.21)*
222.414	4 _{3,2,2} –3 _{2,2,2}	22	2.1(–5)	<0.22	~0.20 (0.10)	<0.73 (0.15):
222.423	4 _{3,2,1} –3 _{2,2,1}	22	1.6(–5)	<0.22	~0.56 (0.30)	<2.09 (0.42):
222.427	4 _{3,1,3} –3 _{2,2,3}	22	4.9(–5)	<0.22	~0.23 (0.12)	<0.85 (0.11):
222.434	4 _{3,1,1/0} –3 _{2,2,1/0}	22	4.2(–5)	<0.22	1.68 (0.63)	<1.86 (0.52):
222.435	4 _{3,1,2} –3 _{2,2,2}	22	2.8(–5)	<0.22	~0.22 (0.08)	<0.34 (0.10):
225.202	24 _{4,21,2/3} –24 _{3,22,2/3}	296	4.7(–5)	<0.19	<0.30	0.55 (0.18)*
225.204	24 _{4,21,1} –24 _{3,22,1}	296	4.7(–5)	<0.19	~0.15 (0.14)*	0.87 (0.28)*
225.205	24 _{4,21,0} –24 _{3,22,0}	296	4.7(–5)	<0.19	~0.09 (0.08)*	0.55 (0.18)*
238.975	29 _{5,25,2/3/1/0} –28 _{6,22,2/3/1/0}	432	1.6(–5)	<0.21	<0.22	<1.64 (0.64):
239.020	24 _{5,19,5/3} –24 _{4,20,5/3}	309	6.1(–5)	<0.21	<0.63 (0.12):	<0.68 (0.14):
239.021	24 _{5,19,1} –24 _{4,20,1}	309	6.1(–5)	<0.21	<1.67 (0.33):	<1.82 (0.36):
239.021	24 _{5,19,0} –24 _{4,20,0}	309	6.1(–5)	<0.21	<0.63 (0.13):	<0.68 (0.14):
240.978	5 _{3,3,2} –4 _{2,2,2}	26	4.5(–5)	<0.20	~0.11 (0.07)*	2.16 (0.66)*
240.983	5 _{3,3,3} –4 _{2,2,3}	26	5.4(–5)	<0.20	~0.06 (0.04)*	0.19 (0.06)*
240.985	5 _{3,3,1} –4 _{2,2,1}	26	5.1(–5)	<0.20	~0.49 (0.30)*	1.49 (0.46)*
240.990	5 _{3,3,0} –4 _{2,2,0}	26	5.4(–5)	<0.20	~0.19 (0.12)	<0.64 (0.28):

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line. * Calculated flux (see Eq. (4)).

Table A.11. Observed line fluxes $\int T_{\text{MB}}dV$ (K km s⁻¹) for CH₂CO.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources		
				IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
CH ₂ CO						
222.120	11 _{4,8/7} –10 _{4,7/6}	273	1.1(–4)	<0.22	<0.27	<0.54 (0.10):
222.198	11 _{0,11} –10 _{0,10}	65	1.2(–4)	<0.22	<0.81 (0.26):	<2.34 (0.75):
222.201	11 _{3,9/8} –10 _{3,8/7}	181	1.1(–4)	<0.22	<0.81 (0.26):	<2.34 (0.75):
222.229	11 _{2,10} –10 _{2,9}	116	1.2(–4)	<0.22	<0.28 (0.06):	<2.22 (0.32):
222.315	11 _{2,9} –10 _{2,8}	116	1.2(–4)	<0.22	~0.24 (0.22)	~0.91 (0.57)
363.937!	18 _{2,16} –17 _{2,15}	218	5.5(–4)	<4.65 (1.08):	<0.22	<20.29 (3.31):

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line. ! Frequency from CDMS.

Table A.12. Observed line fluxes $\int T_{\text{MB}}dV$ (K km s⁻¹) for CH₃CHO.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources		
				IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
CH ₃ CHO						
330.822	5 _{3,3,1} –4 _{2,3,1}	34	1.2(–4)	<0.31	~0.71 (0.80)	~0.97 (0.52)
331.039	17 _{3,14,3} –16 _{3,13,3}	367	1.2(–3)	<1.18 (0.88):	<0.42 (0.29):	<2.50 (1.47):
354.458	18 _{2,16,3} –17 _{2,15,3}	375	1.5(–3)	<0.18	<3.11 (1.11):	<3.25 (1.26):
354.525	19 _{0,19,5} –18 _{0,18,5}	377	1.5(–3)	<0.17	<0.19	<0.34
354.813	18 _{2,16,2} –17 _{2,15,2}	170	1.5(–3)	<0.18	<0.23	~1.54 (0.92)
354.844	18 _{2,16,0} –17 _{2,15,0}	170	1.5(–3)	<0.18:	<1.13 (0.55):	<4.92 (1.34):

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line.

Table A.13. Observed line fluxes $\int T_{\text{MB}}dV$ (K km s⁻¹) for HCOOH and its isotopic species.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources		
				IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
HCOOH						
222.110	7 _{2,6} –7 _{1,7}	43	1.9(–6)	<0.22	<0.27	<0.24
225.086	10 _{4,7} –9 _{4,6}	110	1.0(–4)	<0.18	<0.18	<1.97 (1.70):
225.091	10 _{4,6} –9 _{4,5}	110	1.0(–4)	<0.18	<0.18	<1.97 (1.70):
225.238	10 _{3,8} –9 _{3,7}	88	1.1(–4)	<0.19	<0.39 (0.24):	<2.59 (0.78):
330.931	4 _{3,1} –4 _{2,2}	39	5.1(–6)	<0.31	<0.32	~0.47 (0.63)
331.145	3 _{3,0} –3 _{2,1}	35	3.6(–6)	<0.29	<0.29:	<2.77 (1.80):
338.109	15 _{1,15} –14 _{0,14}	127	1.1(–5)	—	—	<5.11 (3.10):
338.202	15 _{3,13} –14 _{3,12}	158	4.1(–4)	—	—	<13.55 (6.28):
345.031	16 _{0,16} –15 _{0,15}	143	4.5(–4)	<0.26	~0.30 (0.30)	~1.18 (0.70)
345.253	14 _{3,12} –14 _{2,13}	142	8.3(–6)	<0.29	<0.19:	<1.59 (1.01):
354.448	17 _{0,17} –16 _{1,16}	161	1.3(–5)	<0.18	<1.70 (0.57):	<6.42 (1.60):

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line.

Table A.14. Observed line fluxes $\int T_{\text{MB}} dV$ (K km s⁻¹) for CH₃CCH.

Frequency [GHz]	Transition	E_{up} [K]	A [s ⁻¹]	Sources		
				IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
CH ₃ CCH						
222.099	13 ₄ –12 ₄	190	3.4(–5)	<0.22	<0.65 (0.35):	<2.79 (0.56):
222.129	13 ₃ –12 ₃	140	3.6(–5)	~0.29 (0.17)	1.14 (0.43)	3.15 (0.94)
222.150	13 ₂ –12 ₂	104	3.7(–5)	~0.41 (0.19)	1.25 (0.46)	<4.25 (0.85):
222.163	13 ₁ –12 ₁	82	3.8(–5)	0.74 (0.29)	2.20 (0.68)	3.14 (1.01)
222.167	13 ₀ –12 ₀	75	3.8(–5)	0.93 (0.35)	2.13 (0.70)	3.27 (0.97)
239.088	14 ₆ –13 ₆	346	3.9(–5)	<0.21	<0.22 (0.05):	<2.40 (0.48):
239.138	14 ₅ –13 ₅	267	4.1(–5)	<0.21	<2.38 (0.48):	<6.39 (1.28):
239.179	14 ₄ –13 ₄	202	4.3(–5)	0.71 (0.14)	<0.68 (0.14):	~0.39 (0.21)
239.211	14 ₃ –13 ₃	151	4.5(–5)	~0.40 (0.24)	~0.75 (0.41)	2.47 (0.64)
239.234	14 ₂ –13 ₂	115	4.6(–5)	0.36 (0.16)	~0.89 (0.89)	2.06 (0.63)
239.248	14 ₁ –13 ₁	93	4.7(–5)	0.79 (0.23)	1.62 (0.52)	2.91 (0.74)
239.252	14 ₀ –13 ₀	86	4.7(–5)	0.87 (0.24)	1.68 (0.43)	3.46 (1.07)

Notes. The notation $a(-b)$ stands for $a \times 10^{-b}$. < Means upper limit. ~ Means uncertain detection, $S/N \lesssim 2$. – Means frequency not observed. : Means blended line.

Appendix B: Rotation diagrams

RTD diagrams for H_2CO , CH_3OH , $\text{C}_2\text{H}_5\text{OH}$, HNCO , NH_2CHO , CH_3CN , $\text{C}_2\text{H}_5\text{CN}$, HCOOCH_3 , CH_3OCH_3 , CH_2CO and CH_3CCH . Lines from different frequency bands are corrected for differential beam dilution assuming a $R_{7=100}$ K source size for the warm species and $14''$ source size for the cold species, as indicated in the plots. Optically thin, unblended lines with $S/N \gtrsim 2$ are marked with filled circles. Lines with high S/N ($\lesssim 2$) are marked with diamonds, those included in the fit with filled diamonds. Optically thick lines and blended lines are marked with open squares and triangles, respectively. Upper limits are marked with arrows. Upper limits used to constrain the fit are marked with filled stars. The error bars are calculated using Eq. (A.2).

Fig. B.1. RTD fits for H_2CO . All lines included into the fit belong to para- H_2CO .

Fig. B.2. RTD fits for CH_3OH . Lines with $E_{\text{up}} < 100$ K are considered contaminated with the cold CH_3OH . Lines with $S/N \gtrsim 1$ are included in the fit.

Fig. B.3. RTD fits for $\text{C}_2\text{H}_5\text{OH}$. No $\text{C}_2\text{H}_5\text{OH}$ was detected in IRAS 20126+4104. For IRAS 18089 lines with $S/N \gtrsim 1$ have been used.

Fig. B.4. RTD fits for HNCO . No HNCO was detected in IRAS 20126+4104.

Fig. B.5. RTD fits for NH_2CHO . No NH_2CHO was detected in IRAS 20126+4104.

Fig. B.6. RTD fits for CH_3CN .

Fig. B.7. RTD fits for $\text{C}_2\text{H}_5\text{CN}$. No $\text{C}_2\text{H}_5\text{CN}$ was detected in IRAS 20126+4104. For IRAS 18089 lines with $S/N \lesssim 2$ have been included in the fit.

Fig. B.8. RTD fits for HCOOCH_3 . No HCOOCH_3 was detected in IRAS 20126+4104.

Fig. B.9. RTD fits for CH_3OCH_3 . No CH_3OCH_3 was detected in IRAS 20126+4104.

Fig. B.10. RTD fits for CH_2CO . No CH_2CO was detected in IRAS 20126+4104.

Fig. B.11. RTD fits for CH_3CHO .

Fig. B.12. RTD fits for HCOOH .

Fig. B.13. RTD fits for CH_3CCH .

Appendix C: Weeds model parameters

Table C.2. Weeds model parameters for IRAS 18089-1732.

Table C.1. Weeds model parameters for IRAS 20126+4104.

Species	N_S [cm ⁻²]	T_{ex} [K]	θ_S ["]	Δv [km s ⁻¹]
CH ₃ OH	1.5E+17	300	2.2	6
CH ₃ OH	4.8E+16	20	3.0	5
¹³ CH ₃ OH	—	—	—	—
H ₂ CO	6.0E+15	150	2.2	6
H ₂ ¹³ CO	<2.0E+15	150	2.2	6
CH ₃ CN ¹	1.3E+15	200	2.2	6
CH ₃ ¹³ CN	<2.0E+13	200	2.2	6
HNCO	1.0E+15	200	2.2	6
HN ¹³ CO	<1.0E+14	200	2.2	6
C ₂ H ₅ OH	<3.0E+15	100	2.2	6
C ₂ H ₅ CN	<5.0E+14	80	2.2	5
CH ₃ OCH ₃	<1.0E+16	100	2.2	6
HCOOCH ₃	<2.0E+15	200	2.2	6
NH ₂ CHO	3.0E+14	300	2.2	6
CH ₂ CO	<5.0E+13	50	14.0	6
CH ₃ CHO	<1.0E+14	50	8.0	6
HCOOH	<5.0E+13	40	8.0	6
CH ₃ CCH	7.0E+14	35	14.0	2
Additional species				
CH ₃ COCH ₃	<0.1E+17	300	2.2	5
HCCCN	1E+13	100	14	7
SO	0.3E+15	50	14	6
³⁴ SO	0.2E+14	50	14.0	6
SO ₂	0.3E+15	50	14.0	6
³³ SO ₂	0.05E+15	50	14.0	5
HCN	0.4E+14	50	14.0	20
HCN	4E+13	50	14.0	5
H ¹³ CN	0.1E+13	50	14.0	6
CN, $v = 0, 1$	0.2E+15	50	14	2

Notes. ¹³CH₃OH is not available in the JPL database used for Weeds modeling. ⁽¹⁾ Based on 21'' beam spectra.

Species	N_S [cm ⁻²]	T_{ex} [K]	θ_S ["]	Δv [km s ⁻¹]
CH ₃ OH	3.5E+17	300	2.4	5
CH ₃ OH	2.6E+17	20	3.0	5
¹³ CH ₃ OH	—	—	—	—
H ₂ CO	1.5E+16	150	2.4	6
H ₂ ¹³ CO	3.5E+15	150	2.4	6
CH ₃ CN	3.5E+15	200	2.4	6
CH ₃ ¹³ CN	4.0E+14	200	2.4	6
HNCO	4.0E+15	200	2.4	6
HN ¹³ CO	7.0E+14	200	2.4	6
C ₂ H ₅ OH	2.5E+16	150	2.4	6
C ₂ H ₅ CN	4.0E+15	80	2.4	5
CH ₃ OCH ₃	1.0E+17	100	2.4	6
HCOOCH ₃	3.0E+16	200	2.4	6
NH ₂ CHO	5.0E+14	100	2.4	6
CH ₂ CO	1.5E+14	50	14.0	5
CH ₃ CHO	<1.0E+14	50	8.0	6
HCOOH	<5.0E+13	40	8.0	6
CH ₃ CCH	1.8E+15	40	14.0	3.5
Additional species				
CH ₃ COCH ₃	0.5E+17	300	2.4	5
HCCCN	3E+13	150	14	6
SO	1E+15	50	14	6
³⁴ SO	0.5E+14	50	14.0	6
SO ₂	0.5E+15	50	14.0	6
³³ SO ₂	0.5E+15	50	14.0	5
HCN	2E+14	50	14.0	7
HCN	-5E+13	50	14.0	4
H ¹³ CN	1.5E+13	50	14.0	6
CN, $v = 0, 1$	0.4E+15	50	14	5
NH ₂ CN	0.1E+14	50	14	7
OC ³⁴ S	0.5E+15	50	14.0	7
CP	0.5E+15	50	14.0	4
C ³⁴ S	1.4E+14	50	14.0	5
HCOCH ₂ OH	0.05E+17	300	2.0	6

Notes. ¹³CH₃OH is not available in the JPL database used for Weeds modeling.

Table C.3. Weeds model parameters for G31.41+0.31.

Species	N_S [cm ⁻²]	T_{ex} [K]	θ_S [']	Δv [km s ⁻¹]
CH ₃ OH	2.0E+18	300	2.0	7
CH ₃ OH	2.7E+17	20	5.0	7
¹³ CH ₃ OH	—	—	—	—
H ₂ CO	6.0E+16	150	2.0	6
H ₂ ¹³ CO	7.0E+15	150	2.0	6
CH ₃ CN ⁽¹⁾	2.0E+16	300	2.0	7
CH ₃ ¹³ CN	2.0E+15	300	2.0	7
HNCO	6.0E+16	200	2.0	7
HN ¹³ CO	2.0E+15	200	2.0	7
C ₂ H ₅ OH	2.0E+17	100	2.0	7
C ₂ H ₅ CN	2.0E+16	80	2.0	5
CH ₃ OCH ₃	1.0E+18	100	2.0	5
HCOOCH ₃ ⁽²⁾	1.0E+18	300	2.0	7
NH ₂ CHO	4.0E+15	300	2.0	7
CH ₂ CO	6.5E+14	50	14.0	6
CH ₃ CHO	1.0E+15	50	8.0	6
HCOOH	0.2E+16	40	8.0	6
CH ₃ CCH	2.8E+15	60	14.0	4
Additional species				
CH ₃ COCH ₃	1E+17	100	2.0	7
HCCCN	7E+13	150	14	7
SO	7E+14	50	14	8
³⁴ SO	1E+14	50	14.0	6
SO ₂	0.1E+16	50	14.0	7
³³ SO ₂	1E+15	50	14.0	6
HCN	2E+14	50	14.0	12
HCN	-6E+13	50	14.0	6
H ¹³ CN	2E+13	50	14.0	6
CN, $v = 0, 1$	0.8E+15	50	14	6
NH ₂ CN	0.3E+14	50	14	7
O ¹³ CS	5E+14	50	14	5
OC ³⁴ S	1E+15	50	14.0	7
CP	1E+15	50	14.0	6
C ³⁴ S	2E+14	50	14.0	7
HCOCH ₂ OH	0.2E+17	300	2.0	7
S ₃	1.5E+16	15	14	6
CH ₃ NH ₂	1E+17	150	2.0	7

Notes. ¹³CH₃OH is not available in the JPL database used for Weeds modeling. ⁽¹⁾ Based on 21'' beam spectra. ⁽²⁾ Based on 21'' beam spectra, 1.5E+17 in the 14'' beam spectra.

Appendix D: Additional detections

Several lines of other species were found in the observed frequency ranges, particularly in the line-rich G31.41+0.31. Table D.1 lists additional detections with column densities obtained from the Weeds analysis with the best T_{ex} when this could be derived. Several transitions of acetone, CH₃COCH₃, are detected. In G31, the best agreement is obtained at 100 K, whereas for IRAS 18089, this is $T_{\text{ex}} = 300$ K. No CH₃COCH₃ was detected in IRAS 20126, and the tabulated value is an upper limit at 300 K. Figure D.1 shows the strongest acetone lines in G31 together with the Weeds model.

Glycoaldehyde, HCOCH₂OH, has previously been detected in G31 (Beltrán et al. 2009), has several lines in the covered ranges. All lines are however blended with other transitions, and only upper limit could therefore be derived reliably. The tabulated column density is for a temperature of 300 K, constrained by non-detection of lines with low E_{up} .

Two transitions of HC₃N are detected at 218.325 GHz (E_{up} 131.0 K) ($J = 24 \rightarrow 23$) and 354.697 GHz (E_{up} 340.5 K) ($J = 39 \rightarrow 38$). The Weeds model on the two lines, well separated in E_{up} , gives a beam averaged column density of

Fig. D.1. Beam-averaged spectra of CH₃COCH₃ ($23_{1/2,22,1}-22_{1/2,21,1}$), ($23_{1/2,22,0}-22_{1/2,21,1}$), ($28_{7/8,21,0}-27_{8/7,20,0}$) and ($28_{7/8,21,0}-27_{8/7,20,1}$) towards G31.41+0.31 hot core. Dashed lines indicate Gaussian fits to the lines and red solid line show the Weeds model on the CH₃COCH₃ lines at 100 K.

$7 \times 10^{13} \text{ cm}^{-2}$ and a temperature of 150 K for G31. The HC_3N emission cannot be matched with emission contained within a $2.0''$ volume as the low- E_{up} transition becomes optically thick. For IRAS 20126, the tabulated column density is at 100 K, constrained by the two transitions. Only the high E_{up} transition was covered for IRAS 18089 giving a column density of $3 \times 10^{13} \text{ cm}^{-2}$ at 150 K and $6 \times 10^{13} \text{ cm}^{-2}$ at 100 K.

Several CH_3NH_2 transitions are observed with a T_{ex} of 150 K. Blends with other species make the rotation temperature and column density inaccurate, however. Several unidentified lines coincide with $\text{NH}_2\text{CH}_2\text{CH}_2\text{OH}$, but only upper limits of $1.0 \times 10^{16} \text{ cm}^{-2}$ at 100–300 K could be derived reliably using Weeds.

Several weak lines of NH_2CN are detected, with a T_{ex} of 50 K derived from the Weeds model, but line blends make the rotation temperature and column density inaccurate. For CN, SO, ^{34}SO , O^{13}CS , OC^{34}S , SO_2 , $^{33}\text{SO}_2$, C^{34}S , and CP not enough lines were observed to derive T_{ex} values, and tabulated column densities are assuming a temperature of 50 K.

One line of HCN, $J = 4-3$, was observed. HCN emission in G31 is composed of a broad (12 km s^{-1}) and a narrow (6 km s^{-1}) component (redshifted by 2.5 km s^{-1}) causing self-absorption. For IRAS 18089, the broad component has a width of 7 km s^{-1} and narrow 4 km s^{-1} (redshifted by 1 km s^{-1}). For IRAS 20126 the HCN emission has a very broad 20 km s^{-1} component blueshifted by 2 km s^{-1} and a narrow (5 km s^{-1}) component redshifted by 1 km s^{-1} . The tabulated column densities are those derived from the H^{13}CN column density.

Table D.1. Additional detections. Column densities given are source averaged for species with $T_{\text{ex}} > 100 \text{ K}$ and beam averaged for species with $T_{\text{ex}} < 100 \text{ K}$ (except for HCCCN, the emission of which is warm and arises from extended volume). The tabulated values are those obtained from Weeds analysis with the best T_{ex} .

Species	IRAS 20126+4104	IRAS 18089-1732	G31.41+0.31
Source-averaged column densities			
CH_3COCH_3	$<0.1\text{E}+17$	$0.5\text{E}+17$	$1.0\text{E}+17$
HCOCH_2OH	–	$<0.5\text{E}+16$	$<0.2\text{E}+17$
CH_3NH_2	–	–	$2.4\text{E}+14$
$\text{NH}_2\text{CH}_2\text{CH}_2\text{OH}$	–	–	$<1.0\text{E}+16$
Beam-averaged column densities			
NH_2CN	–	$<0.1\text{E}+14$	$7.0\text{E}+14$
OC-13-S	–	–	$5.0\text{E}+14$
OC-34-S	–	$0.5\text{E}+15$	$1.0\text{E}+15$
HCCCN	$1.0\text{E}+13$	$3.0\text{E}+13$	$7.0\text{E}+13$
S_3	–	–	$1.5\text{E}+16$
$\text{CN}, v = 0, 1$	$0.2\text{E}+15$	$0.4\text{E}+15$	$0.8\text{E}+15$
SO_2	$0.3\text{E}+15$	$0.5\text{E}+15$	$0.1\text{E}+16$
SO	$0.3\text{E}+15$	$1.0\text{E}+15$	$7.0\text{E}+14$
CS-34	–	$1.4\text{E}+14$	$2.4\text{E}+14$
CP	–	–	$1.0\text{E}+15$
S-33-O ₂	$<0.5\text{E}+14$	$0.5\text{E}+15$	$1.0\text{E}+15$
S-34-O	$0.2\text{E}+14$	$0.5\text{E}+14$	$1.0\text{E}+14$
HCN	$8\text{E}+13$	$2.0\text{E}+14$	$2.0\text{E}+14$
HC-13-N	$0.1\text{E}+13$	$1.5\text{E}+13$	$2.0\text{E}+13$