Dust emission in star-forming dwarf galaxies: General properties and the nature of the submm excess[★]

Y. I. Izotov^{1,2,3}, N. G. Guseva^{1,2}, K. J. Fricke^{1,4}, E. Krügel¹, and C. Henkel^{1,5}

¹ Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, 53121 Bonn, Germany
 e-mail: izotov@mao.kiev.ua

² Main Astronomical Observatory, Ukrainian National Academy of Sciences, Zabolotnoho 27, 03680 Kyiv, Ukraine

³ LUTH, Observatoire de Paris, CNRS, Université Paris Diderot, Place Jules Janssenn, 92190 Meudon, France

⁴ Institut für Astrophysik, Göttingen Universität, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany

⁵ Astronomy Department, King Abdulaziz University, PO Box 80203, 21589 Jeddah, Saudi Arabia

Received 30 January 2014 / Accepted 5 August 2014

ABSTRACT

We studied the global characteristics of dust emission in a large sample of emission-line star-forming galaxies. The sample consists of two subsamples. One subsample (SDSS sample) includes ~4000 compact star-forming galaxies from the Sloan Digital Sky Survey (SDSS), which were also detected in all four bands at 3.4 μm , 4.6 μm , 12 μm , and 22 μm of the WISE all-sky survey. The second subsample (*Herschel* sample) is a sample of 28 compact star-forming galaxies observed with *Herschel* in the FIR range. Data of the *Herschel* sample were supplemented by the photometric data from the *Spitzer* observations, GALaxy Evolution Explorer (GALEX) survey, SDSS, Wide-Field Infrared Survey Explorer (WISE), Two Micron All Sky Survey (2MASS), NRAO VLA Sky Survey (NVSS), and Faint Images of the Radio Sky at Twenty-cm (FIRST) survey, as well as optical and *Spitzer* spectra and data in submm and radio ranges. It is found that warm dust luminosities of galaxies from the SDSS sample and cold and warm dust luminosities of galaxies from the *Herschel* sample are strongly correlated with H β luminosities, which implies that one of the main sources of dust heating in star-forming galaxies is ionising UV radiation of young stars. Likely, a significant fraction of dust is located inside H II and surrounding regions. We found tight correlations between masses of cold and warm dust, again implying the same source of dust heating. Using the relation between warm and cold dust masses for estimating the total dust mass in star-forming galaxies with an accuracy better than ~0.5 dex is proposed. On the other hand, it is shown for both samples that dust temperatures do not depend on the metallicities. The dust-to-neutral gas mass ratio strongly declines with decreasing metallicity, similar to that found in other studies of local emission-line galaxies, high-redshift gamma-ray burst hosts, and damped Ly- α absorbers. On the other hand, the dust-to-ionised gas mass ratio is about one hundred times as high implying that most of dust is located in the neutral gas. It is found that thermal free-free emission of ionised gas in compact star-forming galaxies is important in the submm and mm ranges, and it might be responsible for the submm emission excess. This effect is stronger in galaxies with lower metallicities and is also positively affected by an increased star-formation rate.

Key words. galaxies: fundamental parameters – galaxies: starburst – galaxies: ISM – galaxies: abundances

1. Introduction

Dust plays an important role in the thermal and dynamical evolution of galaxies at different spatial scales from protostars to large molecular complexes. Observations with ground-based and space-born telescopes, Infrared Astronomical Satellite (IRAS), Infrared Space Observatory (ISO), Submillimetre Common-User Bolometer Array (SCUBA), *Spitzer*, and *Herschel*, revealed significant populations of dusty luminous and ultraluminous infrared galaxies at redshifts $z \gtrsim 1$ (e.g. [Hauser & Dwek 2001](#); [Elbaz et al. 2002](#); [Dole et al. 2006](#); [Frayser et al. 2006](#)), indicating that dust grains are being formed in intense and short star-formation episodes and implying a rapid transition in high-redshift galaxies from the formation of zero-metal population III stars to the formation of stellar populations more typical for present-day galaxies (e.g. [Nozawa et al. 2003](#); [Schneider et al. 2012](#); [Chiaki et al. 2014](#)). However, the dust formation in the early universe is debated. In particular, [Valiante et al. \(2011\)](#)

find that population III stars do not contribute much to the dust budget. They claim that the dust is formed by population II/I, SNe, and asymptotic giant branch (AGB) stars at the later stage.

Studies of high-redshift low-metallicity galaxies are very difficult because of their faintness and small angular sizes. We have a chance to study star formation and role of dust in low-metallicity environments by investigating properties of nearby star-forming dwarf galaxies, such as irregular, blue compact dwarf (BCD), and green pea (GP) galaxies. These galaxies can be considered as local counterparts or “analogs” of high-redshift galaxies, such as Lyman-break galaxies (LBGs) and Lyman- α emitting galaxies, because of their high star-formation activity and low metallicity (e.g. [Cardamone et al. 2009](#); [Izotov et al. 2011a](#)). Due to their proximity they can be studied in greater detail than it is possible in distant galaxies. Studying the properties of a metal-poor interstellar medium (ISM), we may be able to better understand the physical conditions in high-redshift galaxies.

The properties of radiation in the mid- and far-infrared ranges of star-forming dwarf galaxies were studied in many

[★] Figure 6 is available in electronic form at <http://www.aanda.org>

papers by considering samples of galaxies and individual galaxies (e.g. [Draine et al. 2007](#); [Engelbracht et al. 2005, 2008](#); [Cannon et al. 2006](#); [Hunt et al. 2005a, 2010](#); [Galliano et al. 2008](#); [Galamez et al. 2009, 2011](#); [Madden et al. 2006, 2013](#); [Rémy-Ruyer et al. 2013, 2014](#); [Wu et al. 2006, 2007, 2008](#); [Hao et al. 2009](#); [O'Halloran et al. 2010](#); [Cormier et al. 2012](#); [Indebetouw et al. 2009, 2013](#); [Karczewski et al. 2013](#)). Spectral energy distributions (SEDs) in these ranges are dominated by dust emission resulting from reprocessing of UV radiation from young massive stars, in particular in the most metal-deficient BCDs I Zw 18 and SBS 0335–052E ([Wu et al. 2007](#); [Houck et al. 2004](#)).

It was concluded in numerous papers that the main dust heating source in dwarf emission-line galaxies is the radiation from young stellar populations. Dust in these galaxies is heated to higher temperatures of ~ 30 – 40 K as compared to ~ 20 K in spirals (e.g. [Engelbracht et al. 2008](#); [Hirashita et al. 2008](#)). The polycyclic aromatic hydrocarbon (PAH) emission in dwarf star-forming galaxies is weak or absent, and it decreases with decreasing metallicity (e.g. [Wu et al. 2007](#); [Hunt et al. 2010](#)). Different mechanisms were invoked to explain the weakness of PAH features in these galaxies: destruction by the intense UV radiation ([Madden et al. 2006](#)), destruction by shocks ([O'Halloran et al. 2006](#)), delayed formation by AGB stars ([Galliano et al. 2008](#)), and formation of molecular clouds ([Sandstrom et al. 2010](#)). It was also found that the dust emission in dwarf galaxies cannot be characterised by radiation with a single temperature. Warm and hot dust with temperatures of up to several hundred degrees must be present in many of these galaxies in addition to cold dust to reproduce the SED in the mid- and far-infrared (FIR) ranges (e.g. [Griffith et al. 2011](#); [Izotov et al. 2011b, 2014](#)). It is most likely that dust temperature gradients are present in the ISM around massive and compact stellar clusters (e.g. [Hunt et al. 2014](#)).

The mid- and far-infrared emission in dwarf galaxies results from absorption of radiation at shorter wavelengths in the UV and optical ranges. It includes not only the stellar component and nebular emission lines but also the nebular continuum (free-bound and free-free emission). The necessity to include the latter in young starbursts was discussed by [Izotov et al. \(2011a\)](#) as otherwise stellar masses may be overestimated by a factor of up to ~ 3 .

It also was established that the fraction of nebular continuum increases with wavelength. Nebular continuum dominates at $\lambda 1$ – $3\ \mu\text{m}$ with a flux fraction of 70–90% in extremely young starbursts, which are characterised by high equivalent widths $EW(\text{H}\beta) > 100\ \text{\AA}$ of the $\text{H}\beta$ emission line. Ignoring this result would lead to wrong conclusions. For example, [Engelbracht et al. \(2008\)](#) suggested that dust already dominates at $1.8\ \mu\text{m}$ in SBS 0335–052E, Haro 11, and SHOC 391, while it is clear from detailed fitting of optical spectra that the main contributor to the continuum in the near-infrared (NIR) range is ionised gas.

Therefore, photometric data are not sufficient for correct SED fitting in the optical range; spectra are needed as well. Using spectra and the $\text{H}\beta$ emission line flux, one can also predict the flux of free-free emission in the radio range ([Caplan & Deharveng 1986](#)) and can estimate how much star formation is hidden in the UV and optical ranges.

Among the first studies, where detailed fitting of UV and optical SEDs, which included both stellar and gaseous continua, was applied for analysis of optical and infrared properties of large samples of compact star-forming galaxies were those performed by [Izotov et al. \(2011b\)](#) and [Izotov et al. \(2014\)](#). In particular, [Izotov et al. \(2014\)](#) considered a sample

of $\sim 14\,000$ SDSS galaxies with strong emission lines and found that mid- and FIR luminosities are strongly correlated with the luminosity of ionising stellar radiation.

This study is the continuation of the work by [Izotov et al. \(2011b\)](#) and [Izotov et al. \(2014\)](#), where relations between luminosities and optical and NIR colour characteristics were mainly discussed. Here we concentrate on the determination of the dust characteristics, such as temperature and mass. We use two samples. The first sample selected from the SDSS consists of compact star-forming galaxies with available mid-infrared data from the WISE all-sky survey. This sample was discussed by [Izotov et al. \(2014\)](#). The second sample consists of nearby star-forming galaxies observed with *Herschel* by [Madden et al. \(2013\)](#) and [Rémy-Ruyer et al. \(2013\)](#).

The criteria for selection and properties of both samples are discussed in Sect. 2. We describe the determination of galaxy parameters in Sect. 3. Results of this study are discussed in Sect. 4. Our findings are summarised in Sect. 5.

2. The samples

2.1. SDSS sample

We use a sample of $\sim 14\,000$ star-forming galaxies selected by [Izotov et al. \(2014\)](#) from the spectroscopic database of the SDSS data release 7 (DR7; [Abazajian et al. 2009](#)). The details of data selection can be found in [Izotov et al. \(2014\)](#). Out of this sample, we select ~ 4000 compact galaxies with angular diameters $\leq 6''$, which were detected in all four WISE bands at $\lambda 3.4\ \mu\text{m}$, $\lambda 4.6\ \mu\text{m}$, $\lambda 12\ \mu\text{m}$, and $\lambda 22\ \mu\text{m}$, allowing the determination of the warm and hot dust properties. The SDSS spectra and SDSS u , g , r , i , and z magnitudes of all selected galaxies are available in the SDSS database. Finally, we restrict the sample to only those ~ 1000 galaxies out of the ~ 4000 , where the $[\text{O III}] \lambda 4363\ \text{\AA}$ emission line was measured with an accuracy better than 50%. This allows for a reliable determination of the oxygen abundance. Hydrogen emission lines in these galaxies are strong allowing for a correction for dust extinction from the decrement of several hydrogen lines.

2.2. Herschel sample

We use a sample of 48 star-forming galaxies studied by [Rémy-Ruyer et al. \(2013\)](#) with the *Herschel* space observatory and adopt their *Herschel* FIR fluxes and distances. This sample was supplemented by the data on the total fluxes from other photometric surveys, namely from the GALEX¹ ([Martin et al. 2005](#)), SDSS² ([Abazajian et al. 2009](#)), WISE³ ([Wright et al. 2010](#)), 2MASS⁴ ([Skrutskie et al. 2006](#)), NVSS⁵ ([Condon et al. 1998](#)) and FIRST⁶ ([Kimball & Ivezić 2008](#)) surveys. Furthermore, *Spitzer* spectra⁷ ([Lebouteiller et al. 2011](#)) in the mid-infrared (MIR) range, spectra in the optical range, and, whenever it was possible, data in the submm and radio ranges at different wavelengths were collected. Out of 48 galaxies by

¹ <http://galex.stsci.edu/GR6/>

² <http://skyserver.sdss3.org/dr10/en/tools/chart/chartinfo.aspx>

³ <http://irsa.ipac.caltech.edu/cgi-bin/Gator/nph-scan?mission=irsa&submit=Select&projshort=WISE>

⁴ <http://irsa.ipac.caltech.edu/cgi-bin/Gator/nph-scan?mission=irsa&submit=Select&projshort=2MASS>

⁵ <http://www.cv.nrao.edu/nvss/>

⁶ <http://sundog.stsci.edu/cgi-bin/searchfirst>

⁷ <http://cassis.sirtf.com/atlas/index.shtml>

Table 1. References to the data for galaxies from the *Herschel* sample.

Object	RA (J2000)	Dec (J2000)	Distance	References				
				H β flux	Optical spectra	<i>Spitzer</i> photometry	submm and radio	H I 21 cm
Haro 11	00h36m52.7s	−33d33m17.0s	92.1	1	2	3	4, 1, 5, 6	53
Haro 3	10h45m22.4s	+55d57m37.0s	19.3	7	8	9	5, 54, 57	10
HS 0017+1055	00h20m21.4s	+11d12m21.0s	79.1	11	11	...	57	...
HS 0052+2536	00h54m56.4s	+25d53m08.0s	191.0	11	11	...	...	...
HS 0822+3542	08h25m55.5s	+35d32m32.0s	11.0	7	12	3	57	13
HS 1222+3741	12h24m36.7s	+37d24m37.0s	181.7	12	12	...	57	...
HS 1304+3529	13h06m24.2s	+35d13m43.0s	78.7	12	12	...	57	...
HS 1330+3651	13h33m08.3s	+36d36m33.0s	79.7	12	12	...	57	...
I Zw 18	09h34m02.0s	+55d14m28.0s	18.2	7	12	3	14, 16, 57	10
II Zw 40	05h55m42.6s	+03d23m32.0s	12.1	17	18	3	19, 20, 21, 22, 23, 5, 54	24
Mrk 1089	05h01m37.7s	−04d15m28.0s	56.6	25	26	...	1, 5, 27, 54	29
Mrk 1450	11h38m35.7s	+57d52m27.0s	19.8	7	30	3	5, 57	31
Mrk 153	10h49m05.0s	+52d20m08.0s	40.3	7	12	3	5, 57	32
Mrk 209	12h26m15.9s	+48d29m37.0s	5.8	7	12	33	5, 55, 56, 57	10
Mrk 930	23h31m58.3s	+28h56m50.0s	77.8	7	8	3	34, 5	35
NGC 1140	02h54m33.6s	−10d01m40.0s	20.0	7	18	3	19, 20, 28, 5	24
Pox 186	13h25m48.6s	−11d36m38.0s	18.3	37	37	...	...	...
SBS 0335−052E	03h37m44.0s	−05d02m40.0s	56.0	38	39	3	40, 41, 5, 54	42
SBS 1159+545	12h02m02.4s	+54d15m50.0s	57.0	30	30	...	57	...
SBS 1211+540	12h14m02.5s	+53d45m17.0s	19.3	30	12	...	57	35
SBS 1249+493	12h51m52.5s	+49d03m28.0s	110.8	43	43	...	57	52
SBS 1415+437	14h17m01.4s	+43d30m05.0s	13.6	44	43	...	57	35
SBS 1533+574	15h34m13.8s	+57d17m06.0s	54.2	45	45	...	57	35
Tol 1214−277	12h17m17.1s	−28d02m33.0s	120.5	46	47	3	...	...
UGC 4483	08h37m03.0s	+69d46m31.0s	3.2	7	30	3	...	48
UM 448	11h42m12.4s	+00d20m03.0s	87.8	7	39	3	28, 49, 5, 57, 58	10
UM 461	11h51m33.3s	−02d22m22.0s	13.2	7	12	3	57	50
VII Zw 403	11h27m59.9s	+78d59m39.0s	4.5	7	45	3	51, 59	10

Notes. Distances and *Herschel* photometric data are from Rémy-Ruyer et al. (2013). GALEX photometric data are from <http://galex.stsci.edu/GR6/>. SDSS photometric data are from <http://skyserver.sdss3.org/dr10/en/tools/chart/chartinfo.aspx>. WISE data are from <http://irsa.ipac.caltech.edu/cgi-bin/Gator/nph-scan?mission=irsa&submit=Select&projshort=WISE>. 2MASS data are from <http://irsa.ipac.caltech.edu/cgi-bin/Gator/nph-scan?mission=irsa&submit=Select&projshort=2MASS>. NVSS and FIRST data in the continuum at 20 cm are from <http://www.cv.nrao.edu/nvss/> and <http://sundog.stsci.edu/cgi-bin/searchfirst>, respectively; low-resolution *Spitzer* spectra are from <http://cassis.sirtf.com/atlas/index.shtml>

References. (1) Schmitt et al. (2006); (2) Guseva et al. (2012); (3) Engelbracht et al. (2008); (4) Galametz et al. (2009); (5) Condon et al. (1998); (6) Madden et al. (2006); (7) Moustakas & Kennicutt (2006); (8) Izotov & Thuan (2004); (9) Hunt et al. (2006); (10) Huchtmeier & Richter (1989); (11) Ugryumov et al. (2003); (12) Izotov et al. (2006a), <http://das.sdss.org/spectro/>; (13) Chengalur et al. (2006); (14) Leroy et al. (2007); (15) Cannon et al. (2005); (16) Hunt et al. (2005b); (17) Lagos et al. (2007); (18) Guseva et al. (2000); (19) Galliano et al. (2005); (20) Albrecht et al. (2004); (21) Hirashita (2011); (22) Hunt et al. (2005b); (23) Kepley et al. (2014); (24) Springob et al. (2005); (25) Vacca & Conti (1992); (26) Izotov & Thuan (1999); (27) Galametz et al. (2009); (28) Bica et al. (1995); (29) Williams et al. (1991); (30) Izotov et al. (1994); (31) Huchtmeier et al. (2005); (32) Thuan & Martin (1981); (33) Dale et al. (2009); (34) Rosa-Gonzalez et al. (2007); (35) Thuan et al. (1999b); (36) Hunter et al. (1994); (37) Guseva et al. (2004); (38) Izotov et al. (2006b); (39) Izotov & Thuan (1998); (40) Johnson et al. (2009); (41) Hunt et al. (2014); (42) Ekta et al. (2009); (43) Thuan et al. (1995); (44) Gil de Paz et al. (2003); (45) Izotov et al. (1997); (46) Guseva et al. (2011); (47) Izotov et al. (2001); (48) de Vaucouleurs et al. (1991); (49) Dressel & Condon (1978); (50) Courtois et al. (2011); (51) Thuan et al. (2004); (52) Pustilnik et al. (2002); (53) Cormier et al. (2014); (54) Rabidoux et al. (2014); (55) Viallefond & Thuan (1983); (56) Klein et al. (1991); (57) Kimball & Ivezić (2008); (58) James et al. (2002); (59) Leroy et al. (2005).

Rémy-Ruyer et al. (2013), we excluded galaxies without optical or *Spitzer* spectra, galaxies with large angular radii $\geq 60''$, and galaxies, which were not detected by *Herschel* in any of the PACS or SPIRE photometric bands. Furthermore, we excluded the compact object UM 311 because this is not a galaxy, but an H II region within a spiral galaxy.

In the end, the *Herschel* sample consists of 28 compact galaxies with coordinates and distances shown in Table 1.

References on available photometric and spectroscopic data are also given in the table.

3. The determination of galaxy parameters

Most global galaxy parameters (metallicities, stellar masses, H β luminosities and UV, optical, infrared and radio monochromatic luminosities) for the SDSS sample were derived by

Izotov et al. (2014), which we adopt in this paper. We apply the same technique to derive galaxy parameters for the *Herschel* sample. The technique is briefly described in this section.

3.1. Element abundances and H β luminosity

The SDSS spectra of the galaxies from the SDSS sample are used to derive emission-line fluxes and equivalent widths, the extinction coefficient $C(\text{H}\beta)$, the H II region luminosity in the H β emission line, and chemical element abundances. The observed line fluxes were obtained using the IRAF⁸ SPLOT routine. The line flux errors included statistical errors in addition to errors introduced by the standard star absolute flux calibration, which we set to 1% of the line fluxes. These errors are later propagated into the calculation of abundance errors. The line fluxes were corrected for the Milky Way and internal reddening using the extinction curve of Cardelli et al. (1989) and for underlying hydrogen stellar absorption. The extinction coefficients are defined as $C(\text{H}\beta) = 1.47E(B - V)$, where $E(B - V) = A(V)/R_V$ and $R_V = 3.2$ (Aller 1984).

The oxygen abundances were derived according to the procedures described by Izotov et al. (1994, 1997) and Thuan et al. (1995), where a two-zone photoionised H II region model was adopted: a high-ionisation zone with temperature $T_e(\text{O III})$, and a low-ionisation zone with temperature $T_e(\text{O II})$. In the H II regions with a detected [O III] $\lambda 4363$ emission line, the temperature $T_e(\text{O III})$ was calculated using the direct method based on the [O III] $\lambda 4363/(\lambda 4959 + \lambda 5007)$ line ratio. The electron temperature $T_e(\text{O III})$ in H II regions with non-detected [O III] $\lambda 4363$ emission line was derived by a semi-empirical method (Izotov & Thuan 2007; Izotov et al. 2014), which is based on the relation between $T_e(\text{O III})$ and the total flux of strong [O II] $\lambda 3727$, [O III] $\lambda 4959$, and [O III] $\lambda 5007$ emission lines.

We use the relation between the electron temperatures $T_e(\text{O III})$ and $T_e(\text{O II})$ obtained by Izotov et al. (2006a) from the H II region models of Stasińska & Izotov (2003) to derive $T_e(\text{O II})$. Ionic and total oxygen abundances were derived using expressions for ionic abundances and ionisation correction factors (ICFs) obtained by Izotov et al. (2006a).

The extinction-corrected luminosity $L(\text{H}\beta)$ was obtained from the observed H β emission-line flux by adopting the total extinction, which included the Milky Way, internal galaxy extinctions, and the distance derived from the redshift. For distance determination, Izotov et al. (2014) used the relation $D = f(z, H_0, \Omega_M, \Omega_\Lambda)$ from Refsdal et al. (1967), where the Hubble constant $H_0 = 67.3 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and cosmological parameters $\Omega_M = 0.273$, $\Omega_\Lambda = 0.682$ were obtained from the *Planck* mission data (Planck Collaboration XV 2014). The equivalent widths $\text{EW}(\text{H}\beta)$ were reduced to the rest frame.

The SDSS spectra were obtained with a small aperture of $3''$ in diameter. To derive integrated characteristics of the galaxies from their spectra and to make a comparison with photometric data more accurate, Izotov et al. (2014) corrected spectroscopic data for the aperture using the relation $2.5^{r(3'')-r}$, where r and $r(3'')$ are the SDSS r -band total magnitude and the magnitude within the $3''$ spectroscopic aperture, respectively.

The SDSS spectra for some galaxies from the *Herschel* sample were also available. Izotov et al. (2014) used these

data to derive the same parameters as those for SDSS sample galaxies. In other cases, we used optical spectra obtained with different telescopes to derive extinction, physical conditions and element abundances. The H β fluxes were obtained from SDSS spectra only for three *Herschel* galaxies, HS 1222+3741, HS 1304+3529, and HS 1330+3651. For other galaxies, we adopted the observed H β fluxes from the literature, which were obtained with the largest available apertures and corrected them for the extinction derived from the optical spectra. References on the optical spectra and H β fluxes are given in Table 1.

3.2. Stellar masses

Izotov et al. (2014) derived galaxy stellar masses by modelling the galaxy SEDs in the optical range for each galaxy from the SDSS sample. They took into account both the stellar and ionised gas emission.

The method is based on fitting a series of model SEDs to the observed one and finding the best fit. The fit, described in more detail by Guseva et al. (2006, 2007) and Izotov et al. (2011a, 2014), was performed for each SDSS spectrum over the whole observed spectral range of $\lambda\lambda 3900\text{--}9200 \text{ \AA}$. As each SED is the sum of both stellar and ionised gas emission, its shape depends on the relative contribution of these two components. The contribution of gaseous emission relative to stellar emission can be parameterized by the equivalent width $\text{EW}(\text{H}\beta)$ of the H β emission line. The shape of the spectrum depends also on reddening and the star-formation history of the galaxy.

Izotov et al. (2014) approximated the star-formation history in each galaxy by a recent short burst with age $< 10 \text{ Myr}$, which accounts for the young stellar population, and a prior continuous star formation responsible for the older stars with age $\geq 10 \text{ Myr}$. The contribution of each stellar population to the SED was parameterized by the varying ratio of the masses of the young to old stellar populations, M_y/M_o . Then the total stellar mass is equal to $M_* = M_y + M_o$. We adopted M_* and M_y , calculated by Izotov et al. (2014) for SDSS sample galaxies, where details of the SED fitting can be found.

For the galaxies from the *Herschel* sample, we derive stellar masses, using the same method of SED fitting and spectra as cited in Table 1.

3.3. Integrated dust characteristics

To derive integrated dust characteristics in our SDSS and *Herschel* samples we use the modified blackbody fitting technique, similar to that used by Rémy-Ruyer et al. (2013).

Rémy-Ruyer et al. (2013) fitted the *Herschel* data by adopting a single dust temperature T . This is sufficient to fit far-infrared fluxes and to derive the mass of the cold dust. However, there is evidence for warm and hot dust in star-forming dwarf galaxies (e.g. Griffith et al. 2011; Izotov et al. 2011b, 2014; Hunt et al. 2014).

Therefore, we adopted a three-component model with the temperatures T_{cd} , T_{wd} , and T_{hd} to fit dust emission in the wavelength range $\sim 3\text{--}500 \text{ }\mu\text{m}$ covered by WISE, *Spitzer*, and *Herschel* observations:

$$F_\nu = \frac{a_{\text{cd}}\nu^{3+\beta}}{\exp(h\nu/kT_{\text{cd}}) - 1} + \frac{a_{\text{wd}}\nu^{3+\beta}}{\exp(h\nu/kT_{\text{wd}}) - 1} + \frac{a_{\text{hd}}\nu^{3+\beta}}{\exp(h\nu/kT_{\text{hd}}) - 1}, \quad (1)$$

⁸ IRAF is the Image Reduction and Analysis Facility distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy (AURA) under cooperative agreement with the National Science Foundation (NSF).

where h and k are Planck and Boltzman constants, respectively, β is the emissivity index, a_{cd} , and a_{wd} , and a_{hd} are scaling coefficients.

The cold dust component with the temperature T_{cd} contributes mainly at wavelengths $\geq 70 \mu\text{m}$ covered by *Spitzer*/MIPS $70 \mu\text{m}$ and $160 \mu\text{m}$ and *Herschel* observations. The warm dust component with the temperature T_{wd} strongly contributes in the wavelength range $\sim 10\text{--}50 \mu\text{m}$ covered mainly by the WISE $12 \mu\text{m}$ and $22 \mu\text{m}$, *Spitzer*/MIPS $24 \mu\text{m}$ and *Spitzer*/IRS observations. Finally, the hot dust component with the temperature T_{hd} contributes in the wavelength range $< 10 \mu\text{m}$, which is covered by the WISE $3.4 \mu\text{m}$ and $4.6 \mu\text{m}$, and *Spitzer*/IRAC observations. In reality, there is a gradient in dust temperature. However, we show later that a three-component model is sufficient to fit dust emission in the entire mid- and FIR ranges.

We use Monte Carlo simulations to derive T_{cd} , T_{wd} , and T_{hd} , and scaling coefficients a_{cd} , a_{wd} , and a_{hd} by minimization of χ^2 :

$$\chi^2 = \sum_{j=1}^m \frac{[F_j(\text{obs}) - F_j(\text{mod})]^2}{\sigma_j^2(\text{obs})}, \quad (2)$$

where $F_j(\text{obs})$ and $F_j(\text{mod})$ are observed and modelled fluxes, σ_j is the uncertainty of the observed flux, and m is the number of observed fluxes.

The 1σ errors (68.3% confidence level) of T_{cd} , T_{wd} , T_{hd} , a_{cd} , a_{wd} , and a_{hd} are estimated by adopting $\Delta\chi^2 = \chi^2 - \chi_{\text{min}}^2 = 7.04$ for six degrees of freedom. For some objects, we considered SED modelling with a two-component dust, excluding the hot component. In this case, $\Delta\chi^2 = 4.72$ for four degrees of freedom. Derived errors of temperatures and scaling coefficients were then propagated to obtain errors for dust masses and luminosities.

The dust masses M_{cd} , M_{wd} , and M_{hd} of each component are given by

$$M_{\text{dust}} = \frac{D^2}{\kappa(\lambda_0)} \left(\frac{\lambda}{\lambda_0} \right)^\beta \frac{F_\nu(\lambda)}{B_\nu(\lambda, T_{\text{dust}})}, \quad (3)$$

where $\lambda_0 = 100 \mu\text{m}$, $\kappa(\lambda_0) = 34.7 \text{ cm}^2 \text{ g}^{-1}$ is the dust emissivity cross section per unit mass at $100 \mu\text{m}$ for the best fit SMC (Small Magellanic Cloud) dust (Weingartner & Draine 2001), D is the distance to the galaxy, $T_{\text{dust}} = T_{\text{cd}}$, T_{wd} or T_{hd} is the dust temperature, $B_\nu(\lambda, T_{\text{dust}})$ is the Planck function, and $F_\nu(\lambda)$ is the monochromatic flux of dust emission.

Rmy-Ruyer et al. (2013) varied the emissivity index β from 0 to 2.5 to achieve the best agreement between the observed and modelled cold dust emission. At wavelengths $\geq 20 \mu\text{m}$, the dust absorption coefficient is fairly well approximated by a power law ν^β fit with $\beta = 2.0$ (Draine 2003; Krgel 2008), and as this is the range where most of the energy is emitted, we use this value to calculate the dust luminosity of the cold component from

$$L(T_{\text{dust}}) = 4\pi M_{\text{dust}} \kappa(\lambda_0) \int \left(\frac{\lambda}{\lambda_0} \right)^{-\beta} B(\lambda, T_{\text{dust}}) d\lambda. \quad (4)$$

At shorter wavelengths, β is smaller and more variable, but, for simplicity, we use the same β also for the warm and hot component. Nevertheless we also discuss the expected changes if β is lower.

The photometric WISE and *Spitzer* fluxes at $\lesssim 15 \mu\text{m}$ prior to fitting were corrected for contributions from the stellar and gaseous (continuum and emission lines) emission. The stellar and gaseous continuum can be important at $\lambda \lesssim 5 \mu\text{m}$. We subtract it by using extrapolations of SED fits obtained from optical

spectra. However, if observed WISE $\lambda 3.4 \mu\text{m}$, $\lambda 4.6 \mu\text{m}$ fluxes and *Spitzer* $\lambda 3.6 \mu\text{m}$, $\lambda 4.5 \mu\text{m}$ fluxes are smaller than the extrapolated fluxes, we excluded them from the hot dust emission fitting. The stellar and gaseous continuum is negligible at longer wavelengths and does not affect the determination of warm and cold dust parameters.

The grid of CLOUDY spherical ionisation-bounded H II region models calculated in a wide range of input parameters (version c13.01, Ferland et al. 1998, 2013) is used to predict and subtract the strongest MIR emission lines, [Ar III] $\lambda 9.0 \mu\text{m}$, [S IV] $\lambda 10.51 \mu\text{m}$, and [Ne III] $\lambda 15.55 \mu\text{m}$ at their respective redshifted wavelengths. The range of parameters and a grid are described by Izotov et al. (2013). The input CLOUDY chemical abundances were derived from the optical spectra of each object. The equivalent widths of MIR emission lines ($\lesssim 0.1 \mu\text{m}$) are small as compared to the widths of the WISE photometric bands ($\sim 10 \mu\text{m}$); therefore, corrections for emission lines do not exceed $\sim 1\%$. The CLOUDY models also predict fluxes of PAH lines at $\lambda 3.3 \mu\text{m}$, $\lambda 6.2 \mu\text{m}$, $\lambda 7.9 \mu\text{m}$, $\lambda 11.3 \mu\text{m}$, $\lambda 11.8 \mu\text{m}$, and $\lambda 13.3 \mu\text{m}$. However, we do not subtract PAH emission from the observed fluxes due to the complexity of PAH features fitting. Our inspection of the *Spitzer*/IRS spectra shows that PAH emission is visible only in the six highest-metallicity galaxies from our *Herschel* sample with $12 + \log \text{O}/\text{H} \geq 8.1$. Neglecting corrections for PAH emission would overestimate the temperature of the hot dust component by at most 10%. Parameters of the warm and cold components will not be changed.

Since only SDSS optical, 2MASS NIR and WISE MIR data are available for the most of the SDSS sample galaxies, we cannot fit the cold dust component. Therefore, for this sample, we adopted a one-component dust model to derive characteristics of the warm dust from the flux ratio at $\lambda 12 \mu\text{m}$ and $\lambda 22 \mu\text{m}$. For a small number of 58 SDSS galaxies with red $3.4 \mu\text{m}\text{--}4.6 \mu\text{m}$ colours of ≥ 1.5 mag, implying the presence of hot dust, we also adopted a two-component dust model to derive characteristics of the hot and warm dust from the $\lambda 3.4 \mu\text{m}/\lambda 4.6 \mu\text{m}$ and $\lambda 12 \mu\text{m}/\lambda 22 \mu\text{m}$ flux ratios, respectively.

For the *Herschel* sample we use available data in the optical, NIR, MIR, FIR, submm and mm ranges to fit the observed data with the three-component dust model. However, the two-component dust model is sufficient for six galaxies. We note that we discuss the parameters of only cold and warm dust emission in all cases below. This is because hot dust emission may not be in thermal equilibrium. Therefore, its characteristics, such as apparent temperature and mass, may be misleading and do not correctly represent its true parameters.

4. Results

4.1. SDSS sample of compact galaxies

We use WISE $12 \mu\text{m}$ and $22 \mu\text{m}$ fluxes to derive the temperatures T_{wd} , the luminosities L_{wd} , and the masses M_{wd} of the warm dust.

In Fig. 1, we show the dependence of the warm dust temperature T_{wd} on the oxygen abundance $12 + \log \text{O}/\text{H}$ that is calculated with the emissivity index $\beta = 2.0$. Only ~ 1000 galaxies with reliably derived oxygen abundances are shown in the Figure. Galaxies with the H β equivalent width $EW(\text{H}\beta) \geq 50 \text{ \AA}$ and $< 50 \text{ \AA}$ are also represented. Most of the galaxies are spread in the T_{wd} interval of $\sim 80\text{--}150 \text{ K}$ with an average value of $\sim 120 \text{ K}$. The average temperature would be higher by $\sim 20 \text{ K}$ if $\beta = 1.0$ is adopted.

No clear dependence of dust temperature on $12 + \log \text{O}/\text{H}$ is found, implying that the metallicity is not a factor, which

Fig. 1. Dependence of the warm dust temperature T_{wd} on the oxygen abundance $12 + \log \text{O}/\text{H}$ in ~ 1000 galaxies from the SDSS sample. Temperatures are calculated from the WISE fluxes at $12 \mu\text{m}$ and $22 \mu\text{m}$ for galaxies with $\text{EW}(\text{H}\beta) \geq 50 \text{ \AA}$ (red filled circles) and $\text{EW}(\text{H}\beta) < 50 \text{ \AA}$ (blue filled circles). The solid line corresponds to average T_{wd} .

regulates dust emission. We also do not find differences between warm dust temperatures in galaxies with high-excitation and low-excitation H II regions.

Figures 2a and b respectively show dependencies of warm dust luminosities L_{wd} on total galaxy luminosities L_{tot} and on H β luminosity $L(\text{H}\beta)$ for ~ 1000 compact galaxies with reliable oxygen abundances. The total luminosity L_{tot} is calculated in the wavelength range $0.1\text{--}22 \mu\text{m}$ (Izotov et al. 2013). Linear most likelihood fits of the relations are shown by solid lines.

We find that the relation in Fig. 2b is tighter than that in Fig. 2a. The difference is more obvious from the comparison of Figs. 2c and d, where we show the entire SDSS sample of ~ 4000 compact star-forming galaxies. This difference indicates that warm dust is heated by the UV radiation of star-forming regions with a significant fraction of ionising stellar radiation. Thus much of the warm dust is likely associated with H II regions and neutral gas clouds surrounding these H II regions.

This conclusion is supported by the result that there is no offset between galaxies with high and low $\text{EW}(\text{H}\beta)$ (Figs. 2b, d), implying that the warm dust luminosity is determined by the $L(\text{H}\beta)$ luminosity which in turn is proportional to the luminosity of ionising radiation. The relations on the total luminosities, which include both ionising and non-ionising radiation, are broader with clear separation of galaxies with high and low $\text{EW}(\text{H}\beta)$ (Figs. 2a, c).

The dependence of the warm dust mass M_{wd} , calculated with $\beta = 2.0$, on total stellar mass M_* is shown in Fig. 3a for galaxies with high and low $\text{EW}(\text{H}\beta)$, respectively. The warm dust mass would be increased by a factor of ~ 2 if $\beta = 1.0$ is adopted. At a variance with the $L_{\text{wd}} - L_{\text{tot}}$ relation, the correlation $M_{\text{wd}} - M_*$ is much weaker. This can be due to the presence of a faint old stellar population, not participating in the heating of warm dust. Its contribution to the stellar mass can be high, while it does not contribute much to the total luminosity because of the steeper than linear relation between stellar mass and luminosity.

The dependence of M_{wd} on the mass of the young stellar population M_y is much tighter (Fig. 3b) supporting the conclusion that warm dust in SDSS compact dwarf galaxies is heated to a large extent by the radiation from young stars. Similarly to Fig. 3a, the warm dust mass is increased by a factor of ~ 2 , if an emissivity index β of 1.0 is adopted. However, a clear offset is present in both panels of Fig. 3 – the mass of warm dust in

galaxies with lower $\text{EW}(\text{H}\beta)$ is lower than that in galaxies with high $\text{EW}(\text{H}\beta)$.

Relations L_{wd} vs. $L(\text{H}\beta)$ (Fig. 2b) and M_{wd} vs. M_y (Fig. 3b) are steeper than linear relations. We may assume that the non-linearity of the L_{wd} vs. $L(\text{H}\beta)$ relation is due to the result that the warm dust in more luminous and more massive starbursts is warmer and is thus characterised by higher emissivity (Fig. 4a). However, in this case, one would expect a shallower than linear relation M_{wd} vs. M_y because of the positive correlation between T_{wd} and M_y (Fig. 4b) although it is steeper. Probably, other factors such as the dust-to-gas mass ratios play a role in steepening relations in Figs. 2b and 3b. The dust-to-gas mass ratios are smaller for lower-metallicity galaxies, which are also fainter and less massive. Possible deviations of warm dust emission from the thermal equilibrium may also play a role.

We produce dependencies of the M_{wd}/M_y ratios on $\text{EW}(\text{H}\beta)$ for ~ 1000 SDSS galaxies with best derived oxygen abundances and for the entire sample of ~ 4000 SDSS compact galaxies in Figs. 5a and b. The clear decrease of the mass ratio with decreasing $\text{EW}(\text{H}\beta)$ or respective increasing age is seen. It is more evident for the entire sample (Fig. 5b). This decrease is caused mainly by the decrease of M_{wd} , because M_y is almost not changed at $\text{EW}(\text{H}\beta) \geq 10 \text{ \AA}$, corresponding to instantaneous burst ages $< 10 \text{ Myr}$.

We fit the distributions for both samples by quadratic maximum-likelihood regressions. For comparison, we also show dependencies of M_{wd}/M_y ratios on $\text{EW}(\text{H}\beta)$, which are proportional to the luminosities of ionising radiation and to the total luminosities of the instantaneous burst (Schaerer & Vacca 1998; Leitherer et al. 1999). It is seen that these dependencies are not as steep, but the dependencies, which are proportional to the luminosities of ionising radiation, are in closer agreement with the likelihood regressions. The differences between the dependencies are likely due to that the M_{wd}/M_y ratios increase with the luminosity of ionising radiation more steeply than linearly. Thus, this result and the tight correlation between the luminosities L_{wd} and $L(\text{H}\beta)$ suggest that warm dust is associated with H II regions and is heated in significant part by ionising radiation of hot O-stars beyond the Ly-limit at $912 \text{ \AA}$.

4.2. Herschel sample

The original *Herschel* sample of 48 galaxies by Rémy-Ruyer et al. (2013) is more heterogeneous as compared to the SDSS galaxies considered in the previous subsection and includes extended irregular galaxies and BCDs. Large angular sizes of some galaxies in the *Herschel* sample make a comparison of data obtained at different wavelengths uncertain as large aperture corrections are needed. On the other hand, these are nearby objects, some of them are bright, and most of them were extensively observed in different wavelength ranges. Applying selection criteria discussed in Sect. 2.2, we reduced the sample to 28 compact galaxies. We collect the available data for the reduced *Herschel* sample from the UV-range to the radio-range using mainly databases of all-sky surveys. References to these observational data are given in Table 1.

4.2.1. SED fits

The SED fits of stellar and ionised gas emission at $\lambda \leq 15 \mu\text{m}$, of dust emission at $\lambda \sim 2\text{--}1000 \mu\text{m}$ and of free-free emission at $\lambda \geq 100 \mu\text{m}$ and the observed data are shown in Fig. 6. We note that modelled fluxes in this Figure are redshifted to observed

Fig. 2. Dependence of the warm dust luminosity L_{wd} of the sample of ~ 1000 SDSS compact galaxies with reliably derived oxygen abundance on **a)** total luminosity L_{tot} within $0.1\text{--}22 \mu\text{m}$ and **b)** $H\beta$ luminosity $L(H\beta)$. Red and blue filled circles are for galaxies with $EW(H\beta) \geq 50 \text{ \AA}$ and $EW(H\beta) < 50 \text{ \AA}$, respectively. **c)** and **d)** Same as in **a)** and **b)** but for the entire sample of ~ 4000 SDSS compact star-forming galaxies. In all panels, black solid lines are fits to the data where L_{wd} is calculated by adopting $\beta = 2$. Green solid lines in **b)** and **d)** are one-to-one relations between warm dust luminosities and luminosities of ionising radiation $L_{\text{ion}} \sim 100L(H\beta)$.

Fig. 3. Dependence of the warm dust mass M_{wd} of the sample of ~ 1000 SDSS compact galaxies with reliably derived oxygen abundance on **a)** total galaxy stellar mass M_* and **b)** mass of the young stellar population M_y . Red and blue filled circles are for galaxies with $EW(H\beta) \geq 50 \text{ \AA}$ and $EW(H\beta) < 50 \text{ \AA}$, respectively. In both panels, solid lines are fits to the data where M_{wd} is calculated adopting $\beta = 2$.

wavelengths and reddened adopting the extinction coefficient $C(H\beta)$, which is derived from optical spectra and the reddening law by Cardelli et al. (1989) with $R_V = 3.2$ (cf. Sect. 3.1). This is done to make a comparison with observed fluxes.

First, the optical spectra were used to derive line intensities and equivalent widths of emission lines, physical conditions, redshifts and extinction. We produce SEDs, which are the sum of stellar SED and ionised gas SED. These fits were

extrapolated to the UV and IR ranges and cover a wavelength range of $0.1\text{--}15 \mu\text{m}$. Observed fluxes in optical spectra prior to the fitting were corrected for aperture. The aperture correction for nine compact galaxies with SDSS spectra was done comparing the photometric r magnitude of the galaxy with r magnitude measured within the spectroscopic slit of $3''$ in diameter. Typical aperture corrections for compact *Herschel* galaxies are of order $\sim 1.5\text{--}2.0$. For other galaxies, we scale the observed spectra by

Fig. 4. Dependence of the warm dust temperature T_{wd} of the sample of ~ 1000 SDSS compact galaxies with a reliably derived oxygen abundance on **a)** $H\beta$ luminosity $L(H\beta)$ and **b)** mass of the young stellar population M_y . Red and blue filled circles are for galaxies with $EW(H\beta) \geq 50 \text{ \AA}$ and $EW(H\beta) < 50 \text{ \AA}$, respectively. In both panels, T_{wd} is calculated adopting $\beta = 2$.

Fig. 5. **a)** Dependence of the warm dust mass M_{wd} to the young stellar population mass M_y ratio on the $H\beta$ equivalent width $EW(H\beta)$ for the sample of ~ 1000 SDSS compact galaxies with a reliably derived oxygen abundance. **b)** Same as in **a)** but for the entire sample of ~ 4000 SDSS compact star-forming galaxies. In both panels, the warm dust mass M_{wd} is calculated by adopting $\beta = 2$. Solid lines are quadratic fits to the data, while dashed and dotted lines correspond to M_{wd}/M_y ratios, which are proportional to the luminosity L_{ion} of ionising radiation and total luminosity L_{tot} , respectively.

comparing $H\beta$ or $H\alpha$ line fluxes inside the slit with those derived from narrow-band imaging and from spectra, which were obtained with large apertures.

Second, the extinction- and aperture-corrected flux of the $H\beta$ emission line was used to calculate the SED of free-free emission in the submm and radio ranges (at $\lambda \geq 100 \mu\text{m}$ in Fig. 6), according to Caplan & Deharveng (1986).

Third, we find that the three-component model with cold, warm, and hot dust is generally required to fit the SED in the mid- and FIR ranges. For this, *Spitzer*/IRS spectra with varying wavelength ranges, but typically at $\lambda 5.5\text{--}35 \mu\text{m}$, and available *Herschel*, *Spitzer*, and WISE photometric data, are shown by line and different symbols in Fig. 6. The two-component model without hot dust is sufficient only in six galaxies (HS 0052+2536, HS 0822+3542, I Zw 18, Mrk 209, UGC 4483, and VII Zw 403). The SED fits of individual dust components are shown for hot, warm, and cold dust, respectively. The total SED of dust emission is also shown, which additionally includes the free-free emission component at $\lambda \geq 100 \mu\text{m}$. The wavelength at which dust emission and free-free emission are equal are also indicated. In all cases, a dust emissivity index β of 2.0 is adopted.

4.2.2. Comparison with Rémy-Ruyer et al. (2013) results

In Fig. 6, we use fits to derive temperatures, luminosities, and masses of cold, warm, and hot dust components in 28 compact galaxies from the *Herschel* sample by Rémy-Ruyer et al. (2013).

Derived temperatures, luminosities, and masses of cold and warm dust components are shown in Table 2, and they can be compared to the properties of the cold dust derived by Rémy-Ruyer et al. (2013). We do not show parameters for the hot dust component, because the equilibrium conditions for this component may not be valid. Additionally, the table shows oxygen abundances, extinction-corrected $H\beta$ luminosities, masses of the young stellar population M_y and total stellar masses M_* .

We note that Rémy-Ruyer et al. (2013) did not derive parameters of the cold dust in objects, which were not detected at $160 \mu\text{m}$ and longer wavelengths. This refers to seven galaxies, HS 0822+3542, HS 1222+3741, I Zw 18, SBS 1159+545, SBS 1249+493, Tol 1214–277, and UGC 4483, out of the 28 galaxies from our sample. We also note, that we had to correct the luminosity of cold dust in SBS 0335–052E, mistakenly given by (Rémy-Ruyer et al. 2013, their Table 4) upwards by a factor of ~ 100 .

Table 2. Global parameters of galaxies from the *Herschel* sample.

Object	Dust parameters						12+	log	log	log
	Temperature (K)		log Mass ($M_{\odot}$)		log Luminosity ($L_{\odot}$)		log O/H	$L(\text{H}\beta)^a$	M_{y}	M_{*}
	cold	warm	cold	warm	cold	warm		($L_{\odot}$)	($M_{\odot}$)	($M_{\odot}$)
Haro 11	34 $^{+1}_{-2}$	104 $^{+4}_{-9}$	6.54 $^{+0.09}_{-0.03}$	3.73 $^{+0.28}_{-0.12}$	10.74 $^{+0.06}_{-0.12}$	10.82 $^{+0.10}_{-0.23}$	8.36	9.37	9.45	10.51
Haro 3	26 $^{+1}_{-1}$	93 $^{+6}_{-6}$	5.89 $^{+0.03}_{-0.08}$	2.32 $^{+0.08}_{-0.13}$	9.39 $^{+0.11}_{-0.04}$	9.13 $^{+0.16}_{-0.16}$	8.28	6.74	5.97	7.76
HS 0017+1055	38 $^{+12}_{-7}$	121 $^{+11}_{-4}$	4.11 $^{+0.42}_{-0.49}$	1.07 $^{+0.12}_{-0.29}$	8.55 $^{+0.70}_{-0.55}$	8.56 $^{+0.22}_{-0.09}$	7.63	6.96	6.96	7.22
HS 0052+2536	36 $^{+1}_{-9}$	90 $^{+9}_{-1}$	5.73 $^{+0.57}_{-0.04}$	2.85 $^{+0.02}_{-0.30}$	10.06 $^{+0.07}_{-0.71}$	9.58 $^{+0.25}_{-0.01}$	8.04	7.90	6.90	8.83
HS 0822+3542	33 $^{+5}_{-2}$	107 $^{+12}_{-1}$	2.83 $^{+0.18}_{-0.47}$	-0.92 $^{+0.03}_{-0.29}$	6.94 $^{+0.04}_{-0.08}$	6.24 $^{+0.03}_{-0.02}$	7.45	5.28	4.60	5.97
HS 1222+3741	54 $^{+7}_{-12}$	111 $^{+11}_{-7}$	3.87 $^{+0.55}_{-0.43}$	1.70 $^{+0.25}_{-0.29}$	9.25 $^{+0.09}_{-0.12}$	8.96 $^{+0.05}_{-0.08}$	7.79	7.25	6.87	8.13
HS 1304+3529	32 $^{+2}_{-3}$	72 $^{+6}_{-6}$	4.96 $^{+0.20}_{-0.18}$	2.53 $^{+0.30}_{-0.29}$	9.02 $^{+0.05}_{-0.07}$	8.68 $^{+0.06}_{-0.07}$	7.93	7.24	6.76	8.19
HS 1330+3651	27 $^{+3}_{-1}$	82 $^{+8}_{-5}$	5.29 $^{+0.10}_{-0.18}$	2.01 $^{+0.16}_{-0.28}$	8.89 $^{+0.05}_{-0.03}$	8.48 $^{+0.01}_{-0.05}$	7.98	6.81	7.42	9.34
I Zw 18	45 $^{+2}_{-1}$	118 $^{+3}_{-3}$	2.38 $^{+0.09}_{-0.11}$	-0.67 $^{+0.09}_{-0.07}$	7.31 $^{+0.02}_{-0.02}$	6.75 $^{+0.02}_{-0.02}$	7.17	6.13	5.94	6.42
II Zw 40	28 $^{+2}_{-1}$	101 $^{+2}_{-6}$	5.25 $^{+0.05}_{-0.11}$	1.87 $^{+0.19}_{-0.03}$	8.95 $^{+0.18}_{-0.05}$	8.88 $^{+0.06}_{-0.17}$	8.23	7.37	7.93	8.00
Mrk 1089	25 $^{+3}_{-2}$	77 $^{+6}_{-5}$	6.84 $^{+0.17}_{-0.25}$	3.56 $^{+0.21}_{-0.29}$	10.23 $^{+0.06}_{-0.03}$	9.88 $^{+0.03}_{-0.10}$	8.10	7.93	8.02	8.89
Mrk 1450	32 $^{+3}_{-1}$	98 $^{+5}_{-2}$	4.01 $^{+0.04}_{-0.14}$	1.00 $^{+0.08}_{-0.20}$	8.07 $^{+0.12}_{-0.03}$	7.95 $^{+0.03}_{-0.07}$	7.84	6.48	5.77	7.55
Mrk 153	33 $^{+2}_{-2}$	88 $^{+2}_{-1}$	4.70 $^{+0.12}_{-0.11}$	1.61 $^{+0.02}_{-0.04}$	8.77 $^{+0.14}_{-0.16}$	8.29 $^{+0.04}_{-0.01}$	7.86	6.95	7.84	8.60
Mrk 209	32 $^{+3}_{-2}$	96 $^{+9}_{-1}$	3.11 $^{+0.13}_{-0.15}$	0.13 $^{+0.13}_{-0.30}$	7.14 $^{+0.08}_{-0.05}$	7.01 $^{+0.01}_{-0.07}$	7.74	5.83	5.56	5.83
Mrk 930	29 $^{+3}_{-2}$	89 $^{+8}_{-5}$	6.24 $^{+0.13}_{-0.22}$	2.99 $^{+0.19}_{-0.23}$	9.98 $^{+0.04}_{-0.06}$	9.66 $^{+0.03}_{-0.02}$	8.03	7.93	7.78	9.46
NGC 1140	25 $^{+1}_{-1}$	80 $^{+3}_{-1}$	5.97 $^{+0.04}_{-0.06}$	2.38 $^{+0.02}_{-0.12}$	9.33 $^{+0.02}_{-0.01}$	8.79 $^{+0.01}_{-0.04}$	8.38	7.12	6.98	8.72
Pox 186	25 $^{+2}_{-6}$	80 $^{+9}_{-6}$	3.85 $^{+0.59}_{-0.13}$	0.89 $^{+0.25}_{-0.28}$	7.24 $^{+0.06}_{-0.11}$	7.29 $^{+0.01}_{-0.01}$	7.70	5.65	4.98	6.33
SBS 0335–052E	57 $^{+7}_{-12}$	129 $^{+3}_{-8}$	3.06 $^{+0.25}_{-0.16}$	1.37 $^{+0.19}_{-0.06}$	8.61 $^{+0.14}_{-0.34}$	9.03 $^{+0.02}_{-0.01}$	7.30	7.19	6.78	7.26
SBS 1159+545	33 $^{+15}_{-8}$	88 $^{+9}_{-1}$	3.78 $^{+0.62}_{-0.70}$	1.34 $^{+0.04}_{-0.29}$	7.85 $^{+0.31}_{-0.16}$	7.99 $^{+0.03}_{-0.01}$	7.44	6.35	5.76	5.77
SBS 1211+540	42 $^{+15}_{-7}$	99 $^{+7}_{-9}$	2.46 $^{+0.35}_{-0.45}$	-0.48 $^{+0.30}_{-0.28}$	7.22 $^{+0.31}_{-0.13}$	6.50 $^{+0.04}_{-0.10}$	7.58	5.79	5.67	7.06
SBS 1249+493	28 $^{+10}_{-6}$	87 $^{+12}_{-6}$	5.10 $^{+0.62}_{-0.70}$	1.97 $^{+0.15}_{-0.30}$	8.79 $^{+0.09}_{-0.04}$	8.60 $^{+0.04}_{-0.03}$	7.68	6.90	6.53	7.41
SBS 1415+437	37 $^{+5}_{-5}$	89 $^{+6}_{-6}$	3.27 $^{+0.25}_{-0.22}$	0.59 $^{+0.24}_{-0.26}$	7.65 $^{+0.09}_{-0.12}$	7.27 $^{+0.07}_{-0.09}$	7.55	6.11	5.35	6.06
SBS 1533+574	26 $^{+2}_{-1}$	91 $^{+9}_{-9}$	5.34 $^{+0.04}_{-0.16}$	2.11 $^{+0.30}_{-0.29}$	8.83 $^{+0.04}_{-0.02}$	8.87 $^{+0.04}_{-0.04}$	8.05	6.70	6.30	8.08
Tol 1214–277	37 $^{+17}_{-9}$	87 $^{+11}_{-9}$	4.18 $^{+0.62}_{-0.65}$	1.93 $^{+0.28}_{-0.30}$	8.56 $^{+0.32}_{-0.19}$	8.57 $^{+0.01}_{-0.01}$	7.52	7.54	6.94	6.99
UGC 4483	30 $^{+11}_{-5}$	89 $^{+8}_{-3}$	2.31 $^{+0.58}_{-0.70}$	-1.17 $^{+0.10}_{-0.30}$	6.21 $^{+0.09}_{-0.09}$	5.52 $^{+0.01}_{-0.07}$	7.46	4.67	4.46	5.16
UM 448	29 $^{+3}_{-1}$	87 $^{+8}_{-1}$	6.84 $^{+0.10}_{-0.20}$	3.74 $^{+0.02}_{-0.24}$	10.65 $^{+0.07}_{-0.03}$	10.37 $^{+0.02}_{-0.02}$	8.32	8.23	8.59	10.45
UM 461	31 $^{+3}_{-1}$	88 $^{+1}_{-5}$	3.57 $^{+0.10}_{-0.16}$	0.66 $^{+0.16}_{-0.02}$	7.52 $^{+0.06}_{-0.01}$	7.33 $^{+0.02}_{-0.01}$	7.73	5.83	5.45	6.94
VII Zw 403	29 $^{+2}_{-1}$	78 $^{+7}_{-1}$	3.33 $^{+0.07}_{-0.12}$	-0.20 $^{+0.01}_{-0.25}$	7.12 $^{+0.04}_{-0.04}$	6.15 $^{+0.03}_{-0.01}$	7.66	5.22	4.83	5.98

Notes. ^(a) Extinction-corrected.

In Fig. 7a, we show a comparison between the cold dust temperatures in 21 galaxies, for which data are available in Rémy-Ruyer et al. (2013) with those in this paper. For most of the galaxies the agreement is good. However, there are some outliers. These are objects, for which Rémy-Ruyer et al. (2013) adopted $\beta \sim 0.0$. The most deviant object is HS 0017+1055. We derive $T_{\text{cd}} = 38^{+12}_{-7}$ K, using a three-component dust model. Our fit reproduces all available WISE, *Spitzer*/IRS and *Herschel* observations of this galaxy well, and the derived T_{cd} is similar to that obtained for other galaxies (Table 2). Rémy-Ruyer et al. (2013) found $T_{\text{cd}} = 98^{+34}_{-36}$ K, using a one-component dust model with $\beta = 0.0^{+1.34}_{-0.00}$.

On the other hand, the luminosities of the cold dust are in good agreement (Fig. 7b). This is expected, because fits in both cases reproduce the observed FIR fluxes. Derived masses of cold dust differ to a larger extent (Fig. 7c), despite the overall agreement between Rémy-Ruyer et al. (2013) and our determinations.

4.2.3. Dust temperatures

In Fig. 8, relations among the oxygen abundances $12 + \log \text{O/H}$ and the temperatures of cold, and warm dust for galaxies from the *Herschel* sample are shown. Temperatures vary in the range

of $25 \text{ K} < T_{\text{cd}} < 57 \text{ K}$ with the average value of $\sim 30 \text{ K}$ for the cold component and in the range of $72 \text{ K} < T_{\text{wd}} < 129 \text{ K}$ with the average value of $\sim 90 \text{ K}$ for the warm component. No obvious trends with metallicity are found, excluding the two most deviant and most metal-deficient BCDs, I Zw 18 and SBS 0335–052E. This result confirms that found for the temperatures of warm dust in SDSS sample galaxies (Fig. 1). We note, however, that the average temperatures for the SDSS sample are higher, $\sim 120 \text{ K}$ for the warm component. The cause for this discrepancy remains unclear. However, the temperatures derived for the warm component in the *Herschel* sample are more robust, because SED fitting in these objects is controlled by the photometric data from WISE, *Spitzer*, and *Herschel* and by *Spitzer*/IRS spectra, while only WISE photometric data are available for the SDSS sample. We also cannot exclude imperfect correction of WISE fluxes for the contribution of the nebular emission lines, which was done with the use of Cloudy photoionised H II region models.

4.2.4. Dust luminosities

Dependencies of cold and warm dust luminosities on the $\text{H}\beta$ luminosity $L(\text{H}\beta)$ are shown in Fig. 9. The relation in Fig. 9a is almost as tight as that in Fig. 9b for only compact galaxies with *Spitzer*/IRS spectra. This suggests that likely the same

Fig. 7. Relation between the **a)** temperatures, **b)** luminosities, and **c)** masses of cold dust obtained for the *Herschel* sample in this paper and by Rémy-Ruyer et al. (2013). In all panels, solid lines mark a one to one correspondence.

sources heat the warm and cold dust in compact galaxies from the *Herschel* sample.

4.2.5. Dust masses

In Fig. 10, dust mass to total stellar mass ratios on oxygen abundance $12 + \log \text{O}/\text{H}$ for cold and warm dust components are shown. No clear dependence is present. The two most-metal deficient galaxies, I Zw 18 and SBS 0335–052E are among the galaxies of the *Herschel* sample with the lowest cold dust mass (Table 2). In Fig. 10a, they have low M_{cd}/M_* mass ratios of $\sim 10^{-4}$, which are lower by a factor of ~ 10 than the average value for the sample. This, however, does not deviate more than by 1σ from other sample galaxies.

Fig. 8. Dependencies of **a)** cold and **b)** warm dust temperatures for the *Herschel* sample on oxygen abundances $12 + \log \text{O}/\text{H}$. The galaxies I Zw 18 and SBS 0335–052E are encircled.

Deviations of I Zw 18 and SBS 0335–052E are much larger in the M_{cd}/M_y vs. $12 + \log \text{O}/\text{H}$ diagram (Fig. 11a). This is presumably because of the low cold dust mass and high fraction of $\sim 30\%$ of the young stellar population in the total mass, indicating that both galaxies are relatively unevolved and had no time to produce dust in large quantities. Although the old stellar population with age ≥ 1 Gyr is present in I Zw 18, its mass of $\geq 2 \times 10^6 M_\odot$ (Annibali et al. 2013) is small compared to the H I mass of $10^8 M_\odot$ (Lelli et al. 2012). There is no direct evidence for an old stellar population in SBS 0335–052E. Pustilnik et al. (2004) inferred from the integrated light of SBS 0335–052E that the stellar-to-total baryon mass fraction in this galaxy is 0.035.

Therefore, M_{cd}/M_y in these galaxies is only by ~ 3 times higher than M_{cd}/M_* (Table 2). On the other hand, the young-to-total stellar mass ratio is much smaller in many other galaxies from the *Herschel* sample, elevating their dust-to-young stellar mass ratio, as compared to that in Fig. 10a. The difference between I Zw 18 and SBS 0335–052E and other galaxies is lower for the warm dust component.

In Fig. 12, the relation is shown between the warm and cold dust masses. The presence of a correlation between cold and warm dust masses for the *Herschel* sample suggests that these components are linked to young stellar populations. This correlation with dispersion of ~ 0.5 dex opens the opportunity for estimating the cold dust mass from the warm dust mass. The most outlying galaxy in Fig. 12 is SBS 0335–052E with highest $M_{\text{wd}}/M_{\text{cd}}$ ratio. Given the warm dust mass and using the relation in Fig. 12 would overestimate the cold dust mass by one order of magnitude for this galaxy. On the other hand, the relation in the Figure nicely reproduces the cold dust mass in I Zw 18.

Fig. 9. Dependencies of **a)** cold and **b)** warm dust luminosities on $H\beta$ luminosities for galaxies from the *Herschel* sample. Solid lines are linear maximum-likelihood relations, while dotted lines are one-to-one relations between dust luminosities and total luminosities of ionising radiation $L_{\text{ion}} \sim 100L(H\beta)$. I Zw 18 and SBS 0335–052E are encircled.

Only a small fraction of the local star-forming galaxies with low metallicity was observed with *Herschel* (Rémy-Ruyer et al. 2013). On the other hand, there are many more compact star-forming galaxies that were detected with WISE (this paper, Izotov et al. 2011b, 2014). The relation in Fig. 12 allows us to estimate the cold dust mass in these galaxies with an uncertainty better than by a factor of three.

4.2.6. Dust-to-gas mass ratio

The dust-to-gas mass ratio is often considered as a measure of metallicity. In this respect it might be compared to the oxygen abundance $12 + \log O/H$ measured in the ionised gas. One would expect that the dust-to-gas mass ratio and heavy element mass fraction Z_{ion} in the ionised gas should be comparable if the galaxy ISM is well mixed.

However, there is some evidence that the neutral gas metallicity is much lower than that of the ionised gas. Thuan et al. (2002, 2005), Lebouteiller et al. (2004, 2013), Lecavelier des Etangs et al. (2004) using *FUSE* and the Cosmic Origin Spectrograph (COS) onboard HST showed that the oxygen-to-hydrogen abundance ratio in the neutral gas of low-metallicity BCDs is on average several times lower than that in H II regions. It was even suggested in some papers that some fraction of pristine neutral gas might be present in these galaxies.

Based on IRAS observations, Lisenfeld & Ferrara (1998) found that the dust-to-gas mass ratio in dwarf irregulars and BCDs is steeply decreased with decreasing ionised gas

Fig. 10. Dependencies of the dust mass-to-total stellar mass ratio on oxygen abundance are shown for **a)** cold and **b)** warm dust in galaxies from the *Herschel* sample. The black solid lines in **a)** and **b)** indicate average values of M_{cd}/M_* and M_{wd}/M_* , respectively. I Zw 18 and SBS 0335–052E are encircled.

metallicity. Later, Engelbracht et al. (2008) using *Spitzer* observations confirmed this result. A detailed study of the dust-to-gas mass ratio in the galaxies from the *Herschel* sample was done by Rémy-Ruyer et al. (2014). Based on the galaxies from Rémy-Ruyer et al. (2013), they compiled H I data and calculated the mass of the molecular hydrogen by using the CO-to- H_2 conversion factor for galaxies with detected CO emission and by extrapolating the conversion factor for galaxies with low metallicity, where CO emission was not detected. Rémy-Ruyer et al. (2014) found that the dust-to-gas mass ratio is steeply decreased with decreasing metallicity, a result similar to that obtained by Lisenfeld & Ferrara (1998) and Engelbracht et al. (2008).

In Fig. 13a, we show the dust-to-gas mass – oxygen abundance diagram for the *Herschel* sample of compact galaxies together with data from the literature, which include both dwarf and giant galaxies. The mass of molecular hydrogen M_{H_2} for three galaxies from the *Herschel* sample with detected CO emission (Haro 11, Mrk 1089, Mrk 930) was taken from Cormier et al. (2014) and added to the H I mass. The M_{H_2}/M_{HI} ratios for Mrk 1089 and Mrk 930 are low, of ~ 0.2 – 0.3 . However, this ratio of ~ 5 is much higher for Haro 11. For galaxies selected from the literature, we show the data in Fig. 13a, which include the H_2 mass, whenever it was available. We do not use the extrapolation of the CO-to- H_2 conversion factor to low metallicities to derive the H_2 mass in galaxies from the *Herschel* sample with undetected CO emission because this procedure is highly uncertain.

We confirm a sharp decrease of dust-to-gas mass ratios with decreasing metallicity at $12 + \log O/H < 8.5$, while the ratio is

Fig. 11. Dependencies of the dust mass-to-young stellar mass ratio on oxygen abundance are shown for **a)** cold and **b)** warm dust in galaxies from the *Herschel* sample. The black solid lines in **a)** and **b)** indicate average values of M_{cd}/M_y and M_{wd}/M_y , respectively. I Zw 18 and SBS 0335–052E are encircled.

Fig. 12. Mass of warm dust vs. mass of cold dust. The red solid line represents the maximum-likelihood linear regression. I Zw 18 and SBS 0335–052E are encircled.

almost constant in giant galaxies with higher metallicities. The *Herschel* data can be fit by a linear maximum-likelihood relation shown as a black solid line. We note the steep slope of ~ 2.65 , which is steeper than the linear extrapolation of the data for higher-metallicity galaxies by [Draine et al. \(2007\)](#) and the model predictions by [Hirashita et al. \(2002\)](#). We note that the dust-to-gas mass ratios in Fig. 13a are global characteristics of galaxies because they are derived from the total dust and gas masses.

The dotted line in Fig. 13a is the one-to-one line between the dust-to-gas mass ratio and the heavy-element mass fraction Z_{ion} in the ionised gas. Z_{ion} is derived from the relation

$$Z_{ion} = 16 \times \left(\frac{O}{H} \right) \left(\frac{Z}{Z_0} \right), \quad (5)$$

where Z/Z_0 is the ratio by mass of all heavy elements to oxygen, which we adopt to be 2.5. O/H is the oxygen abundance by number derived from the H II region's optical spectrum.

This relation shows that the observed dust-to-gas mass ratio in dwarf galaxies is by a factor of ~ 100 lower than the heavy element mass fraction in the dwarf's H II regions. Keeping in mind that the metallicity of the neutral gas in these galaxies can be lower by a factor of up to ~ 10 than the heavy element mass fraction in H II regions (e.g. [Thuan et al. 2005](#)), we conclude that the global dust-to-gas mass ratio in dwarf galaxies is ~ 10 times lower than the neutral gas metallicity. From the two relations (Fig. 13a), we can derive the dust-to-metal mass ratio, which declines with decreasing metallicity. This is in line with the trends in dust-to-metal ratios (i.e. the mass fraction of metals in dust) found by [De Cia et al. \(2013\)](#) for low-metallicity gamma-ray burst (GRB) hosts and damped Ly- α absorbers (DLAs).

The lowest global dust-to-gas mass ratios are derived for the two lowest-metallicity BCDs, I Zw 18 and SBS 0335–052E, shown in Fig. 13a. They are in nice agreement with dust-to-gas mass ratios obtained for these galaxies by [Rémy-Ruyer et al. \(2014\)](#) and for some other most-metal poor galaxies with $12 + \log O/H \lesssim 7.6$ (this paper, [Lisenfeld & Ferrara 1998](#); [Engelbracht et al. 2008](#)). [Hunt et al. \(2014\)](#) derived a similar dust-to-gas mass ratio for I Zw 18, but at a value ~ 5000 times higher for SBS 0335–052E. The difference in the case of SBS 0335–052E is in part due to that [Hunt et al. \(2014\)](#) derived a dust mass, which is a factor of ~ 40 higher than the value in this paper (Table 2), the latter being consistent with [Rémy-Ruyer et al. \(2013\)](#). Furthermore, [Hunt et al. \(2014\)](#) assumed that the dust is located inside an $\sim 1''$ aperture, which is much smaller than the larger aperture, of $\sim 10''$ in radius, where *Herschel* fluxes were measured. We adopt a large aperture of $\sim 10''$. [Hunt et al. \(2014\)](#) also used the H I mass inside $\sim 1''$ aperture, while we use the total H I mass of $4.2 \times 10^8 M_\odot$ for SBS 0335–052E ([Ekta et al. 2009](#)). Furthermore, hydrogen in the central part of the galaxy within a $1''$ aperture is mostly ionised, and its mass that adopts the electron number density of $\sim 100 \text{ cm}^{-3}$ is by at least a factor of ~ 10 larger than the mass of neutral gas within the same aperture. In these circumstances, ionised gas should be taken into account in the dust-to-gas mass ratio determination. On the other hand, the global dust-to-gas mass ratio is almost insensitive to the presence of ionised gas, because its total mass is much lower than the neutral gas mass.

In this respect, our dust-to-gas mass ratio for SBS 0335–052E can be considered as a global characteristic of the galaxy, while that of [Hunt et al. \(2014\)](#) describes a local characteristic for the central part of the galaxy.

In Fig. 13b, we show the dust-to-ionised-gas mass ratio for our galaxies. The ionised gas mass is derived using relation

$$M(\text{H II}) = \frac{L(\text{H}\beta)m_p}{\epsilon N_e}, \quad (6)$$

where m_p is the proton mass, N_e is the electron number density, $L(\text{H}\beta)$ is the extinction-corrected total H β luminosity, and ϵ is the H β emissivity, as defined by equation ([Aller 1984](#))

$$\epsilon = 1.37 \times 10^{-25} t_e^{-0.982} \exp(-0.104/t_e), \quad (7)$$

$$t_e = 10^{-4} T_e.$$

Fig. 13. a) Dependence of the dust-to-hydrogen mass ratio relative to the oxygen abundance $12 + \log O/H$. The data from this paper are shown by large filled blue circles and are fitted by the linear maximum-likelihood regression (black solid line). The two lowest-metallicity galaxies, I Zw 18 and SBS 0335–052E, are encircled. The dust-to-gas ratio from a compiled sample of dwarf irregulars and blue compact dwarf (BCD) galaxies by Lisenfeld & Ferrara (1998; IRAS data) are denoted by black crosses (note that for GR8 we adopt the oxygen abundance from Morales-Luis et al. 2011). The data for 16 BCDs (IRAS data) by Hirashita et al. (2002) are shown by magenta open squares. The dust-to-gas ratio for 66 star-forming galaxies observed with *Spitzer* by Engelbracht et al. (2008) are shown by light blue triangles. The data compiled from literature for a wide sample of galaxies using different submm ($>160 \mu\text{m}$) observations (Galametz et al. 2011) are denoted by large open black circles. The available data gathered by James et al. (2002) and their observations with SCUBA are shown by red asterisks. b) Dependence of the dust-to-ionised-hydrogen mass ratio relative to the oxygen abundance $12 + \log O/H$. The data from this paper are shown by large filled blue circles and are fitted by the linear maximum-likelihood regression (black solid line). The two lowest-metallicity galaxies, I Zw 18 and SBS 0335–052E, are encircled. The black dotted lines in both panels denote a one-to-one correspondence between the dust-to-gas mass ratio and the heavy-element mass fraction Z_{ion} (see Eq. (5)) in the ionised gas as derived from the oxygen abundance.

It is seen in Fig. 13b that the dust-to-ionised-gas mass ratio is sharply increased with metallicity and generally is higher than the dotted line denoting a ratio of unity between $M_{\text{dust}}/M_{\text{H II}}$ and Z_{ion} . Only in the lowest-metallicity galaxies are they comparable. This suggests that although radiation by a young stellar population is the main source of dust heating, most of dust is located in the neutral gas, and the fraction of dust in the neutral gas is increased with increasing metallicity.

4.2.7. submm excess in star-forming dwarf galaxies

Rémy-Ruyer et al. (2013) extensively discussed the submm excess at wavelengths $\geq 500 \mu\text{m}$, which is detected in a significant fraction of galaxies from the *Herschel* sample. They noted that this excess is more frequent in low-metallicity galaxies. In particular, they discuss two galaxies with this excess, Haro 11 and II Zw 40. Rémy-Ruyer et al. (2013) also detected the submm excess in the galaxies HS 0052+2536 and VII Zw 403 from our *Herschel* sample. Furthermore, Rémy-Ruyer et al. (2013) analyse different mechanisms, which may cause this emission. All of them are related to dust emission.

On the other hand, the presence of cold dust in low-metallicity compact dwarf galaxies with active star formation and intense UV stellar radiation is difficult to understand. We suggest that the free-free emission may play a role in producing submm excess in these galaxies with strong nebular continuum in the optical range and at longer wavelengths, as is indicated by a high equivalent width of the $H\beta$ emission line. It was noted in Sect. 4.2.1 that the fluxes of the free-free emission in submm and radio ranges were calculated from the $H\beta$ emission line flux, according to Caplan & Deharveng (1986). We compare predicted and observed fluxes in the cm range where emission is dominated by free-free processes. For most objects from our *Herschel* sample, no data are available in this wavelength

range. For the remaining galaxies, Haro 11 (Fig. 6a1), II Zw 40 (Fig. 6b1), I Zw 18 (Fig. 6b2), Mrk 1089 (Fig. 6b3), Mrk 209 (Fig. 6b4), Mrk 930 (Fig. 6b7), and SBS 0335–052E (Fig. 6c2), the agreement is good.

Depending on the galaxy properties, the contribution of free-free emission may be significant in the submm or mm wavelength ranges (Fig. 6). We note that the role of free-free emission in producing submm excess was discussed earlier by Galliano et al. (2003) and Galliano et al. (2005) for four high-metallicity star-forming galaxies, NGC 1569, II Zw 40, He 2-10, and NGC 1140. They concluded that free-free emission is insufficient to explain this excess and that emission of very cold dust is needed. However, we point out that Rémy-Ruyer et al. (2013) did not detect the excess in NGC 1140, while it is present with the Galliano et al. (2005) data. We also conclude from our SED fitting (Fig. 6b8) that a submm excess in this galaxy is not apparent.

We note that submm and mm observations for II Zw 40 are controversial (Fig. 6b1). Submillimetre array (SMA) observations at $880 \mu\text{m}$ (Hirashita 2011) and MRT observations at 1.25 mm (Albrecht et al. 2004) are in very nice agreement with our fit, while $850 \mu\text{m}$ SCUBA and 1.3 mm MAMBO observations by Galliano et al. (2005) and Hunt et al. (2005a) are not. It is possible that differences in observed fluxes are due to the fact that II Zw 40 is located at low Galactic latitude with a high Milky Way extinction. This makes subtraction of bright foreground Galactic cirrus somewhat uncertain. We also note that $870 \mu\text{m}$ LABOCA observations by Galametz et al. (2009) are above the fit for Haro 11 (Figs. 6a1). On the other hand, observed and predicted fluxes at $\lambda \geq 850 \mu\text{m}$ for another two galaxies, NGC 1140 (6b8) and SBS 0335–052E (6c2) are in very good agreement and do not show evidence for a very cold dust.

In Fig. 6, blue vertical dashed lines indicate the wavelengths λ_{ff} at which contributions of dust and free-free emission

Fig. 14. Dependence of the wavelength λ_{ff} in submm and mm ranges on the oxygen abundance $12 + \log \text{O}/\text{H}$, at which monochromatic dust emission is equal to monochromatic free-free emission. Red filled circles and blue filled circles are data for galaxies from the *Herschel* sample with $L(\text{H}\beta) \geq 10^7 L_{\odot}$ and $L(\text{H}\beta) < 10^7 L_{\odot}$, respectively. The black solid line is the maximum-likelihood regression for the entire *Herschel* sample. I Zw 18 and SBS 0335–052E are encircled, while Haro 11 and II Zw 40 are surrounded by squares.

are equal. The most extreme cases are the most-metal deficient galaxies from the sample, I Zw 18 and SBS 0335–052E, with $\lambda_{\text{ff}} \sim 370 \mu\text{m}$ and $\sim 340 \mu\text{m}$, respectively (Figs. 6b2 and c2). In particular, ALMA observations of SBS 0335–052E clearly indicate that emission at $870 \mu\text{m}$ is almost totally dominated by free-free emission (Fig. 6c2, Hunt et al. 2014). There is also a general tendency of lowering λ_{ff} with decreasing oxygen abundance, $12 + \log \text{O}/\text{H}$. The wavelength λ_{ff} for galaxies with $12 + \log \text{O}/\text{H} \lesssim 8.0$ is commonly in the submm range, while it is $\geq 1 \text{ mm}$ for higher metallicities (Fig. 6). This agrees with statistics of the submm excess discussed by Rémy-Ruyer et al. (2013).

Regarding Haro 11 and II Zw 40, λ_{ff} are $\sim 700 \mu\text{m}$ and $\sim 960 \mu\text{m}$, respectively (Figs. 6-a1 and 6-b1). Therefore, a submm excess in Haro 11 due to the free-free emission is expected at shorter wavelengths, and it is seen by *Herschel* at $500 \mu\text{m}$, while it is not in II Zw 40.

The dependence of λ_{ff} on oxygen abundance for the entire *Herschel* sample is shown in Fig. 14. A clear increase of λ_{ff} with increasing $12 + \log \text{O}/\text{H}$ is seen, implying that the submm excess in low-metallicity galaxies may be due to the free-free emission from relatively transparent H II regions, which are seen in the optical range. However, metallicity is not the only parameter determining λ_{ff} , which is lower for galaxies with higher luminosity $L(\text{H}\beta)$ of the H β emission line (compare galaxies shown in Fig. 14). In particular, λ_{ff} for the two most deviant high-metallicity galaxies Haro 11 and II Zw 40 with high $L(\text{H}\beta)$ is also in the submm range.

Unfortunately, there are no observational data in the submm range for most of the galaxies shown in Fig. 6. Therefore, we cannot check the importance of free-free emission in producing submm excess in these low-metallicity galaxies. At least, we can conclude that the wavelength, at which the contribution of free-free emission becomes important is progressively decreased with decreasing metallicity, which is consistent with measured statistics of the submm excess (Rémy-Ruyer et al. 2013). Hopefully, forthcoming observations of these galaxies with ALMA will further clarify this issue.

4.3. The components of the SBS 0335–052E SED in the infrared

Thuan et al. (1999a) were the first who noted the existence of highly obscured regions in SBS 0335–052E based on ISO observations. More recently, Plante & Sauvage (2002) based on the same observations concluded that extinction in the obscured region is of order $A_V \sim 30 \text{ mag}$ and that most of the star formation in SBS 0335–052E is hidden and is not seen even in the NIR range. Houck et al. (2004) discovered silicate absorption at $9.7 \mu\text{m}$ in the *Spitzer* spectrum of SBS 0335–052E, which is indicative of the presence of highly obscured regions. Finally, Hunt et al. (2014) modelled the IR SED using the DUSTY code with inclusion of obscured regions.

On the other hand, our fit of free-free emission from the observed H β flux in SBS 0335–052E is in excellent agreement with the observed fluxes in the submm and radio ranges (Fig. 6c2). There are many spectroscopic observations of SBS 0335–052E with H β flux measurements of $\sim (4-10) \times 10^{-14} \text{ erg s}^{-1} \text{ cm}^{-2}$. For the fit of free-free emission in Fig. 6-c2, we adopted the value of $1.05 \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^{-2}$ by Izotov et al. (2006b) for the entire galaxy. However, if, in SBS 0335–052E, most of the star formation is hidden by a foreground absorbing layer as suggested by Plante & Sauvage (2002) then one would expect higher observed radio fluxes, as compared to those predicted from the observed H β flux.

Furthermore, Izotov & Thuan (2011) predicted fluxes of nebular [S IV] $\lambda 10.51 \mu\text{m}$ and [Ne III] $\lambda 15.55 \mu\text{m}$ emission lines from optical spectroscopic observations and Cloudy photoionised H II region models, which are in good agreement with fluxes observed by Houck et al. (2004) in the *Spitzer*/IRS spectrum of SBS 0335–052E. Finally, ALMA continuum observations at $870 \mu\text{m}$ (Hunt et al. 2014) and VLA continuum observations in the cm range (Johnson et al. 2009) do not reveal new spatially distinct star-forming regions in addition to those seen at shorter wavelengths.

Therefore, the origin of silicate absorption at $9.7 \mu\text{m}$ is puzzling as the observations imply that the contribution of obscured star-forming regions to FIR, submm, and thermal radio emission should be lower than that of those star-forming regions, which are visible in the optical range.

According to Starburst99 models for instantaneous bursts (Leitherer et al. 1999), the H β luminosity of visible star-forming regions in SBS 0335–052E, $L(\text{H}\beta) = 1.6 \times 10^7 L_{\odot}$ corresponds to a luminosity of ionising radiation beyond the Lyman limit, $L_{\text{ion}} \sim 100 L(\text{H}\beta) = 1.6 \times 10^9 L_{\odot}$, and a total UV luminosity L_{UV} of $3.9 \times 10^9 L_{\odot}$. The V-band extinction $A(V)$ of 0.3 mag for SBS 0335–052E would correspond to $A(0.1 \mu\text{m}) \sim 3 \times A(V) \sim 0.9 \text{ mag}$ meaning that more than half of the UV radiation is absorbed by dust. This seems to be sufficient to explain our derived L_{FIR} of $1.7 \times 10^9 L_{\odot}$, which includes cold and warm dust components (Table 2).

Figure 15 shows the full spectrum of SBS 0335–052E from the Lyman limit to centimeter radio wavelengths. Between $3 \mu\text{m} < \lambda < 0.5 \text{ mm}$, the radiation comes from dust. At shorter wavelengths, it is stellar and nebular emission including many lines; at longer wavelengths, it is free-free radiation. Because the radio continuum becomes opaque ($\tau_c = 1$) at $\lambda > 3 \text{ cm}$, this implies an emission measure $EM \sim 10^9 \text{ pc cm}^{-6}$ or an electron density of 10^4 cm^{-3} over a length of 1 pc.

More than half of the total luminosity of the galaxy, $L_{\text{gal}} = 3.9 \times 10^9 L_{\odot}$, is emitted at optical wavelengths ($L_{\text{opt}} = 2.2 \times 10^9 L_{\odot}$ for $\lambda < 2 \mu\text{m}$) and has not been attenuated by dust. The remaining luminosity, $L_{\text{FIR}} = 1.7 \times 10^9 L_{\odot}$, has been processed

by dust. If the galaxy is powered by O6 stars with having a luminosity $L_* = 2.5 \times 10^5 L_\odot$, it will take $N_* = 1.56 \times 10^4$ of them to account for L_{gal} .

As one O6 star emits $N_L \sim 1.2 \times 10^{49}$ Lyman continuum photons per s, the output of all O6 stars amounts to $N_* N_L = 1.9 \times 10^{53}$ photons s^{-1} . If the fraction $L_{\text{opt}}/L_{\text{gal}} \sim 22/39$ leads to ionisation, one comes up with a number of ionising photons $N_{\text{ion}} = 1.1 \times 10^{53}$ which agrees very well with the number derived from the optically thin radio emission, for example, at $\lambda = 1$ cm (Johnson et al. 2009).

By modelling the dust radiation from a source in a radiative transfer calculation for simple geometries, one can sometimes derive good estimates of the spatial distribution, mass, and temperature range of the dust. Here, however, the geometry is obviously much too complex, and we restrict ourselves to a decomposition of the infrared part of the SED into components and their semi-quantitative analysis. Therefore, most of the following numbers (where we use the correct emissivity $\kappa_\nu B_\nu(T_{\text{dust}})$ and not the approximation $\nu^2 B_\nu(T_{\text{dust}})$) are rough but indicative of the range of possible values.

In analysing the SED, we start by looking at its coldest component defined by the FIR emission. The data points at 70 and 160 μm imply a colour temperature of about $T_1 = 65$ K in which the dust acquires at a distance of $R_1 \sim 100$ pc from a central source of luminosity L_{gal} . Of course, when L and R are fixed, the temperature depends on the size and mineralogy of the grain, but T_1 is a mean value for the grain's standard size range (0.03 to 0.3 μm) and standard composition (silicate, amorphous carbon and graphite). The mass of this dust component is $M_1 \sim 2000 M_\odot$ or ~ 1.5 times higher than the mass derived adopting $\beta = 2$ (Table 2), and it is located at the edge of or just outside the H II region. If cold dust forms a shell of width ΔR_1 around the powering star cluster and if ρ is the density of dust per cm^3 , then the visual extinction optical depth through this shell is $\tau_{1,V} = \kappa_V \Delta R_1 \rho = \kappa_V M_1 / 4\pi R_1^2 \sim 0.13$ for an extinction coefficient at V of $\kappa_V \sim 4 \times 10^4 \text{ cm}^2$ per g of dust. This cool shell is therefore almost transparent to radiation from inside.

To explain the SED, one needs more warmer components. An important guide to their constraint plays the 9.7 μm silicate feature. It is claimed to be seen in absorption by Houck et al. (2004) with an optical depth $\tau_{9.7 \mu\text{m}} = 0.49$. One way to produce it would be by a layer of such an optical depth that lies in front of a background source with an intrinsic, undiminished flux of ~ 0.02 Jy at 9.7 μm . The foreground layer must be cooler than 150 K, or it would show the 9.7 μm resonance in emission. The background source itself, unless it is optically thick to stellar radiation, should for similar reasons be cooler than 150 K.

The visual extinction optical depth of the foreground layer is $\tau_V \sim 0.49 \times \tau_{9.7 \mu\text{m}} \sim 7.5$. If it were in front of the stellar superclusters it would strongly suppress their optical and UV emission which would then be intrinsically brighter by a factor $\sim e^{7.5}$ than what is observed. That, however, is excluded because the IR part of the SED accounts for less than half of the total luminosity. The layer also cannot lie in front of the optically visible H II region, which covers most or a considerable fraction of it, because its short wavelength emission is well detected and the extinction over it does not show much variation.

We, therefore, favour a configuration in which the near- and mid-IR emission come from one, or more likely, several optically hidden sources. Let there be N_{hid} of them with a luminosity L_{hid} with $N_{\text{hid}} L_{\text{hid}} \sim 0.27 L_{\text{gal}}$ for each. For each, one has to do a radiative transfer calculation to compute their spectra. They can then be added independently to the optical and FIR fluxes. These

Fig. 15. Full spectrum of SBS 0335-052E. Green points: WISE. Blue points: *Spitzer*. Cyan points: *Herschel*. Black points: submm and radio. The IR part of the SED can be deconvolved into three components: a cold one of $T_1 = 65$ K (blue line), a warm one of $T_2 = 160$ K (green line), and emission from dust surrounding the still enshrouded star clusters (red line). Their sum is shown by dot-dashed line.

dust hidden source(s) may be O star clusters, which are still dust enshrouded and at an earlier evolutionary stage than the visible super-star clusters. Although there must be some general trigger for the star burst in SBS 0335-052E, one sees six separated clusters and their ignition is not likely to have occurred simultaneously within a period much less than the lifetime of an O6 star.

The hidden sources of total luminosity $N_{\text{hid}} L_{\text{hid}}$ may be anywhere within the field of view of the detector: within or outside the H II region, in front of or behind it. It does not make much difference whether one imagines a few or a greater number of (spherical) hidden sources. They are in many ways quite similar: they are small (< 10 pc), so that they would never noticeably obscure the H II region and have a visual optical depth $\tau_V \sim 6$. The model that enters Fig. 15 has 40 such hidden sources, with each having a radius of 1 pc and a luminosity $L_{\text{hid}} = 2.5 \times 10^7 L_\odot$, representing a very dense cluster of about 100 O stars.

In Fig. 15, the contribution from the hidden sources emitting in the infrared is given by the thin red curve peaking near 25 μm . The cold component is also shown. To improve the fit, another warm component with $T_2 = 160$ K and a dust mass $M_2 \sim 15 M_\odot$ is shown. One would find a temperature of ~ 160 K at a distance of 10 pc from an unobscured source of $10^9 L_\odot$.

We mention that a satisfactory overall fit to the SED can, in principle, be achieved if silicate dust is absent, so that there is only carbon dust and no 10 μm feature. Then one does not need the hidden sources, and the dip near 10 μm would result from a crossover of two temperature components. However, the heavy elements in this galaxy come from exploding massive stars, and it is hard to avoid the creation of silicon and oxygen.

The existence of an additional very cold component ($\ll 60$ K) can never be excluded, but it sets limits on its maximum mass because the emission at 0.8 mm is clearly free-free radiation. For example, the mass of a 20 K component, which would be about 3 kpc away from the galaxy center, could not be much more massive than $10^4 M_\odot$.

5. Summary

We studied global characteristics of dust emission in a large sample of emission-line star-forming galaxies. The sample consists of two subsamples. One subsample includes ~ 4000 compact star-forming galaxies from the SDSS, which were also detected in all four bands at $3.4\ \mu\text{m}$, $4.6\ \mu\text{m}$, $12\ \mu\text{m}$, and $22\ \mu\text{m}$ of the WISE all-sky survey. The second subsample is a sample of 28 star-forming galaxies observed with *Herschel* in the FIR range (Rémy-Ruyer et al. 2013). For the second sample, we also use observations by WISE and *Spitzer* in the MIR range and various observations in submm and radio ranges. Both SDSS and *Herschel* samples are supplemented by available data in the UV, optical, and NIR ranges.

For the SDSS sample, we use $12\ \mu\text{m}$ and $22\ \mu\text{m}$ WISE fluxes to determine temperatures, luminosities, and masses of warm dust. For the *Herschel* sample, we adopt a three component dust model to derive global parameters for cold, warm, and hot components. The parameters of cold and warm dust are compared with other global parameters, such as stellar masses and luminosities in the UV and optical ranges.

Our main results are as follows.

1. We derived temperatures, luminosities, and masses of the warm dust in SDSS compact star-forming galaxies, adopting a modified black-body distribution with a dust emissivity index $\beta = 2.0$. The average temperature of warm dust is ~ 120 K and does not depend on the galaxy metallicity. It is slightly larger for smaller β . Warm dust luminosities are strongly correlated with $H\beta$ luminosities, implying that the main source of warm dust heating is radiation of star-forming regions that include ionising UV radiation, and, thus, warm dust is located inside and/or near H II regions.
2. Warm dust masses in SDSS compact galaxies weakly depend on the total stellar masses, but they are much more tightly correlated with masses of young stellar populations. We find a strong decrease of warm-dust-to-young-stellar-population mass ratio with decreasing equivalent width $EW(H\beta)$ of the $H\beta$ emission line, implying that this ratio is higher in younger star-forming regions and that warm dust is mainly heated by ionising radiation of H II regions. We also varied β and found that warm dust masses would be higher by a factor of ~ 2 with $\beta = 1.0$, as compared to those with $\beta = 2.0$.
3. We used WISE, *Spitzer*, and *Herschel* photometric data, as well as the optical and *Spitzer*/IRS spectra to derive global dust characteristics in 28 compact galaxies from the *Herschel* sample (Rémy-Ruyer et al. 2013) by adopting a three-component dust model and modified blackbody emissivity with the emissivity index $\beta = 2.0$. The mean dust temperatures are ~ 30 K and ~ 90 K for the cold and warm components, respectively, with no clear dependence on metallicity. Since the emission of hot dust may not be in equilibrium, we do not analyse hot dust parameters because they may deliver erroneous results. There are two outliers, the two sources with lowest metallicity, SBS 0335–052E and I Zw 18, which have higher cold dust temperatures. We find general agreement between the cold dust temperatures derived by Rémy-Ruyer et al. (2013) and in this paper.
4. Similarly to the SDSS sample, we find a tight relation between the warm dust and $H\beta$ luminosities for galaxies from the *Herschel* sample. Additionally, correlations are found between cold dust and $H\beta$ luminosities. All these relations suggest that the main source of dust heating in star-forming galaxies is UV radiation of young stellar populations.

5. The masses of the cold dust in the *Herschel* sample galaxies are in the range of $\sim 10^2$ – $10^6\ M_\odot$ and on average they are by a factor of $\sim 10^3$ higher than those of the warm dust. We found a tight correlation between masses of cold and warm dust, implying the same source of dust heating, UV radiation of star-forming regions. It is proposed to use the relation between warm and cold dust masses for estimations of the total dust mass in star-forming galaxies with an accuracy better than ~ 0.5 dex. This can appreciably increase the number of star-forming galaxies with derived dust masses from several tens to several thousands, because the data in the mid-infrared range (e.g. WISE survey) used for the determination of the warm dust characteristics are numerous.
6. We find that dust-to-neutral gas and dust-to-metal mass ratios in *Herschel* star-forming galaxies strongly decline with decreasing metallicity, which is similar to results from other studies of local emission-line galaxies, high-redshift GRB hosts, and damped Ly α absorbers (DLAs). On the other hand, the dust-to-ionised gas mass ratios are about one hundred times as high, indicating that most of dust is located in the neutral gas.
7. It is found that thermal free-free emission of ionised gas in star-forming galaxies is important in the submm range and may cause the submm excess, as discussed by Rémy-Ruyer et al. (2013). This effect is stronger in galaxies with lower metallicity and higher star-formation rate.
8. We showed that the optically thick foreground layer invoked by Plante & Sauvage (2002) and Hunt et al. (2014) to explain silicate dust absorption at $9.7\ \mu\text{m}$ in SBS 0335–052E is in conflict with observations in the UV, optical, and radio ranges and results in much higher radio emission than that observed. We propose a model in which hidden star formation in SBS 0335–052E resides in one or several dense and compact gas clouds, which are optically thick at $9.7\ \mu\text{m}$. These hidden star clusters emit only in the infrared range and do not produce an emission excess in the radio range, leading to satisfactory agreement with all observational data.

Acknowledgements. Y.I.I., N.G.G. and K.J.F. are grateful to the staff of the Max Planck Institute for Radioastronomy (MPIfR) for their warm hospitality. Y.I.I. and N.G.G. acknowledge financial support by the MPIfR. GALEX is a NASA mission managed by the Jet Propulsion Laboratory. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory, California Institute of Technology, funded by the National Aeronautics and Space Administration. Funding for the Sloan Digital Sky Survey (SDSS) and SDSS-II has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Science Foundation, the U.S. Department of Energy, the National Aeronautics and Space Administration, the Japanese Monbukagakusho, and the Max Planck Society, and the Higher Education Funding Council for England. This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

References

- Abazajian, K. N., Adelman-McCarthy, J. K., Agüeros, M. A., et al. 2009, *ApJS*, 182, 543
- Albrecht, M., Chini, R., Krügel, E., Müller, S. A. H., & Lemke, R. 2004, *A&A*, 414, 141
- Aller, L. H. 1984, *Physics of Thermal Gaseous Nebulae* (Dordrecht: Reidel)
- Annibali, F., Cignoni, M., Tosi, M., et al. 2013, *AJ*, 146, 144
- Bica, M. D., Kojoian, G., Seal, J., Dickinson, D. F., & Malkan, M. A. 1995, *ApJS*, 98, 369
- Bourne, N., Dunne, L., Bendo, G. J., et al. 2013, *MNRAS*, 436, 479

- Cannon, J. M., Walter, F., Skillman, E. D., & van Zee, L. 2005, *ApJ*, 621, L21
- Cannon, J. M., Smith, J.-D. T., Walter, F., et al. 2006, *ApJ*, 647, 293
- Caplan, J., & Deharveng, L. 1986, *A&A*, 155, 297
- Cardamone, C., Schawinski, K., Sarzi, M., et al. 2009, *MNRAS*, 399, 1199
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Chengalur, J. N., Pustilnik, S. A., Martin, J.-M., & Kniazev, A. Y. 2006, *MNRAS*, 371, 1849
- Chiaki, G., Schneider, R., Nozawa, T., et al. 2014, *MNRAS*, 439, 3121
- Condon, J. J., Helou, G., Sanders, D. B., & Soifer, B. T. 1996, *ApJS*, 103, 81
- Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, *AJ*, 115, 1693
- Cormier, D., Lebouteiller, V., Madden, S. C., et al. 2012, *A&A*, 548, A20
- Cormier, D., Madden, S. C., Lebouteiller, V. et al. 2014, *A&A*, 564, A121
- Courtois, H. M., Tully, R. B., Makarov, D. I., et al. 2011, *MNRAS*, 414, 2005
- Dale, D. A., Cohen, S. A., Johnson, L. C., et al. 2009, *ApJ*, 703, 517
- De Cia, A., Ledoux, C., Savaglio, S., Schady, P., & Vreeswijk, P. M. 2013, *A&A*, 560, A88
- de Vaucouleurs, G., de Vaucouleurs, A., Corwin, Jr., H. G., et al. 1991, 3rd Reference Catalogue of Bright Galaxies (New York: Springer Verlag)
- Dole, H., Lagache, G., Puget, J.-L., et al. 2006, *A&A*, 451, 417
- Draine, B. T. 2003, *ARA&A*, 41, 241
- Draine, B. T., Dale, D. A., Bendo, G., et al. 2007, *ApJ*, 663, 866
- Dressel, L. L., & Condon, J. J. 1978, *ApJS*, 36, 53
- Ekta, B., Pustilnik, S. A., & Chengalur, J. N. 2009, *MNRAS*, 397, 963
- Elbaz, D., Cesarsky, C. J., Chantal, P., et al. 2002, *A&A*, 384, 848
- Engelbracht, C. W., Gordon, K. D., Rieke, G. H., et al. 2005, *ApJ*, 628, L29
- Engelbracht, C. W., Rieke, G. H., Gordon, K. D., et al. 2008, *ApJ*, 678, 804
- Ferland, G. J., Korista, K. T., Verner, D. A., et al. 1998, *PASP*, 110, 761
- Ferland, G. J., Porter, R. L., van Hoof, P. A. M., et al. 2013, *Rev. Mex. Astron. Astrofis.*, 49, 137
- Freyer, D. T., et al. 2006, *ApJ*, 647, L9
- Galliano, F., Madden, S. C., Jones, A. P., Wilson, C. D., Bernard, J.-P., & Le Peintre, F. 2003, *A&A*, 407, 159
- Galliano, F., Madden, S. C., Jones, A. P., Wilson, C. D., & Bernard, J.-P. 2005, *A&A*, 434, 867
- Galliano, F., Dwek, E., & Chantal, P. 2008, *ApJ*, 672, 214
- Galametz, M., Madden, S. C., Galliano, F., et al. 2009, *A&A*, 508, 645
- Galametz, M., Madden, S. C., Galliano, F., et al. 2011, *A&A*, 532, A56
- Gil de Paz, A., Madore, B. F., & Pevunova, O. 2003, *ApJS*, 147, 29
- Griffith, R. L., Tsai, C.-W., Stern, D., et al. 2011, *ApJ*, 736, L22
- Gruppioni, C., Pozzi, F., Andreani, P., et al. 2010, *A&A*, 518, L27
- Guseva, N. G., Izotov, Y. I., & Thuan, T. X. 2000, *ApJ*, 531, 776
- Guseva, N. G., Papaderos, P., Izotov, Y. I., Noeske, K. G., & Fricke, K. J. 2004, *A&A*, 421, 519
- Guseva, N. G., Izotov, Y. I., & Thuan, T. X. 2006, *ApJ*, 644, 890
- Guseva, N. G., Izotov, Y. I., Papaderos, P., & Fricke, K. J. 2007, *A&A*, 464, 885
- Guseva, N. G., Izotov, Y. I., Stasińska, G., et al. 2011, *A&A*, 529, A149
- Guseva, N. G., Izotov, Y. I., Fricke, K. J., & Henkel, C. 2012, *A&A*, 541, A115
- Gutiérrez, L., & Beckman, J. E. 2010, *ApJ*, 710, L44
- Hao, L., Wu, Y., Charmandaris, V., et al. 2009, *ApJ*, 704, 1159
- Hauser, M. G., & Dwek, E. 2001, *ARA&A*, 39, 249
- Hirashita, H. 2011, *MNRAS*, 418, 828
- Hirashita, H., Tajiri, Y. Y., & Kamaya, H. 2002, *A&A*, 388, 439
- Hirashita, H., Kaneda, H., Onaka, T., & Suzuki, T. 2008, *PASJ*, 60, S477
- Houck, J. R., Charmandaris, V., Brandl, B. R., et al. 2004, *ApJS*, 154, 211
- Huchtmeier, W. K., & Richter, O.-G. 1989, *A General Catalog of HI Observations of Galaxies* (New York: Springer-Verlag)
- Huchtmeier, W. K., Krishna, G., & Petrosian, A. 2005, *A&A*, 434, 887
- Hunt, L., Bianchi, S., & Maiolino, R. 2005a, *A&A*, 434, 849
- Hunt, L. K., Dyer, K. K., & Thuan, T. X. 2005b, *A&A*, 436, 837
- Hunt, L. K., Thuan, T. X., Sauvage, M., & Izotov, Y. I., M. 2006, *ApJ*, 653, 222
- Hunt, L. K., Thuan, T. X., Izotov, Y. I., & Sauvage, M. 2010, *ApJ*, 712, 164
- Hunt, L. K., Testi, L., Casasola, V., et al. 2014, *A&A*, 561, A49
- Hunter, D. A., van Woerden, H., & Gallagher, J. S., III 1994, *ApJS*, 91, 79
- Indebetouw, R., de Messières, G. E., Madden, S., et al. 2009, *ApJ*, 694, 84
- Indebetouw, R., Brogan, C., Chen, C.-H. R., et al. 2013, *ApJ*, 774, 73
- Izotov, Y. I., & Thuan, T. X. 1998, *ApJ*, 500, 188
- Izotov, Y. I., & Thuan, T. X. 1999, *ApJ*, 511, 639
- Izotov, Y. I., & Thuan, T. X. 2004, *ApJ*, 602, 200
- Izotov, Y. I., & Thuan, T. X. 2007, *ApJ*, 665, 1115
- Izotov, Y. I., & Thuan, T. X. 2011, *ApJ*, 734, 82
- Izotov, Y. I., Thuan, T. X., & Lipovetsky, V. A. 1994, *ApJ*, 435, 647
- Izotov, Y. I., Thuan, T. X., & Lipovetsky, V. A. 1997, *ApJS*, 108, 1
- Izotov, Y. I., Chaffee, F. H., & Green, R. F. 2001, *ApJ*, 562, 727
- Izotov, Y. I., Stasińska, G., Meynet, G., Guseva, N. G., & Thuan, T. X. 2006a, *A&A*, 448, 955
- Izotov, Y. I., Schaerer, D., Blecha, A., Royer, F., Guseva, N. G., & North, P. 2006b, *A&A*, 459, 71
- Izotov, Y. I., Guseva, N. G., & Thuan, T. X. 2011a, *ApJ*, 728, 161
- Izotov, Y. I., Guseva, N. G., Fricke, K. J., & Henkel, C. 2011b, *A&A*, 536, L7
- Izotov, Y. I., Stasińska, G., & Guseva, N. G. 2013, *A&A*, 558, A57
- Izotov, Y. I., Guseva, N. G., Fricke, K. J., & Henkel, C. 2014, *A&A*, 561, A33
- James, A., Dunne, L., Eales, S., & Edmunds, M. G. 2002, *MNRAS*, 335, 753
- Johnson, K. E., Hunt, L. K., & Reines, A. E. 2009, *AJ*, 137, 3788
- Karczewski, O. L., Barlow, M. J., Page, M. J., et al. 2013, *MNRAS*, 431, 2493
- Kepley, A. A., Reines, A. E., Johnson, K. E., & Walker, L. M. 2014, *AJ*, 147, 43
- Kimball, A. E., & Ivezić, Z. 2008, *AJ*, 136, 684
- Kirkpatrick, A., Calzetti, D., Galametz, M., et al. 2013, *ApJ*, 778, 51
- Klein, U., Weiland, H., & Brinks, E. 1991, *A&A*, 246, 323
- Krügel, E. 2008, *An introduction to the physics of interstellar dust*, Publ. Taylor & Francis, Series in astronomy and astrophysics
- Lagos, P., Telles, E., & Melnick, J. 2007, *A&A*, 476, 89
- Lebouteiller, V., Kunth, D., Lequeux, J., et al. 2004, *A&A*, 415, 55
- Lebouteiller, V., Barry, D. J., Spoon, H. W. W., et al. 2011, *ApJS*, 196, 8
- Lebouteiller, V., Heap, S., Hubeny, I., & Kunth, D. 2013, *A&A*, 553, A16
- Lecavelier des Etangs, A., Désert, J.-M., Kunth, D., et al. 2004, *A&A*, 413, 131
- Leitherer, C., Schaerer, D., Goldader, J. D., et al. 1999, *ApJS*, 123, 3
- Lelli, F., Verheijen, M., Fraternali, F., & Sancisi, R. 2012, *A&A*, 537, A72
- Leroy, A., Bolatto, A. D., Simon, J. D., & Blitz, L. 2005, *ApJ*, 625, 763
- Leroy, A., Cannon, J., Walter, F., Bolatto, A., & Weiss, A. 2007, *ApJ*, 663, 990
- Lisenfeld, U., & Ferrara, A. 1998, *ApJ*, 496, 145
- Madden, S. C., Galliano, F., Jones, A. P., & Sauvage, M. 2006, *A&A*, 446, 877
- Madden, S. C., Remy-Ruyer, A., Galametz, M., et al. 2013, *PASP*, 125, 600
- Martin, D. C., Fanson, J., Schiminovich, J., et al. 2005, *ApJ*, 619, L1
- Mauch, T., Murphy, T., Buttery, H. J., et al. 2003, *MNRAS*, 342, 1117
- Morales-Luis, A. B., Sánchez Almeida, J., Aguerri, J. A. L., et al. 2011, *ApJ*, 743, 77
- Moustakas, J., & Kennicutt, R. C., Jr. 2006, *ApJS*, 164, 81
- Nozawa, T., Kozasa, T., Umeda, H., Maeda, K., & Nomoto, K. 2003, *ApJ*, 598, 785
- O'Halloran, B., Satyapal, S., & Dudik, R. P. 2006, *ApJ*, 641, 795
- O'Halloran, B., Galametz, M., Madden, S. C., et al. 2010, *A&A*, 518, L58
- Planck collaboration XV 2014, *A&A*, 571, A15
- Plante, S., & Sauvage, M. 2002, *AJ*, 124, 1995
- Pustilnik, S. A., Martin, J.-M., Huchtmeier, W. K., et al. 2002, *A&A*, 389, 405
- Pustilnik, S. A., Pramskij, A. G., & Kniazev, A. Y. 2004, *A&A*, 425, 51
- Rabidoux, K., Pisano, D. J., Kepley, A. A., Johnson, K. E., & Balser, D. S. 2014, *ApJ*, 780, 19
- Refsdal, S., Stabell, R., & de Lange, F. G., 1967, *Mem. RAS*, 71, 143
- Rémy-Ruyer, A., Madden, S. C., Galliano, F., et al. 2013, *A&A*, 557, A95
- Rémy-Ruyer, A., Madden, S. C., Galliano, F., et al. 2014, *A&A*, 563, A31
- Rosa-Gonzalez, D., Schmitt, H. R., Terlevich, E., & Terlevich, R. 2007, *ApJ*, 654, 226
- Sandstrom, K. M., Bolatto, A. D., Draine, B. T., Bot, C., & Stanimirović, S. 2010, *ApJ*, 715, 701
- Schaerer, D., & Vacca, W. D. 1998, *ApJ*, 497, 618
- Schmitt, H. R., Calzetti, D., Armus, L., et al. 2006, *ApJS*, 164, 52
- Schneider, R., Omukai, K., Limongi, M., et al. 2012, *MNRAS*, 423, L60
- Skrutskie, M. F., Cutri, R. M., Stiening, R., et al. 2006, *AJ*, 131, 1163
- Springob, C. M., Haynes, M. P., Giovanelli, R., & Kent, B. R. 2005, *ApJS*, 160, 149
- Stasińska, G., & Izotov, Y. I. 2003, *A&A*, 397, 71
- Thuan, T. X., & Martin, G. E. 1981, *ApJ*, 247, 823
- Thuan, T. X., Izotov, Y. I., & Lipovetsky, V. A. 1995, *ApJ*, 445, 108
- Thuan, T. X., Sauvage, M., & Madden, S. 1999a, *ApJ*, 516, 783
- Thuan, T. X., Lipovetsky, V. A., Martin, J.-M., & Pustilnik, S. A. 1999b, *A&AS*, 139, 1
- Thuan, T. X., Lecavelier des Etangs, A., & Izotov, Y. I. 2002, *ApJ*, 565, 941
- Thuan, T. X., Hibbard, J. E., & Levrier, F. 2004, *AJ*, 128, 617
- Thuan, T. X., Lecavelier des Etangs, A., & Izotov, Y. I. 2005, *ApJ*, 621, 269
- Ugryumov, A. V., Engels, D., Pustilnik, S. A., et al. 2003, *A&A*, 397, 463
- Vacca, W. D., & Conti, P. S. 1992, *ApJ*, 401, 543
- Valiante, R., Schneider, R., Salvadori, S., & Bianchi, S. 2011, *MNRAS*, 416, 1916
- Viallefond, F., & Thuan, T. X. 1983, *ApJ*, 269, 444
- Weingartner, J. C., & Draine, B. T. 2001, *ApJ*, 548, 296
- Williams, B. A., McMahon, P. M., & van Gorkom, J. H. 1991, *AJ*, 101, 1957
- Wright, E. L., Eisenhardt, P. R. M., Mainzer, A. K., et al. 2010, *AJ*, 140, 1868
- Wu, Y., Charmandaris, V., Hao, L., et al. 2006, *ApJ*, 639, 157
- Wu, Y., Charmandaris, V., Hunt, L. K., et al. 2007, *ApJ*, 662, 952
- Wu, Y., Bernard-Salas, J., Charmandaris, V., et al. 2008, *ApJ*, 673, 193

Fig. 6. Spectral energy distribution fits of the galaxies from *Herschel* sample. Observed spectra in the optical and MIR ranges are shown by black and magenta lines. Photometric data are shown from GALEX (blue filled circles), SDSS (open magenta circles), 2MASS (black open circles), WISE (green symbols), *Spitzer* (black filled circles), *Herschel* (red filled circles), and submm and radio (open magenta rhombs). Blue, green and red spectra at $\lambda < 15 \mu\text{m}$ are fitted SEDs for ionised gas, stellar and total emission, which are produced from the optical spectra. Fits of dust emission at $\lambda > 5 \mu\text{m}$ are shown by red, green, and blue dotted lines for different components and by black solid line for total emission. Blue solid line at $\lambda > 100 \mu\text{m}$ is free-free emission extrapolated from the extinction- and aperture-corrected $H\beta$ flux. Blue dashed line indicates the wavelength at which dust flux is equal to extrapolated free-free emission.

Fig. 6. continued.

Fig. 6. continued.

Fig. 6. continued.