

## Hot super-Earths and giant planet cores from different migration histories (*Corrigendum*)

Christophe Cossou<sup>1,2</sup>, Sean N. Raymond<sup>1,2</sup>, Franck Hersant<sup>1,2</sup>, and Arnaud Pierens<sup>1,2</sup>


<sup>1</sup> Univ. Bordeaux, Laboratoire d'Astrophysique de Bordeaux, UMR 5804, 33270 Floirac, France  
e-mail: [christophe.cossou@u-bordeaux1.fr](mailto:christophe.cossou@u-bordeaux1.fr)

<sup>2</sup> CNRS, Laboratoire d'Astrophysique de Bordeaux, UMR 5804, 33270 Floirac, France  
e-mail: [rayray.sean@gmail.com](mailto:rayray.sean@gmail.com)

A&A 569, A56 (2014), DOI: [10.1051/0004-6361/201424157](https://doi.org/10.1051/0004-6361/201424157)

**Key words.** planets and satellites: formation – protoplanetary disks – planet-disk interactions – methods: numerical – errata, addenda

An error occurred during the production process. An incorrect Fig. 14 was published. The correct Fig. 14 is published below.



**Fig. 14.** Comparison between the planets that formed in simulations with dissipating disks and observed extra-solar planets. Both samples are limited to the range  $R > 1.5 R_{\oplus}$  ( $3.3 M_{\oplus}$  assuming Earth density) and  $p < 200$  days.