

Surface brightness profile of the Milky Way's nuclear star cluster (Corrigendum)

R. Schödel¹, A. Feldmeier², D. Kunneriath³, S. Stolovy⁴, N. Neumayer², P. Amaro-Seoane⁵, and S. Nishiyama⁶

¹ Instituto de Astrofísica de Andalucía (CSIC), Glorieta de la Astronomía s/n, 18008 Granada, Spain
e-mail: rainer@iaa.es

² European Southern Observatory, Karl-Schwarzschild-Strasse 2, 85748 Garching bei München, Germany

³ Astronomical Institute, Academy of Sciences, Boční II 1401, 14100 Prague, Czech Republic

⁴ El Camino College, 16007 Crenshaw Blvd., Torrance, CA 90506, USA

⁵ Max-Planck Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, 14476 Potsdam, Germany

⁶ National Astronomical Observatory of Japan, Mitaka, 181-8588 Tokyo, Japan

A&A 566, A47 (2014), DOI: [10.1051/0004-6361/201423481](https://doi.org/10.1051/0004-6361/201423481)

Key words. dust, extinction – Galaxy: center – Galaxy: nucleus – Galaxy: structure – galaxies: nuclei – errata, addenda

In Schödel et al. (2014), the axis labels of Figs. 4, 6–9, 11, and 13 should not refer to “RA” (Right ascension) and “Dec” (Declination) because the orientation of all the images is such that Galactic north is up and Galactic east is to the left. Instead, the axis labels indicate the offsets from Sagittarius A* parallel and perpendicular to the Galactic plane, as is correctly described in Fig. 1 and its caption.

Acknowledgements. We thank Ortwin Gerhard for bringing this to our attention.

References

Schödel, R., Feldmeier, A., Kunneriath, D., et al. 2014, [A&A, 566, A47](https://doi.org/10.1051/0004-6361/201423481)