LETTER TO THE EDITOR

Searching for electromagnetic counterpart of LIGO gravitational waves in the *Fermi* GBM data with ADWO

Zsolt Bagoly¹, Dorottya Szécsi², Lajos G. Balázs^{1,3}, István Csabai¹, István Horváth⁴, László Dobos¹,
János Lichtenberger^{1,5}, and L. Viktor Tóth¹

¹ Dept. of Physics of Complex Systems, Eötvös University, 1053 Budapest, Hungary
e-mail: zsolt.bagoly@elte.hu

² Argelander-Institut für Astronomie der Universität Bonn, 53113 Bonn, Germany

³ Konkoly Observatory, RCAES, Hungarian Academy of Sciences, 1051 Budapest, Hungary

⁴ National University of Public Service, 1083 Budapest, Hungary

⁵ Geodetic and Geophysical Institute, RCAES, Hungarian Academy of Sciences, 1051 Budapest, Hungary

Received 21 March 2016 / Accepted 17 August 2016

ABSTRACT

Aims. The *Fermi* collaboration identified a possible electromagnetic counterpart of the gravitational wave event of September 14, 2015. Our goal is to provide an unsupervised data analysis algorithm to identify similar events in *Fermi*'s Gamma-ray Burst Monitor CTTE data stream.

Methods. We are looking for signals that are typically weak. Therefore, they can only be found by a careful analysis of count rates of all detectors and energy channels simultaneously. Our Automatized Detector Weight Optimization (ADWO) method consists of a search for the signal, and a test of its significance.

Results. We developed ADWO, a virtual detector analysis tool for multi-channel multi-detector signals, and performed successful searches for short transients in the data-streams. We have identified GRB150522B, as well as possible electromagnetic candidates of the transients GW150914 and LVT151012.

Conclusions. ADWO is an independently developed, unsupervised data analysis tool that only relies on the raw data of the *Fermi* satellite. It can therefore provide a strong, independent test to any electromagnetic signal accompanying future gravitational wave observations.

Key words. gamma rays: general – gravitational waves – gamma-ray burst: general – gamma-ray burst: individual: GRB150522

1. Introduction

We present a new method to search for non-triggered, short-duration transients in the data-set of the *Fermi* Gamma-ray Space Telescope's Gamma-ray Burst Monitor (GBM; [Carson 2007](#); [Meegan et al. 2009](#)). The method, called Automatized Detector Weight Optimization (ADWO), combines the data of all available detectors and energy channels, identifying those with the strongest signal. This way, we are able to separate potential events from the background noise and present the statistical probability of a false alarm. Although it is possible to apply our ADWO method to look for non-triggered short gamma-ray bursts (SGRBs), ADWO works the best if a potential event at a given time (and, if available, a given celestial position) is provided as an input. Thus, ADWO is ideal to search for electromagnetic (EM) counterparts of gravitational wave (GW) events, when the time of the event is well known from the GW-detectors' observation.

On September 14, 2015 at 09:50:45.391 UTC the two detectors of the advanced Laser Interferometer Gravitational-Wave Observatory (LIGO) simultaneously observed a transient gravitational-wave signal GW150914 ([Abbott et al. 2016c](#)). The signal is originated from the merger of a binary black hole (BBH) system at low redshift ($z \approx 0.1$; [Abbott et al. 2016a](#)).

GBM observations revealed a weak transient source above 50 keV, 0.4 s after the GW event, with a false alarm probability of 0.0022 ([Connaughton et al. 2016](#)). This weak transient, with a

duration of ≈ 1 s, does not appear to be connected with any other previously known astrophysical, solar, terrestrial, or magnetospheric activity. Its localization is ill-constrained but consistent with the direction of GW150914. The duration and spectrum of the *Fermi* transient event suggest that the radiation was arriving at a large angle relative to the direction where the Fermi Large Area Telescope (LAT) was pointing.

The EM transient was a result of a custom pipeline looking for prompt gamma-ray counterparts in GBM ([Blackburn et al. 2015](#); [Kelley et al. 2013](#)), optimized for LIGO/Virgo GW candidate events. The automatic GBM pipelines (looking for GRBs) did not find any transients.

Neither the *Fermi*-LAT observation ([Ackermann et al. 2016](#)) above 100 MeV nor the partial Swift follow-up ([Evans et al. 2016](#)) in the X-ray, optical and UV bands, nor the INTEGRAL observations ([Savchenko et al. 2016](#)) in the gamma-ray and hard X-ray bands found any potential counterparts to GW150914, they only provide limits on the transient counterpart activity.

However, from a theoretical point of view, EM counterparts such as short duration GRBs associated with GW events are not excluded. Recently, [Perna et al. \(2016\)](#) proposed a scenario where a double black hole merger is accompanied by a SGRB. The evolution of the system starts with two low-metallicity massive stars that are orbiting around each other ([de Mink et al. 2009](#); [Marchant et al. 2016](#); [Mandel & de Mink 2016](#)). Their orbit is so tight initially that their rotational periods are synchronized with the orbital period. Due to the fast rotation, these

stars evolve homogeneously and never expand (as described by Szécsi et al. 2015, for single, homogeneously evolving stars). This way, the stars avoid the supergiant phase and thus a common envelope evolution, which reduces the theoretical uncertainties involved. Assuming that (at least) one of the supernova explosions leaves a long-lived disk behind, Perna et al. (2016) predict that this scenario leads to a relativistic jet to be launched during the merger of the black holes. The burst-duration timescale they derive from their models is in the order of 5 ms. In light of these theoretical models that predict not only the existence of black hole mergers but even the subsequent production of a SGRB, it is quite reasonable to look for EM transients of any possible gravitational wave detection.

LVT151012, the second GW candidate transient event occurred on October 12, 2015 at 09:54:43.555 UTC (Abbott et al. 2016b; The LIGO Scientific Collaboration et al. 2016). They report a false alarm probability of 0.02, and consider it not to be low enough to confidently claim this event as a real GW signal.

This paper is organized as follows: in Sect. 2 we describe our method, in Sect. 3 we test our ADWO method with the short-duration GRB150522B and in Sect. 4 with the SGRB-like signal that accompanied the GW150914 event. We find that our analysis of these signals are in accordance with the results of Connaughton et al. (2016). In Sect. 5 we apply ADWO to look for a potential EM counterpart of the event LVT151012.

2. Input data and methods

2.1. Fermi GBM overview

The *Fermi* GBM includes two sets of detectors: 12 thallium activated Sodium Iodide (NaI(Tl)) and two Bismuth Germanate (BGO) scintillation detectors (Meegan et al. 2009). The NaI(Tl) detectors measure the low-energy spectrum (8 keV to ~ 1 MeV) while the BGO detectors have an energy range of ~ 200 keV to ~ 40 MeV. The effective area of the detectors varies with the photon energy and the angle of incidence, with a maximum of ~ 100 cm² (NaI(Tl)) and ~ 120 cm² (BGO).

Signals from the photomultipliers are analyzed on-board, and the pulse height analysis (PHA) converts the peak heights into 128 PHA channels. The signal distribution in these PHA channels as a function of the incoming photon energy and geometry is described by the detector response matrix (DRM). The DRMs contain the effective detection area as the function of the angular dependence of the efficiency, energy deposition and dispersion, detector non-linearity, as well as the atmospheric and spacecraft scattering. The PHA distribution is usually wide for high-energy photons (especially above ~ 1 MeV), as some photons will scatter prior to detection. The DRMs are provided as a standard data product for each GBM trigger, but neither the program nor the data are public.

It is important to note that the 128 PHA channels have different energy ranges from detector to detector, according to the detector's actual setup. The PHA channels are aggregated into different data products, e.g. CTIME data, which consist of accumulated spectra from each detector with a 8-channel energy and 64/265 ms time resolution.

A GBM trigger occurs when the count rates of two or more detectors exceed the background with a given threshold ($4.5\text{--}7.5\sigma$). The trigger algorithms include four energy ranges (25–50 keV, 50–300 keV, 100–300 keV, and >300 keV) and ten timescales (from 16 ms to 8.192 s). A total of 120 different trigger algorithms can be specified, from which usually ~ 75 operate simultaneously.

2.2. Automatized Detector Weight Optimization (ADWO)

The basic problem of the event analysis is to find the parameters of an event in multi-detector multi-channel time series when the approximate time and direction of the expected signal are given. To calculate the significance of such an event as described by PHA counts, one should take the typical background noise and the spectral model into account. To obtain the background-induced PHA counts, the assumed synthetic spectrum is multiplied by the DRM and binned. This is then compared to the PHA counts derived from the combination of the signal and the background with tools such as XSPEC for fitting Gaussian signals using χ^2 , and C-Stat for Poisson signals (Arnaud 1996).

Contrary to this detection method, here we do not assume the event time, only a possible time interval is given. Our goal is to find the strongest weights and the best time position in this interval using a weighted signal from the multi-detector multi-channel continuous data. The simplest method would be to compare the sum of the count rates within and outside the given time interval. This approach, however, is not the most effective one in a multi-channel multi-detector environment, since for a maximum signal-to-noise ratio usually only those detectors should be summed (selected for the analysis) which produce the strongest signals. Noisy energy channels and not illuminated detectors with very low DRM should either not be taken into account, or only with a low weight. A further complication arises from the fact that we know neither the direction of the event (and, therefore, if a given detector is illuminated or not), nor the spectra.

Our solution for these problems is the following: we give different weights to different energy channels (e_i) and detectors (d_j), and optimize the Signal's Peak to Background's Peak Ratio (SPBPR). The weights are positive and normalized as $\sum e_i = 1$, $\sum d_j = 1$. We do not restrict these weights any further, i.e. we do not include any information about the DRM (which we do not know anyway, without any spectral and directional information).

If the background subtracted intensity in the j th detector's i th energy channel is $C_{ij}(t)$, we define our composite signal as $S(t) = \sum_{i,j} e_i d_j C_{ij}(t)$. The Signal's Peak is the maximum of $S(t)$ within the given time interval, and the Background's Peak is the maximum of $S(t)$ outside this interval. Our goal is to maximize the ratio of these two maximums. The best weights will be built up by iteration, maximizing SPBPR as a function of e_i and d_j . These e_i and d_j weights create an optimal filter among the spectra and detectors. The algorithm will provide not only maximum value of SPBPR, but will search the best weights and the exact time of this maximum, within the pre-defined interval.

We call this algorithm the Automatized Detector Weight Optimization. ADWO is similar to the GRB satellites' triggering mechanism, but includes several improvements. For example, while *Fermi*'s trigger algorithm selects the e_i and d_j factors to be 0 or 1, here we allow intermediate values too. Additionally, the condition that at least two detectors exceed a threshold simultaneously, is not required anymore, since the ADWO algorithm will automatically produce the best d_j weights. For a signal with time-evolving spectrum, ADWO will determine the best trigger time position.

We applied Matlab's/Octave's *fminsearch* routine to find the maximum via the Nelder & Mead Simplex algorithm. The algorithm always started from an equal weights position. The analysis of ≈ 100 signal data points against the $\approx 10^4$ background data points with 80 dimensional data took several minutes on a 4-core Intel i7 processor with 8GB memory, depending on the linear algebra packages used by the programs. After the search converged, the differences of the weights on the final simplex

were below 10^{-4} (the sum of the weights is 1). The sample Matlab/Octave code is available on GitHub¹.

2.3. Analysis of the *Fermi* GBM data

Since November 2012, the *Fermi* GBM continuous time-tagged event (CTTE) data is present for each detector with a time precision of $2 \mu\text{s}$, in all the 128 PHA energy channels (Meegan et al. 2009). Here we use the same CTTE energy channels of Connaughton et al. (2016), with limits of 4.4, 12, 27, 50, 100, 290, 540, 980 and 2000 keV (denoted with $e_1 \dots e_8$, resp.). Since we look for spectrally hard events, we use only the upper 6 energy channels in the 27–2000 keV range ($e_3 \dots e_8$). The exclusion of the low energy channels also reduces the background contamination from soft particle events, such as Cygnus X-1 and other weak variable X-ray sources, since their flux is usually small above 27 keV.

All the 12 NaI(Tl) and both BGO detectors were included in the analysis. Since the BGO detectors' low energy PHA channels start above 100 keV, the corresponding 27–100 keV energy channels are empty. Overall, we have $6 \times 14 - 2 \times 2 = 80$ non-zero time series.

For each detector and for each channel, the CTTE $2 \mu\text{s}$ event data is filtered with a 64 ms wide moving average filter at 1 ms steps, producing the $C_{ij}(t)$ light curve. This filtering is important as the photon event data are quite sparse (the intensity is quite low; for the GW150914 event there is, on average ≈ 5.8 ms between photons in a given detector and energy channel). Our 64 ms window contains 11.2 photons on average, while this window size corresponds to the typical triggered CTTE light curve resolution.

Without filtering, the photon-photon correlation in time that we search for would disappear. Very narrow filters are worthless because of the sparsity constraint, while much wider filters will smooth and filter out short transients, lowering ADWO's sensitivity. As a byproduct, the smoothing also acts as a low-pass filter which reduces the Poisson noise. The moving average filter is the simplest choice here: e.g. using some prior knowledge about the signal's shape, a matching filter tuned to the signal would improve the sensitivity at the expense of generality.

Fermi operates in survey mode most of the time, with slewing at ≈ 4 degrees per minute. This creates a continuously changing background, which should be accounted for, since ADWO would be optimal without directional changes (as it uses the correlation between the detectors and channels). One possibility would be to take the detailed satellite positional information into account and create a physical model to determine the background for a hundreds of seconds (Szécsi et al. 2013). However, we expect that the slow slew will not suppress the sensitivity to the kind of short ($\sim$ s) transients that we are looking for. Therefore, a much simpler, 6th order polynomial background fit was subtracted for each channel and detector, similar to the method of Connaughton et al. (2016). We chose the typical background window to be $\approx (-200, 500)$ s around the search window, depending on the CTTE data availability: this window can contain the majority of GBM's GRB lightcurves and covers approx. 1/7th of *Fermi*'s orbit.

3. GRB150522B

To test ADWO, we analyze the short GRB150522B gamma-ray burst, with $T_{90} = 1.02 \pm 0.58$ s and $2.13 \pm 0.12 \times 10^{-7}$ erg/cm²

Fig. 1. ADWO light curve of GRB150522B in the 27–2000 keV range.

Table 1. Channel weights.

Transient	e_3	e_4	e_5	e_6	e_7	e_8
GRB150522B	0.090	0.297	0.315	0.188	0.000	0.110
GW150914	0.203	0.050	0.056	0.559	0.110	0.022
LVT151012	0.260	0.212	0.010	0.113	0.000	0.406

fluence. These parameters are comparable to the EM companion values of GW150914, as reported by Connaughton et al. (2016). *Fermi* triggered on May 22, 2015 at 22:38:44.068 UTC, and full CTTE data of $(-137, 476)$ s interval relative to the trigger is analyzed, using a 6 s long signal window centered on the trigger. The ADWO obtains a maximal SPBPR of 3.12, and reveals the double pulse shown in the *Fermi* GBM quicklook data (Fig. 1). The detector and energy channel weights are given in Tables 1, 2.

To determine the significance we generated a Poisson-distributed synthetic signal, using the background photon data of the interval, and repeated ADWO for 10^4 Monte-Carlo (MC) simulations with the same window width. There was no simulation with bigger SPBPR value than 3.12, therefore we estimate the false alarm rate to be below 2×10^{-5} Hz, and the false alarm probability to be below 2×10^{-5} Hz $\times$ 0.125 s $\times$ $(1 + \ln(6 \text{ s}/64 \text{ ms})) = 2.8 \times 10^{-5}$, analogously to Connaughton et al. (2016).

4. The GW150914 event

We apply the ADWO method on the *Fermi* CTTE data set covering the event of GW150914: the 6 s long signal window was centered on September 14, 2015 09:50:45 UTC (391 ms before trigger). Here we investigate a $(690-6)$ s time background interval that adds up as 195 s before and 495 s after the time of the possible event. The ADWO has converged (Fig. 2) and the obtained maximal SPBPR is 1.911, 474 ms after the GW trigger.

We repeated ADWO for 10^4 MC simulations using this data: 86 cases had bigger SPBPR than 1.911. The false alarm rate is 0.0014 Hz, giving a false alarm probability of 2.8×10^{-3} Hz $\times$ 0.474 s $\times$ $(1 + \ln(6 \text{ s}/64 \text{ ms})) = 0.0075$, which is higher than 0.0022, the value given by Connaughton et al. (2016).

5. LVT151012

Considering the positional errors of GW150914 on the sky, it can be easily shown that there's a high ($>70-75\%$) probability that a similar error ring will intersect with *Fermi* GBM's field of view in the case of LVT151012 too. Therefore, we apply ADWO on the *Fermi* GBM CTTE data around the event of

¹ <https://github.com/zbagoly/ADWO>

Table 2. Detector weights for the $n0 \dots n9$, na and nb NaI(Tl) and $b0$ and $b1$ BGO detectors (as listed in Meegan et al. 2009).

Transient	$n0$	$n1$	$n2$	$n3$	$n4$	$n5$	$n6$	$n7$	$n8$	$n9$	na	nb	$b0$	$b1$
GRB150522B	0.105	0.106	0.100	0.078	0.146	0.073	0.001	0.031	0.000	0.021	0.009	0.050	0.113	0.167
GW150914	0.000	0.044	0.028	0.151	0.000	0.000	0.035	0.045	0.228	0.090	0.138	0.162	0.000	0.077
LVT151012	0.034	0.062	0.000	0.127	0.073	0.125	0.151	0.000	0.000	0.010	0.234	0.162	0.000	0.022

Fig. 2. ADWO light curve of GW150914 in the 27–2000 keV range. The inset shows the whole time interval where the ADWO search was performed. **Fig. 3.** ADWO light curve of LVT151012 in the 27–2000 keV range. The inset shows the whole time interval where the ADWO search was performed.

LVT151012, covering $(-195, 495)$ s, centered on October 12, 2015 at 09:54:43 UTC. We find a relatively strong signal at 09:54:44.207 UTC in the 6 s signal window, with a SPBPR of 1.805 (Fig. 3). The sum of the 27–290 keV weights is higher than in the case of GW150914, i.e. here the signal is softer than GW150914 at the peak ($E_p \approx 3.5$ MeV), but harder than GRB15522 at the peak ($E_p \approx 130$ keV).

We made 10^4 MC simulations: 308 cases had bigger SPBPR than the original observation. hence the false alarm rate is 0.0051 Hz, and the false alarm probability is estimated to be $0.01 \text{ Hz} \times 0.652 \text{ s} \times (1 + \ln(6 \text{ s}/64 \text{ ms})) = 0.037$.

When cross-checking the lightning detections made by WWLLN (Rodger et al. 2009) with the *Fermi*'s positions and times, we find no TGF candidates (storm activity) within 500 km of the spacecraft position and ± 900 s around the peak.

6. Discussion

Although here we applied our ADWO method to look for particular events, we point out that it is entirely possible to use this unsupervised data analysis method for a general search for non-triggered, short-duration *Fermi* GBM events. Automatized

search processes are important, as the total data-set collected by the *Fermi*'s 8-yr operation is significantly larger than the triggered data-set. It is likely that there are several potential EM events observed but not triggered. For example, based on the CTIME 256 ms data product, Gruber et al. (2012) estimates ≈ 1.6 untriggered SGRB/month in the *Fermi* observations. It is thus a worthwhile future task to identify potential SGRB candidates in the non-triggered *Fermi* GBM data-set using ADWO. Alternatively, we can cross-check those already found by other algorithms.

As our ADWO method is independently developed, and only relies on the raw data of the satellite, it can provide a strong, independent test to any future signal. In regard of the current expectation that LIGO will detect several GW events in the near future, many of which may have a weak EM transient counterpart such as a SGRB, it is of crucial importance to identify those potential EM signals. We therefore expect that ADWO will be successfully applied in the future to find SGRB counterparts of the GW events observed by LIGO. The analysis of the GW151226 event as well as the improvement of ADWO is the topic of a forthcoming paper.

Acknowledgements. This work was supported by OTKA grants NN111016 and NN114560. The authors wish to thank the World Wide Lightning Location Network (<http://wwlln.net>), a collaboration among over 50 universities and institutions, for providing the lightning location data used in this paper. Thanks for the computational resources of the Wigner GPU Laboratory of the Wigner RCP of the H.A.S. A.C. Elbeze for pointing out that GRB150522B was the second GRB of the day and the anonymous referee for valuable comments on the paper.

References

- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016a, *ApJ*, **818**, L22
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016b, *Phys. Rev. D*, **93**, 122003
- Abbott, B. P., Abbott, R., Abbott, T. D., et al. 2016c, *Phys. Rev. Lett.*, **116**, 061102
- Ackermann, M., Ajello, M., Albert, A., et al. 2016, *ApJ*, **823**, L2
- Arnaud, K. A. 1996, in *Astronomical Data Analysis Software and Systems V*, eds. G. H. Jacoby, & J. Barnes, *ASP Conf. Ser.*, **101**, 17
- Blackburn, L., Briggs, M. S., Camp, J., et al. 2015, *ApJS*, **217**, 8
- Carson, J. 2007, *J. Phys. Conf. Ser.*, **60**, 115
- Connaughton, V., Burns, E., Goldstein, A., et al. 2016, *ApJ*, **826**, L6
- de Mink, S. E., Cantiello, M., Langer, N., et al. 2009, *A&A*, **497**, 243
- Evans, P. A., Kennea, J. A., Barthelmy, S. D., et al. 2016, *MNRAS*, **460**, L40
- Gruber, D., et al. (Fermi/GBM Collaboration) 2012, in *Proc. of the Gamma-Ray Bursts 2012 Conference (GRB 2012)*, May 7–11, Munich, Germany, 36
- Kelley, L. Z., Mandel, I., & Ramirez-Ruiz, E. 2013, *Phys. Rev. D*, **87**, 123004
- Mandel, I., & de Mink, S. E. 2016, *MNRAS*, **458**, 2634
- Marchant, P., Langer, N., Podsiadlowski, P., Tauris, T. M., & Moriya, T. J. 2016, *A&A*, **588**, A50
- Meegan, C., Lichti, G., Bhat, P. N., et al. 2009, *ApJ*, **702**, 791
- Perna, R., Lazzati, D., & Giacomazzo, B. 2016, *ApJ*, **821**, L18
- Rodger, C., Brundell, J., Holzworth, R., & Lay, E. 2009, in *Coupling of thunderstorms and lightning discharges to near-Earth space: Proc. Workshop, Corte (France)*, *Am. Inst. Phys. Conf. Proc.*, **1118**, 15
- Savchenko, V., Ferrigno, C., Mereghetti, S., et al. 2016, *ApJ*, **820**, L36
- Szécsi, D., Bagoly, Z., Kóbori, J., Horváth, I., & Balázs, L. G. 2013, *A&A*, **557**, A8
- Szécsi, D., Langer, N., Yoon, S.-C., et al. 2015, *A&A*, **581**, A15
- The LIGO Scientific Collaboration, & the Virgo Collaboration 2016, *ArXiv e-prints* [[arXiv:1606.04856](https://arxiv.org/abs/1606.04856)]