Star-planet interactions

II. Is planet engulfment the origin of fast rotating red giants?

Giovanni Privitera^{1,2}, Georges Meynet¹, Patrick Eggenberger¹, Aline A. Vidotto^{1,3},
Eva Villaver⁴, and Michele Bianda²

¹ Geneva Observatory, University of Geneva, Chemin des Maillettes 51, 1290 Sauverny, Switzerland
e-mail: giovanni.privitera@unige.ch

² Istituto Ricerche Solari Locarno, via Patocchi, 6605 Locarno-Monti, Switzerland

³ School of Physics, Trinity College Dublin, Dublin 2, Ireland

⁴ Department of Theoretical Physics, Universidad Autonoma de Madrid, Modulo 8, 28049 Madrid, Spain

Received 21 April 2016 / Accepted 23 June 2016

ABSTRACT

Context. Fast rotating red giants in the upper part of the red giant branch have surface velocities that cannot be explained by single star evolution.

Aims. We check whether tides between a star and a planet followed by planet engulfment can indeed accelerate the surface rotation of red giants for a sufficiently long time to produce these fast rotating red giants.

Methods. We studied how the surface rotation velocity at the stellar surface evolves using rotating stellar models, accounting for the redistribution of the angular momentum inside the star by different transport mechanisms, the exchanges of angular momentum between the planet orbit and the star before the engulfment, and for the deposition of angular momentum inside the star at the engulfment. We considered different situations with masses of stars in the range between 1.5 and 2.5 $M_{\odot}$, masses of the planets between 1 and 15 M_J (Jupiter mass), and initial semimajor axis between 0.5 and 1.5 au. The metallicity Z for our stellar models is 0.02.

Results. We show that the surface velocities reached at the end of the orbital decay due to tidal forces and planet engulfment can be similar to values observed for fast rotating red giants. This surface velocity then decreases when the star evolves along the red giant branch but at a sufficiently slow pace to allowing stars to be detected with such a high velocity. More quantitatively, star-planet interaction can produce a rapid acceleration of the surface of the star, above values equal to 8 km s⁻¹, for periods lasting up to more than 30% the red giant branch phase. As found already by previous works, the changes of the surface carbon isotopic ratios produced by the dilution of the planetary material into the convective envelope is modest. The increase of the lithium abundance due to this effect might be much more important, however lithium may be affected by many different, still uncertain, processes. Thus any lithium measurement can hardly be taken as a support or argument against any star-planet interaction.

Conclusions. The acceleration of the stellar surface to rotation velocities above limits that depend on the surface gravity does appear at the moment to be the clearest signature of a star-planet interaction.

Key words. planetary systems – planet-star interactions – stars: rotation – stars: low-mass

1. Introduction

It is well known that when stars evolve into the red giant branch after the main-sequence phase, the very large expansion of the envelope imposes very low surface rotation velocities. Indeed, early survey of projected rotational velocities $v \sin i$ (Gray 1981, 1982) showed that giants cooler than about 5000 K are predominantly slow rotators and are characterized by $v \sin i$ of a few km s⁻¹ (see Fig. 7). However, there exists a small percentage of red giants that present much higher $v \sin i$ (Fekel & Balachandran 1993; Massarotti et al. 2008; Carlberg et al. 2011; Tayar et al. 2015).

To explain the high rotation rates of (apparently) single red giants, two kinds of scenarios have been proposed. The first scenario involves a mechanism occurring in the star itself. Simon & Drake (1989) proposed that, at the time of the first dredge-up, the surface could be accelerated by the transfer through convection of angular momentum from the central fast spinning regions to the surface. Fekel & Balachandran (1993) explained the high

rotation and high lithium abundance they observed in their sample of red giants as resulting from such a scenario. The dredge-up episode would bring to the surface not only angular momentum but also freshly synthesized lithium. It is interesting to underline here that this dredge-up scenario is expected to cause rapid rotation at a particular phase in giant stars evolution, namely when the first dredge-up occurs. In the large sample studied by Carlberg et al. (2011), a clustering of the rapid rotators at this phase (between T_{eff} equal to ~4500 and 5500 K or $\log T_{\text{eff}}$ between 3.732 and 3.740) is not seen. Moreover, we showed in Privitera et al. (2016) that the dredge-up actually produces no significant acceleration of the surface and thus cannot be a realistic reason for the high surface rotations that we are discussing here.

A second scenario proposed to explain the high rotation rates in giants involves the swallowing of planet (Peterson et al. 1983; Siess & Livio 1999a; Livio & Soker 2002; Massarotti et al. 2008; Carlberg et al. 2009). The phenomenon of planet and brown dwarf ingestion was studied theoretically by

Sandquist et al. (1998, 2002) for main-sequence (MS) stars and by Soker et al. (1984), Siess & Livio (1999a,b), and recently by Passy et al. (2012) and Staff et al. (2016) for giant stars. Siess & Livio (1999a,b) studied the accretion of a gaseous planet by a red giant and an asymptotic giant branch star (AGB). They considered cases where the planet is destroyed in the stellar envelope. They focused their study on possible consequences of this engulfment on the luminosity and surface composition of the star. Here we want to study the impact on the surface rotation of the star.

In the first paper of this series (Privitera et al. 2016), we followed the orbital evolution of a planet accounting for all the main effects impacting the orbit, that is changes of masses of the star and the planet, frictional and gravitational drags, and tidal forces, and computed the changes of the stellar rotation due to the planet orbital changes. We used rotating stellar models allowing us to follow the angular momentum transport inside the star in a consistent way. In the present paper, we study what happens when the orbital decay causes the planet to be engulfed by the star. More precisely we address the following questions:

- By how much can the surface rotation increase through a planet engulfment process?
- How long is the period during which a rapid surface velocity can be observed after an engulfment?
- Can the increase of the surface velocity trigger some internal mixing?
- Are there other signatures in addition to fast surface rotation that are linked to a planet engulfment event?

In Sect. 2, we discuss the physics included in our models. Section 3 presents post-engulfment evolution of various stellar models. Comparisons with observations are discussed in Sect. 4. Finally in Sect. 5 the main results are listed.

2. Ingredients of the models

2.1. The stellar models

The rotating stellar models are computed using the Geneva stellar evolution code (for a detailed description see Eggenberger et al. 2008). Ekström et al. (2012) provides the details about the ingredients of the models, such as nuclear reaction rates, opacities, mass loss rates, initial composition, overshooting, and diffusion coefficients for rotation, and Privitera et al. (2016) explain how the exchanges of angular momentum between the planetary orbit and the star are treated.

The present models allow us to make predictions of the evolution of the surface and the interior velocity resulting from the following processes: shear turbulence and meridional currents in radiative zones, convection, changes of the structure, loss of angular momentum by stellar winds, changes of angular momentum of the star resulting from tidal forces between the star and the planet, and the process of planet engulfment. In convective zones, solid body rotation is assumed, while radial differential rotation can develop in the radiative zone.

2.2. The planet model

Some knowledge of the structure of the planet is needed to determine where, once engulfed, the planet dissolves and deposits its angular momentum into the star. More precisely, we need to know two quantities (see Eq. (2) below): the planet radius, R_{pl} , and the mean molecular weight of the planet gas, μ_{pl} .

The radius of the planet is determined using the mass-radius relation by Zapolsky & Salpeter (1969),

$$R_{\text{pl}} \approx 0.105 \left(\frac{2x^{1/4}}{1+x^{1/2}} \right) [R_{\odot}], \quad (1)$$

where $x = M_{\text{pl}}/(3.2 \times 10^{-3})$ and M_{pl} is the planetary mass given in solar mass.

The same initial chemical composition is considered for the star and the planet. Here we consider a mass fraction of hydrogen $X = 0.707$, a mass fraction of helium $Y = 0.273$, and a mass fraction of the heavy elements $Z = 0.02$. With that composition, $\mu_{\text{pl}} = 0.614$. Below, we shall consider different compositions for the planet and star for what concerns carbon and lithium. These changes have however very limited impacts on the value of μ_{pl} and are neglected in the estimate of this quantity.

2.3. Physics of the engulfment

In Privitera et al. (2016), given a mass of a star and a mass of the planet, we studied the range of an initial semimajor axis that leads to a planet engulfment along the red giant branch. Now, we need some prescriptions for the fate of the planet inside the star to describe what happens next. According to the work by Soker et al. (1984), planets with masses inferior or equal to about $20 M_{\text{J}}$ are destroyed in the star. In the present work, we consider only planets with masses equal or below $15 M_{\text{J}}$, and therefore we assume that they are destroyed.

We need to know two quantities to be able to compute the impact on the surface velocity of this destruction process. The first quantity is the timescale for the destruction of the planet and the second is the location of this destruction inside the star. For instance, if the destruction occurs in a very short timescale (much shorter than the evolutionary timescale) then, once engulfment occurs, the whole orbital angular momentum of the planet orbit is given to the star in one shot. Moreover, if the destruction occurs in the convective envelope, then this angular momentum can be added in a very simple way to the convective envelope (see below).

2.3.1. Destruction timescale

Livio & Soker (1984) studied the evolution of star-planet systems during the red giant phase. They found that the engulfed planets with masses equal or below $10 M_{\text{J}}$ are destroyed after a few thousand years (see their Fig. 3). This is very short with respect to the evolutionary timescale during the red giant phase. Indeed, the duration of the first ascent of the red giant branch (i.e. before the ignition of helium burning in the core) is 180 Myr for a $1.7 M_{\odot}$ star. Therefore, we can consider that the angular momentum that remains in the orbit of the planet at the moment of the engulfment is delivered to the star in one shot.

2.3.2. Where is the angular momentum deposited?

The location of the planet dissolution (i.e. the dissolution point) depends on the physical mechanism that is responsible for this dissolution. We can consider two main mechanisms: thermal and mechanical destruction of the planet. Thermal dissolution is obtained where the virial temperature of the planet becomes lower than the local stellar temperature. The thermal kinetic energy of the stellar material is higher than the binding energy of the planet more deeply beyond this point.

Fig. 1. Kippenhahn diagram during the red giant phase. The central radiative zone and the convective envelope are represented by a white area and an area with black dots, respectively. The horizontal axis is the age of the star minus its main-sequence lifetime (t_{MS}). The four lines indicate the limit layers that can be reached by the planets of mass $1 M_J$ (red solid line), $5 M_J$ (blue dashed line), $10 M_J$ (green dash-dotted line), and $15 M_J$ (continuous cyan line). The vertical dashed lines labelled A, B, and C indicate when the engulfment occurs for the $15 M_J$ planet at an initial distance of 0.5 au, the $10 M_J$ planet at an initial distance of 1.0 au, and the $1 M_J$ planet at an initial distance of 1.5 au, respectively. All the other cases occur between the lines A and C. The *left panel* corresponds to a model star of $1.5 M_\odot$ and the *right panel* to a mass of $2 M_\odot$.

Knowing the internal structure of the star, it is possible to compare the local temperature in the star and the virial temperature of the planet (Siess & Livio 1999a) as follows:

$$T_{\text{v,pl}} \sim \frac{GM_{\text{pl}}\mu_{\text{pl}}m_{\text{H}}}{kR_{\text{pl}}} \sim 10^5 \mu_{\text{pl}} \frac{M_{\text{pl}}}{M_J} \frac{R_J}{R_{\text{pl}}} [\text{K}], \quad (2)$$

where k is the Boltzmann constant and M_J , R_J are the mass and radius of Jupiter.

Figure 1 indicates the regions where the stellar temperature is equal to the virial temperature of planets of various masses. Also, isothermal lines¹ are indicated in Fig. 1. We also indicated by vertical dashed lines the engulfment time of planets of various masses and initial semimajor axes as obtained in Privitera et al. (2016). As indicated in Sect. 2.3.1, the migration time of the planet is very short, thus the destruction of those planets occurs at the intersection of the isothermal line for the planet mass considered with the corresponding vertical lines indicating the time of engulfment (only a few cases are shown for illustration). A few interesting points can be noted from Fig. 1:

- When the star evolves, the dissolution points reach (in general) deeper layers in the stellar interior. This is a consequence of the expansion of the envelope during the red giant phase, which produces a lowering of the temperature at a given Lagrangian mass. The dissolution point only shifts for a time outwards when the star contracts, for example during the bump. This explains the local minimum that can be seen for instance for the $5 M_J$ mass planet at a time around 0.08 Gyr for the $2 M_\odot$ model (right panel of Fig. 1).
- More massive planets go deeper inside the star. This is of course expected, since a more massive planet requires more extreme conditions to be destroyed than lighter planets.
- Under the assumptions above, and in the light of Fig. 1, it is reasonable to assume that the planets are destroyed in the convective envelope of the star.

Siess & Livio (1999a) also examined the possibility that when the planet gets closer to the stellar core, tidal effects induce strong distortions of the planet that can destroy it. Using the elongation stress at the centre of the planet as approximated by Soker et al. (1987), we find in most cases that the planet would be destroyed in the convective envelope as is the case when the criterion based on the virial temperature is used. The planet would only be destroyed by tides just below the convective envelope in a few cases. In the following we consider only the criterion based on the virial temperature and thus assume that all our engulfed planets deliver their angular momentum in the convective envelope.

We call $\mathcal{L}_{\text{pl}}$ the angular momentum of the planet orbit at the time of engulfment and $\mathcal{L}_\star(\text{ce})$ the angular momentum of the external convective envelope (ce) of the star just before engulfment. To obtain the new angular velocity of the envelope after engulfment, $\bar{\Omega}$, we write

$$\bar{\Omega}(\text{ce}) = \Omega(\text{ce}) \left(1 + \frac{\mathcal{L}_{\text{pl}}}{\mathcal{L}_\star(\text{ce})} \right), \quad (3)$$

where $\Omega(\text{ce})$ is the angular velocity of the convective envelope just before the engulfment.

Equation (3) assumes that the moment of inertia of the convective envelope is not changed by the planet engulfment process. Actually the engulfment process might add thermal energy in the upper layers of the stellar envelope, making these layers expand for a while. But the excess of energy is rapidly radiated away and thus the star rapidly evolves back to its initial state. Just as a numerical example, in cases in which all the kinetic energy in the planetary orbit of a $15 M_J$ initially orbiting a $1.7 M_\odot$ at 0.5 au would be added as an increase of the internal energy in the outer layers, then the excess of energy would be radiated away in about 55 yr, thereby in a very short time compared to the evolutionary timescale.

2.4. Initial conditions considered

We consider stars with initial masses in the range between 1.5 and $2.5 M_\odot$, with a metallicity $Z = 0.02$ and an initial rotation equal to $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1$ and 0.5 , where Ω_{ini} is the initial angular velocity and Ω_{crit} the critical angular velocity on the zero

¹ These lines would not be isothermal lines in the case in which the structure of the planet were to change during its journey inside the star. Here it is assumed that during the very short migration time the planet structure does not change significantly. The only change occurs at the very end, when the planet dissolves.

age main sequence (ZAMS). These correspond to initial surface velocities of ~ 30 and 160 km s^{-1} , respectively.

As explained in Privitera et al. (2016), we focused here on stellar masses that are larger or equal to $1.5 M_{\odot}$ because stars in this mass range do not have a sufficiently extended outer convective zone to activate a dynamo during the main sequence, so that unless they host a fossile magnetic field, they do not undergo any significant surface magnetic braking. The results obtained in the present work are thus not sensitive to the uncertainties related to the modelling of this braking during the main sequence. In contrast, lower initial mass stars have an extended outer convective zone during the main sequence. Therefore they can activate a dynamo and suffer strong magnetic braking. The evolution of the rotational properties of these lower initial mass stars is then different and will be the topic of another paper in this series (Privitera et al. 2016).

Planets with masses equal to 1, 5, 10, and $15 M_J$ were considered². The initial semimajor axes (a_0) were taken as equal to 0.5, 1, and 1.5 au and the eccentricities of the orbits are equal to 0. In Privitera et al. (2016), the evolution of the planetary orbit and the star were computed in a consistent way up to the point of engulfment. The end of the computation in Privitera et al. (2016) represents the initial conditions for the present work (see Table A.1).

3. Impact of planet engulfment on surface velocities of red giants

The impact of tides and engulfment on the surface velocity of a red giant depends at least on the following parameters: the mass of the planet, its initial distance to the star, and the mass of the star. Other parameters such as metallicity (which is not discussed here) and rotation (which is discussed here) of the star also affect the results.

Figure 2 shows the evolution of the surface equatorial velocities for the present stellar models with and without engulfment of a planet. Tables A.2 and A.3 present some characteristics of the star after an engulfment.

3.1. Planet engulfment by a $1.5 M_{\odot}$ star

We begin by discussing the case of the $1.5 M_{\odot}$ stellar model (upper left diagram in Fig. 2). The case of an engulfment of a $15 M_J$ initially at a distance of 0.5 au is considered for the purpose of the discussion. We note several features.

Planet engulfment may have a very strong effect on the surface rotation of a red giant. Compare for instance in the upper left panel, the (black) dotted line that shows the evolution of the surface velocity of a $1.5 M_{\odot}$ red giant that began its evolution with an angular velocity equal to 50% of the critical velocity (142 km s^{-1}) and with no planet engulfment, with the (blue) dashed line showing the evolution of the surface velocity of a similar star engulfing a planet. While the isolated star has a surface velocity that remains below 7 km s^{-1} for $\log g$ below 2.8, the star that engulfed the planet shows an extremely rapid increase of the surface velocity up to a maximum value of about 40 km s^{-1} .

² In our models, we take planets with masses that are not larger than $15 M_J$, which means below the limiting value $M_{\text{cri}} \approx 20 M_J$ estimated by Soker et al. (1984), above which the planet is no longer completely destroyed when engulfed by the stellar envelope. This limit is also consistent with the value found in the numerical simulations by Staff et al. (2016).

The left panel of Fig. 3 shows the variation of the angular velocity inside the $1.5 M_{\odot}$ with an initial angular velocity equal to 10% of the critical velocity just before and just after the engulfment of a planet. The angular velocity in the convective zone extending from a mass equal to about $0.3 M_{\odot}$ up to the surface has its angular velocity increased by about 1.7 dex (a factor 50). At engulfment (around a $\log g$ equal to 2.5), the ratio between the angular velocity of the core and that of the envelope decreases from a value of about 32 000 to a value equal to 1000 (see the right panel of Fig. 3). Then the contrast increases again as a result of the expansion of the envelope and contraction of the core.

The present models cannot account for the small contrast of about one dex between the core and envelope rotation rates that have been obtained by asteroseismology for red giants at the base of the giant branch (Beck et al. 2012; Mosser et al. 2012; Deheuvels et al. 2012, 2014). In that respect, an additional internal angular momentum transport process is missing in the radiative interior of the present models (Eggenberger et al. 2012; Marques et al. 2013; Cantiello et al. 2014). This weakness has however no major impact on the process studied here. The missing transport mechanism would add a fraction of the angular momentum of the core to the envelope, but this does not accelerate the envelope much because the angular momentum in the core is quite small with respect to that contained in the envelope. Therefore the angular momentum coming from the planetary orbit remains in any case that is much larger than the angular momentum of the envelope, and the final result of a planet engulfment would remain nearly unchanged.

After the engulfment, the surface velocity decreases because the stellar envelope continues to expand, but it remains above 8 km s^{-1} for a time, i.e. nearly 36 Myr (see Table A.3), which means for about 15% of the total duration of the red giant branch phase. No significant changes of the structure of the star (i.e. change of the extent of the convective envelope and/or of the mixing of the chemical elements) is produced by the increase of the rotational speed of the convective envelope. So in our models, any changes of the surface abundances can only be attributed to the dilution of the planet material into the convective envelope. This is briefly investigated in the next section when comparisons with observations are discussed.

In Fig. 2 we also indicated the maximum value for the surface rotation that is expected during the red giant phase (see the black long-dashed line) for a star with no planet. This line has been obtained assuming that the star rotates near the critical velocity on the ZAMS and that a solid body rotation is maintained at every time. This is clearly an extremum. This case of solid body represents an extremum case in terms of efficiency of angular momentum transport and also in terms of surface velocities³. This means that any observed star with initial masses of about $1.5 M_{\odot}$ and with a surface velocity that is higher than that given by the curve V_{max} in the left upper panel of Fig. 2 needs some acceleration process, which might come either from tidal forces or from an engulfment. We see that for a $1.5 M_{\odot}$ star at a metallicity $Z = 0.02$, this maximum velocity is well below

³ One could argue that, once the solid body rotation is reached, some processes might still advect angular momentum from the centre to the surface producing a negative gradient of Ω , i.e. a core rotating slower than the envelope, and increasing the surface velocity. However, this is not supported by asteroseismic determinations of the interior rotation of red giants (e.g. Beck et al. 2012; Mosser et al. 2012; Deheuvels et al. 2012, 2014), which show that red giants have a clear increase of rotation towards the central regions. In that respect, the hypothesis of solid body rotation gives a very conservative upper limit for the surface rotation of red giants.

Fig. 2. Evolution of the surface velocity as a function of the logarithm of the surface gravity. Dotted (black) lines are for stars without engulfment. Cases for $\Omega/\Omega_{\text{ini}} = 0.1$ and 0.5 are shown. The black long-dashed line (labelled V_{max}) corresponds to solid body rotation without engulfment starting with an initial rotation near the critical velocity (see text). Continuous lines correspond to cases with planet engulfment for stars with an initial velocity equal to 10% the critical value and the short-dashed lines correspond to the same cases but for an initial rotation equal to 50% of the critical value. The different colours for solid and dashed lines are for planet masses of 5 (red), 10 (magenta), and 15 M_J (blue) and initial orbital distances of 0.5, 1, and 1.5 au from left to right (see labels). The different panels are for different values of the mass of the star. In the case of $2.5 M_{\odot}$, the evolution is pursued until the end of the core He-burning phase, while for lower initial mass stars, the evolution is stopped at the tip of the red giant branch.

the 40 km s^{-1} reached by the engulfment of a $15 M_J$ planet with an initial distance to the star equal to 0.5 au. Therefore planet engulfment may be a sufficiently strong mechanism to produce surface acceleration beyond any reasonable process that can occur in single stars. Of course not just any planet engulfment will produce such strong acceleration, but at least some events can produce surface velocities beyond what can be reasonably explained by single star evolution.

We see that decreasing the initial distance between the star and the planet produces higher surface velocities and longer periods during which high surface rotations can be observed. The main reason for this is that the smaller the initial distance, the engulfment occurs earlier along the red giant branch. The earlier the engulfment occurs, the smaller the moment of inertia of the convective envelope, and the effect produced by the injection of the planetary angular momentum in the stellar convective

Fig. 3. *Left panel:* variation of the angular velocity as a function of the Lagrangian mass inside a $1.5 M_{\odot}$ model along the red giant branch just before (blue continuous line) and after the engulfment (red dashed line) of a $15 M_J$ mass planet at an initial distance of 0.5 au from the star. *Right panel:* evolution of the ratio between the angular velocity of the core and surface angular velocity as a function of the surface gravity along the red giant branch for the same models as in the left panel.

envelope is thus more important. We could argue however that the larger the distance, the larger the amount of angular momentum in the planet orbit. However the effect indicated above (the small moment of inertia of the envelope) is the most important and clearly overcomes the effect of the larger angular momentum associated with a larger orbital radius. An interesting consequence of that point was noted already by [Carlberg et al. \(2009, 2011\)](#); planet engulfment produces the largest increases of the surface rotation when the engulfment occurs at the beginning of the red giant branch.

We see also that the higher the mass of the planet, the higher the maximum surface velocity reached (assuming identical initial distance of the planet to the star) and longer the period during which the surface velocity is superior to a given limit (see Tables A.2 and A.3).

If keeping all the other parameters equal, we change the initial rotation of the star, passing from $\Omega_{ini}/\Omega_{crit} = 0.1$ to 0.5, we obtain very similar behaviors. This indicates that the outcome at least in the case of the $1.5 M_{\odot}$ does not depend much on the initial rotation of the star.

3.2. Planet engulfment by stars with masses between 1.7 and $2.5 M_{\odot}$

In Fig. 4, the increase of the surface velocity due to a planet engulfment is compared for stars with different initial masses. We see that in general, for a given planet mass and a given initial distance between the star and the planet, the engulfment occurs at a higher surface gravity when the mass of the star decreases. The maximum surface velocities that are reached are higher for lower initial mass stars. We added a track for a one solar mass model to provide a clear illustration of this. As explained in Sect. 2, such a star suffers magnetic braking during the main-sequence phase. This is introduced for the one solar mass model using a solar-calibrated value for the efficiency of this braking. As a result, the surface velocity at the base of the red-giant branch is much lower for the one solar mass model than for stars with higher initial masses. When the engulfment occurs, the acceleration is stronger for this model than for the more massive stars because of the larger ratio between the planetary orbital angular momentum and the stellar envelope angular momentum. Since the $2.5 M_{\odot}$ star does not evolve through the He flash, we pursued the evolution during the core He-burning phase. This is why

Fig. 4. Evolution of the surface velocity of various initial mass stellar models during the red giant phase as a function of the surface gravity. The initial rotations are between 30 and 50 km s^{-1} . Planets have a mass equal to $15 M_J$ and the orbits have an initial semimajor axis equal to 0.5 au.

its evolution looks so different from the lower initial mass stars (compare the lower right panel of Fig. 2 with the other panels). Actually, just after the engulfment, we can see a very similar behaviour as those we could see in the other panels. The surface velocity increases nearly vertically at engulfment and after engulfment decreases with surface gravity. When the tip of the red giant branch is reached, the star contracts and the track evolves again at high surface gravities, following first the same path as that used when the star was evolving up the red giant branch, passing again through the same point where engulfment occurred, and evolving to higher surface velocities still when it continues to contract. Most of the core He-burning phase occurs along the extremity of the loop at surface gravities between 2.7 and 3.0. After the core He-burning phase, the envelope of the star expands, the surface gravity decreases, and the star joins the asymptotic giant branch.

Interestingly, the surface velocity during the core He-burning phase remains quite high after an engulfment (above the curve V_{max} in the lower right panel of Fig. 2). Therefore, the signature of planet engulfment may remain visible for the whole core He-burning phase. This would likely be the case for lower initial mass stars too that are evolving through a He flash, although this remains to be confirmed by computations.

In general, as was the case for the $1.5 M_{\odot}$ star, changing the initial rotation rate of the star does not have a very large impact on the results. We note, however, the very thin loop at engulfment in the lower left panel of Fig. 2 in the case of $2 M_{\odot}$ models with an initial rotation equal to 50% the critical angular velocity and with a $15 M_J$ at initial distance of 0.5 au. For this model, the time of engulfment coincides with the time when the bump occurs. During the bump, the star contracts (hence the increase of the surface gravity seen on Fig. 2). This contraction occurs when the H shell enters the domain whose composition has been enriched in hydrogen by the downwards extension of the external convective zone. As a consequence, the H-burning shell expands and reduces its temperature. By a mirror effect, the envelope contracts provoking an additional increase of the surface velocity and the thin loop that appears at the end of the engulfment process.

3.3. How long does a red giant star remain fast rotating after an engulfment?

In the literature, a value of $v \sin i$ higher than 8 km s^{-1} is sometimes used as a criterion to qualify a red giant as fast rotating (see e.g. Carlberg et al. 2012). We see on Fig. 2 that indeed higher values than 8 km s^{-1} would definitively require some interaction with a second body for surface gravities that are smaller than $\log g$ equal 2.0 ($1.5 M_{\odot}$), 2.1 ($1.7 M_{\odot}$), 2.15 ($2 M_{\odot}$), and 2.1 ($2.5 M_{\odot}$). Of course, this does not exclude that stars showing surface velocities below this limit have not undergone any interaction, but in that case, alternative single star models could be proposed to explain the high velocity as well.

Figure 5 shows a synthetic view of the durations of the high surface velocity periods. We see that, for a given stellar model, the duration increases when the initial semimajor axis decreases and the mass of the planet increases (see also Tables A.2 and A.3). The conditions for, let us say, having a high surface velocity during about 10% of the red giant phase are less restricted when the mass of the star passes from 1.5 to $2 M_{\odot}$ ⁴. The same occurs when, keeping the stellar mass constant; the initial rotation passes from 10 to 50% of the critical velocity, although the impact is less important than changing the initial mass of the star.

3.4. Impact of tidal forces with respect to engulfment

The physics of what happens during the engulfment is not so well known and thus we may wonder about the reliability of the present results. An interesting point to mention in that context is the fact that the tidal forces may already lead to significant accelerations of the stellar surface. Figure 6 shows the evolution of the surface rotation as a function of the luminosity for a few selected models of $2 M_{\odot}$ stars. The small dots along the curves with engulfment indicate the surface velocity reached just before engulfment. The acceleration up to this point is only due to tidal forces. The acceleration after that point is due to the engulfment process itself.

We see that tides alone may be sufficient to accelerate the surface rotation of the star significantly. Since the acceleration due to tides is very rapid (see Privitera et al. 2016), as is the acceleration due to the engulfment, there is no chance that using the

rotational velocity of the star to distinguish observationally between tides- and engulfment-acceleration processes, would both occur.

4. Comparisons with observations

The observed sample of Carlberg et al. (2011) (1287 stars) is shown in Fig. 7 superposed on the evolution of the surface velocities for different $2.5 M_{\odot}$ models as computed in the present work. A significant number of stars are rotating much faster than what is predicted by these models. An extreme case is the star Tyc5854-011526-1 that has a $v \sin i$ of 84 km s^{-1} , which corresponds to two-thirds of the critical velocity of a $2.5 M_{\odot}$ at the considered effective temperature. In the remainder of this section, we compare our theoretical predictions with the observations of red giants performed by Carlberg et al. (2012). We choose this sample since it gives surface velocities and surface gravities for a significant number of red giants (91 stars). Moreover, it also provides indications for some stars on the $^{12}\text{C}/^{13}\text{C}$ ratio at the surface and for lithium abundance. This allows us to check whether stars that are believed to have acquired their high surface rotation as a result of planet engulfment also show some special features in their surface composition.

We do not perform a very detailed analysis since that analysis was already carried out in Carlberg et al. (2012), but we want to address the following questions:

- Are there among the observed stars, cases that cannot be explained by single star models?
- Can the observed surface velocities of such cases be reproduced by invoking a planet engulfment?
- If yes, is it possible to deduce the initial conditions required to explain these systems by a planet engulfment?
- Is there any chance of seeing some signatures of the planet engulfment in the surface composition?

Before investigating these points based on the present computations, we must see whether the metallicities and mass range of our stellar models are adequate for a comparison with the sample of Carlberg et al. (2012). For what concerns the metallicity, the correspondence between the metallicity of the observed sample and metallicity of the models is marginal. The $[\text{Fe}/\text{H}]$ of the sample studied by Carlberg et al. (2012) is on average lower than solar, while the present computations were performed with a heavy mass fraction of $Z = 0.020$ which is over solar ($Z_{\odot} = 0.014$). However, whatever the metallicity, the engulfment would produce a strong acceleration of the surface. It may be that a given surface velocity would be reached with slightly different initial conditions, but the main outcomes are not changed.

In the left panel of Fig. 8, the observed sample is shown together with the tracks of the single (no engulfment) stellar models computed in the present work (case with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1$). In case we had made our computations of stellar models with a lower initial metallicity, the tracks would have been slightly shifted to the left, i.e. to the blue. Keeping this in mind, we see that the range of masses followed here, would more or less go through the averaged observed positions. It may be that some stars have lower initial mass than $1.5 M_{\odot}$ and some may have higher initial mass than $2.5 M_{\odot}$, but on the whole the mass range of the models is relatively well representative of most of the masses of the sample.

⁴ The duration of the period during which the surface velocity remains superior to a given limit after an engulfment is larger in lower mass stars. However, when this period is normalized to the total duration of the red giant branch, then the fraction spent above a given limit decreases when the mass decreases.

Fig. 5. Fraction of the total red giant branch duration spent with a surface equatorial rotation velocity that is higher than 8 km s^{-1} after an engulfment. *Upper left panel:* case of a $1.5 M_{\odot}$ with an initial rotation $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1$. *Upper right panel:* same as the *upper left panel*, but with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$. *Lower left panel:* same as the *upper left panel*, but for a star of mass $2.0 M_{\odot}$. *Lower right panel:* same as the *lower left panel*, but with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$.

Fig. 6. Evolution of the surface velocity of a $2 M_{\odot}$ stellar model during the red giant phase as a function of the luminosity. The initial rotation of the star is 10% of the critical velocity (on the ZAMS). When evolution proceeds, the luminosity in general increases except at the bump visible in the left panel under the form of a small hook. Magenta lines indicate cases without planet engulfment, while blue, green, red, and cyan lines correspond to the engulfment of a planet with a mass equal to 1, 5, 10, and $15 M_J$, respectively. The surface velocity reached just before engulfment, i.e. resulting only from tidal forces, is indicated by a small dot. *Left panel:* the initial semimajor axis is equal to 0.5 au. *Right panel:* the initial semimajor axis is equal to 1.5 au.

4.1. Stars with a clear signature of a past interaction

In many works, a red giant is considered as fast rotating and thus as a possible candidate for having gone through a planet engulfment process if its $v \sin i$ is larger than 8 km s^{-1} . This criterion is however too schematic. To understand this point, we point out

the right panel of Fig. 8. We see that, actually, many of the red-filled points ($v \sin i > 8 \text{ km s}^{-1}$) are not very far from the line corresponding to the track for the $1.5 M_{\odot}$ with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$, allowing such stars to be possibly explained without invoking any interaction with a companion. On the contrary, much lower

surface velocities can be the indication of a planet engulfment. This can be seen looking at the magenta-dotted line, which is in the lower right corner in the right panel of Fig. 8. The tracks with no engulfment pass well below the slowly rotating points observed with surface gravities below about 1.6, while the track with an engulfment of a one Jupiter mass planet initially at a distance of 0.5 au of a $1.5 M_{\odot}$ star with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$ would pass through at least part of these observed points. Thus, the surface rotation above which some interaction may be needed depends strongly on the surface gravity and cannot be given by only one number.

How can we distinguish stars that would nearly certainly require some interaction as the cause for their high surface velocities? The continuous (blue) line shows the evolution of the surface velocity for a $2.5 M_{\odot}$ that would begin its evolution at the critical velocity and evolves as a solid body rotating model. This evolution, as explained in the previous section, represents an extremum in terms of efficiency of the internal angular momentum transport and surface rotation. Actually, this is upper limit is very generous because we know that red giants are not rotating as solid body.

Turning now to stars that are found above this limit, we can be quite confident that, for these cases, an interaction with a companion must have occurred. These stars are shown as circled blue magenta filled points and individually labelled with a capital letter. Hence, this answers our first question above. We see that stars are indeed observed with surface velocities that cannot be explained by single star evolution⁵.

As a final remark, as already stated in the introduction, the bulk of stars with $v \sin i$ lower than 8 km s^{-1} are in between the single star evolutionary tracks with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1$ and $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$. This shows that single star models can well account for the surface rotation of the bulk of red giants.

4.2. Can planet engulfment explain the cases likely resulting from an interaction?

Can planet engulfment reproduce the surface velocities observed for those red giants that have such large surface rotations that no single star models can reproduce them (see the points labelled by upper-case letters A to F in Figs. 9 and 8)? The answer is yes, as can be seen by comparing the locations of these stars in the right panel of Fig. 8 with the black continuous tracks. Only a few cases are shown for purpose of illustration, but there is no doubt that playing with the initial conditions (mass of the star, mass of the planet, and initial distance of the planet), it is indeed possible to produce a sufficient surface acceleration to reach these observed surface velocities for a sufficiently long duration to allow these high surface velocities to be observed.

4.3. Can the initial conditions be deduced from observations?

Can we determine the initial conditions needed to reproduce these systems? The solutions are unfortunately not unique. For instance, the positions of the points E and F could be reproduced by the engulfment of typically a $10 M_J$ planet initially at a distance of 0.5 au of a $2.5 M_{\odot}$ with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$, or by the engulfment of a $10 M_J$ planet initially at a distance of 0.5 au from a

Fig. 7. Surface velocities $v \sin i$ as a function of the effective temperature for the sample of red giants observed by Carlberg et al. (2011). The red dots correspond to stars with $v \sin i$ larger than 8 km s^{-1} . The lines indicate the evolutionary tracks for $2.5 M_{\odot}$ stellar models for different initial rotations (from bottom to top, the time-averaged velocities during the MS phases are ~ 20 , 100 , and 280 km s^{-1}). The upper dotted line shows the critical velocity.

$1.5 M_{\odot}$ with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$, and there are likely other solutions. Of course, additional information can provide some further constraints. For instance, from the position in the HR diagram of the E and F stars (see left panel), the solution with the $2.5 M_{\odot}$ would be favoured with respect to that involving a $1.5 M_{\odot}$. In that case the star would be likely in the clump and not along the red giant branch.

4.4. The surface compositions after an engulfment

The left panel of Fig. 9 shows isotopic ratios at the surface of the observed stars and predictions of the theoretical models. The continuous lines show the predictions of the present stellar models without any planet engulfment. The predictions depend on the initial rotation. Higher initial rotations lead to lower $^{12}\text{C}/^{13}\text{C}$ ratios. Models with an initial rotation equal to 50% of the critical value go roughly through the middle of the observed points⁶.

We can deduce the following points looking at the left panel of Fig. 9:

- The planet engulfment process decreases the shear between the core and the convective envelope and thus weakens the shear mixing in that region. However this effect is negligible because even in models without engulfment, which have a stronger shear in that region, we see no effect. Indeed, such an effect would produce a continuous decrease of the $^{12}\text{C}/^{13}\text{C}$ ratio once the convective envelope has reached its deepest point. This is obviously not the case as shown in the left panel of Fig. 9.

⁵ We may wonder, however, whether a more massive star (like a $3 M_{\odot}$) could explain these points. At least this does not appear to be the case for point A which has, according to its position in the theoretical HR diagram, an initial mass clearly below $2.5 M_{\odot}$.

⁶ The present models do not account for thermohaline mixing for instance, which has been proposed by Charbonnel & Zahn (2007) as causing an additional decrease of the $^{12}\text{C}/^{13}\text{C}$ ratio and of the lithium along the red giant branch (see also Charbonnel & Lagarde 2010).

Fig. 8. *Left panel:* surface gravities vs. effective temperatures. The continuous lines correspond to evolutionary tracks for single star models (no engulfment) for (from bottom to top) $1.5 M_{\odot}$ (red), $1.7 M_{\odot}$ (magenta), $2.0 M_{\odot}$ (black), and $2.5 M_{\odot}$ (blue), starting with an initial angular velocity equal to 50% of the critical velocity. The dots show the observations by Carlberg et al. (2012), the black circles are for stars with $v \sin i < 8 \text{ km s}^{-1}$, the filled red and circled blue magenta points are for stars with $v \sin i > 8 \text{ km s}^{-1}$. The velocities of the circled blue magenta points cannot be explained by any reasonable model for single stars. The high rotation of these stars results with great probability from an interaction with an additional body. These points are labelled by letters (see text). The filled blue triangles are observations by Adamów et al. (2014), the filled blue square is the Li-rich star BD+48 740 discussed by Adamów et al. (2012a), and the blue star is the super Li-rich giant HD 107028 studied by Adamów et al. (2015). The two empty squares correspond to fast rotators obtained by Tayar et al. (2015) for stars with masses equal to 1.43 and $2.07 M_{\odot}$ along the red giant branch. *Right panel:* surface equatorial rotation vs. surface gravities for $1.5 M_{\odot}$ models. The two red tracks correspond to no planet engulfment cases starting with different initial rotations. The black curves show the evolution for stars with planet engulfment. The dots are the same observations as those shown in the left panel.

Fig. 9. Comparison between surface abundances and theoretical predictions. The points are observations (see caption of Fig. 8 for the meaning of the different symbols). *Left panel:* surface isotopic ratios $^{12}\text{C}/^{13}\text{C}$ vs. surface gravities. The continuous lines are the same evolutionary tracks as those shown in the left panel of Fig. 8. The dotted lines show the new isotopic ratios obtained when account is made for the dilution in the convective envelope of the planet material (see text for more details) starting from the models with $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$. *Right panel:* surface lithium abundance vs. surface gravities. The continuous (black) horizontal line shows the averaged lithium abundance observed at the surface of stars with $v \sin i$ smaller than 8 km s^{-1} . The dotted lines show the results of dilution of the planet material into the convective envelope.

- Another process can change the isotopic ratio: the dilution of the planet material into the convective envelope. To estimate the importance of this process, we followed the same line of reasoning as presented in [Carlberg et al. \(2012\)](#). We applied their Eq. (3), which gives the new isotopic ratio, $(^{12}\text{C}/^{13}\text{C})_{\text{new}}$ accounting for the dilution of the planetary material into the convective envelope

$$(^{12}\text{C}/^{13}\text{C})_{\text{new}} = \frac{10^{A(\text{C})_p} \frac{r_p q_e}{1+r_p} + 10^{A(\text{C})_*} \frac{r_*}{1+r_*}}{10^{A(\text{C})_p} \frac{q_e}{1+r_p} + 10^{A(\text{C})_*} \frac{1}{1+r_*}},$$

where $A(\text{C})_p$ is $\text{Log}(N(\text{C})/N(\text{H})) + 12$ in the planet and $A(\text{C})_*$ is the same quantity for the star ($N(\text{C})$ is the number of carbon per unit volume accounting for the two isotopes ^{12}C and ^{13}C). The ratio r is equal to $N(^{12}\text{C})/N(^{13}\text{C})$ and q_e is the ratio of the mass of the planet to the mass of the convective envelope. We took the values directly from our stellar models for the quantities concerning the star (*). We considered the same values for the planet as in [Carlberg et al. \(2012\)](#) namely $A(\text{C})_p = 8.87$ dex (about three times the solar carbon abundance as determined by [Wong et al. 2004](#)) and $r_p = 89$, which is a standard value for the solar system ([Lodders & Fegley 1998](#)). Using the above formula, we obtain after engulfment the level of $^{12}\text{C}/^{13}\text{C}$ shown by the dashed segments in Fig. 9. We see that since the planetary material has a higher $^{12}\text{C}/^{13}\text{C}$ ratio than the stellar envelope; this effect makes the surface isotopic ratio a bit larger. The effect is however small and thus the measure of this isotopic ratio appears as a poor indicator of a planet engulfment, especially that changes of the initial mass, rotation, and likely other possible mixing processes have stronger effects than the engulfment.

- Looking only at the observed points, we see that the fast rotators and those stars whose surface velocities cannot be explained by single star evolution do not present different carbon isotopic ratios than the rest of the sample, confirming that this isotopic ratio does not appear to be very sensitive to the engulfment.

In the right panel of Fig. 9, observed lithium abundances are shown. It seems as indicated by [Carlberg et al. \(2012\)](#), that the fast rotators present on average higher lithium abundances. However, if we consider the Li-rich giants observed by [Adamów et al. \(2012a, 2014, 2015\)](#), none of these stars have a $v \sin i > 8 \text{ km s}^{-1}$ (see the right panel of Fig. 8), which somewhat blurs the trend.

Similar estimates as that for the carbon isotopic ratio can be performed for the lithium abundance. Using Eq. (2) of [Carlberg et al. \(2012\)](#)

$$A(\text{Li})_{\text{new}} = \text{Log}(q_e 10^{A(\text{Li})_p} + 10^{A(\text{Li})_*}) - \text{Log}(1 + q_e),$$

we estimate the new surface lithium abundance, $A(\text{Li})_{\text{new}}$, after accretion of the planetary material. In our stellar models, the abundance of lithium is not followed explicitly, thus we did as in [Carlberg et al. \(2012\)](#), considering for $A(\text{Li})_*$ a value equal to -0.18 dex corresponding to the averaged abundance of lithium observed at the surface of slow rotators; see the horizontal continuous line in the right panel of Fig. 9. For $A(\text{Li})_p$, we took a value equal to 3.3 ([Lodders & Fegley 1998](#)), while for q_e , we used the stellar models quantities. The results are shown by the dotted tracks in the right panel of Fig. 9. The lines represent the surface lithium abundances that would be observed in case the engulfment occurred at various surface gravities. When the outer convective zone begins to appear, the convective envelope is very small and the Li abundance that results from a

planet dilution is very high. This makes the vertical part of the track. Then when the envelope increases in mass, the evolution goes to lower Li abundances and to lower gravities.

From the right panel of Fig. 9, we see that indeed the dilution of the planetary material into the convective envelope can have a strong effect on the surface Li abundance. The conditions at the base of the convective of our stellar models are not favourable to a rapid destruction of that element by proton captures, thus it is burned unless some mixing occurs below the convective envelope (see also [Aguilera-Gómez et al. 2016](#)).

The strong effect on lithium abundance found here after the planet engulfment confirms what was suggested by many authors in the past (see e.g. [Alexander 1967](#); [Siess & Livio 1999a](#); [Carlberg et al. 2012](#); [Adamów et al. 2012b](#)), namely that the planet engulfment process can indeed produce lithium-rich red giants. However, the lithium signature of planet engulfment remains difficult to disentangle from other processes. Lithium is a fragile element that is destroyed at about 2.6 million degrees, so any surface enrichment might disappear rapidly in some stellar models. Lithium can also be produced in some red giant models, which completely blurs the picture for what concerns the origin of a high lithium abundance.

In that respect it is interesting to see the lithium surface abundances of those stars whose high surface velocities cannot be explained by single star models (see the circled blue magenta filled points in Fig. 9). Their positions are not very peculiar in the sense that slowly rotating stars are also found with similar levels of Li abundances. All these stars, however, are above the averaged level for the (apparently) slowly rotating stars, leaving open the possibility that they could have received some lithium from a planet.

On the other hand, as noted already above, none of the Li-rich giants observed by [Adamów et al. \(2012a, 2014, 2015\)](#) are apparently fast rotators. These kinds of stars are therefore difficult to be explained by a planet engulfment process unless some mechanism, not accounted for in the present models, such as a strong wind magnetic braking for instance, has occurred. In that case those red giants would have once been fast rotators, but the wind magnetic braking was efficient enough for the star to have lost memory of its post-engulfment fast rotating stage.

5. Perspectives and conclusions

As was already shown in [Privitera et al. \(2016\)](#), there are observed red giant stars whose high surface velocities cannot be explained by single star evolution. The upper velocities that can be reached by single star models of the considered mass are indicated as a function of the surface gravity in Fig. 2 by the long-dashed black line labelled V_{max} . The use of such gravity dependent limits to isolate red giants that have likely engulfed a planet, or at least interacted with, it is probably more realistic than the commonly used limit of 8 km s^{-1} .

The present models show that tidal interactions followed by planet engulfment can reproduce the high surface velocities of these stars during sufficiently long periods that allow these high rotations to be observable. Surface velocities beyond the upper limit allowed by single star evolution can be reached already just by tidal interaction.

We obtain also that the high rotation obtained after the engulfment is maintained beyond the end of the red giant branch for the $2.5 M_{\odot}$. It would be extremely interesting to check whether such a conclusion would hold for our lower initial mass models that go through a He-flash episode.

Conditions also exist for producing engulfments with much less spectacular impacts. The engulfment of a one M_J planet by a $1.5 M_\odot$ star would produce surface velocities of only a few km per seconds at gravities below about 1.6. This is quite small and these stars would hardly be recognized as having engulfed a planet.

The above conclusions make the link between observed fast rotating red giants and red giants that have engulfed a planet much less clear in the sense that not all fast rotating red giants need an interaction to have reached their surface velocities; this may reflect simply a high initial rotation rate of the progenitor. Also, not all slowly rotating red giants may be explained by single star evolution that is typically in the upper part of the red giant branch.

We discussed in a simple way the consequences of an engulfment on the surface composition of red giants and confirmed the results obtained by [Carlberg et al. \(2012\)](#) that the effect on the carbon isotopic ratios are very small and the impact on lithium might be quite large, although remaining difficult to interpret (see below).

We showed that the chemical signatures of an engulfment are still quite ambiguous because planet engulfment either produce too small signal as in the case of the carbon isotopic ratios, or produce signals that cannot be attributed in a non-ambiguous way to a planet engulfment. This makes the surface velocity of a red giant the most stringent observable feature indicating a past star-planet interaction.

The present results also show that the evolution after an engulfment is not very different from that obtained without any engulfment. The main difference is in the surface rotation (and of course the rotation as a whole of the external convective zone). A planet engulfment would somewhat lower the contrast between the angular velocity of the core and that of the envelope, but not at a level that could be useful to reproduce the small contrast as obtained by asteroseismology for a few red giants (e.g. [Beck et al. 2012](#); [Mosser et al. 2012](#); [Deheuvels et al. 2012, 2014](#)).

There are many other points that should be addressed in further works. In particular, we shall investigate whether the acceleration of the surface due to a planet engulfment may trigger a magnetic field.

Acknowledgements. This research has made use of the Exoplanet Orbit Database and the Exoplanet Data Explorer at exoplanets.org. The project has been supported by Swiss National Science Foundation grants 200021-138016, 200020-160119 and 200020-15710.

References

- Adamów, M., Niedzielski, A., Villaver, E., Nowak, G., & Wolszczan, A. 2012a, *ApJ*, **754**, L15
- Adamów, M., Niedzielski, A., Villaver, E., Nowak, G., & Wolszczan, A. 2012b, *ApJ*, **754**, L15
- Adamów, M., Niedzielski, A., Villaver, E., Wolszczan, A., & Nowak, G. 2014, *A&A*, **569**, A55
- Adamów, M., Niedzielski, A., Villaver, E., et al. 2015, *A&A*, **581**, A94
- Aguilera-Gómez, C., Chanamé, J., Pinsonneault, M., & Carlberg, J. 2016, *ApJ*, accepted [[arXiv:1605.05332](#)]
- Alexander, J. B. 1967, *The Observatory*, **87**, 238
- Beck, P. G., Montalbán, J., Kallinger, T., et al. 2012, *Nature*, **481**, 55
- Cantiello, M., Mankovich, C., Bildsten, L., Christensen-Dalsgaard, J., & Paxton, B. 2014, *ApJ*, **788**, 93
- Carlberg, J. K., Majewski, S. R., & Arras, P. 2009, *ApJ*, **700**, 832
- Carlberg, J. K., Majewski, S. R., Patterson, R. J., et al. 2011, *ApJ*, **732**, 39
- Carlberg, J. K., Cunha, K., Smith, V. V., & Majewski, S. R. 2012, *ApJ*, **757**, 109
- Charbonnel, C., & Lagarde, N. 2010, *A&A*, **522**, A10
- Charbonnel, C., & Zahn, J.-P. 2007, *A&A*, **467**, L15
- Deheuvels, S., García, R. A., Chaplin, W. J., et al. 2012, *ApJ*, **756**, 19
- Deheuvels, S., Doğan, G., Goupil, M. J., et al. 2014, *A&A*, **564**, A27
- Eggenberger, P., Meynet, G., Maeder, A., et al. 2008, *Ap&SS*, **316**, 43
- Eggenberger, P., Montalbán, J., & Miglio, A. 2012, *A&A*, **544**, L4
- Ekström, S., Georgy, C., Eggenberger, P., et al. 2012, *A&A*, **537**, A146
- Fekel, F. C., & Balachandran, S. 1993, *ApJ*, **403**, 708
- Gray, D. F. 1981, *ApJ*, **251**, 155
- Gray, D. F. 1982, *ApJ*, **262**, 682
- Livio, M., & Soker, N. 1984, *MNRAS*, **208**, 763
- Livio, M., & Soker, N. 2002, *ApJ*, **571**, L161
- Lodders, K., & Fegley, B. 1998, *The planetary scientist's companion* (Oxford University Press)
- Marques, J. P., Goupil, M. J., Lebreton, Y., et al. 2013, *A&A*, **549**, A74
- Massarotti, A., Latham, D. W., Stefanik, R. P., & Fogel, J. 2008, *AJ*, **135**, 209
- Mosser, B., Goupil, M. J., Belkacem, K., et al. 2012, *A&A*, **540**, A143
- Passy, J.-C., Mac Low, M.-M., & De Marco, O. 2012, *ApJ*, **759**, L30
- Peterson, R. C., Tarbell, T. D., & Carney, B. W. 1983, *ApJ*, **265**, 972
- Privitera, G., Meynet, G., Eggenberger, P., et al. 2016, *A&A*, **591**, A45
- Sandquist, E., Taam, R. E., Lin, D. N. C., & Burkert, A. 1998, *ApJ*, **506**, L65
- Sandquist, E. L., Dokter, J. J., Lin, D. N. C., & Mardling, R. A. 2002, *ApJ*, **572**, 1012
- Siess, L., & Livio, M. 1999a, *MNRAS*, **304**, 925
- Siess, L., & Livio, M. 1999b, *MNRAS*, **308**, 1133
- Simon, T., & Drake, S. A. 1989, *ApJ*, **346**, 303
- Soker, N., Livio, M., & Harpaz, A. 1984, *MNRAS*, **210**, 189
- Soker, N., Regev, O., Livio, M., & Shara, M. M. 1987, *ApJ*, **318**, 760
- Staff, J. E., De Marco, O., Wood, P., Galaviz, P., & Passy, J.-C. 2016, *MNRAS*, **458**, 832
- Tayar, J., Ceillier, T., García-Hernández, D. A., et al. 2015, *ApJ*, **807**, 82
- Wong, M. H., Mahaffy, P. R., Atreya, S. K., Niemann, H. B., & Owen, T. C. 2004, *Icarus*, **171**, 153
- Zapolsky, H. S., & Salpeter, E. E. 1969, *ApJ*, **158**, 809

Appendix A: Detailed properties of the stellar models before and after engulfment

This Appendix contains three tables that list information on the characteristics of the star before and after an engulfment. Table A.1 indicates the properties of the stellar models just before the engulfment. The first column is the mass of the planet. Columns 2 to 5 give the mass of the star, its radius, surface angular velocity, mass loss rate, and mass of the convective envelope at this time.

Tables A.2 and A.3 present the changes occurring owing to the tides/engulfment processes. The first column is the mass of the planet. Column 2 gives the equatorial surface velocity before the engulfment. This velocity is already significantly higher than that of the corresponding single star at that stage. Actually, the velocity indicated in that column would be the velocity acquired by the tidal forces alone. Column 3 gives the duration of the phases during which, starting from the velocity indicated in Col. 2, the surface velocity would be higher than 8 km s^{-1} , $\Delta t_t(\text{FR})$. To compute this duration we used the results shown in Fig. A.1. This figure shows the evolution as a function of time of the surface velocity just after an engulfment. We see that roughly the surface velocity varies as $v(\Delta t) \sim v_{\text{max}}(1 - 0.04\Delta t)$, where Δt (in Myr) is $t - t_{\text{MS}}$, with t the age of the star, t_{MS} the age of the star at the end of the MS phase, and v_{max} the surface velocity reached just after the engulfment. This relation does not much depend on the mass of the planet, or equivalently on the value of v_{max} . It means that after 2.5 Myr, whatever the initial velocity, the surface velocity decreases by 10% with respect to its initial value⁷. We can use such a plot to estimate the time during which a star maintains a certain velocity after the engulfment evolving up along the red giant branch. This is how we estimated the time, $\Delta t_t(\text{FR})$, indicated in Col. 3 of Tables A.2 and A.3 showing the duration of the period when the star has a surface velocity

Fig. A.1. Evolution of the surface velocity (normalized to the velocity reached just after the engulfment, v_{max}) as a function of time for a $2 M_{\odot}$ stellar model with an initial rotation equal to 10% of the critical velocity. Various masses for the planets are considered. All planets started at a distance equal to 0.5 au from their parent star.

higher than 8 km s^{-1} when only the acceleration due to tides is accounted for.

In Tables A.2 and A.3, Cols. 4 to 6 indicate the surface velocity right after the engulfment; the duration of the phases during which the surface velocity, after an engulfment, is higher than 8 km s^{-1} ($\Delta t_{t+c}(\text{FR})$), and finally the fraction of the red giant branch phase spent with a surface velocity superior to 8 km s^{-1} . Tables A.2 show the results obtained with stellar models that have an initial angular velocity equal to 10% of the critical velocity, and Tables A.3 show the results for models that have an initial angular velocity equal to 50% of the critical velocity.

⁷ Actually, this timescale changes a bit with the mass of the planet as can be seen in Fig. A.1, which is slightly shorter for the 5 than for the 15 M_J planet. On the other hand, the differences between the various planet masses are not very large.

Table A.1. Characteristics of the star at the time of engulfment; see also [Privitera et al. \(2016\)](#) for additional pieces of information.

M_{pl} [M_{J}]	$M_{\star}(\text{Age}_{\text{eng}})$ [$M_{\odot}$]	$R_{\star}(\text{Age}_{\text{eng}})$ [$R_{\odot}$]	$\Omega_{\text{surf}}(\text{Age}_{\text{eng}})$ [10^{-7} s^{-1}]	$\dot{M}_{\star}(\text{Age}_{\text{eng}})$ [$10^{-9} M_{\odot} \text{ yr}^{-1}$]	$M_{\text{env}}(\text{Age}_{\text{eng}})$ [% $M_{\star}$]	$M_{\star}(\text{Age}_{\text{eng}})$ [$M_{\odot}$]	$R_{\star}(\text{Age}_{\text{eng}})$ [$R_{\odot}$]	$\Omega_{\text{surf}}(\text{Age}_{\text{eng}})$ [10^{-7} s^{-1}]	$\dot{M}_{\star}(\text{Age}_{\text{eng}})$ [$10^{-9} M_{\odot} \text{ yr}^{-1}$]	$M_{\text{env}}(\text{Age}_{\text{eng}})$ [% $M_{\star}$]
	$\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1, Z = 0.02$									
	$M_{\star,\text{ini}} = 1.5 M_{\odot}, R_{\star,\text{ini}} = 1.506 R_{\odot}, v_{\text{surf,ini}} = 25.6 \text{ km s}^{-1}$					$M_{\star,\text{ini}} = 1.7 M_{\odot}, R_{\star,\text{ini}} = 1.535 R_{\odot}, v_{\text{surf,ini}} = 25.1 \text{ km s}^{-1}$				
	$a_{\text{ini}} = 0.5 \text{ [AU]}$									
1	1.485274	29.3838	1.01	−0.77	67.62	1.687587	30.5011	0.84	−0.79	67.09
5	1.490112	20.8069	9.55	−0.32	69.45	1.691796	22.2625	5.58	−0.35	68.87
10	1.492073	17.4992	23.50	−0.21	70.46	1.693976	17.6225	15.97	−0.19	69.85
15	1.493253	15.4550	38.69	−0.16	71.03	1.694331	17.1070	29.72	−0.18	70.01
	$a_{\text{ini}} = 1.0 \text{ [AU]}$									
1	1.458215	70.2482	0.26	−6.75	61.94	1.663381	70.2862	0.13	−6.38	61.41
5	1.472279	50.4068	2.10	−3.02	64.37	1.675412	52.1931	1.73	−3.08	63.64
10	1.477916	42.1161	5.87	−2.00	65.56	1.680259	44.1649	4.67	−2.07	64.88
15	1.481351	37.0043	10.69	−1.48	66.52	1.683111	38.9341	9.03	−1.55	65.66
	$a_{\text{ini}} = 1.5 \text{ [AU]}$									
1	1.422241	109.5148	0.15	−20.61	57.76	1.626031	115.7902	0.12	−21.88	56.75
5	1.448623	81.8077	1.06	−9.98	60.60	1.652720	85.0667	0.99	−10.21	59.84
10	1.459680	68.0247	2.00	−6.38	62.19	1.663087	71.0516	1.14	−6.56	61.42
15	1.465830	62.7179	5.29	−5.69	63.21	1.668403	64.7079	4.11	−5.42	62.29
	$M_{\star,\text{ini}} = 2.0 M_{\odot}, R_{\star,\text{ini}} = 1.620 R_{\odot}, v_{\text{surf,ini}} = 26.5 \text{ km s}^{-1}$					$M_{\star,\text{ini}} = 2.5 M_{\odot}, R_{\star,\text{ini}} = 1.785 R_{\odot}, v_{\text{surf,ini}} = 28.2 \text{ km s}^{-1}$				
	$a_{\text{ini}} = 0.5 \text{ [AU]}$									
1	1.989876	34.5306	0.52	−0.97	65.38	−	−	−	−	−
5	1.993967	24.2140	4.58	−0.39	66.90	2.498326	31.0165	2.32	−0.65	63.77
10	1.995228	20.4300	11.68	−0.26	67.48	2.498676	28.5510	3.78	−0.52	63.77
15	1.995974	18.1078	17.46	−0.19	67.86	2.498689	28.4553	6.89	−0.52	63.77
	$a_{\text{ini}} = 1.0 \text{ [AU]}$									
1	1.969336	73.7458	0.19	−6.52	60.02	−	−	−	−	−
5	1.979736	55.3260	1.41	−3.26	62.39	−	−	−	−	−
10	1.983725	47.8862	3.63	−2.30	63.30	−	−	−	−	−
15	1.986025	43.0656	5.95	−1.76	64.05	−	−	−	−	−
	$a_{\text{ini}} = 1.5 \text{ [AU]}$									
1	1.936279	119.2380	0.06	−21.53	55.27	−	−	−	−	−
5	1.959307	89.1782	0.68	−10.47	58.46	−	−	−	−	−
10	1.967829	76.6386	1.72	−7.29	59.86	−	−	−	−	−
15	1.972431	68.9721	3.21	−5.66	60.75	−	−	−	−	−
	$\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5, Z = 0.02$									
	$M_{\star,\text{ini}} = 1.5 M_{\odot}, R_{\star,\text{ini}} = 1.473 R_{\odot}, v_{\text{surf,ini}} = 142 \text{ km s}^{-1}$					$M_{\star,\text{ini}} = 1.7 M_{\odot}, R_{\star,\text{ini}} = 1.580 R_{\odot}, v_{\text{surf,ini}} = 130 \text{ km s}^{-1}$				
	$a_{\text{ini}} = 0.5 \text{ [AU]}$									
1	1.485476	30.4342	1.93	−0.86	67.47	1.688074	31.4697	1.56	−0.87	66.58
5	1.490038	22.1058	8.21	−0.38	69.16	1.691756	23.5966	6.98	−0.42	68.03
10	1.491929	18.6698	19.48	−0.25	70.01	1.693187	20.1127	14.86	−0.28	68.68
15	1.492999	16.3263	35.18	−0.19	70.59	1.694140	17.6355	28.11	−0.21	69.17
	$a_{\text{ini}} = 1.0 \text{ [AU]}$									
1	1.459109	70.9117	0.44	−7.05	61.77	1.664231	72.3823	0.40	−7.01	61.06
5	1.472818	51.5472	2.34	−3.24	64.21	1.676070	54.2353	1.92	−3.45	63.29
10	1.482561	36.0369	8.27	−1.36	66.49	1.680738	45.5303	4.98	−2.30	64.37
15	1.481257	37.5313	10.94	−1.60	66.23	1.683347	43.9700	12.58	−2.42	65.15
	$a_{\text{ini}} = 1.5 \text{ [AU]}$									
1	1.418420	116.2842	0.20	−24.21	57.16	1.627502	118.5560	0.18	−23.82	56.38
5	1.449313	83.4278	0.63	−10.54	60.43	1.654111	87.8077	0.96	−11.17	59.47
10	1.460203	69.9438	1.32	−6.84	62.03	1.663785	73.3716	1.13	−7.28	60.90
15	1.466107	61.3007	5.41	−5.40	62.91	1.669147	65.6610	1.88	−5.55	61.94
	$M_{\star,\text{ini}} = 2.0 M_{\odot}, R_{\star,\text{ini}} = 1.667 R_{\odot}, v_{\text{surf,ini}} = 138 \text{ km s}^{-1}$					$M_{\star,\text{ini}} = 2.5 M_{\odot}, R_{\star,\text{ini}} = 1.833 R_{\odot}, v_{\text{surf,ini}} = 147 \text{ km s}^{-1}$				
	$a_{\text{ini}} = 0.5 \text{ [AU]}$									
1	1.991163	33.5930	1.41	−0.93	64.96	−	−	−	−	−
5	1.994217	25.0269	5.53	−0.44	66.10	2.498400	32.5785	3.10	−0.75	62.02
10	1.996013	18.3343	16.65	−0.21	66.67	2.498741	29.5638	6.37	−0.59	62.02
15	1.997273	21.2124	17.83	−0.29	68.44	2.498893	28.3750	9.41	−0.54	61.77
	$a_{\text{ini}} = 1.0 \text{ [AU]}$									
1	1.970755	75.3894	0.37	−7.06	59.60	−	−	−	−	−
5	1.980597	57.0608	1.93	−3.61	61.77	−	−	−	−	−
10	1.984246	49.5622	3.61	−2.56	62.70	−	−	−	−	−
15	1.986321	46.0991	6.57	−2.15	63.25	−	−	−	−	−
	$a_{\text{ini}} = 1.5 \text{ [AU]}$									
1	−	−	−	−	−	−	−	−	−	−
5	1.960723	92.4787	0.78	−11.56	57.84	−	−	−	−	−
10	1.968937	78.6476	1.86	−7.91	59.24	−	−	−	−	−
15	1.973119	71.6140	2.24	−6.30	60.14	−	−	−	−	−

Notes. A small – indicates that no engulfment occurs during the red giant phase for the considered initial conditions.

Table A.2. Effect of planet engulfment on the surface rotation velocities of red giants.

M_{pl} [M_J]	$v_{\text{surf}}(\text{Be})$ [km s $^{-1}$]	$\Delta t_{\text{f}}(\text{FR})$ [Myr]	$v_{\text{surf}}(\text{Af})$ [km s $^{-1}$]	$\Delta t_{\text{f+e}}(\text{FR})$ [Myr]	$\Delta t_{\text{f+e}}(\text{FR})/t(\text{RG})$ [% t_{rgb}]	$v_{\text{surf}}(\text{Be})$ [km s $^{-1}$]	$\Delta t_{\text{f}}(\text{FR})$ [Myr]	$v_{\text{surf}}(\text{Af})$ [km s $^{-1}$]	$\Delta t_{\text{f+e}}(\text{FR})$ [Myr]	$\Delta t_{\text{f+e}}(\text{FR})/t(\text{RG})$ [% t_{rgb}]
	$\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1, Z = 0.02$									
	$1.5 M_{\odot}, \mathcal{L}_{0,\star} = 0.41 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 0.37 [\text{Gyr}]$					$1.7 M_{\odot}, \mathcal{L}_{0,\star} = 0.46 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 0.18 [\text{Gyr}]$				
	$a_{\text{ini}} = 0.5 [\text{AU}]$									
1	1.44	–	2.06	0.0	0.00	1.35	–	1.8	0.0	0.00
5	7.26	–	13.80	15.3	4.29	6.26	–	8.64	2.8	1.55
10	17.00	26.24	28.90	29.6	8.30	14.60	19.65	19.60	23.9	13.13
15	28.20	32.36	42.30	39.7	11.15	23.30	26.28	35.70	31.2	17.20
	$a_{\text{ini}} = 1 [\text{AU}]$									
1	0.83	–	1.20	0.0	0.00	0.78	–	1.18	0.0	0.00
5	4.81	–	7.37	0.0	0.00	4.19	–	6.28	0.0	0.00
10	11.90	4.28	17.40	7.6	2.13	10.30	3.03	14.4	5.7	3.13
15	12.50	5.60	28.00	11.5	3.22	16.50	7.03	24.8	9.7	5.35
	$a_{\text{ini}} = 1.5 [\text{AU}]$									
1	0.66	–	1.04	0.0	0.00	0.60	–	0.95	0.0	0.00
5	4.81	–	6.02	0.0	0.00	3.42	–	5.87	0.0	0.00
10	9.22	1.14	13.40	3.2	0.91	7.99	–	12.50	2.8	1.55
15	15.80	4.12	23.60	5.7	1.61	12.8	3.07	18.70	4.8	2.64
	$2.0 M_{\odot}, \mathcal{L}_{0,\star} = 0.61 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 0.08 [\text{Gyr}]$					$2.5 M_{\odot}, \mathcal{L}_{0,\star} = 0.92 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 10 [\text{Myr}]$				
	$a_{\text{ini}} = 0.5 [\text{AU}]$									
1	0.99	–	1.30	0.0	0.00	–	–	–	–	–
5	4.90	–	7.71	0.0	0.00	3.33	–	18.4	0.7	6.82
10	11.80	8.69	16.70	14.7	19.29	7.00	–	37.9	0.7	6.82
15	16.18	13.51	22.00	19.9	26.01	7.94	–	47.0	0.7	6.82
	$a_{\text{ini}} = 1.0 [\text{AU}]$									
1	0.70	–	0.99	0.0	0.00	–	–	–	–	–
5	3.60	–	5.42	0.0	0.00	–	–	–	–	–
10	7.87	–	12.10	3.8	4.98	–	–	–	–	–
15	8.42	0.49	17.90	6.8	8.95	–	–	–	–	–
	$a_{\text{ini}} = 1.5 [\text{AU}]$									
1	0.53	–	0.82	0.0	0.00	–	–	–	–	–
5	2.82	–	4.22	0.0	0.00	–	–	–	–	–
10	6.33	–	9.18	0.9	1.14	–	–	–	–	–
15	10.30	1.61	15.50	3.7	4.83	–	–	–	–	–

Notes. The cases for different initial mass stars beginning their evolution with a rotation equal to $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.1$ are considered.

Table A.3. Same as Table A.2 but for $\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5$.

M_{pl} [M_{J}]	$v_{\text{surf}}(\text{Be})$ [km s $^{-1}$]	$\Delta t_{\text{f}}(\text{FR})$ [Myr]	$v_{\text{surf}}(\text{Af})$ [km s $^{-1}$]	$\Delta t_{\text{f+e}}(\text{FR})$ [Myr]	$\Delta t_{\text{f+e}}(\text{FR})/t(\text{RG})$ [% t_{rgb}]	$v_{\text{surf}}(\text{Be})$ [km s $^{-1}$]	$\Delta t_{\text{f}}(\text{FR})$ [Myr]	$v_{\text{surf}}(\text{Af})$ [km s $^{-1}$]	$\Delta t_{\text{f+e}}(\text{FR})$ [Myr]	$\Delta t_{\text{f+e}}(\text{FR})/t(\text{RG})$ [% t_{rgb}]
	$\Omega_{\text{ini}}/\Omega_{\text{crit}} = 0.5, Z = 0.02$									
	$1.5 M_{\odot}, \mathcal{L}_{0,\star} = 2.16 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 0.24 [\text{Gyr}]$					$1.7 M_{\odot}, \mathcal{L}_{0,\star} = 2.39 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 0.12 [\text{Gyr}]$				
	$a_{\text{ini}} = 0.5 [\text{AU}]$									
1	3.21	–	4.09	0.0	0.00	2.94	–	3.42	0.0	0.00
5	9.45	6.51	12.60	11.9	4.96	8.24	0.88	11.50	8.2	6.76
10	18.60	20.92	25.50	25.1	10.51	15.60	16.20	20.80	18.3	15.09
15	28.90	27.52	40.40	35.9	15.00	24.00	20.78	34.50	26.8	22.15
	$a_{\text{ini}} = 1 [\text{AU}]$									
1	1.69	–	2.19	0.0	0.00	1.53	–	2.04	0.0	0.00
5	5.93	–	8.39	0.6	0.23	5.15	–	7.23	0.0	0.00
10	14.80	7.03	20.90	9.8	4.08	11.10	3.01	15.80	5.6	4.61
15	20.80	9.27	28.70	10.7	4.48	18.30	5.89	31.90	9.6	7.97
	$a_{\text{ini}} = 1.5 [\text{AU}]$									
1	1.15	–	1.64	0.0	0.00	1.10	–	1.49	0.0	0.00
5	4.74	–	6.81	0.0	0.00	3.97	–	5.84	0.0	0.00
10	9.96	1.69	15.60	3.5	1.48	8.54	0.44	12.00	2.3	1.94
15	16.40	3.97	22.90	5.3	2.22	13.60	3.01	19.80	4.5	3.72
	$2.0 M_{\odot}, \mathcal{L}_{0,\star} = 3.06 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 0.06 [\text{Gyr}]$					$2.5 M_{\odot}, \mathcal{L}_{0,\star} = 4.65 [10^{50} \text{ g cm}^2 \text{ s}^{-1}], t_{\text{rgb}} \simeq 8 [\text{Myr}]$				
	$a_{\text{ini}} = 0.5 [\text{AU}]$									
1	2.87	–	3.3	0.0	0.00	–	–	–	–	–
5	5.40	–	9.6	3.6	6.19	5.20	–	24.60	0.5	6.42
10	17.10	11.62	21.3	18.6	31.86	9.22	6.42	42.1	0.5	6.42
15	18.70	16.07	33.7	26.3	45.16	12.95	6.42	57	0.5	6.42
	$a_{\text{ini}} = 1.0 [\text{AU}]$									
1	1.52	–	1.96	0.0	0.00	–	–	–	–	–
5	4.72	–	7.64	0.0	0.00	–	–	–	–	–
10	8.97	0.96	12.50	3.5	6.07	–	–	–	–	–
15	13.80	3.81	21.30	6.1	10.39	–	–	–	–	–
	$a_{\text{ini}} = 1.5 [\text{AU}]$									
1	–	–	–	–	–	–	–	–	–	–
5	3.39	–	5.03	0.0	0.00	–	–	–	–	–
10	7.01	–	10.20	1.3	2.22	–	–	–	–	–
15	11.00	1.74	15.50	2.3	3.95	–	–	–	–	–