

Observing transiting planets with JWST

Prime targets and their synthetic spectral observations (*Corrigendum*)

P. Mollière¹, R. van Boekel¹, J. Bouwman¹, Th. Henning¹, P.-O. Lagage^{2,3}, and M. Min^{4,5}

¹ Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany
e-mail: molliere@mpia.de

² Irfu, CEA, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

³ AIM, Université Paris Diderot, 91191 Gif-sur-Yvette, France

⁴ SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584 CA Utrecht, The Netherlands

⁵ Astronomical institute Anton Pannekoek, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands

A&A 600, A10 (2017), DOI: [10.1051/0004-6361/201629800](https://doi.org/10.1051/0004-6361/201629800)

Key words. methods: numerical – planets and satellites: atmospheres – radiative transfer – errata, addenda

In Sect. 6.2 it was incorrectly stated that NIRISS provides little diagnostic power to distinguish between the various cases of strongly irradiated planets without heat re-distribution. In fact, from radiometric considerations alone, NIRISS is as good or better than NIRSpec in these cases, as is apparent in Fig. 7.