

Period spacings in red giants

III. Coupling factors of mixed modes[★]

(Corrigendum)

B. Mosser¹, C. Pinçon¹, K. Belkacem¹, M. Takata², and M. Vradar³

¹ LESIA, Observatoire de Paris, PSL Research University, CNRS, Université Pierre et Marie Curie, Université Paris Diderot, 92195 Meudon, France cedex, France
e-mail: benoit.mosser@obspm.fr

² Department of Astronomy, School of Science, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, 113-0033 Tokyo, Japan

³ Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, 4150-762 Porto, Portugal

A&A 600, A1 (2017), DOI: [10.1051/0004-6361/201630053](https://doi.org/10.1051/0004-6361/201630053)

Key words. stars: oscillations – stars: interiors – stars: evolution – errata, addenda

A table with the coupling factors and their uncertainties, computed as in Sect. 3 of the original paper, has been appended to the article as a CDS/VizieR document. This table downloadable at <http://vizier.u-strasbg.fr/viz-bin/VizieR?-source=J/A+A/600/A1> contains 5166 stars identified with their KIC numbers, coordinates, large separations $\Delta\nu$, period spacings $\Delta\Pi_1$, and seismic estimates of the mass.

* A table is only available at the CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/A+A/600/A1>