

Exploring the climate of Proxima B with the Met Office Unified Model (Corrigendum)

Ian A. Boutle^{1,2}, Nathan J. Mayne¹, Benjamin Drummond¹, James Manners^{1,2}, Jayesh Goyal¹,
F. Hugo Lambert³, David M. Acreman¹, and Paul D. Earnshaw²

¹ Physics and Astronomy, College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, EX4 4QL, UK
e-mail: i.boutle@exeter.ac.uk

² Met Office, FitzRoy Road, Exeter, EX1 3PB, UK

³ Mathematics, College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, EX4 4QF, UK

A&A, 601, A120 (2017), <https://doi.org/10.1051/0004-6361/201630020>

Key words. stars: individual: Proxima Cen – planets and satellites: individual: Proxima B – planets and satellites: atmospheres – planets and satellites: detection – planets and satellites: terrestrial planets – errata, addenda

It has been brought to our attention that there is a minor typo in the x -axis title of the second panel of Fig. 5, which should have units of “ kg kg^{-1} ” rather than “ g kg^{-1} ”. The corrected figure appears below.




Fig. 5. Ten orbit mean profiles of temperature and water vapour (*left and right panels*, respectively) from the sub-stellar point on the dayside of the tidally locked planet (solid) and its antipode (dashed). Filled triangles show the surface temperature, and crosses indicate the position of the model levels.